

# Twenty Years of Progress in Physiological and Biochemical Investigation of RALF Peptides<sup>1</sup>[OPEN]

Matthew R. Blackburn,<sup>a</sup> Miyoshi Haruta,<sup>a,2</sup> and Daniel S. Moura<sup>b,2,3</sup>

<sup>a</sup>Department of Biochemistry, University of Wisconsin-Madison, Madison, Wisconsin 53706

<sup>b</sup>Laboratório de Bioquímica de Proteínas, Departamento de Ciências Biológicas, Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo (ESALQ/USP), Piracicaba, SP 13418-900, Brazil

ORCID IDs: 0000-0001-6067-4478 (M.R.B.); 0000-0003-1686-1860 (M.H.); 0000-0003-2042-1403 (D.S.M.)

A family of secreted peptides known as the RALFs (rapid alkalization factors) induce rapid alkalization of the extracellular compartment of plant cells. RALFs are perceived by a family of *Catharanthus roseus* Receptor Like Kinase 1 Like proteins (CrRLK1Ls), a structurally related family of plasma membrane-enriched receptors. RALF perception coincides with the downregulation of plasma membrane H<sup>+</sup>-ATPase function, which reduces energy for solute uptake and results in growth inhibition. In this review, we discuss the initial discovery of this ubiquitous family of peptide growth regulators and their cognate receptors. We describe recent detailed structural studies of RALF-CrRLK1Ls interactions and the identification of a subset of downstream signaling components. We then summarize the current state of RALFs-CrRLK1L research with emphasis on the Arabidopsis (*Arabidopsis thaliana*) RALFs (AtRALFs) and their roles in cellular regulation of plant growth and development and in biotic and abiotic stress responses.

## RALF DISCOVERY, STRUCTURE, AND DIVERSITY

The origins of the RALF field lie in the work of Clarence “Bud” Ryan’s group at Washington State University, with the discovery of the first RALF peptide using an activity assay originally designed to detect systemins, which are small, ~18 amino acid peptide hormones involved in the wound response in tobacco (*Nicotiana tabacum*) plants. This assay targeted the alkalization caused by systemins and was used to test HPLC-fractionated tobacco leaf extracts. Surprisingly, an HPLC fraction eluting later than systemins elicited an unexpectedly fast and prominent increase in the pH of the culture medium. Edman degradation later

confirmed that this activity was a larger peptide of approximately 5 kDa (Pearce et al., 2001). This new peptide failed to induce proteinase inhibitors when applied to cut stems of small tomato (*Solanum lycopersicum*) plants, suggesting it was not involved in the wound response. However, germinating seeds of tomato and Arabidopsis experienced immediate root growth arrest when transferred to media spiked with the peptide versus unspiked media, and this effect was mainly perceived in the elongation zone of the root.

A search of sequence databases using the original 17 N-terminal amino acids obtained from peptide sequencing revealed that RALFs are ubiquitous in the plant kingdom and identified conserved motifs. For example, a pair of Arg residues (RR), a feature typically

## ADVANCES

- RALF peptide-induced alkalization of extracellular media or the cell wall reduces the proton electrochemical potential that is required for solute uptake, leading to cell growth suppression.
- RALFs are more widespread than originally thought as they have been found in fungi and even in bacteria.
- RALF precursors are preproteins or preproteins, and their primary structures and precursor processing may be the determinant of their physiological roles.
- RALF signaling components include GPI-anchored proteins, G-protein subunits, and soluble protein kinases; they also interact with other hormone pathways including those for brassinolide and ABA.
- Unbound AtRALF peptides do not appear to be structured in solution, but AtRALF-receptor complexes can be co-crystallized, and their interaction domains have been modeled.

<sup>1</sup>This work was supported by the National Science Foundation (MCB MSN203469 to M.H. and PGRP MSN187204 to M.R.B.) and the São Paulo Research Foundation (2014/14487-1 to D.S.M.).

<sup>2</sup>Senior authors.

<sup>3</sup>Author for contact: danielmoura@usp.br.

All authors participated in writing the article.

[OPEN] Articles can be viewed without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.19.01310

found in animal prohormones processed by convertases, is located just upstream from the N terminus of the active peptide in almost all RALFs (Pearce et al., 2001). Furthermore, the dibasic site RR not only functions as the cleavage site for prohormone processing enzymes but is also central to the involvement of AtRALFs in coordinating the immune response (Matos et al., 2008; Srivastava et al., 2009; Stegmann et al., 2017).

It was established early on that the N-terminal YISY motif of the mature peptide is required for productive binding of the peptide to its target receptor; this was experimentally demonstrated in truncation and mutagenesis experiments on synthetic RALF (Pearce et al., 2010). Importantly, the flanking tyrosines and internal I/L of this motif show the strongest conservation, whereas the internal Ser shows moderate conservation, with occasional substitution by Gly or Thr. GASYY and the C-terminal RCRR(S) motifs also occur in many mature AtRALF sequences, but their role remains to be experimentally determined. However, truncation experiments that demonstrated the requirement of the C-terminal portion of the AtRALF1 peptide for activity suggest that these motifs may play a role in peptide structure or binding (Pearce et al., 2010).

Notably, many AtRALF sequences contain four Cys residues that are positionally conserved and absolutely required for biological activity (Pearce et al., 2001; Haruta et al., 2008). For these peptides, oxidative folding of the cleaved peptides also occurs to generate the mature AtRALF peptide, and mutation or blocking (alkylation) of these cysteines is known to disrupt physiological activity. Despite the presence of two functionally important disulfide bridge sites, the oxidatively folded mature peptide appears to be largely unstructured due to the strikingly minimal length of the peptide loop between each pair of bridged cysteines (Pearce et al., 2001). This configuration was independently validated with the recently solved solution structure of AtRALF8, which indicates an entirely unstructured peptide except where movement of the peptide backbone is constrained in the immediate vicinity of each disulfide bridge (Frederick et al., 2019; Fig. 1A, inset). It is possible that all Cys-rich AtRALFs adopt this configuration because the relative amino acid positions of those Cys residues are conserved among AtRALF peptide sequences.

Given these conserved motifs, it has been possible to demonstrate that RALF-related sequences are much more universal than originally thought, as peptide sequences with RALF motifs have been isolated from phytopathogenic fungi and even from bacteria (Masachis et al., 2016; Thynne et al., 2017). On the other hand, the diversity of RALF sequences within and among species suggests that the RALF signaling paradigm is complex. In *Arabidopsis* alone, the *AtRALF* genes comprise a multigene family of more than 30 members. The exact number varies according to the criteria used to establish AtRALF identity, and current reported numbers range from 33 to 37 (Cao and Shi, 2012; Campbell and Turner, 2017).

Dozens of other AtRALF homologs are present in other species of terrestrial plants. Phylogenetic analysis of nearly 800 identified RALF peptides across the plant kingdom found that much of the sequence variability of the AtRALF peptides occurs in the sequence of mature RALF peptides, and this variability further divides the mature peptides into four distinct clades (Campbell and Turner, 2017).

Since their discovery, an enormous amount of research has extended our understanding of the RALF family, its interactions with other proteins, and its biological functions across the plant kingdom. Despite the ubiquity of the RALF peptides, *Arabidopsis* emerged as the natural model to study their functions and contains nine AtRALF sequences closely related to the originally isolated tobacco RALF. For this reason, the following sections will largely discuss the RALF family in the context of *Arabidopsis* biology.

## THE DISCOVERY OF RALF RECEPTORS AND ANALYSIS OF RALF-RECEPTOR INTERACTIONS


The first study to investigate AtRALF-receptor interactions used a synthetic, radiolabeled tomato RALF (LeRALF) peptide modified with a photolabile azido handle to probe for LeRALF receptors. When cross linking was induced in *Lycopersicon peruvianum* cell suspension cultures spiked with this synthetic LeRALF, two membrane proteins at approximately 25 and 120 kDa were identified (Scheer et al., 2005). The RALF receptor was first identified in 2014 in the laboratory of Michael R. Sussman at the University of Wisconsin-Madison in a phosphoproteomic study that screened for AtRALF1-dependent changes using stable-isotope labeling combined with mass spectrometry of proteins that could be coimmunoprecipitated with AtRALF1 (Haruta et al., 2014).

This phosphoproteomic screen detected an increased abundance of phosphopeptides derived from a plasma membrane  $H^+$ -ATPase 2 (AHA2) and calcium-dependent protein kinase 9 (CPK9). This coincided with increases in the phosphorylation of FERONIA (FER), a member of CrRLK1L family of receptor-like kinases (Haruta et al., 2014). AtRALF1 was also shown to elevate cytoplasmic calcium and suppress root growth in *Arabidopsis* seedlings in a FER-dependent manner, because *fer* mutants were insensitive to AtRALF1 in root growth suppression and cytoplasmic calcium mobilization, further supporting the hypothesis that the AtRALF1 peptide is sensed by the FER receptor kinase. These results suggested the FER signaling axis as a molecular mechanism for AtRALF-induced alkalinization of the extracellular media and elevation of cytoplasmic calcium concentrations.

FER is localized at the plasma membrane, where it functions as an AtRALF receptor in root cell growth regulation and is involved in many diverse pathways in plant physiology, including female gamete control of

pollen tube rupture upon arrival of a female gametophyte to release sperm for fertilization (Huck et al., 2003; Rotman et al., 2003), rosette morphology, and guard cell movement (Yu and Assmann, 2018), immune responses (Stegmann et al., 2017), and abiotic stress responses (Chen et al., 2016; Xu et al., 2019). FER interacts with a RAC/ROP GTPase in the cellular signaling pathway regulating root hair development (Duan et al., 2010). FER also interacts with the GPI-anchored protein (GPI-AP) LORELEI (LRE) and LRE-LIKE GPI-AP1 (LLG1), proteins that act as chaperones that correctly target FER to the plasma membrane (Li et al., 2015). Besides their chaperone role, LLG proteins also bind AtRALF23 and function as a coreceptor for FER (Li et al., 2015; Xiao et al., 2019). In retrospect, after more than 15 years of ongoing research in the AtRALF field, it is possible that the original proteins identified by Scheer et al. (2005) are tomato homologs of Arabidopsis LLG proteins and CrRLK1Ls.

The presence of a protein complex containing AtRALF1, FER, and LLG1 was shown *in vivo*, and LLG1 was proposed to function as a coreceptor for FER in AtRALF1 perception (Li et al., 2015). Recent detailed structural studies report a submicromolar affinity ( $K_D$ ) for AtRALF binding in AtRALF-receptor complexes, including AtRALF1 or AtRALF23-FER interactions, indicating that this diverse family of peptides interacts with target receptors with high specificity (Liu et al., 2018; Xiao et al., 2019). An analysis of AtRALF23 cocrystallized with FER and LLG2 demonstrated that AtRALF23 adopts a helical fold when complexed with FER and LLG2 (Xiao et al., 2019; Fig. 1A). Unstructured, fully reduced AtRALF peptide (with four free thiols) was as active as correctly oxidized AtRALF peptide, suggesting the unusual possibility that the disulfide configuration itself can be induced during or right before receptor binding (Pearce et al., 2010).



**Figure 1.** RALF peptides adopt an induced fold when complexed with a CrRLK1L protein. A, the left shows the crystal structure of the heterotypic complex among LLG2 (pale blue), RALF23 (red), and the ectodomain of FERONIA (pale orange). Density for the full-length RALF23 peptide was obtained for residues from RYISY to the first conserved Cys of the mature peptide. The right shows 900 pivot along the y axis by an  $\sim 90^\circ$  tilt out of the page and zoomed in, with  $\sim 40$  residues of the N-terminal domain of ectoFER removed to clear the view of RALF23. The mature peptide hormone intercalates tightly between the juxtamembrane domain of FERONIA (light orange) and the flexible loops of LLG2 (light blue), adopting an induced alpha helix. YISY sidechains are shown as sticks for emphasis. PDB ID: 6A5E (Xiao et al., 2019). The inset shows the solution NMR structure of RALF8 confirms a mostly unstructured mature peptide with a minimal loop-spanning disulfide configuration formed from the four invariant cysteines. Bridged Cys sidechains are represented as yellow sticks. YITY sidechains are shown as sticks for emphasis. PDB ID: 6NU4 (Frederick et al., 2019). B, Chemical cross linking and footprinting data from RALF1 complexed with the extracellular domain of FERONIA (gray spheres) *in vitro*, in the absence of an LLG protein, mapped to the crystal structure 6A5E. LLG2 (cyan cartoon) is included in the image for spatial reference but was not part of the experiment. Dark blue, residues that gain protection from the bulk solvent in the presence of full-length RALF1 (EDC-GEE labeling); red, Lys 60 (K60) on FERONIA, which cross linked to the C-terminal Lys of RALF1 (BS3 cross linking); yellow, the peptide containing K60, identified by mass spectrometry (Liu et al., 2018).

In the study of Xiao et al. (2019), AtRALF23 was found to directly interact with the coreceptor LLG2 through tight interactions between its N terminus, which forms an alpha helix in crystallo, and disordered loops in LLG2 (Fig. 1A). Notably, this complex packs against the juxtamembrane domain of FER, and the YISY motif of AtRALF23 intercalates within the loops of LLG2 through a combination of polar and hydrophobic interactions. This suggests a general mechanism for how YISY, within the N-terminal  $\alpha$ -helical domain, is minimally required for AtRALF peptides to bind their target receptors and coreceptors and supports the notion that AtRALF peptides become structured upon binding to their receptors (Xiao et al., 2019).

AtRALF peptide binds its receptor independently of a GPI-anchored coreceptor like LLG2. Recent in vitro chemical cross linking studies of AtRALF1 complexed with the extracellular domain of FER have suggested that the AtRALF1 peptide occupies a region spanning a large hydrophobic cleft on FER and that the C terminus of AtRALF1 is involved in binding to FER (Liu et al., 2018). Specifically, chemical cross linking with the amine-reactive cross linker *bis*-sulfo-succinimidyl suberate (BS3) identified an N-terminal Lys on AtRALF1 that cross links with a Lys on the ectodomain of FER near the hydrophobic cleft, and protein footprinting demonstrated that changes in solvent exposure occurred on residues in the vicinity of the cleft and this Lys on FER in response to AtRALF1 binding (Fig. 1B). Similar AtRALF1-FER interprotein cross linking was also observed in vitro when other amine reactive chemical cross linkers, disuccinimidyl sulfoxide, or dithiobis (succinimidylpropionate) were tested. Fluorescence polarization experiments examining the interaction of Alexa 488 labeled AtRALF1 and various truncated forms of FER ectodomain (ectoFER) showed increases in fluorescence polarization values with wild-type ectoFER-bound form of Alexa 488 AtRALF1, but not in those with truncated forms. Thus, the experiments demonstrated that both malectin-like domains of FER are required for AtRALF1 binding, suggesting that the ligand binding site of AtRALF1 spans the cleft and both malectin-like domains (Liu et al., 2018).

It is worth noting that other CrRLK1L ectodomain structures have also been published for ANXUR1 (ANX1) and ANX2, a pair of CrRLK1L receptor kinases most closely related to FER (Du et al., 2018; Moussu et al., 2018) and that these models indicate that the cleft between the malectin-like domains of these CrRLK1Ls have a different residue composition than the intermalectin-like cleft of FER, suggesting that the AtRALF1-FER interaction is not necessarily mirrored by ANX1/2 and AtRALF4/19 interactions. Since three-dimensional structures of ANX1 and ANX2 were obtained as unbound forms, it is yet to be examined how closely AtRALF23-FER and AtRALF4/19-ANX1/2 complex structures are related.

These recent studies provide evidence about how an AtRALF peptide—specifically, the behavior of the N- and C-terminal portions of the peptide—binds its

receptor and is an important first step to understanding the biochemical and structural basis of RALF perception. However, additional high-resolution structural and biochemical studies of these complex interactions remain to be conducted to resolve apparent inconsistencies between the findings of Liu et al. (2018) and Xiao et al. (2019).


As with any in vitro experiment, the observed interactions in these structural studies do not necessarily reflect the physiological environment and could be prone to mixing or crystallization artifacts. Furthermore, the role of correct disulfide configurations was not explicitly addressed in either study, and it remains to be determined how the conserved Cys of each AtRALF contributes to ligand-receptor interactions. In addition, N-Hydroxysuccinimide-based cross linkers, such as BS3 or disuccinimidyl sulfoxide, rely on the availability of Lys residues located at or near the protein interaction interface, and thus AtRALF1-FER interaction sites modeled by the cross linking technique may not necessarily reflect the domains or amino acid residues that could be located at the closest contact of the two proteins. Ultimately, a unifying mechanism governing the specificity of RALF-receptor interactions remains to be determined and is a major open question in the RALF field (Fig. 2).

How do individual AtRALF peptides coordinate so many complex and specific responses, especially when considering the CrRLK1L family of receptors that bind the structurally similar AtRALF peptides? This question remains the subject of intense research efforts, and there is particular interest in solving the structure of a mature AtRALF peptide in a complex with its receptor. To date, RALF ligand interactions with CrRLK1Ls ectodomains are demonstrated for AtRALF1 and FER, AtRALF23 and FER, BUDDHA'S PAPER SEAL1/2 (BUPS1/2) and AtRALF4/19, and ANX1/2 and AtRALF4/19 (Haruta et al., 2014; Ge et al., 2017; Xiao et al., 2019). AtRALF ligand perception is mediated by the CrRLK1L family and LLG coreceptor family, but their signaling output is likely a complex interplay of different AtRALFs playing roles as agonists and antagonists (Ge et al., 2017; Stegmann et al., 2017).

## THE RALF SIGNALING AXIS EXTENDS TO OTHER DIVERSE PROTEINS AND FUNCTIONS

Both RALFs and CrRLK1Ls interact with numerous proteins that are implicated in RALF signaling. A pollen-specific RALF from tomato (SIPRALF) was first identified as a potential interactor of the LRR domain of a tomato pollen LEUCINE-RICH REPEAT EXTENSIN (LRX) protein in a Y2H screen (Covey et al., 2010). Pollen-expressed AtRALF4 and 19 interact with pollen-expressed proteins of the LRX family, which regulates pollen tube integrity and growth (Mecchia et al., 2017; Ge et al., 2019a). In roots, the amino terminus of LRX4 interacts with both AtRALF1 and the amino terminus of





**Figure 2.** Two proposed models considering available data on the induced fold of AtRALF and AtRALF-CrRLK1L protein binding interactions. To date, the solution NMR structure of AtRALF8 (Frederick et al., 2019) remains the only structure of an unbound AtRALF peptide and suggests that most Cys-rich peptides are similarly disordered. Note that LLG is shown here associated with FER before AtRALF binding for simplicity, but direct experimental evidence of AtRALF-independent LLG-FER interactions has not yet been demonstrated. Model 1 shows the crystal structure of the ectodomain of FERONIA complexed with LLG2 and AtRALF23 is currently the only structure of an AtRALF-receptor complex and suggests that the N terminus of AtRALF23 assumes an induced fold during binding and that its YISY motif is highly involved in AtRALF binding (Xiao et al., 2019). Meanwhile, the chemical cross linking and protein footprinting study of the AtRALF1/FER complex in the absence of an LLG protein suggests that the C terminus of AtRALF1 spans the intermalectin-like cleft. In the case of AtRALF23, binding occurs in an LLG-dependent manner near the juxtamembrane domain while AtRALF1 binding occurs independently of LLG at the malectin-like domains, suggesting two distinct, AtRALF-specific binding mechanisms. Model 2 shows, alternatively, these two binding modes reflect an interaction that is generalizable to many mature AtRALF peptides. Until more comprehensive data are available on AtRALF-receptor interactions, this remains a possibility.

FER to regulate cell elongation by controlling vacuolar expansion (Dünser et al., 2019). Similarly, the LRR domains of LRX3 and LRX4 can interact with both AtRALF22/23 and FER to regulate salt stress response (Zhao et al., 2018). Notably, it was recently found that AtRALF1 also interacts in a  $\text{Ca}^{2+}$  and pH-dependent manner with calmodulin-like 38 (CML38), another secreted protein that is now understood to be an important component of the AtRALF1 root growth signaling pathway, further complicating the current understanding of AtRALF1 specificity (Campos et al., 2018).

Molecular genetic studies have shown that FER is involved in plant cell growth and reproduction, mechanosensing, pathogen defense, salt, and nutrient stress responses (Table 1; Li et al., 2016; Stegmann et al., 2017; Feng et al., 2018; Xu et al., 2019). Since *fer* knockout mutants display a wide range of phenotypes, it has been proposed that FER may be acting as a scaffold in an array of protein complexes. It is possible that CrRLK1L receptors act as heterodimers in such complexes. This idea is supported by the detection of BUPS1/2-ANX1/2 or BUPS1-BUPS2 complex formation in both an in vitro protein interaction assay and a yeast two-hybrid assay; specifically, these results suggested that BUPS and ANXs form heterodimers to regulate pollen tube elongation (Ge et al., 2017; Zhu et al., 2018). A more recent study further suggested that LLG2/LLG3 are also acting as coreceptors for RALF4 and RALF19 perception by BUPS1, BUPS2, ANX1, and ANX2 receptors in maintaining pollen tube integrity (Ge et al., 2019c).

## REGULATION OF CELL PHYSIOLOGY, GROWTH, AND DEVELOPMENTAL PROCESSES BY RALF PEPTIDES

In the immediate aftermath of the discovery of the RALF peptides in tomato, tobacco, alfalfa (*Medicago truncatula*), and hybrid poplar (*Populus trichocarpa* × *Populus deltoides*), only two cellular activities—alkalinization and MAP kinase activation—were known in addition to the inhibition of root growth as physiological roles (Pearce et al., 2001; Haruta and Constabel, 2003). All of the original RALF peptides, whether biochemically isolated, recombinantly produced, or chemically synthesized, exhibit alkalization activity in cell suspension cultures (Pearce et al., 2001, 2010; Haruta and Constabel, 2003; Haruta et al., 2008; Mingossi et al., 2010; Morato do Canto et al., 2014). Additionally, AtRALFs were shown to transiently increase the concentration of the intracellular second messenger  $\text{Ca}^{2+}$  in Arabidopsis seedlings expressing the intracellular calcium reporter, aequorin; this suggested that AtRALFs exhibit growth factor-like coupling (Haruta et al., 2008; Morato do Canto et al., 2014).

Strikingly, alkalization activity does not appear to be universally essential for the AtRALFs' physiological roles, as some of these peptides elicit activities that are independent or partially independent of alkalization (Dressano et al., 2017; Stegmann et al., 2017; Campos et al., 2018). Intriguingly, when *NaRALF* of *Nicotiana attenuata* was silenced by an inverted repeat RNA construct, the *irRALF* plants unexpectedly retained the ability to acidify the external media (Wu et al., 2007). In

**Table 1.** Physiological overlap of AtRALF signaling and other biological pathways

Physiological Conditions	Interaction with RALF Signaling	Cellular Mechanisms	References
Brassinolide (BL) (DWF4)	Antagonistic effect between BR and AtRALF1 (or AtRALF23)	EXPANSIN5 (AtEXPA5), DWARF gene expression BR downregulates RALF1 or 23 expression	(Srivastava et al., 2009; Bergonci et al., 2014a, 2014b)
Absciscic acid (ABA)	Cross talk between AtRALF1 and ABA signaling and stress responses	ABI2 protein phosphatase dependence FER interaction with G protein $\beta$ -subunit (AGB1)	(Chen et al., 2016) (Yu and Assmann, 2018)
<i>Pseudomonas syringae</i> infection (elicitors flg22 or elf18)	Negative regulation of innate immunity by AtRALF23	Reduced protein interaction of FLS2 or EFR with BAK1	(Stegmann et al., 2017)
Carbon/nitrogen nutrient ratio	AtRALF1 and FER suppress stress response caused by high carbon/nitrogen	Phosphorylation of ATL6 E3 ubiquitin ligase and reduced stability of 14-3-3 protein	(Xu et al., 2019)

Arabidopsis cell cultures, AtRALFs tested so far all show robust extracellular pH alkalization activities, except the pollen-expressed AtRALF4 (Morato do Canto et al., 2014). In contrast, AtRALF19 is also expressed in pollen but is a strong alkalization factor, showing a 4.6 nM half-maximal activity in the assay (Morato do Canto et al., 2014). Those observations may suggest that seven amino acid substitutions between AtRALF4 and AtRALF19 contribute to their differential alkalization activity as well as that endogenous AtRALF19 or other AtRALFs' expression levels influence AtRALF4 effect on cells. AtRALF alkalization of the extracellular media or apoplast may be a prerequisite for other AtRALFs to associate with their receptors, as has been suggested by the pH dependence of AtRALF34 for binding to the ectodomain of the THESEUS1 (THE1) receptor (Gonneau et al., 2018). The pH dependence of AtRALF binding could also be related to the affinity of AtRALFs binding to the receptors, but this possibility has yet to be investigated.

The initial discovery of RALFs identified the inhibition of root elongation as an important function (Pearce et al., 2001). A clearer role for AtRALFs in the inhibition of cell expansion emerged from the work on AtRALF1, which remains one of the most studied RALF peptides. Overexpression of AtRALF1 causes bushy, semidwarf plants with small leaves, short roots, and a reduction in the number of both lateral roots and small cells in roots (Matos et al., 2008; Bergonci et al., 2014a). On the other hand, knockdown or knockout mutants of the *AtRALF1* gene showed the opposite phenotype: plants with long roots and long hypocotyls, increased lateral root number, and large root cells (Bergonci et al., 2014a). Biochemically isolated and *Escherichia coli*-produced AtRALF1 was active in the typical cytoplasmic  $\text{Ca}^{2+}$ -assay and inhibited cell expansion and root growth (Haruta et al., 2008; Mingossi et al., 2010; Morato do Canto et al., 2014).

Other RALF peptides are also involved in regulating growth (overexpression of two other AtRALF sequences, AtRALF8 and AtRALF23, also resulted in semidwarf plants; Srivastava et al., 2009; Atkinson et al., 2013). The *AtRALF23* gene was found to be downregulated by brassinolide (BL) treatment, and its overexpression also

resulted in semidwarf plants with reduced root growth and a compromised BL response (Nemhauser et al., 2004; Srivastava et al., 2009). Antagonism between AtRALFs and BL was also demonstrated when AtRALF-induced genes involved in cell wall rearrangement were identified (Table 1; Bergonci et al., 2014a, 2014b). Brassinosteroids (BRs) are important regulators of growth and development and play a critical role in cell expansion. BRs are mainly perceived by the cell membrane receptor kinase BRASSINOSTEROID INSENSITIVE1 (BRI1) that, upon binding of BR, associates with its coreceptor BRI1-ASSOCIATED KINASE1 (BAK1) to up-regulate BR cellular responses (Belkhadir and Jaillais, 2015). AtRALF1 also binds BAK1, and BAK1 is critical for the inhibition of root growth and cell expansion. The antagonistic relationship between AtRALF and BRs could well be regulated by switching both signaling pathways on and off through BAK1 (Dressano et al., 2017).

The specific developmental stages or environmental conditions in which AtRALF1-FER signaling contributes to root growth regulation remains unclear. One plausible hypothesis states that the AtRALF1-FER pair controls growth suppression of roots under dim light, because *fer* seedlings display a faster growth phenotype than wild type in these conditions. FER and other members of CrRLK1Ls may also be involved in cell growth regulation during photomorphogenesis, because those genes were shown to be transcriptionally induced by BL, a growth regulator that overlaps with light signaling (Li et al., 1996; Guo et al., 2009a, 2009b; Clouse, 2011; Zhu et al., 2013). Additionally, *fer-4* root elongation was not completely suppressed when tested at higher AtRALF1 concentrations (Haruta et al., 2014), suggesting that another member of the CrRLK1L family is also involved in sensing AtRALF1 during root growth and development.

FER was originally discovered as the female determinant of sperm cell release from the pollen tube as it penetrates the synergid cell of the ovule (Escobar-Restrepo et al., 2007). The discovery of FER as the AtRALF1 receptor and the abundance of RALF peptides in the reproductive tissues led to a concerted effort in the research community to unravel the roles of

these peptides in plant reproduction. The interest in reproduction-associated RALFs began when a family of AtRALF-like sequences was identified in the expression sequence tag library derived from the ovule of *Solanum chacoense* (Germain et al., 2005), which led to the critical finding that silencing *ScRALF3* expression by RNA interference caused developmental arrest in pollen mitosis and abnormal nuclear distribution in the embryo sac (Chevalier et al., 2013; Loubert-Hudon et al., 2020). *ScRALF3* mRNA expression and promoter activity assay results suggested that *ScRALF3* functions as a secreted signal regulating cell-to-cell communication between the embryo sac and the sporophytic tissue surrounding the ovule, demonstrating the first example of RALF physiology occurring outside of the roots. Additional research efforts in *Arabidopsis* led to the discovery of another pair of RALF-CrRLK1L interactors: pollen-expressed AtRALF4 and AtRALF19 are peptide ligands of the pollen-expressed CrRLK1L BUPS1/2 and ANX1/2 receptors. These receptor-bound complexes are responsible for maintaining cell integrity during pollen tube germination and growth (Ge et al., 2017). As pollen tubes approach ovules, ovule-expressed AtRALF34 peptide competes with pollen-expressed AtRALF4 and AtRALF19 for binding to ANX1/2 or BUPS1/2, allowing sperm cell discharge (Ge et al., 2017).

Recently, two CrRLK1L receptor-like kinases, HERCULES1 (HERK1) and ANJEA (ANJ), were identified as female determinants of pollen tube reception (Galindo-Trigo et al., 2020). Both HERK1 and ANJ interact with FER and, although not yet proved, an analogous mechanism was proposed where pollen-derived RALFs would act upon receptor heterocomplexes made of FER, HERK1, and ANJ assembled on female tissues to control pollen tube reception (Galindo-Trigo et al., 2020). Fertilization is the result of an interplay between signal molecules, their receptors, second messengers, and cell wall components. Establishing RALFs, CrRLK1Ls, and LLG coreceptor pairs involved in each part of this process and their relation to each of the intracellular events is presently a matter of intense research interest (Ge et al., 2019a). It is also essential to understand whether a RALF-CrRLK1L-LLG ternary complex multimerizes into higher-order complexes that may be composed of different sets of RALFs, CrRLK1Ls, and LLGs (Ge et al., 2019b).

## REGULATION OF BIOTIC AND ABIOTIC STRESS RESPONSES BY RALF PEPTIDES

Recently, several physiological studies have implicated AtRALFs-FER signaling in biological pathways including innate immune responses, stress responses mediated by abscisic acid (ABA), and a nutrient stress condition (Table 1).

It has been suggested that FER may act as a scaffold protein for the assembly of an immune receptor complex (Chakravorty et al., 2018; Duan and Cheung, 2018; Haruta et al., 2018a). An AtRALF23-FER interaction

potentially coordinates the assembly of an immune complex involving EF-TU RECEPTOR (EFR), FLAGELLIN SENSING 2 (FLS2), and BAK1 (Stegmann et al., 2017). Interestingly, a search for mutants that could recover the immune function lost in *bak1-5* mutants resulted in the isolation of the subtilase SITE-1 PROTEASE (S1P)/SBT6.1, the prohormone convertase that processes AtRALF23 (Srivastava et al., 2009; Stegmann et al., 2017). This finding is striking because it shows that proteolytic processing of AtRALF23 from its precursor inhibits plant immune responses that are activated by cellular perception of flg22 or elf18 elicitor peptides by FLS2 or EFR. RALF was known to affect the level of reactive oxygen species (ROS) production in *N. attenuata*, and decreased levels of ROS accumulation were found in *irRALF* plants, although that observation was related to root hair development (Wu et al., 2007). AtRALF23 and AtRALF33 are suppressors of ROS production when elicited by elf18, a peptide derived from the bacterial elongation factor Tu (Table 1); however AtRALF17, a peptide whose precursor lacks a canonical convertase processing site, induces ROS production instead of suppressing it, further suggesting the importance of S1P in RALF-mediated immune responses (Stegmann et al., 2017). AtRALF32 neither induces nor suppresses ROS production elicited by elf18 and also lacks a conserved processing site yet has a propeptide region (Stegmann et al., 2017). This further suggests that different AtRALFs have different biological effects, which might be contingent on the biological context (such as tissue type) and how the mature peptide is processed.

MYC2 transcription factor, which was identified from a transcriptome analysis of the *fer* mutant and promoter motif analysis, was also shown to be a FER kinase substrate and act as a negative regulator of immunity to *Pseudomonas syringae* pv. *tomato* DC3000 (pst DC3000; Guo et al., 2018). *Arabidopsis* plants treated with AtRALF23 suppress the induction of immune responses, while plants overexpressing the *AtRALF23* gene are more susceptible to pathogenic infection. Thus, AtRALF23 ligand itself is a negative regulator of immune response. Meanwhile, *Arabidopsis fer* mutants showed hypersensitivity to this pathogen, indicating that the normal function of FER is to up-regulate the immune response by acting as a scaffold for proteins involved in pathogen-associated molecular pattern-triggered immunity and by suppressing jasmonate-responsive signaling.

As a Ser/Thr receptor kinase, FER has been proposed to activate itself by autophosphorylation (Escobar-Restrepo et al., 2007), and it exhibits kinase activity in vitro on both itself and on synthetic substrates (Minkoff et al., 2017; Haruta et al., 2018a). To date, there is evidence of additional substrates of FER, including ARABIDOPSIS TOXICOS EN LEVADURA6 (ATL6) and ERBB3-BINDING PROTEIN1 (EBP1; Li et al., 2018; Xu et al., 2019).

AtRALF-CrRLK1L signaling also plays a role in stomatal opening and stress responses. AtRALF8 was

### OUTSTANDING QUESTIONS

- How does AtRALF-induced phosphorylation of the proton pump AHA2 influence H<sup>+</sup>-ATPase function and lead to the inhibition of cell elongation?
- Which FERONIA co-receptors participate in AtRALF1 perception at the cell surface and initiate protein phosphorylation signaling to suppress root growth?
- The AtRALF peptide family is diverse and binds to a large array of receptors and coreceptors. What is the mechanism governing the specificity of AtRALF-receptor interactions and the assembly of distinct signaling complexes?
- Do biological conditions affect whether AtRALF members act as agonists or antagonists?
- If extracellular alkalinization, the canonical hallmark of AtRALF activity, is not essential for AtRALF functions, then what is the biological role(s) of pH increase?

originally identified as one of the genes that were transcriptionally upregulated in Arabidopsis plants exposed to a combination of water deficit and nematode stresses (Atkinson et al., 2013). FER was identified in a screen for protein interactors of AGB1, a G protein  $\beta$ -subunit that is involved in ABA-mediated regulation of stomatal movements. AtRALF1 was then shown to regulate stomatal aperture, which is important in both immune and drought responses, through FER (Table 1; Yu and Assmann, 2018). This work was the first to demonstrate that AtRALF1 inhibits stomatal opening and induces its closure through FER, which functions like a G protein-coupled receptor. Because FER and AGB1 are both known to be involved in salt stress responses mediated by ABA, the mutant phenotypes of *fer* and *agb1* were compared in two different cell types, guard cells and roots, to examine epistatic relationship of the two genes (Yu and Assmann, 2018). Both *fer* and *agb1* mutants perform poorly in high-salt conditions and the double mutant demonstrated an even more severe phenotype of hypersensitivity to the salt stress (Yu and Assmann, 2018). When salt was combined with AtRALF1 treatment, these plants also demonstrated a hypersensitivity phenotype. This phenotype was observed again when AtRALF22 was overexpressed (Zhao et al., 2018). Interestingly, in a different study, *atralf1* mutant lines did not show any phenotypic difference from control plants when exposed to high-salt conditions, and AtRALF1 overexpressors showed increased tolerance to salt (Feng et al., 2018). However,

contrasting results for AtRALF-treated plants or for plants in which endogenous levels of expression are genetically manipulated have been reported (Wiegheas et al., 2019).

### CONCLUSION

The initial biochemical screening for extracellular alkalinization activity led to the discovery of the first RALF peptide nearly 20 years ago. As the proton gradient at the cell surface is key to cell growth, this peptide family has long been thought to regulate cell expansion. Consistent with this prediction, both biochemical and genetic experiments showed that a plasma membrane H<sup>+</sup>-ATPase acts downstream of AtRALF1 and FER action in cellular signaling suppressing root growth (Haruta et al., 2018b). Moreover, an increasing body of literature supports cell growth regulation by RALF ligands and their cognate receptor complexes in other tissues. As demonstrated by molecular, genetic, and biochemical studies of the RALF peptide and CrRLK1L receptor families, many physiological responses interact with cell elongation signaling in the plant. The wealth of knowledge that has been produced on RALF biology over the last two decades has also opened many new questions about this complex area of plant physiology (see Outstanding Questions). To fully understand how this complex signaling mechanism is regulated, experimental approaches with high spatial and/or temporal resolution will be needed. We believe that many additional fruitful RALF-CrRLK1L studies will follow to understand plant cell growth regulation mechanisms initiated at the interface of plasma membrane and cell wall.

### ACKNOWLEDGMENTS

The authors thank the reviewers for critical reading of the manuscript and suggesting substantial improvements. The authors apologize for not citing other important studies in this review due to page limitations.

Received October 25, 2019; accepted February 16, 2020; published February 18, 2020.

### LITERATURE CITED

- Atkinson NJ, Lilley CJ, Urwin PE (2013) Identification of genes involved in the response of Arabidopsis to simultaneous biotic and abiotic stresses. *Plant Physiol* 162: 2028–2041
- Belkhadir Y, Jaillais Y (2015) The molecular circuitry of brassinosteroid signaling. *New Phytol* 206: 522–540
- Bergonci T, Ribeiro B, Ceciliato PHO, Guerrero-Abad JC, Silva-Filho MC, Moura DS (2014a) *Arabidopsis thaliana* RALF1 opposes brassinosteroid effects on root cell elongation and lateral root formation. *J Exp Bot* 65: 2219–2230
- Bergonci T, Silva-Filho MC, Moura DS (2014b) Antagonistic relationship between AtRALF1 and brassinosteroid regulates cell expansion-related genes. *Plant Signal Behav* 9: e976146
- Campbell L, Turner SR (2017) A comprehensive analysis of RALF proteins in green plants suggests there are two distinct functional groups. *Front Plant Sci* 8: 37
- Campos WF, Dressano K, Ceciliato PHO, Guerrero-Abad JC, Silva AL, Fiori CS, Morato do Canto A, Bergonci T, Claus LAN, Silva-Filho MC, et al (2018) *Arabidopsis thaliana* rapid alkalinization factor 1-mediated



- root growth inhibition is dependent on calmodulin-like protein 38. *J Biol Chem* **293**: 2159–2171
- Cao J, Shi F (2012) Evolution of the AtRALF gene family in plants: Gene duplication and selection patterns. *Evol Bioinform Online* **8**: 271–292
- Chakravorty D, Yu Y, Assmann SM (2018) A kinase-dead version of FERONIA receptor-like kinase has dose-dependent impacts on rosette morphology and RALF1-mediated stomatal movements. *FEBS Lett* **592**: 3429–3437
- Chen J, Yu F, Liu Y, Du C, Li X, Zhu S, Wang X, Lan W, Rodriguez PL, Liu X, et al (2016) FERONIA interacts with ABI2-type phosphatases to facilitate signaling cross-talk between abscisic acid and RALF peptide in *Arabidopsis*. *Proc Natl Acad Sci USA* **113**: E5519–E5527
- Chevalier E, Loubert-Hudon A, Matton DP (2013) SCRALF3, a secreted RALF-like peptide involved in cell-cell communication between the sporophyte and the female gametophyte in a solanaceous species. *Plant J* **73**: 1019–1033
- Clouse SD (2011) Brassinosteroid signal transduction: From receptor kinase activation to transcriptional networks regulating plant development. *Plant Cell* **23**: 1219–1230
- Covey PA, Subbiah CC, Parsons RL, Pearce G, Lay FT, Anderson MA, Ryan CA, Bedinger PA (2010) A pollen-specific RALF from tomato that regulates pollen tube elongation. *Plant Physiol* **153**: 703–715
- Dressano K, Ceciliato PHO, Silva AL, Guerrero-Abad JC, Bergonci T, Ortiz-Moreno FA, Bürger M, Silva-Filho MC, Moura DS (2017) BAK1 is involved in AtRALF1-induced inhibition of root cell expansion. *PLoS Genet* **13**: e1007053
- Du S, Qu LJ, Xiao J (2018) Crystal structures of the extracellular domains of the CrRLK1L receptor-like kinases ANXUR1 and ANXUR2. *Protein Sci* **27**: 886–892
- Duan Q, Cheung AY (2018) Context-specific dependence on FERONIA kinase activity. *FEBS Lett* **592**: 2392–2394
- Duan Q, Kita D, Li C, Cheung AY, Wu HM (2010) FERONIA receptor-like kinase regulates RHO GTPase signaling of root hair development. *Proc Natl Acad Sci USA* **107**: 17821–17826
- Dünser K, Gupta S, Herger A, Feraru MI, Ringli C, Kleine-Vehn J (2019) Extracellular matrix sensing by FERONIA and Leucine-Rich Repeat Extensins controls vacuolar expansion during cellular elongation in *Arabidopsis thaliana*. *EMBO J* **38**: e100353
- Escobar-Restrepo JM, Huck N, Kessler S, Gagliardini V, Gheyselinck J, Yang WC, Grossniklaus U (2007) The FERONIA receptor-like kinase mediates male-female interactions during pollen tube reception. *Science* **317**: 656–660
- Feng W, Kita D, Peaucelle A, Cartwright HN, Doan V, Duan Q, Liu M-C, Maman J, Steinhorst L, Schmitz-Thom I, et al (2018) The FERONIA receptor kinase maintains cell-wall integrity during salt stress through Ca<sup>2+</sup> signaling. *Curr Biol* **28**: 666–675.e5
- Frederick RO, Haruta M, Tonelli M, Lee W, Cornilescu G, Cornilescu CC, Sussman MR, Markley JL (2019) Function and solution structure of the *Arabidopsis thaliana* RALF8 peptide. *Protein Sci* **28**: 1115–1126
- Galindo-Trigo S, Blanco-Tourrián N, DeFalco TA, Wells ES, Gray JE, Zipfel C, Smith LM (2020) CrRLK1L receptor-like kinases HERK1 and ANJEA are female determinants of pollen tube reception. *EMBO Rep* **21**: e48466
- Ge Z, Bergonci T, Zhao Y, Zou Y, Du S, Liu MC, Luo X, Ruan H, García-Valencia LE, Zhong S, et al (2017) *Arabidopsis* pollen tube integrity and sperm release are regulated by RALF-mediated signaling. *Science* **358**: 1596–1600
- Ge Z, Cheung AY, Qu LJ (2019a) Pollen tube integrity regulation in flowering plants: insights from molecular assemblies on the pollen tube surface. *New Phytol* **222**: 687–693
- Ge Z, Dresselhaus T, Qu LJ (2019b) How CrRLK1L receptor complexes perceive RALF signals. *Trends Plant Sci* **24**: 978–981
- Ge Z, Zhao Y, Liu MC, Zhou LZ, Wang L, Zhong S, Hou S, Jiang J, Liu T, Huang Q, et al (2019c) LLG2/3 are coreceptors in BUPs/ANX-RALF signaling to regulate *Arabidopsis* pollen tube integrity. *Curr Biol* **29**: 3256–3265.e5
- Germain H, Chevalier E, Caron S, Matton DP (2005) Characterization of five RALF-like genes from *Solanum chacoense* provides support for a developmental role in plants. *Planta* **220**: 447–454
- Gonneau M, Desprez T, Martin M, Doblas VG, Bacete L, Miart F, Sormani R, Hématy K, Renou J, Landrein B, et al (2018) Receptor kinase THESEUS1 is a rapid alkalization factor 34 receptor in *Arabidopsis*. *Curr Biol* **28**: 2452–2458.e4
- Guo H, Li L, Ye H, Yu X, Algreen A, Yin Y (2009a) Three related receptor-like kinases are required for optimal cell elongation in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **106**: 7648–7653
- Guo H, Nolan TM, Song G, Liu S, Xie Z, Chen J, Schnable PS, Walley JW, Yin Y (2018) FERONIA receptor kinase contributes to plant immunity by suppressing jasmonic acid signaling in *Arabidopsis thaliana*. *Curr Biol* **28**: 3316–3324.e6
- Guo H, Ye H, Li L, Yin Y (2009b) A family of receptor-like kinases are regulated by BES1 and involved in plant growth in *Arabidopsis thaliana*. *Plant Signal Behav* **4**: 784–786
- Haruta M, Constabel CP (2003) Rapid alkalization factors in poplar cell cultures. Peptide isolation, cDNA cloning, and differential expression in leaves and methyl jasmonate-treated cells. *Plant Physiol* **131**: 814–823
- Haruta M, Gaddameedi V, Burch H, Fernandez D, Sussman MR (2018a) Comparison of the effects of a kinase-dead mutation of FERONIA on ovule fertilization and root growth of *Arabidopsis*. *FEBS Lett* **592**: 2395–2402
- Haruta M, Monshausen G, Gilroy S, Sussman MR (2008) A cytoplasmic Ca<sup>2+</sup> functional assay for identifying and purifying endogenous cell signaling peptides in *Arabidopsis* seedlings: Identification of AtRALF1 peptide. *Biochemistry* **47**: 6311–6321
- Haruta M, Sabat G, Stecker K, Minkoff BB, Sussman MR (2014) A peptide hormone and its receptor protein kinase regulate plant cell expansion. *Science* **343**: 408–411
- Haruta M, Tan LX, Bushey DB, Swanson SJ, Sussman MR (2018b) Environmental and genetic factors regulating localization of the plant plasma membrane H<sup>+</sup>-ATPase. *Plant Physiol* **176**: 364–377
- Huck N, Moore JM, Federer M, Grossniklaus U (2003) The *Arabidopsis* mutant feronia disrupts the female gametophytic control of pollen tube reception. *Development* **130**: 2149–2159
- Li C, Liu X, Qiang X, Li X, Li X, Zhu S, Wang L, Wang Y, Liao H, Luan S, et al (2018) EBP1 nuclear accumulation negatively feeds back on FERONIA-mediated RALF1 signaling. *PLoS Biol* **16**: e2006340
- Li C, Wang L, Cui Y, He L, Qi Y, Zhang J, Lin J, Liao H, Lin Q, Yang T, et al (2016) Two FERONIA-like receptor (FLR) genes are required to maintain architecture, fertility, and seed yield in rice. *Mol Breed* **36**: 151
- Li C, Yeh FL, Cheung AY, Duan Q, Kita D, Liu MC, Maman J, Luu EJ, Wu BW, Gates L, et al (2015) Glycosylphosphatidylinositol-anchored proteins as chaperones and co-receptors for FERONIA receptor kinase signaling in *Arabidopsis*. *eLife* **4**: e06587
- Li J, Nagpal P, Vitart V, McMorris TC, Chory J (1996) A role for brassinosteroids in light-dependent development of *Arabidopsis*. *Science* **272**: 398–401
- Liu P, Haruta M, Minkoff BB, Sussman MR (2018) Probing a plant plasma membrane receptor kinase's three-dimensional structure using mass spectrometry-based protein footprinting. *Biochemistry* **57**: 5159–5168
- Loubert-Hudon A, Mazin BD, Chevalier É, Matton DP (2020) The SCRALF3 secreted peptide is involved in sporophyte to gametophyte signalling and affects pollen mitosis I. *Plant Biol (Stuttg)* **22**: 13–20
- Masachis S, Segorbe D, Turrà D, Leon-Ruiz M, Fürst U, El Ghalid M, Leonard G, López-Berges MS, Richards TA, Felix G, et al (2016) A fungal pathogen secretes plant alkalizing peptides to increase infection. *Nat Microbiol* **1**: 16043
- Matos JL, Fiori CS, Silva-Filho MC, Moura DS (2008) A conserved dibasic site is essential for correct processing of the peptide hormone AtRALF1 in *Arabidopsis thaliana*. *FEBS Lett* **582**: 3343–3347
- Mecchia MA, Santos-Fernandez G, Duss NN, Somoza SC, Boisson-Dernier A, Gagliardini V, Martínez-Bernardini A, Fabrice TN, Ringli C, Muschietti JP, et al (2017) RALF4/19 peptides interact with LRX proteins to control pollen tube growth in *Arabidopsis*. *Science* **358**: 1600–1603
- Mingossi FB, Matos JL, Rizzato AP, Medeiros AH, Falco MC, Silva-Filho MC, Moura DS (2010) SacRALF1, a peptide signal from the grass sugarcane (*Saccharum* spp.), is potentially involved in the regulation of tissue expansion. *Plant Mol Biol* **73**: 271–281
- Minkoff BB, Makino SI, Haruta M, Beebe ET, Wrobel RL, Fox BG, Sussman MR (2017) A cell-free method for expressing and reconstituting membrane proteins enables functional characterization of the plant receptor-like protein kinase FERONIA. *J Biol Chem* **292**: 5932–5942
- Morato do Canto A, Ceciliato PHO, Ribeiro B, Ortiz Morea FA, Franco Garcia AA, Silva-Filho MC, Moura DS (2014) Biological activity of nine recombinant AtRALF peptides: implications for their perception and function in *Arabidopsis*. *Plant Physiol Biochem* **75**: 45–54

- Moussu S, Augustin S, Roman AO, Broyart C, Santiago J** (2018) Crystal structures of two tandem malectin-like receptor kinases involved in plant reproduction. *Acta Crystallogr D Struct Biol* **74**: 671–680
- Nemhauser JL, Mockler TC, Chory J** (2004) Interdependency of brassinosteroid and auxin signaling in *Arabidopsis*. *PLoS Biol* **2**: E258
- Pearce G, Moura DS, Stratmann J, Ryan CA Jr.** (2001) RALF, a 5-kDa ubiquitous polypeptide in plants, arrests root growth and development. *Proc Natl Acad Sci USA* **98**: 12843–12847
- Pearce G, Yamaguchi Y, Munske G, Ryan CA** (2010) Structure-activity studies of RALF, Rapid Alkalinization Factor, reveal an essential—YISY—motif. *Peptides* **31**: 1973–1977
- Rotman N, Rozier F, Boavida L, Dumas C, Berger F, Faure JE** (2003) Female control of male gamete delivery during fertilization in *Arabidopsis thaliana*. *Curr Biol* **13**: 432–436
- Scheer JM, Pearce G, Ryan CA** (2005) LeRALF, a plant peptide that regulates root growth and development, specifically binds to 25 and 120 kDa cell surface membrane proteins of *Lycopersicon peruvianum*. *Planta* **221**: 667–674
- Srivastava R, Liu JX, Guo H, Yin Y, Howell SH** (2009) Regulation and processing of a plant peptide hormone, AtRALF23, in *Arabidopsis*. *Plant J* **59**: 930–939
- Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, Holton N, Belkhadir Y, Zipfel C** (2017) The receptor kinase FER is a RALF-regulated scaffold controlling plant immune signaling. *Science* **355**: 287–289
- Thynne E, Saur IML, Simbaqueba J, Ogilvie HA, Gonzalez-Cendales Y, Mead O, Taranto A, Catanzariti AM, McDonald MC, Schwessinger B, et al** (2017) Fungal phytopathogens encode functional homologues of plant rapid alkalinization factor (RALF) peptides. *Mol Plant Pathol* **18**: 811–824
- Wiegand A, Prüfer D, Schulze Gronover C** (2019) Loss of function mutation of the Rapid Alkalinization Factor (RALF1)-like peptide in the dandelion *Taraxacum kok-saghyz* entails a high-biomass taproot phenotype. *PLoS One* **14**: e0217454
- Wu J, Kurten EL, Monshausen G, Hummel GM, Gilroy S, Baldwin IT** (2007) NaRALF, a peptide signal essential for the regulation of root hair tip apoplastic pH in *Nicotiana attenuata*, is required for root hair development and plant growth in native soils. *Plant J* **52**: 877–890
- Xiao Y, Stegmann M, Han Z, DeFalco TA, Parys K, Xu L, Belkhadir Y, Zipfel C, Chai J** (2019) Mechanisms of RALF peptide perception by a heterotypic receptor complex. *Nature* **572**: 270–274
- Xu G, Chen W, Song L, Chen Q, Zhang H, Liao H, Zhao G, Lin F, Zhou H, Yu F** (2019) FERONIA phosphorylates E3 ubiquitin ligase ATL6 to modulate the stability of 14-3-3 proteins in response to the carbon/nitrogen ratio. *J Exp Bot* **70**: 6375–6388
- Yu Y, Assmann SM** (2018) Inter-relationships between the heterotrimeric G $\beta$  subunit AGB1, the receptor-like kinase FERONIA, and RALF1 in salinity response. *Plant Cell Environ* **41**: 2475–2489
- Zhao C, Zayed O, Yu Z, Jiang W, Zhu P, Hsu CC, Zhang L, Tao WA, Lozano-Durán R, Zhu JK** (2018) Leucine-rich repeat extensin proteins regulate plant salt tolerance in *Arabidopsis*. *Proc Natl Acad Sci USA* **115**: 13123–13128
- Zhu JY, Sae-Seaw J, Wang ZY** (2013) Brassinosteroid signalling. *Development* **140**: 1615–1620
- Zhu L, Chu LC, Liang Y, Zhang XQ, Chen LQ, Ye D** (2018) The *Arabidopsis* CrRLK1L protein kinases BUPS1 and BUPS2 are required for normal growth of pollen tubes in the pistil. *Plant J* **95**: 474–486