

RT-MAP-8406

**ARE THERE FREE GROUPS
IN DIVISION RINGS?**

Jairo Z. Gonçalves
Arnaldo Mandel

Setembro 1984

ARE THERE FREE GROUPS IN DIVISION RINGS?

Jairo Z. Gonçalves and Arnaldo Mandel

Instituto de Matemática e Estatística,
Universidade de São Paulo,
São Paulo, Brazil

September, 1984

Address: IMEUSP
Caixa Postal 20570
01000 - SÃO PAULO - SP
BRAZIL

MR subject classification (1980): Primary: 16A39. Secondary:
16A25, 16A70, 20E05.

ABSTRACT

The following conjecture is investigated: a noncentral subnormal subgroup of the multiplicative group of a division ring contains a noncyclic free subgroup. Special cases are proved, entailing several known commutativity theorems. Also a new framework is presented for some kinds of commutativity theorems, based on the existence of (group) words for which one can always find an appropriate substitution by elements of such a subnormal subgroup that yields a noncentral element. Several families of such words are given; one gets commutativity theorems imposing some restrictions (like periodicity) to the image of these words.

ARE THERE FREE GROUPS IN DIVISION RINGS?

J.Z. Gonçalves

A. Mandel

1) INTRODUCTION

We are concerned with a problem posed by Lichtman [8], which, at least for discussion purposes, we formulate as:

CONJECTURE 1: *The multiplicative group of a noncommutative division ring contains a noncyclic free subgroup.*

A stronger conjecture will actually be approached:

CONJECTURE 2: *Any noncentral subnormal subgroup of the multiplicative group of a division ring contains a noncyclic free subgroup.*

An affirmative answer to either of these would be an umbrella for several known theorems. For instance, in [5], Chapter 3, there are several group-theoretic conditions whose occurrence in the multiplicative group of a division ring, entail commutativity of the ring; those conditions, to which we return later, are impossible in presence of a noncommutative free subgroup, hence commutativity would follow from Conjecture 1. (There are other commutativity theorems based on algebraic conditions,

however, which are apparently unrelated to this conjecture). Further work of Herstein [6,7], Stuth [12] and others extend some of these results to conditions on a subnormal subgroup which imply it is central; these theorems would easily follow from Conjecture 2 (actually, these papers are probably stepping stones in any attempt to settle the Conjecture). As direct evidence towards these conjectures, we quote Gonçalves [2], Lichtman [9,10], and the present work.

On this line, the main results appear in section 3, and they can be put together as:

THEOREM: *Conjecture 2 holds for a subnormal subgroup which contains a noncentral element x such that either (a) x is algebraic over the center Z of the ring, and the field $Z(x)$ admits a nontrivial Z -automorphism, or (b) $x^p \in Z$, where $p = 2$ or $\text{char} Z (> 0)$.*

The conjectures suggest a new perspective on some commutativity theorems. Let w be a word on the free group on n generators. For any group G , w induces, by substitution, a map of the n -fold cartesian product $G \times \dots \times G$ to G , which for short we call the *word map* w on G . Say that w is *noncentral* for G if its image is not contained in the center $C(G)$; equivalently, if the relation $w=1$ does not hold in $G/C(G)$. Clearly, if G contains a nonabelian free subgroup, every word is noncentral for G . Say that w is an N -word (SN-word) provided it is noncentral for

every G that is (a noncentral subnormal subgroup of) the multiplicative subgroup of a noncommutative division ring. Sections 4 and 5 present several such words (actually, all words should be SN). Many of the aforementioned commutativity theorems make statements about the image of an SN word map; they are covered in section 4.

We fix some notation throughout: D is a division ring with center Z , their multiplicative subgroups are D^* , Z^* ; N is a subnormal subgroup of D^* . We abbreviate "contains a noncyclic free group" to "contains a free group". The group theoretic commutator $xyx^{-1}y^{-1}$ will be denoted (x,y) , while $[x,y] = xy-yx$. We also use the iterated inner derivation $[x,y]^{(k)}$ given by: $[x,y]^{(0)} = x$, $[x,y]^{(k)} = [[x,y]^{(k-1)}, y]$.

2) A COMPUTATIONAL TOOL KIT

The combined use of the lemmas at this section is at the core of most results in this paper. The first one is a takeoff from Herstein [7]. We denote by $C_X(p)$ the coefficient of X in p , where p is a polynomial in the indeterminate X .

DESCENDING LEMMA: Let $a, u \in D^*$, and suppose that

$$u^n = \alpha \in Z \text{ for some } n > 0. \text{ Consider}$$

the sequences of polynomials $p_i = p_i(a, u; X)$, $\bar{p}_i \in D[X]$, $q_i \in Z[X^n]$, $i=0, 1, \dots$ defined by:

$$p_0 = (1-uX)(1-\alpha X^n), \bar{p}_0 = 1 + uX + u^2X^2 + \dots + u^{n-1}X^{n-1}, q_0 = 1 - \alpha X^n, \text{ and}$$

for $i > 0$, $p_i = p_{i-1} a \bar{p}_{i-1} a^{-1}$, $\bar{p}_i = a p_{i-1} a^{-1} \bar{p}_{i-1}$, $q_i = q_{i-1}^2$. Then:

$$a) \quad p_i(0) = \bar{p}_i(0) = q_i(0) = 1$$

$$b) \quad C_X(p_i) = -[u, a]^{(i)} a^{-i} = -C_X(\bar{p}_i),$$

Further, suppose that $N_k \triangleleft N_{k-1} \triangleleft \dots \triangleleft N_0 = D^*$ is a normal series, that $a \in N_k$ and that $\lambda \in Z$ is such that $\lambda^n \cdot \alpha \neq 1$, and define elements $d_i = d_i(\lambda) = d_i(a, u; \lambda) = p_i(\lambda) / q_i(\lambda)$. Then:

$$c) \quad d_i(\lambda) \in N_i, \quad i=0, 1, \dots, k$$

$$d) \quad d_i(\lambda)^{-1} = \bar{p}_i(\lambda) / q_i(\lambda).$$

Proof: All the proofs are by induction on i . The case $i=0$ is trivial for (a), (b), (c), while (d) follows

from the identity $1 - \lambda^n \alpha = (1 - \lambda)(1 + \lambda u + \dots + \lambda^{n-1} u^{n-1})$.
 So assume $i > 0$. Again (a) is trivial, and using (a) we compute:

$C_x(p_i) = C_x(p_{i-1}) + a C_x(\bar{p}_{i-1}) a^{-1} = -u^{(i-1)} a^{-(i-1)} + a u^{(i-1)} a^{-(i-1)} a^{-1} = -u^{(i)} a^{-i}$, and a similar computation yields $C_x(\bar{p}_i)$. Further, observing that $q_i(\lambda) \in Z^*$, with a liberal use of the inductive hypothesis we get:

$d_i(\lambda) = p_i(\lambda)/q_i(\lambda) = (p_{i-1}(\lambda)/q_{i-1}(\lambda) a(\bar{p}_{i-1}(\lambda)/q_{i-1}(\lambda)) a^{-1}) = d_{i-1} a d_{i-1}^{-1} a^{-1} \in N$
 and $d_i(\lambda)^{-1} = (d_{i-1} a)^{-1} = (a, d_{i-1}) = \bar{p}_i(\lambda)/q_i(\lambda)$. \square

The next results form a piece of folklore we have found useful to make explicit.

LEMMA 2.1: *If $a, u \in D$ are such that $[u, a] \neq 0$ and there exists a $k > 0$ such that $[u, a]^{(k)} = 0$, then there exists an $x \in D$ such that $a^{-1} x a = x + 1$.*

Proof: Choose $r > 0$ such that $[u, a]^{(r)} \neq 0$ but $[u, a]^{(r+1)} = 0$ and set $x = a [u, a]^{(r-1)} ([u, a]^{(r)})^{-1}$. Then $[x, a] = a$ and the result follows. \square

LEMMA 2.2: *Suppose that the polynomial $p \in D[X]$ maps infinitely many elements of Z to Z . Then $p \in Z[X]$.*

Proof: With $p = a_0 + a_1 X + \dots + a_n X^n$, let $\lambda_0, \dots, \lambda_n$ be distinct elements of Z and let $p(\lambda_i) = z_i$. Then $(a_0, a_1, \dots, a_n) = (z_0, z_1, \dots, z_n) ((\lambda_i^j))^{-1}$, where $((\lambda_i^j))$ is a Vandermonde matrix. \square

LEMMA 2.3: *For every $a \in N-Z$, there is an N -conjugate u of a such that $(u, a) \neq 1$.*

Proof: Deny. Let N_1 be the group generated by the N -conjugates of a thus a is in the center of N_1 . Since $N_1 \triangleleft N \triangleleft D^*$, Stuth [12, Thm. 2] implies that $a \in Z$, a contradiction. \square

We can now present a simplified proof of a theorem of Herstein [6], which also illustrates the use of the Lemmas.

THEOREM 2.4: *If $N \triangleleft D^*$ is periodic, then $N \subseteq Z^*$*

Proof: We begin with a quick run through some of Herstein's arguments. The proof is by contradiction, hence we assume that N contains a noncentral element a .

Since $a^n = 1$ for some $n > 2$ (as $a^2 = 1$ implies $a = \pm 1$), we can choose $r > 1$ such that the map $a \rightarrow a^r$ induces a non-trivial automorphism of $Z(a)$. By the Noether-Skolem theorem, there is a $w \in D^*$ such that $w^{-1} a w = a^r$. Thus, for some $k > 0$ $w^k a w^k = a$, and $D_1 = \{ \sum \alpha_{ij} a^i w^j \mid 0 \leq i < n, 0 \leq j < r, \alpha_{ij} \in Z(w^r) \}$ is a finite dimensional $Z(w^r)$ -division-algebra (see also

Lemma 3.2), Now $N \cap D_1^*$ is subnormal in D_1^* , periodic, and contains the noncentral element a . Thus, by substituting D_1 for D , N_1 for N , we obtain the additional hypothesis that $[D:Z] < \infty$.

By Lemma 2.3, there exists $u \in N$ such that $(a,u) \neq 1$. Let G be the subgroup of N generated by $\{a,u\}$. As $[D:Z] < \infty$, G is isomorphic to a group of matrices; also it is finitely generated and periodic, hence by the affirmative answer to the Burnside problem, G is finite. Denote by F the prime field of Z and let k be the linear span of G over F . Then k is a finite dimensional F -division-algebra, and Wedderburn's Theorem implies that $F=\mathbb{Q}$

We refresh notation again by substituting k for D , $k \cap N$ for N , so that now D is a finite dimensional \mathbb{Q} -division algebra..

Let $m = [D:\mathbb{Q}]$, so that D^* can be identified with a subgroup of $GL(m,\mathbb{Q})$. If $x \in D^*$ is periodic of order n , the degree $\phi(n)$ of the cyclotomic polynomial of index n must be at most m . As $\lim_{r \rightarrow \infty} \phi(r) = \infty$ there exist an $n > 0$ such that every periodic element of D^* satisfies $x^n = 1$.

We now apply the Descending Lemma to a,u . For some k , for each $\lambda \in \mathbb{Q} - \{\pm 1\}$, we have that $p_k(\lambda)/q_k(\lambda) \in N$ so that $p_k(\lambda)^n/q_k(\lambda)^n = 1$. Hence, we may apply Lemma 2.2 to conclude that $p_k^n \in Z[X]$. As $p_k(0) = 1$, we obtain $C_x(p_k^n) = n C_x(p_k) = -n[u,a]^{(k)} a^{-k} \in Z$, thus $[u,a]^{(k+1)} = 0$. By Lemma 2.1, there is an x such that $axa = x+1$, therefore, $x = a^{-n} x a^n = x+n$, the final contradiction as $n > 0$. \square

3) SPECIAL ELEMENTS ENTAILING FREE GROUPS

One of the authors has proved [2] that Conjecture 2 is true when $[D:Z]$ is finite. This will be used as a basic tool, so we restate it in a convenient form, and present the proof for completeness.

LEMMA 3.1. *Suppose that D_1 is a subdivision ring of D , finite dimensional over its center Z_1 . If $N \cap D_1$ is not in the center of D_1 , then N contains a free group.*

Proof: Indeed, $N \cap D_1^*$ is a noncentral subnormal subgroup of D_1^* , hence we retrieve our original D, N, Z setting with the additional hypothesis that $[D:Z]=r<\infty$. Thus N is isomorphic to a group of matrices and, if it contains no free group, Tits' Theorem [13] implies that there is a normal solvable subgroup L of N such that N/L is locally finite. Theorem 4 of Stuth [12] implies that $L \subseteq Z^*$. Let η be the norm on D to Z and let N' be the commutator subgroup of N . If $x \in N'$, $\eta(x)=1$, and $x^n = \lambda \in Z$ for some n , hence $\lambda^n = \eta(\lambda) = \eta(x^n) = \eta(x)^n = 1$. It follows that N' is a periodic subnormal subgroup of D^* . By Theorem 2.4, $N' \subseteq Z^*$, whence N is solvable and again by the same theorem of Stuth, $N \subseteq Z^*$, a contradiction. \square

Using this Lemma at appropriate places, one can change the proof of Lichtman [9, Thm. 2] extending

his result to subnormal subgroups of D^* ,

Our main division subring producing tool is:

LEMMA 3.2: Let $x \in D^*$ and suppose that conjugation by x induces a nontrivial automorphism of finite order of a commutative subring R of D . If $y \in R$ does not commute with x , then $Z(x,y)$ is contained in a division subring of D which is finite dimensional over its center.

Proof: Let $y_i = x^{-i}yx^i$, $i=1,2, \dots$; by hypothesis,

$y_k=y$ for some $k>1$. Consider the polynomial

$$p(X) = \prod_{i=1}^k (X - y_i) = \sum_{i=0}^k w_i X^i \in R[X].$$

Let Z_1 be the division

subring of D generated over Z by $\{w_0, w_1, \dots, w_k, x^k\}$, and

let D_1 be the Z_1 -module spanned by the monomials

$$x^i \cdot y_1^{i_1} \dots y_k^{i_k}, \quad i, i_1, \dots, i_k \geq 0.$$

Since $x^{-1}px=p$, x commutes with each w_i , and as $x^{-k}yx^k=y$, x^k commutes with each y_i . Hence Z_1 is a field and it commutes with each monomial. The relations

$$y_i y_j = y_j y_i, \quad \text{and} \quad y_i x = x y_{i+1} \quad (i+1 \text{ mod } k)$$

show that the set of monomials is closed under multiplication, whence D_1

is a Z_1 -algebra. As x and each y_i is algebraic of degree

$\leq k$ over Z_1 , D_1 is spanned by the monomials $x^i \cdot y_1^{i_1} \dots y_k^{i_k}$,

$0 \leq i, i_1, \dots, i_k < k$, thus D is a finite-dimensional Z_1 -algebra,

and is therefore also a division ring. \square

THEOREM 3.3.: If $a \in N$ is algebraic over Z and the field $Z(a)$ has a nontrivial Z -automorphism then N

contains a free group.

Proof: By the Noether-Skolem theorem, there exists an element $b \in D^*$ such that the given automorphism is induced from conjugation by b . Nontriviality of the automorphism means that $(b, a) \neq 1$, and a algebraic implies that the automorphism has finite order. By Lemma 3.2, $Z(a, b)$ is contained in a finite-dimensional division ring D_1 , and since $a \in N \cap D_1$, Lemma 3.1 yields the result. \square

COROLLARY 3.4[2] : *If N contains a noncentral torsion element, then it also contains a free group.*

Proof: If $a \in N - Z$ and $a^n = 1$, then $n > 2$ and for r coprime with n , $a \rightarrow a^r$ induces a nontrivial automorphism of $Z(a)$. \square

A further improvement of 3.4, suggested by several results in the literature, and which is the gist of most of section 5 is stated as a conjecture, weaker than Conjecture 2.

CONJECTURE 3: *If $N/N \cap Z^*$ contains a torsion element, then N contains a free group.*

PROPOSITION 3.5: *If Z is an absolute field, or if it contains all roots of 1, then Conjecture 3 holds for N .*

Proof: In the first case, $a^n \in Z^*$ implies that a is torsion. In the second, if $a^n = \alpha \in Z$, $Z(a)$ is a splitting field of $X^n - \alpha$, whence the result follows from 3.3. \square

PROPOSITION 3.6: *If $\text{char} D = p > 0$ and N contains a non-central element x such that x^p is central, then N contains a free group.*

Proof: Choose y such that $[y, x] \neq 0$. From a well known calculation, $[y, x]^{(p)} = [y, x^p] = 0$. Lemma 2.1 yields a w such that $x^{-1} w x = w + 1$. Applying Lemma 3.2 to x and $R = Z(w)$ and then Lemma 3.1 yields the result. \square

The exponent 2 yields the desired result irrespective of characteristic. Indeed, a slightly better fact is true (and compare [6, Thm. 4]):

PROPOSITION 3.7: *If N contains a noncentral element satisfying a quadratic polynomial over Z , then N contains a free group.*

Proof: Call that element x . If x is separable over Z , clearly $Z(x)$ has a nontrivial automorphism, and the result follows from Theorem 3.3. Otherwise we are under the hypothesis of 3.6 with $p=2$. \square

As observed in the last paragraph of [6]:

COROLLARY 3.8: *D^*/Z^* contains no nontrivial subnormal p -subgroup, when $p = \text{char } D$ or $p=2$.*

Proof: If there was such a group it would be of form N/Z^* with $Z^* \subseteq N \triangleleft D^*$. But 3.6 or 3.7 would imply that N , hence N/Z^* would contain a free group.

We use an argument of Lichtman [9] to prove:

PROPOSITION 3.9: *If $Z^* \subset N$ and N/Z^* contains a nonabelian finite group, the N contains a free group.*

Proof: Let G/Z^* be that group, with $G \leq D^*$. Then the linear span $Z[G]$ is a finite dimensional Z -algebra, spanned by a set of coset representatives of G/Z^* . Thus it is a subdivision ring of D , which allows the application of Lemma 3.1. \square

4) SN-WORDS

Recall that a word w in the free group on n generators is said to be *noncentral* on a group G if the word map $G^n \rightarrow G$ defined by substitution into w is not into the center of G . And w is said to be an N -word (SN-word) if for every division ring D (any noncentral $N \triangleleft D^*$), w is noncentral on D^* (N). The meaning of being an N -word or SN-word for some class of rings shall be understood in the obvious way.

THEOREM 4.1.: *If $N \triangleleft D^*$ and a word w is noncentral on N , then for some $g_1, \dots, g_n \in N$, $w(g_1, \dots, g_n)$ has infinite order.*

Proof.: Choose a noncentral element in the image of w .

There is nothing to prove unless it is torsion. In this case, Corollary 3.4 shows that N contains a free group on infinitely many generators. Substituting distinct generators into w yields the desired result. \square

This Theorem leads to a variety of specializations, since we are able to produce several SN-words, or N -words.

THEOREM 4.2: *Any word that is not in the commutator subgroup of the free group is SN. Further, let u, v be SN-word, and let w be any word. Then the*

following are SN-words:

a) uw , provided u and w have no letters in common

b) $w^{-1}uw$, u^{-1}

c) (u,v) , provided u and v have no letters in common

d) u^2

e) u^p , for division rings of characteristic $p > 0$.

Proof: Let w be a word not in the commutator of the free group. Suppose that some noncentral $N \triangleleft D^*$, is mapped by w into its center. There exists a letter x such that substituting in w all other letters by 1, yields the word x^n , for some nonzero integer n . It follows that $a^n \in Z^*$ every $a \in N$, and that contradicts Theorem 5.2, which we prove later.

Parts (a) and (b) are trivial. To prove (c), let $N \triangleleft D^*$ as usual. Denote by U the image of the map u on N and by V the image of v on N . The subgroups $\langle U \rangle$, $\langle V \rangle$ these sets generate are invariant in N , hence subnormal in D^* , and noncentral as u and v are SN. By Lemma 3 of [12], the group $(\langle U \rangle, \langle V \rangle)$ is noncentral. As U and V are normal sets in N , $(\langle U \rangle, \langle V \rangle)$ is generated by the commutators (a,b) with $a \in U$, $b \in V$; hence, for some $a \in U$, $b \in V$, (a,b) is noncentral, and (a,b) is in the image of (u,v) .

To prove (d), let a be a noncentral image of u on N ;

if a^2 is noncentral we are done, else we apply u to generators of an infinitely generated free subgroup of N , which exists by Theorem 3.7. The proof of (e) is similar. \square

The following is due to Amitsur [1, Thm. 19]:

THEOREM 4.3: *If Z is infinite then D^* satisfies no non-trivial relation $w=1$; that is, the word map w on D^* is not constant.*

COROLLARY 4.4: *Any word is noncentral for division rings with infinite center.*

Proof: If w is central for D^* , and x is a new letter, then (w,x) maps D^* to 1. \square

We close the section with a couple of more contrived examples of SN-words. The first one is a small-perturbation in an argument of Herstein [4].

PROPOSITION 4.4: *The word $(xy)^n x^{-n} y^{-n}$, $n \geq 1$ is SN.*

Proof: Suppose false, and let us consider a group $N \triangleleft D^*$ on which this word maps into the center. Without loss, we may assume that $Z^* \subseteq N$. Thus, we have that in N/Z^* , $(xy)^n = y^n x^n$ for every x, y . Therefore, for every x, y :
 $y^{n+1} x^{n+1} = (yx)^{n+1} = yx(yx)^n = yx x^n y^n = yx^{n+1} y^n$, hence
 $y^n x^{n+1} y^{-n} x^{-(n+1)} = 1$ in N/Z^* . This is impossible, since by Theorem 4.2, $y^n x^{n+1} y^{-n} x^{-(n+1)}$ is SN. \square

PROPOSITION 4.5: Let w be an SN-word, r, n positive integers and let x be a letter not occurring in w . Then $(w, x^r)^n$ is SN, provided $\text{char} D = p > 0$ and $p \nmid r$.

Proof: It is enough to consider the case where $p \nmid n$ since the general case will follow from (4.2.d). For a contradiction, suppose that $(w, x^r)^n$ maps N to its center, N being noncentral.

Let w_1 be a copy of w with entirely new letters, also distinct from x . By 4.2, $(w_1, (w, x))$ is noncentral, hence there exist $a, u \in N$, a in the image of w (hence of w_1), u in the image of (w, x^r) such that $(a, u) \neq 1$. By our temporary hypothesis $u^n \in Z$. If Z is finite, Proposition 3.5 implies that N contains a free group, whence $(w, x)^n$ cannot map N into its center. Hence Z is infinite.

We apply the Descending Lemma to a, u , and obtain suitable polynomials, p_k, \bar{p}_k, q_k and elements $d_k(\lambda) \in N$, of which the subscript k will be dropped. Further, since a is in the image of w , $(a, d^r(\lambda))^n \in Z$ for each possible λ . Therefore, $s = a p^r a^{-1} \bar{p}^r$ is a polynomial mapping infinitely many central elements to the center, and by Lemma 2.2, its coefficients are central. In particular,

$$\begin{aligned} c_x(s) &= a c_x(p^r) a^{-1} + c_x(p^{-r}) = r a [u, a]^{(k)} a^{-k} a^{-1} + r c_x [u, a]^{(k)} = \\ &= r [u, a]^{(k+1)} a^{-(k+1)} \in Z, \text{ and as } r \neq 0, [u, a]^{(k+2)} = 0. \end{aligned}$$

By Lemma 2.1, there is a t such that $a^{-1}ta=t+1$. Hence conjugation by a induces a nontrivial automorphism of $Z(t)$, of finite order p . Combining Lemmas 3.2 and 3.1 with the fact that $a \in N-Z^*$ yields that N contains a free group, and the original word has to be noncentral. \square

A slight modification at this proof's end would make it work in characteristic 0 provided D was algebraic over Z . In general for characteristic 0 we only have the following result, whose proof, on the lines of Theorem 5.2, is left to the reader.

PROPOSITION 4.5: *If m, n, r are nonzero integers, then $(x^m, y^n)^r$ is SN for division rings of characteristic 0.*

5) COMMUTATIVITY THEOREMS

We begin with a restatement of Theorem 4.1.

PROPOSITION 5.1: *If w is an SN-word and for every substitution $w(x)$ in $N \triangleleft D^*$, there is a $n > 1$ such that $w(x)^n = 1$, then $N \leq Z^*$. \square*

If Conjecture 3 is true, the stronger result will hold that if a SN word maps a subnormal subgroup of D^*/Z^* to torsion elements, then that subgroup is trivial. Special cases of Conjecture 3 yield special cases of the extension: it holds when Z is absolute or when the period of each $w(x) \bmod Z^*$ has as prime divisors only 2 and char Z . Another special case has been announced [11], that the extension holds for D^* provided Z is uncountable.

A special case of the proposed extension is that periodic subnormal subgroup of D^*/Z^* must be trivial; this was proved by Herstein [7] in case Z is uncountable. Without cardinality restrictions, we show it for groups of bounded period.

THEOREM 5.2: *Suppose that $N \triangleleft D^*$ and there exists an integer $n > 0$ such that $x^n \in Z$ for every $x \in N$. Then $N \leq Z^*$.*

Proof: By 3.5, it is enough to prove this when Z is infinite (and non-absolute). Further, we choose n minimal such that the hypothesis is satisfied, and then,

if $\text{char} D = p > 0$, Proposition 3.6. implies that $p \nmid n$.

Suppose that N is noncentral, for a contradiction. Then, there exist $a, u \in N$ such that $(a, u) \neq 1$. Applying the Descending Lemma, we obtain polynomials $p = p_k(a, u, X) \in D[X]$, $q \in Z[X^n]$ such that for infinitely many $\lambda \in Z$, $p(\lambda)/q(\lambda) \in N$. Then, for infinitely many $\lambda \in Z$, $p^n(\lambda) \in Z$, whence by Lemma 2.2, $C_X p^n = n C_X(p) \in Z$, and as $n \neq 0$, $C_X(p) = -[u, a]^{(k)} a^{-k} \in Z$. It follows that $[u, a]^{(k+1)} = 0$, hence by Lemma 2.1, there is a w such that $a^{-1} w a = w + 1$. As $a^n \in Z$, $w = a^{-n} w a^n = w + n$, whence $n = 0$ in D , a contradiction. \square

The same use of the Descending Lemma, coupled with the pigeonhole principle will yield another proof of the aforementioned theorem of Herstein. Indeed, it is easy to extend [7, Thm. 1]:

PROPOSITION 5.3: *If Z is uncountable and the word (x, y) maps N/Z^* to torsion elements, where $Z^* \subseteq N \ll D^*$, then $N = Z^*$.*

Proof: Suppose that $N \not\subseteq Z^*$. Again, by 3.6, the order of a noncentral element of $N \text{ mod } Z^*$ cannot be divisible by $\text{char} D$. Choose a noncentral commutator a , and an N -conjugate u such that $(a, u) \neq 1$. By hypothesis, there is an $m > 1$, such that $a^m = u^m \in Z$. Using the Descending Lemma we find torsion $\text{mod } Z^*$ elements $d_{k+1}(\lambda) = (d_k(\lambda), a)$, with $d_k(\lambda) \in N$, for uncountably many λ . Thus, some $n \geq 1$

is the order of $d_{k+1}(\lambda) \bmod Z^*$ for infinitely many λ . As in the preceding proof, we get a contradiction. \square

A further approximation to either of the conjectures would be a possible generalization of Amitsur's Theorem 4.3. for noncentral subnormal subgroups of D^* . We did not advance much in this direction. The next results could be contrived to statements that certain words are noncentral for division rings satisfying very restrictive conditions. To avoid misunderstandings, we consider the trivial group to be torsion free.

PROPOSITION 5.4: *Suppose that $\text{char}D=0$. Let n, m be relatively prime integers, and let r, s be integers such that $0 \neq r \cdot n \neq s \cdot m$. Then, if the relation $(x^n, y^n)^r = (x^m, y^m)^s$ holds in N/Z^* , N/Z^* is torsion free.*

Proof: We shall assume that the relation holds, and that there is an $a \in N-Z$ such that $a^p \in Z$ for some $p > 1$.

By Proposition 4.6, $ms \neq 0$. Now, the given relation is equivalent to both $(x^n, y^n)^{-r} = (x^m, y^m)^{-s}$ and $(x^{-n}, y^{-n})^r = (x^{-m}, y^{-m})^s$, hence there is no loss in supposing that $n, r > 0$. To avoid more notation, we split the proof in four separate cases, given by the four combinations of signs of m and s . We shall do completely the case m, s both positive.

As in earlier cases, apply the Descending Lemma, choosing an appropriate u , thus obtaining polynomials

$p_k(a, u; X)$, and so on. The given relation can be rewritten as: $(x^n, y^n)^r (y^m, x^m)^s \in Z$ for every $x, y \in N$; thus, upon the substitution $x=a, y=d_k(\lambda)$ and the now familiar reasoning, we deduce that:

$$r(-na^n[u, a]^{(k)} a^{-k} a^{-n} + n[u, a]^{(k)} a^{-k}) + r(-m[u, a]^{(k)} a^{-k} + ma^m[u, a]^{(k)} a^{-k} a^{-m}) \in$$

and a new derivation yields, with $w=[u, a]^{(k+1)}$:

$$sm a^m w a^{-m} - r n a^n w a^{-n} - (sm - rn) = 0.$$

If $w=0$, we get a contradiction in the usual way. Otherwise, let V be the Z -submodule of D spanned by $\{w, awa^{-1}, \dots, a^{p-1} w a^{-(p-1)}\}$, and let T be the automorphism of V given by $T(x) = axa^{-1}$. Clearly $T^p=1$, while the last equation implies that $rmT^m - snT^n - (rm - sn) = 0$. Now, when m, n are coprime, the greatest common divisor of X^{p-1} and $smX^m - rnX^n - (sm - rn)$ is, as we see below, $X-1$, whence $T=1$. That is, $awa^{-1}=w$, and we conclude that $[u, a]^{(k+2)} = [w, a]=0$, and this gives a contradiction via Lemma 2.1.

As for the polynomials, let θ be a common complex root of these. Thus we have that $\theta^p=1$ and $\theta^n(\theta^{m-n} - rn/sm) = 1 - rn/sm$. Then $|\theta^{m-n} - rn/sm| = |1 - rn/sm|$, and this implies $\theta^{m-n}=1$ (just draw a picture). Substituting the value of θ^{m-n} yields $\theta^n=1$, and as m, n are coprime, $\theta=1$.

In the other cases one proceeds similarly, and obtains the same pair of equations for T . When $m < 0$, the large equation must be multiplied by T^{-m} in order to give

a polynomial, but the reasoning is still the same. \square

The same method yields:

PROPOSITION 5.5: Let sequences $n_1, \dots, n_\ell, m_1, \dots, m_\ell,$
 r_1, \dots, r_ℓ of nonzero integers be
 given and suppose that the only complex root of the
 rational function $\sum_i r_i m_i X^{n_i} - \sum_i r_i m_i$ which is also
 a root of 1 is $X=1$. Then if $\text{char} D=0$, and the word
 $\prod_i (x^{n_i}, y^{m_i})^{r_i}$ maps N to its center, then N/Z^* is torsion
 free.

REFERENCES:

- [1] S. A. Amitsur, *Rational identities and applications to algebra and geometry*, J. Algebra 3(1966), 304-359.
- [2] J.Z. Gonçalves, *Free groups in subnormal subgroups and the residual nilpotence of the group of units of group rings*, Can. Math. Bull. (to appear).
- [3] I.N. Herstein and W.R. Scott, *Subnormal subgroups of division rings*, Can. J. Math. 15(1963), 80-83.
- [4] I.N. Herstein, *Power maps in rings*, Mich. Math. J. 8(1961), 29-32.
- [5] I.N. Herstein, *Noncommutative rings*, The Carus Math. Monographs n. 15, The Math. Assoc. of America, 1968.
- [6] I.N. Herstein, *Multiplicative commutators in division rings*, Israel J. Math. 31(1978), 180-188.
- [7] I.N. Herstein, *Multiplicative commutators in division rings II*, Rend. Circ. Mat. di Palermo, Serie II, Tomo XXIX (1980), 485-489.
- [8] A.I. Lichtman, *On subgroups of the multiplicative group of skew fields*, Proc. Amer. Math. Soc. 63(1977), 15-16.
- [9] A.I. Lichtman, *Free subgroups of normal subgroups of the multiplicative group of skew fields*, Proc. Amer.

Math. Soc. 71(1978), 174-178.

- [10] A.I. Lichtman, *On normal subgroups of the multiplicative group of skew fields generated by a polycyclic-by-finite group*, J. Algebra, 78(1982), 548-577.
- [11] L. Makar-Limanov and P. Malcolmson, *Words periodic over the center of a division ring*. Abstract 810-16-11 Abs. Amer. Math. Soc. 2(1984), 160.
- [12] C. Stuth, *A generalization of the Brauer - Cartan-Hua theorem*, Proc. Amer. Math. Soc. 15(1964), 211-217.
- [13] J. Tits, *Free subgroups in linear groups*, J. Algebra 20(1972), 250-270.