

DEPARTAMENTO DE CIÊNCIA DA COMPUTAÇÃO

Relatório Técnico

RT-MAC-2001-10

THE SYMMETRIC EIGENVALUE
COMPLEMENTARITY PROBLEM

MARCELO QUEIROZ
JOAQUIM JÚDICE
CARLOS HUMES JÚNIOR

Novembro de 2001

The Symmetric Eigenvalue Complementarity Problem

Marcelo Queiroz ^{*} Joaquim Júdice [†] Carlos Humes Jr.

November 16, 2001

Abstract

In this paper the Eigenvalue Complementarity Problem (EiCP) with real symmetric matrices is addressed. It is shown that the symmetric (EiCP) is equivalent to finding an equilibrium solution of a differentiable optimization problem in a compact set. A necessary and sufficient condition for feasibility is obtained which, when verified, gives a convenient starting point for any gradient-ascent local optimization method to converge to a solution of the (EiCP). It is further shown that similar results apply to the Symmetric Generalized Eigenvalue Complementarity Problem (GEiCP). Computational tests show that these reformulations improve the speed and robustness of the solution methods.

1 Introduction

The computation of eigenvalues is of crucial importance in a variety of practical problems in physics and engineering. Eigenvalues are related to the resonance frequency of structures,

^{*}Research supported by FAPESP Grant No. 97/06227-2.

[†]Research supported by FCT project POCTI/35059/MAT/2000.

and to stability of dynamical systems [Chatelin]. Real symmetric matrices appear in a large portion of these practical problems [Parlett].

A classical approach to solve large-scale eigenvalue problems with real symmetric matrices is to apply unconstrained optimization techniques to the so-called Rayleigh quotient. This function has the property that every equilibrium point is an eigenvector with a corresponding eigenvalue given by the Rayleigh quotient.

The Eigenvalue Complementarity Problem (EiCP) appeared in the study of static equilibrium states of finite dimensional mechanical systems with unilateral frictional contact [Costa & alii] and takes the following form

$$(\text{EiCP}): \text{Find } \lambda > 0, x \neq 0 \text{ such that } \begin{cases} w = (\lambda B - A)x \\ w \geq 0 \\ x \geq 0 \\ w^t x = 0. \end{cases}$$

The reader is referred to [Costa & alii] for further details.

The purpose of this paper is to study the (EiCP) when A and B are symmetric matrices. This particular problem is shown to be equivalent to finding an equilibrium point of a generalized Rayleigh quotient satisfying $x \geq 0$ and $\lambda > 0$, which can be solved using any gradient ascent method, from a convenient starting point. This result is akin to a classical result for Nonlinear Complementarity Problems (NCP) with symmetric jacobians, although the classical result is not applicable to a NCP reformulation of the (EiCP) (section 4). The result is extended to a generalization of the symmetric (EiCP) of practical interest, the symmetric (GEiCP). Computational tests show that these reformulations improve the speed and robustness of the solution methods. Some preliminary research on the asymmetric case is also presented.

The structure of the paper is as follows. In section 2 the classical eigenvalue problem is discussed, with emphasis to symmetric matrices. The relationship of this problem with the optimization of the Rayleigh quotient is established, showing that the eigenpairs can be computed with gradient related methods.

In section 3 the Eigenvalue Complementarity Problem (EiCP) and the Generalized Eigenvalue Problem (GEiCP) are introduced. Some general results concerning the feasibility of these problems and the number of solutions are presented. It is shown that in general these problems are NP-hard. In this paper the term feasibility is a synonym for the existence of a solution, which means the existence of a complementary solution in the case of (EiCP) or (GEiCP).

The symmetric case is studied in section 4. It is shown that in this case both the (EiCP) and the (GEiCP) are equivalent to finding equilibrium points of optimization problems, and therefore are relatively easy to solve. Computational experience is reported in section 5.

Some preliminary work on the asymmetric case is presented in section 6, and the last section (section 7) presents some conclusions and future work.

2 Eigenvalue Problems and Optimization

Given a matrix $A \in \mathbb{R}^{n \times n}$, the classical Eigenvalue Problem (EiP) is:

$$(\text{EiP}): \text{Find } \lambda \in \mathbb{R}, x \in \mathbb{R}^n \setminus \{0\} \text{ such that } Ax = \lambda x.$$

A necessary and sufficient condition for the existence of a solution with $x \neq 0$ is that the kernel of $(A - \lambda I)$ be non-trivial, and this occurs if and only if λ is a root of the polynomial equation $\det(A - \lambda I) = 0$. Such values of λ are referred to as the solutions of the (EiP), and correspond to the eigenvalues of A . Since the polynomial above has degree n , the (EiP) has

at most n distinct solutions.

If $\bar{\lambda}$ is an eigenvalue of A , computing a corresponding eigenvector amounts to finding a non-trivial solution of the linear system $(A - \bar{\lambda}I)x = 0$. Since this system is homogeneous, x is an eigenvector corresponding to $\bar{\lambda}$ if and only if αx is also an eigenvector corresponding to $\bar{\lambda}$, for any $\alpha \neq 0$. An extensive treatment of properties and algorithms for the (EiP) can be found in [Parlett] and [Chatelin].

A classical approach to solve large-scale (EiP) problems with real symmetric matrices is to apply unconstrained optimization techniques to the Rayleigh quotient

$$\lambda(x) = \frac{x^t A x}{x^t x}.$$

This quotient is only defined for $x \neq 0$. The gradient of the Rayleigh quotient is

$$\nabla \lambda(x) = \frac{2}{x^t x} [A - \lambda(x)I]x.$$

Since $\nabla \lambda(x) = 0$ if and only if $[A - \lambda(x)I]x = 0$, any equilibrium point $(x, \lambda(x))$ of the Rayleigh quotient is a solution of the (EiP). The expression of the gradient of the Rayleigh quotient above is only valid when A is symmetric. If $A \neq A^t$, the correct expression is $\nabla \lambda(x) = \frac{1}{x^t x} [A + A^t - 2\lambda(x)I]x$, and the relationship between equilibrium points and solutions of the (EiP) ceases to hold.

The Generalized Eigenvalue Problem (GEiP) is often referred to in the literature: given $A, B \in \mathbb{R}^{n \times n}$,

(GEiP): Find $\lambda \in \mathbb{R}, x \in \mathbb{R}^n \setminus \{0\}$ such that $Ax = \lambda Bx$.

The λ -solutions of this problem are referred to as general eigenvalues of (A, B) . This problem is in general more difficult than the (EiP) [Parlett]. When B is symmetric positive definite

and A is symmetric, a generalized Rayleigh quotient can be defined as

$$\lambda(x) = \frac{x^t A x}{x^t B x}$$

and the (GEiP) can also be solved by optimization. The gradient in this case is

$$\nabla \lambda(x) = \frac{2}{x^t B x} [A - \lambda(x) B] x,$$

and the equilibrium points of $\lambda(x)$ correspond to general eigenvectors and eigenvalues of (A, B) .

The lemma below presents two fundamental properties of the generalized Rayleigh quotient.

Lemma 1 *For all $x \neq 0$ the following equalities hold:*

1. $\lambda(\alpha x) = \lambda(x), \forall \alpha > 0;$
2. $x^t \nabla \lambda(x) = 0.$

Proof :

1. $\lambda(\alpha x) = \frac{(\alpha x)^t A (\alpha x)}{(\alpha x)^t B (\alpha x)} = \frac{\alpha^2 x^t A x}{\alpha^2 x^t B x} = \lambda(x).$
2. $x^t \nabla \lambda(x) = x^t \left[\frac{2}{x^t B x} [A - \lambda(x) B] x \right] = \frac{2}{x^t B x} [x^t A x - \lambda(x) x^t B x] = \frac{2}{x^t B x} [x^t A x - x^t A x] = 0.$

■

Since the case $B = I$ and B symmetric positive definite are subject to the same approach with respect to the optimization of the Rayleigh quotient, all the complementary problems introduced in the next section are formulated with two matrices (A, B) .

Other functions whose equilibrium points correspond to pairs of eigenvalues and eigenvectors exist for special classes of matrices. For instance, the functions $S_A(x) = x^t x - 2\sqrt{x^t A x}$, $P_A(x) = (x^t x)^2 - 2x^t A x$ and $L_A(x) = x^t x - \ln(x^t A x)$, for a symmetric positive definite matrix A , have their equilibrium points corresponding to eigenvectors of A , whereas $S_A(\cdot)$ and $P_A(\cdot)$ also have the origin as an equilibrium point. The papers [Mongeau & Torki, Auchmuty 89, Auchmuty 91] present theoretical results and computational experience comparing these functions to the Rayleigh quotient for symmetric positive definite matrices. The present work does not address these functions in order to keep the framework more general.

3 The Eigenvalue Complementarity Problem

As stated in the Introduction, the Eigenvalue Complementarity problem (EiCP) appeared in the study of static equilibrium states of mechanical systems with unilateral friction, in [Costa & alii]. It may be expressed as

$$(\text{EiCP}): \text{Find } \lambda > 0, x \neq 0 \text{ such that } \begin{cases} w = (\lambda B - A)x \\ w \geq 0 \\ x \geq 0 \\ w^t x = 0. \end{cases}$$

where B is a positive definite matrix.

Note that any solution with $w = 0$ correspond to a solution of the (GEiP). By allowing $w_J \geq 0$ when $x_J = 0$, the (EiCP) has a larger solution set than the (GEiP), and is therefore considered a relaxation of that problem.

The Generalized Eigenvalue Complementarity Problem (GEiCP)_J is

$$(\text{GEiCP})_J: \text{Find } \lambda > 0, x \neq 0 \text{ such that } \begin{cases} w = (\lambda B - A)x \\ w_J = 0 \\ w_J \geq 0 \\ x_J \geq 0 \\ w_J^t x_J = 0 \end{cases}$$

where B is a positive definite matrix, $J \subseteq \{1, 2, \dots, n\}$ is given, and $\bar{J} = \{1, 2, \dots, n\} \setminus J$. The (EiCP) is clearly the particular case of the $(\text{GEiCP})_J$ with $J = \{1, 2, \dots, n\}$.

For any solution (λ, x, w) , the value λ is called a *(general) complementary eigenvalue* of (A, B) , and x is a corresponding *(general) complementary eigenvector*. Since the set of complementary eigenvectors of a given complementary eigenvalue is a cone, there is no loss of generality in restricting the problem to finding solutions satisfying $\|x\|_2 = 1$, which replaces the constraint $x \neq 0$. In the case of the (EiCP) the linear constraint $\|x\|_1 = e^t x = 1$ can be considered instead of $\|x\|_2 = 1$, since $x \geq 0$.

It is easy to see that any solution of the (EiCP) (or (GEiCP)) with $w = 0$ is a positive eigenvalue of (A, B) with a corresponding eigenvector satisfying some sign constraints. In general, the following property holds.

Proposition 2 *For any solution (λ, x) of $(\text{GEiCP})_J$, there is a set I satisfying $\bar{J} \subseteq I \subseteq \{1, 2, \dots, n\}$, such that λ is a positive eigenvalue of (A_{II}, B_{II}) and x_I is a corresponding eigenvector satisfying $x_{J \cap I} \geq 0$.*

Proof: Set $I = \{i \mid w_i = 0\}$. Then $I \supseteq \bar{J}$, and $x_I = 0$, by complementarity.

Therefore $A_{II}x_I = \lambda B_{II}x_I$. ■

For the (EiCP), this result says that given a solution (λ, x) , λ is a positive eigenvalue of (A_{II}, B_{II}) and x_I is a corresponding nonnegative eigenvector. A direct corollary says that

the solution sets of the (EiCP) and (GEiCP) are finite.

Proposition 3 *The (EiCP) has at most $2^{n+1} - n - 2$ solutions. The (GEiCP) $_J$ has at most $(n - |J| + 2)2^{|J|} - |J| - 2$ solutions.*

Proof: Consider first the (GEiCP) $_J$. There are $2^{|J|}$ possible subsets I such that $\bar{J} \subset I \subset \{1, 2, \dots, n\}$. For each possible subset I there are at most $|I|$ distinct solutions, namely the eigenvalues of $B_{II}^{-1}A_{II}$. Therefore the total number of solutions is at most

$$\begin{aligned} \sum_{i=0}^{|J|} (|\bar{J}| + i) \binom{|J|}{i} &= (n - |J|) \sum_{i=0}^{|J|} \binom{|J|}{i} + \sum_{i=1}^{|J|} \sum_{k=i}^{|J|} \binom{|J|}{k} \\ &= (n - |J|)2^{|J|} + \sum_{i=1}^{|J|} (2^{|J|-i+1} - 1) \\ &= (n - |J|)2^{|J|} + 2^{|J|+1} - |J| - 2 \\ &= (n - |J| + 2)2^{|J|} - |J| - 2. \end{aligned}$$

Since the (EiCP) is the particular instance of the (GEiCP) $_J$ with $J = \{1, 2, \dots, n\}$, the number of solutions is at most

$$(n - |J| + 2)2^{|J|} - |J| - 2 = 2^{n+1} - n - 2.$$

■

The complete set of solutions of both (EiCP) and (GEiCP) can be obtained via complete enumeration, as follows.

Complete Set of Solutions of (GEiCP) $_J$

1. For each subset I such that $\bar{J} \subseteq I \subseteq \{1, 2, \dots, n\}$ compute the set $\Lambda(I)$ of eigenvalues of $B_{II}^{-1}A_{II}$.

2. For each $\bar{\lambda} \in \Lambda(I)$ such that $\bar{\lambda} > 0$, try to compute x_I such that

$$\begin{cases} A_{II}x_I = \bar{\lambda}B_{II}x_I \\ A_{JI}x_I \geq \bar{\lambda}B_{JI}x_I \\ x_{I \cap J} \geq 0 \\ \|x_I\|_2 = 1. \end{cases}$$

If such a solution exists, then $x = (x_I, 0)$ is a general complementary eigenvector corresponding to the general complementary eigenvalue $\bar{\lambda}$.

The soundness of this method is a direct consequence of proposition 2. This algorithm also provides a necessary and sufficient test for feasibility of a given (GEiCP). The following proposition shows that any necessary and sufficient test for the feasibility of a general (GEiCP) is doomed to be computationally expensive.

Proposition 4 *The feasibility of the (GEiCP)_J is a NP-complete decision problem.*

Proof : Any solution (λ, x) of the (GEiCP)_J satisfies $x_J \geq 0$ and

$$x^t Ax = x^t (\lambda Bx - w) = \lambda x^t Bx - x^t w = \lambda x^t Bx > 0,$$

since $x \neq 0$, $\lambda > 0$ and B is positive definite. Therefore, deciding if the (GEiCP)_J is feasible is at least as difficult as deciding if there exists a x such that $x_J \geq 0$ and $x^t Ax > 0$, for a given matrix A . The latter problem is NP-complete [Murty, theorem 2.20]. Since the feasibility of the (GEiCP)_J is in NP (i.e. one can polynomially check whether a given pair (λ, x) is a solution of the (GEiCP)_J), the result of the proposition follows. ■

It follows that solving the (GEiCP) is in general a NP-hard problem. Despite this fact, for some classes of matrices the feasibility of the corresponding (EiCP) or (GEiCP) can be answered easily.

Lemma 5 *If A is negative semi-definite (i.e. $x^t A x \leq 0$ for all x), the corresponding (GE-
iCP) is infeasible.*

Proof : Since A is negative semi-definite, it follows that A_{II} is negative semi-definite for all $I \subseteq \{1, 2, \dots, n\}$. Therefore $x_I^t A_{II} x_I \leq 0$, for any x , and the problem is infeasible. ■

Lemma 6 *If A is positive ($A_{ij} > 0, \forall i, j$), and $B = I$, then the (EiCP) has a solution $\bar{\lambda} > 0$, with a corresponding eigenvector $\bar{x} > 0$. Moreover, if A is symmetric, then the solution $\bar{\lambda}$ is unique.*

Proof : Since A is positive, any feasible solution $\bar{x} \neq 0$ has to satisfy $\bar{x} > 0$, for otherwise $\bar{x}_i = 0$ and $\bar{x} \neq 0$ implies $\bar{w}_i = \bar{\lambda} \bar{x}_i - A_i \bar{x} = -A_i \bar{x} < 0$. By complementarity, $\bar{w} = 0$, and therefore this (EiCP) is equivalent to the (EiP). The existence of a solution of the (EiP) with a positive matrix is Perron's Theorem [Ortega, theorem 6.1.2]. The uniqueness in the symmetric case is verified as follows. Let $\hat{\lambda} \neq \bar{\lambda}$, and consider a vector \hat{x} such that $A\hat{x} = \hat{\lambda}\hat{x}$. Then,

$$(\hat{\lambda} - \bar{\lambda})\hat{x}^t \bar{x} = (A\hat{x})^t \bar{x} - \hat{x}^t (A\bar{x}) = \hat{x}^t A^t \bar{x} - \hat{x}^t (A\bar{x}) = 0,$$

and since $\hat{\lambda} \neq \bar{\lambda}$, it follows that $\hat{x}^t \bar{x} = 0$. Since $\bar{x} > 0$, there is an index i such that $\hat{x}_i < 0$, and therefore \hat{x} is not a solution of the (EiP). ■

Lemma 7 *If there is some index j such that $A_{jj} > 0$ and $A_{ij} \leq 0, \forall i \neq j$, then (λ, x) is a solution of the (EiCP), with $\lambda = A_{jj}$ and $x = e^j$.*

Proof : Trivial. ■

In particular this property holds for the well-known class of nonsingular M-matrices, which are defined as P-matrices (all principal minors are positive) with nonpositive off-diagonal elements.

The following section is devoted to the special case of the Eigenvalue Complementarity Problems with symmetric matrices. Some additional properties of the general (GEiCP) are presented in section 6.

4 The Symmetric Eigenvalue Complementarity Problem

In this section the symmetric versions of (EiCP) and (GEiCP) are studied. Some properties of these problems are derived, including a necessary and sufficient condition for feasibility. When this condition is verified, the problem can be solved by a variety of well-known methods in differentiable nonlinear programming.

For clarity of exposition, the results are presented for the (EiCP) and the (GEiCP) separately.

The Symmetric (EiCP)

The symmetric (EiCP) is closely related to the classical eigenvalue problem. Since $x \neq 0$ for any solution, the complementarity condition $w^t x = 0$ may be rewritten as $x^t(\lambda Bx - Ax) = 0$, or equivalently,

$$\lambda(x) = \frac{x^t Ax}{x^t Bx}.$$

This is the generalized Rayleigh quotient. Analogously to the classical case, equilibrium points of the Rayleigh quotient in the nonnegative orthant with $\lambda(x) > 0$ are solutions of

the (EiCP).

Proposition 8 *The symmetric (EiCP) is equivalent to*

$$(OEiCP) \begin{cases} \max & \lambda(x) \\ \text{s.t.} & x \geq 0 \\ & e^t x = 1, \end{cases}$$

in the sense that every equilibrium solution x of (OEiCP) with $\lambda(x) > 0$ is a solution of the (EiCP).

Proof : The optimization problem (OEiCP) is equivalent to

$$\max_{x \in \mathbb{R}^n} \inf_{\substack{u \geq 0 \\ \alpha \in \mathbb{R}}} \frac{x^t A x}{x^t B x} + u^t x + \alpha(e^t x - 1).$$

Any equilibrium solution of this problem satisfies the Kuhn-Tucker conditions

$$\begin{cases} u + \alpha e = \frac{2}{x^t B x} [\lambda(x) B - A] x \\ u \geq 0 \\ x \geq 0 \\ u^t x = 0 \\ e^t x = 1. \end{cases}$$

By performing the scalar product of the first equation with x and noticing that $u^t x = 0$, $e^t x = 1$, and $\lambda(x) = \frac{x^t A x}{x^t B x}$, it follows that $\alpha = 0$. The resulting system of equations corresponds closely to the (EiCP), with $u = \frac{2}{x^t B x} w$, and without the constraint $\lambda > 0$. It follows that any equilibrium solution of the (OEiCP) with $\lambda(x) > 0$ is a solution of the (EiCP). ■

The conclusion that $\alpha = 0$ in the proof of proposition 8, though apparently nonintuitive, may be motivated by the homogeneity of $\lambda(\cdot)$ and the orthogonality between x and $\nabla\lambda(x)$ (lemma 1). Consider, for instance, an equilibrium point of the (OEiCP) satisfying $x > 0$. By the homogeneity of $\lambda(\cdot)$, the projection of $\nabla\lambda(x)$ on the half-line $\{\alpha x \mid \alpha \geq 0\}$ vanishes. By the equilibrium property with respect to the constraint $e^t x = 1$, the projection of $\nabla\lambda(x)$ on the affine manifold $\{x \mid e^t x = 1\}$ also vanishes. Therefore, $\nabla\lambda(x) = 0$, and so the Lagrange multiplier α in the Kuhn-Tucker conditions becomes superfluous.

Proposition 8 resembles a classical result for nonlinear complementarity problems of the form

$$\begin{cases} x \geq 0 \\ F(x) \geq 0 \\ x^t F(x) = 0, \end{cases}$$

with $\nabla F(x)$ symmetric for all x . In fact, such problems can be solved as optimization problems because $F(x)$ is shown to correspond to the gradient of a primitive function $f(x)$, whose equilibrium points are solutions of the NCP. In the present case the constraint $e^t x = 1$ (or $x \neq 0$) breaks the symmetry of the enlarged NCP, obtained by considering an additional complementary variable α satisfying

$$\begin{cases} w = (B - \mu A)x \\ \alpha = -1 + e^t x \\ x, w \geq 0 \\ \mu, \alpha \geq 0 \\ w^t x = \alpha \mu = 0, \end{cases}$$

where $\mu = \frac{1}{\lambda}$ with respect to the original problem. In this setting, the hypothesis of the symmetric jacobians do not hold. Therefore, it is not possible to use the classical theorem

to conclude that the NCP above is solvable as an optimization problem.

From the reformulation of the (EiCP) as a nonlinear programming problem several conclusions are drawn. The first one is a characterization of feasibility of the (EiCP).

Proposition 9 *(EiCP) is feasible if and only if there exists some $x \geq 0$ such that $x^t Ax > 0$.*

Proof: First suppose that the (EiCP) is feasible. If (λ, x) is any solution of the (EiCP), then $x \geq 0$ and $x^t Ax = \lambda x^t Bx > 0$, since $x \neq 0$ and B is positive definite.

Suppose, on the other hand, that there is a $\hat{x} \geq 0$ such that $\hat{x}^t A \hat{x} > 0$. Since $\lambda(x)$ is a continuous function (for $x \neq 0$) and the set $X = \{x \geq 0 \mid e^t x = 1\}$ is compact, there exists a $\bar{x} \in X$ such that $\lambda(\bar{x}) \geq \lambda(x)$, $\forall x \in X$. This is an equilibrium point for the (OEiCP), and in particular, $\lambda(\bar{x}) \geq \lambda(\hat{x}) > 0$. Therefore, \bar{x} is a solution of the (EiCP). ■

For the practical solution of the symmetric (EiCP), any ascent gradient method, with an initial solution $\hat{x} \in X$ such that $\hat{x}^t A \hat{x} > 0$, obtains an equilibrium point which is a solution of the problem. The nonconvexity of the objective function is actually not a problem, since the global optimization problem (OEiCP) need not be solved. However, finding such an initial point is a NP-complete problem (proposition 4). But for a very large class of matrices this problem is trivial, as the next proposition shows. For these matrices, an initial point can be found by inspection, and a solution of the (EiCP) is thereafter easily obtained.

Proposition 10 *Suppose that the matrix A satisfies one of the following conditions*

1. $\exists i : A_{ii} > 0$;
2. $\exists i, j : A_{ii} = 0, A_{jj} \leq 0$ and $A_{ij} > 0$;

3. $A \geq 0$, $A \neq 0$;

4. A is a S -matrix ($\exists x \geq 0 : Ax > 0$);

Then a point $\hat{x} \in X$ such that $\hat{x}^t A \hat{x} > 0$ can be obtained easily and the corresponding (EiCP) is feasible.

Proof :

1. Let $\hat{x} = e^i$; then $\hat{x}^t A \hat{x} = A_{ii} = 1$.

2. Let $\hat{x}_j = 1$, $\hat{x}_i = \frac{1-A_{ii}}{2A_{ij}}$, and $\hat{x}_k = 0$, $\forall k \neq i, j$; then $\hat{x}^t A \hat{x} = A_{jj}x_j^2 + 2A_{ij}x_i x_j = 1$.

3. If $A \geq 0$, $A \neq 0$, then $x = e$ is such that $x^t A x > 0$.

4. If A is a S -matrix, then there exists a $\hat{x} \geq 0$ such that $A \hat{x} > 0$. It follows that $\hat{x}^t A \hat{x} > 0$. Such a \hat{x} can be found by solving the following linear program:

$$(LP) \left\{ \begin{array}{l} \max \quad y \\ \text{s.t.} \quad Ax - ye \geq 0 \\ \quad \quad e^t x = 1 \\ \quad \quad x \geq 0 \\ \quad \quad y \in \mathbb{R}. \end{array} \right.$$

This problem is always feasible, and since the variable x lies in a compact set, and $y \leq \min\{(Ax)_i\}$, an optimal solution exists. Now since A is a S -matrix, the optimal solution has $y > 0$, and the corresponding $x \geq 0$ satisfies $x^t A x \geq y x^t e = y > 0$.

■

Note that condition 1 of the previous result (A has at least one positive diagonal entry) defines a very large class of matrices, that includes non-trivial positive semi-definite matrices (i.e. $x^t Ax \geq 0$ for all x) and strictly copositive matrices (i.e. $x^t Ax > 0$ for all $0 \neq x \geq 0$). Condition 2 includes non-trivial copositive matrices (i.e. $x^t Ax \geq 0$ for all $x \geq 0$).

For the design of an initialization algorithm, besides applying the tests indicated above, additional heuristics may be implemented. One of these is to find local maxima of the nonconvex quadratic problem

$$(QP) \begin{cases} \max & x^t Ax \\ \text{s.t.} & e^t x = 1 \\ & x \geq 0 \end{cases}$$

from a few random starting points. Other tests are given by lemmas 5, 6 and 7 in the previous section. The definitive answer can be given by solving a general LCP as below.

Proposition 11 *Given a matrix A , there exists a vector $x \geq 0$ such that $e^t x = 1$ and $x^t Ax \geq \varepsilon > 0$ if and only if the system*

$$\begin{cases} u + \alpha e + Ax = 0 \\ -\alpha \geq \varepsilon \\ u \geq 0 \\ x \geq 0 \\ u^t x = 0 \\ e^t x = 1 \end{cases}$$

is feasible.

Proof : This system corresponds to the Kuhn-Tucker conditions of the (QP) and the additional constraint $-\alpha \geq \varepsilon$. This additional constraint is equivalent

to $x^t Ax \geq \epsilon$, which can be seen by taking the inner product of the first equation with x , and noticing that $u^t x = 0$ and $e^t x = 1$. Since the feasible set of the (QP) is compact, the existence of a $x \geq 0$ such that $e^t x = 1$ and $x^t Ax \geq \epsilon > 0$ is equivalent to the existence of an equilibrium point of the (QP) satisfying the same conditions, which is equivalent to the general LCP above. ■

An interesting property of the symmetric (EiCP) regards the number of solutions. In fact it is possible to find a smaller upper bound than the one presented in the general case.

Proposition 12 *The number of solutions of the symmetric (EiCP) is at most $2^n - 1$.*

Proof : Consider a solution (λ, x) with $I = \{i \mid x_i > 0\}$. It is proved next that λ is the only eigenvalue of A_{II} with a corresponding positive eigenvector. Suppose, for the sake of contradiction, that there is some other solution (μ, y) such that $I_y = \{i \mid y_i > 0\} = I$ and $\mu \neq \lambda$. Since A is symmetric, $(\lambda - \mu)x_I^t y_I = y_I^t Ax - x_I^t Ay = 0$. But this contradicts $x_I > 0$ and $y_I > 0$. Therefore, for each possible set I at most one solution is possible, which means that the number of solutions is bounded above by $2^n - 1$. ■

The symmetric (GEiCP)

The main results from this section can be rephrased in the context of the symmetric (GEiCP). Once again the complementarity constraint $x^t w = 0$ allows λ to be expressed as a function of x ,

$$\lambda(x) = \frac{x^t Ax}{x^t Bx}.$$

Since x is not necessarily nonnegative (only x_J is nonnegative), the condition $x \neq 0$ cannot be replaced by $e^t x = 1$. The cumbersome constraint $\|x\|_2 = 1$ is adopted for the sake of theory only. It is later dropped from the computational model, which is able to handle this constraint implicitly. The symbol $\|\cdot\|$ without sub-index is used to denote the euclidean norm.

Proposition 13 *The symmetric (GEiCP) is equivalent to*

$$(OGEiCP) \begin{cases} \max & \lambda(x) \\ \text{s.t.} & x_J \geq 0 \\ & \|x\| = 1, \end{cases}$$

in the sense that every equilibrium solution x of (OGEiCP) with $\lambda(x) > 0$ is a solution to the (GEiCP).

Proof : The optimization problem (OGEiCP) is equivalent to

$$\begin{aligned} \max_{x \in \mathbb{R}^n} \quad & \inf_{\substack{u_J \geq 0 \\ u_J = 0 \\ \alpha \in \mathbb{R}}} \frac{x^t A x}{x^t B x} + u^t x + \alpha(x^t x - 1). \end{aligned}$$

Any equilibrium solution of this problem satisfies the Kuhn-Tucker conditions

$$\begin{cases} u + \alpha x = \frac{2}{x^t B x} [\lambda(x) B - A] x \\ u_J \geq 0 \\ x_J \geq 0 \\ u_J^t x_J = 0 \\ u_J = 0 \\ x^t x = 1. \end{cases}$$

As before, taking the inner product of the first equation with x , it can be seen that the Lagrangian multiplier α corresponding to the constraint $x^t x = 1$ has value 0. The system corresponds closely to the (GEiCP), with $u = \frac{2}{x^t B x} w$, and without the constraint $\lambda > 0$, and the result follows. ■

The difficulty of the optimization problem (OGEiCP) is revealed by the nonconcavity of the objective function (maximization problem) and by the nonconvex constraint $\|x\| = 1$. The nonconvexity of the objective function is not really a problem, since one looks after equilibrium solutions, and not globally optimal solutions, exactly as in the (EiCP) case. The feasible region of (OGEiCP) is a spherical surface with sign constraints, and therefore projection methods can be easily implemented via renormalization of iterates. See, for instance, the projected steepest ascent method below, and proposition 15.

The feasibility condition for the (GEiCP) is analogous to the one for the (EiCP).

Proposition 14 *(GEiCP)_J is feasible if and only if there exists some $x \in \mathbb{R}^n$ such that $x_J \geq 0$ and $x^t A x > 0$.*

It is clear that if the (EiCP) is feasible, then the (GEiCP)_J is feasible. But the feasibility of the (GEiCP)_J holds for a larger class of matrices. For instance, the (GEiCP)_J with $n = 2$, $J = \{1\}$, $B = I$ and

$$A = \begin{bmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{2} & -1 \end{bmatrix}$$

is feasible, since $x = (2, -1)^t$ is such that $x_J = 2 \geq 0$ and $x^t A x = 1 > 0$. A solution is obtained with $x = (0.8844, -0.3663)$ and $\lambda = 0.2071$. However, the corresponding (EiCP) is infeasible, since $-A$ is copositive.

For the practical solution of the symmetric (GEiCP)_J, a starting point satisfying $x_J \geq 0$ and $x^t A x > 0$ is needed, as well as an ascent gradient method for the optimization problem

(OG*Ei*CP).

The initialization for the solution of the (GEiCP)_J is akin to the (EiCP) case. The heuristics of lemmas 10, 5, 6 and 7 are applicable in this framework as well, and the feasibility question can be definitively answered by solving the general LCP below.

$$\begin{cases} u + \alpha e + Ax = 0 \\ -\alpha \geq \varepsilon \\ u_J \geq 0 \\ x_J \geq 0 \\ u_J^t x_J = 0 \\ u_J = 0 \\ e^t x_J = 1. \end{cases}$$

In this problem $x \neq 0$ is replaced by $e^t x_J = 1$. This is not done in the optimization problem (OG*Ei*CP) to ensure compactness of the feasible region, which is needed in the convergence proof of the method presented below.

Given a good starting point, a gradient-ascent method can be easily adapted to handle the constraint $\|x\| = 1$ implicitly, by renormalizing every iterate of the sequence. As an example, the Projected Steepest Ascent Method for the solution of the symmetric (GEiCP) is presented next, using Armijo's step-length choice.

Projected Steepest Ascent Method for the Symmetric (GEiCP)_J

0. Let $\varepsilon \in (0, 1)$.

Let x^0 satisfy $x_J \geq 0$, $\|x\| = 1$ and $x^t Ax > 0$.

Let $k \leftarrow 0$.

1. Compute $\nabla\lambda(x^k) = \frac{2}{(x^k)^t B x^k} [A - \lambda(x^k)B]x^k$, and d^k as

$$d_j^k = \begin{cases} 0 & \text{if } j \in J, \ x_j^k = 0 \text{ and } \nabla\lambda(x^k)_j < 0 \\ \nabla\lambda(x^k)_j & \text{otherwise.} \end{cases}$$

2. Let

$$\begin{aligned} t_k &= \min \left\{ t \in \mathbb{N} \mid (x^k + \frac{1}{2^t} d^k)_J \geq 0 \text{ and } \lambda \left(x^k + \frac{1}{2^t} d^k \right) - \lambda(x^k) \geq \frac{\varepsilon}{2^t} \|d^k\|^2 \right\}, \\ \mu_k &= \frac{1}{2^{t_k}}, \\ y^{k+1} &= x^k + \mu_k d^k, \\ x^{k+1} &= \frac{y^{k+1}}{\|y^{k+1}\|}. \end{aligned}$$

3. Set $k \leftarrow k + 1$ and go back to step 1.

Proposition 15 *The Projected Steepest Ascent Method converges to a solution of the symmetric (GEiCP).*

Proof : First notice that the direction d^k computed in step 2 is either an ascent direction or zero. This happens since $\nabla\lambda(x^k)^t d^k = (d^k)^t d^k > 0 \iff d^k \neq 0$.

The renormalization in step 3 can be done, since $y^{k+1} \neq 0$. This follows from the fact that $(x^k)^t d^k = (x^k)^t \nabla\lambda(x^k) = \frac{2}{(x^k)^t B x^k} [(x^k)^t A x^k - \lambda(x^k)(x^k)^t B x^k] = 0$. Furthermore, the renormalization preserves the value of the objective function,


$$\lambda(x^{k+1}) = \lambda \left(\frac{y^{k+1}}{\|y^{k+1}\|} \right) = \lambda(y^{k+1}),$$

as well the constraint $x_j^k \geq 0$. Therefore the sequence $\{x^k\}$ is well-defined and belongs to the compact set $\{x \mid x_J \geq 0, \|x\| = 1\}$.

For any convergent subsequence $x^k \rightarrow x^*$, it is clear that $\nabla\lambda(x^k) \rightarrow \nabla\lambda(x^*)$ and $d^k \rightarrow d^*$ where

$$d_j^* = \begin{cases} 0 & \text{if } j \in J, \ x_j^* = 0 \text{ and } \nabla\lambda(x^*)_j < 0 \\ \nabla\lambda(x^*)_j & \text{otherwise.} \end{cases}$$

It is proved in the sequel that $d^* = 0$. In fact, by the Armijo step-length choice there exists a constant $C > 0$ such that $\lambda(y^{k+1}) - \lambda(x^k) \geq C \|d^k\|^2$. This is a canonical proof [Bazaraa, Sherali & Shetty, theorem 8.6.3] that uses the Mean-Value Theorem, the Cauchy-Schwarz inequality and the Lipschitz-continuity of $\nabla\lambda(\cdot)$. Since $\|x^{k+1} - x^k\| \rightarrow 0$, it follows that $\theta^k \rightarrow 0$, where θ^k is the angle between x^{k+1} and x^k .



Therefore

$$\cos \theta^k = \frac{\|y^{k+1} - x^k\|}{\|x^k\|} = \|y^{k+1} - x^k\| \rightarrow 0.$$

Since $\lambda(\cdot)$ is continuous, $\lambda(y^{k+1}) - \lambda(x^k) \rightarrow 0$. Since $\|d^k\|^2 < \frac{1}{C}(\lambda(y^{k+1}) - \lambda(x^k))$, it follows that $\|d^k\|^2 \rightarrow 0$, that is, $d^* = 0$.

Since $\nabla\lambda(x^*)_J = d_J^* = 0$ and $\nabla\lambda(x^*)_J \leq 0$, it follows that $x = x^*$, $\alpha = 0$ and $u = -\nabla\lambda(x^*)$ solve the Kuhn-Tucker conditions for the $(\text{OGEiCP})_J$, and since $\lambda(x^*) \geq \lambda(x^0) > 0$, this is a solution of the $(\text{GEiCP})_J$. ■

The above algorithm has been chosen for clarity of exposition. Other line search strategies may be used, as well as other ascent directions d^k (such that $\nabla\lambda(x^k)^t d^k > 0$). It must be added that the line search in step 2 can be solved by exact maximization, since the derivative of the function $\hat{\lambda}(\mu) = \lambda(x^k + \mu d^k)$ is zero if and only if μ is a root of a polynomial of order

3, as can easily be seen. Therefore at most 4 evaluations of the function $\hat{\lambda}(\cdot)$ are needed, namely the 3 roots mentioned above, and the point $x^k + \bar{\mu}d^k$ corresponding to the maximal step-length $\bar{\mu}$ such that $(x^k + \bar{\mu}d^k)_J \geq 0$.

5 Computational Experience

In order to compare the original formulation of the (GEiCP) as a mixed NCP, and its reformulation as a nonlinear programming problem (OGEiCP), with respect to the relative difficulty of solution, two commercial packages have been chosen inside the GAMS 2.5E system: GAMS/PATH 4.3 for mixed NCPs, and GAMS/MINOS 5.5 for NLPs.

The test problems were given by 40 randomly generated matrices $A \in \mathbb{R}^{n \times n}$, 10 matrices for each value of $n = 10, 20, 30, 40$ ¹. Each matrix was forced to satisfy $A_{11} > 0$, which gives a convenient starting solution $x^0 = (1, 0, \dots, 0)^t \geq 0$ such that $\lambda(x^0) > 0$. The corresponding problems, taking $B = I$ and $J = \{1, 2, \dots, n\}$, were given as input for both software packages. The tests were run in a Pentium II 333 MHz, running MS-Windows 98.

The PATH algorithm is an implementation of a Nonsmooth Damped Newton Method [Dirkse & Ferris]. This method is applied to a reformulation of the mixed NCP as a generalized equation. The MINOS algorithm combines a Reduced-Gradient Method with a Quasi-Newton Method.

The computational experience is summarized by the following table. The measures for the number of iterations and running time are average measures over the 10 problems in the corresponding dimension. For the PATH algorithm, major iterations correspond to Newton steps, and minor iterations are pivot steps in Lemke's Method used to construct a piecewise-linear path from an iterate to the corresponding Newton point. Columns 4,5 and 6 are the

¹maximum dimension allowed by the free version of the software.

number of problems solved without restarts, the number of problems solved using up to 3 restarts from distinct initial points, and the number of problems not solved by the PATH algorithm, respectively. Note that the convergence of the PATH method is only established under the hypothesis of local inversibility of the jacobians defining the NCP, which does not hold for the (GEiCP).

Dimension	PATH						MINOS	
	# Major Iter.	# Minor Iter.	# Successes	# Suc. w/ Restart	# Failures	Time (sec)	# Iter.	Time (sec)
10	33.6	165.5	7	2	1	0.30	16.8	0.04
20	47.9	588.7	7	3	0	0.39	35.6	0.05
30	77.1	958.8	4	6	0	0.71	52.5	0.09
40	93.9	2418.7	3	6	1	1.30	67.8	0.15

Note that the PATH algorithm, without restarts, fails in 30% to 70% of the tests. Thanks to the restart mechanism and the heuristics employed, a considerable portion of the problems has been solved. In another test, 50 random problems were generated as above for $n = 40$; the PATH algorithm, with up to 3 restarts, failed for 16% of them. The MINOS algorithm, on the other hand, has always found a solution, with running times ranging from 7.5 to 8.7 times faster.

It is possible to conclude from the above experience that the MINOS algorithm applied to the (OGEiCP) performs better than the PATH algorithm applied to the original formulation

of the (GEiCP), with respect to running time and robustness (in finding a solution). It is important to add that the PATH algorithm is considered to be the most robust procedure for solving mixed complementarity problems. So the numerical results of this experience clearly indicate the relevance of the optimization formulation for finding a solution of the (GEiCP) in the symmetric case.

6 The Asymmetric Case for $n = 2$

In this section the (EiCP) with $B = I$ is considered, without the symmetry assumption on the matrix A . It is shown that the existence of a solution when all the diagonal elements of A are positive and the number of solutions established for symmetric matrices also hold for asymmetric matrices of order 2.

Proposition 16 *If A is a 2×2 -matrix, then the (EiCP) has at most 3 solutions.*

Proof : The upper bound given by proposition 3 is 4. If $A_{ii} \leq 0$ for some i , or if a column of A is positive, then the solutions corresponding to $I = \{i\}$ or $I = \{j\}$ are infeasible, and so the result follows.

On the other hand, if $A_{ii} > 0$ and $A_{ij} \leq 0$, for all $i \neq j$, then the solutions corresponding to $I = \{i\}$ or $I = \{j\}$ are feasible. The remaining case is $I = \{1, 2\}$, for which it is shown next that at most one solution is feasible.

The characteristic polynomial of A

$$f(\lambda) = \det \begin{pmatrix} \lambda - A_{11} & -A_{12} \\ -A_{21} & \lambda - A_{22} \end{pmatrix} = \lambda^2 - (A_{11} + A_{22})\lambda + [A_{11}A_{22} - A_{12}A_{21}]$$

has at most 2 positive roots. Furthermore,

$$f(A_{11}) = f(A_{22}) = -A_{12}A_{21} \leq 0;$$

If $A_{12}A_{21} = 0$, then either A_{11} or A_{22} is an eigenvalue of A , with the same solution of the cases $I = \{i\}$ or $I = \{j\}$. Therefore the (EiCP) has at most 3 solutions.

Now suppose that $A_{12}, A_{21} < 0$. Then $f(A_{11}) = f(A_{22}) < 0$, and there is an eigenvalue $\lambda > \max\{A_{11}, A_{22}\} > 0$. The corresponding eigenvector x must satisfy

$$(\lambda - A_{11})x_1 = A_{12}x_2$$

$$x_1 + x_2 = 1$$

$$x_1, x_2 \geq 0.$$

This is impossible, since $A_{12} < 0$ and $\lambda > A_{11}$. Therefore this λ is not a solution of the (EiCP), and the maximal number of solutions is 3. ■

Proposition 17 *If A is a matrix of order 2 with positive diagonal elements, then the (EiCP) is feasible.*

Proof : If there is some $j \neq i$ such that $A_{ij} \leq 0$, then the result follows. Otherwise, $A_{ij} > 0, \forall i, j$, and $f(A_{11}) = f(A_{22}) = -A_{12}A_{21} < 0$, where, as before, $f(\lambda)$ is the characteristic polynomial of A . Then there is some eigenvalue $\lambda > \max\{A_{11}, A_{22}\} > 0$. Since $A_{12} > 0$, the following system has a solution (x_1, x_2) ,

$$(\lambda - A_{11})x_1 = A_{12}x_2$$

$$x_1 + x_2 = 1$$

$$x_1, x_2 \geq 0.$$

This solution satisfy $A_{21}x_1 = (\lambda - A_{22})x_2$, since λ is an eigenvalue of A . Therefore, the (EiCP) is feasible. ■

This result implies that the (EiCP) is feasible when A is a strictly copositive or a P-matrix (in particular, when A is positive definite) of order 2. Notice that all these classes of

matrices satisfy the necessary condition for the feasibility of the (EiCP)

$$\exists x \geq 0 \text{ such that } x^t A x > 0$$

stated before. Furthermore they are sufficient for the symmetric case. More research is required to investigate whether these classes of matrices are sufficient for the feasibility of the (EiCP) in the case of a general asymmetric matrix.

7 Conclusions and Future Work

In this paper the Eigenvalue Complementarity Problems introduced in [Costa & alii] have been considered. It has been shown that the symmetric cases are solvable with optimization techniques applied to the Rayleigh quotient subject to linear constraints. The initialization for the optimization process is a NP-complete problem, but for the majority of practical cases the initial point can be found by inspection.

Some preliminary work on the general case with asymmetric matrices of order 2 concerning the characterization of feasibility and the number of solutions has been presented. The generalization of these results for asymmetric matrices of any order is the subject of ongoing research.

References

- [Auchmuty 89] AUCHMUTY, G.; Unconstrained Variational Principles for Eigenvalues of Real Symmetric Matrices. *SIAM Journal on Mathematical Analysis*, 20(5):1186-1207, 1989.

- [Auchmuty 91] AUCHMUTY, G.; Globally and Rapidly Convergent Algorithms for Symmetric Eigenproblems. *SIAM Journal on Matrix Analysis and Applications*, 12(4):690-706, 1991.
- [Bazaraa, Sherali & Shetty] BAZARAA, M. S.; SHERALI, H. D.; SHETTY, C. M.; Nonlinear Programming: Theory and Algorithms, 2nd Edition. John Wiley and Sons, New York, 1993.
- [Chatelin] CHATELIN, F.; Eigenvalues of Matrices. John Wiley & Sons, 1993.
- [Costa & alii] COSTA, A. P.; FIGUEIREDO, I. N.; MARTINS, J. A. C.; JÚDICE, J.; A Complementarity Eigenproblem in the Stability Analysis of Finite Dimensional Elastic Systems with Frictional Contact. In.: Complementarity: Applications, Algorithms and Extensions, edited by M. Ferris, J. S. Pang and O. Mangasarian, Kluwer, New York, pp. 67-83, 2001.
- [Dirkse & Ferris] DIRKSE, S.; FERRIS, M.; The PATH Solver: A Nonmonotone Stabilization Scheme for Mixed Complementarity Problems. *Optimization and Software*, 5:123-156, 1995.
- [Harker & Pang] HARKER, P. T.; PANG, J.-S.; Finite-Dimensional Variational Inequality and Nonlinear Complementarity Problems: A Survey of Theory, Algorithms and Applications. *Mathematical Programming - Series B - Variational Inequality Problems*, 48(2):161-220, 1990.

- [Júdice & Faustino] JÚDICE, J. J.; FAUSTINO, A. M.; A Sequential LCP Method for Bilevel Linear Programming. *Annals of Operations Research*, 34:89-106, 1992.
- [Mongeau & Torki] MONGEAU, M.; TORKI, M.; Computing Eigenelements of Real Symmetric Matrices via Optimization. *Technical Report MIP 99-54*, Université Paul Sabatier, Toulouse, 1999.
- [Murty] MURTY, K. G.; Linear Complementarity, Linear and Nonlinear Programming. Heldermann Verlag, Berlin, 1988.
- [Ortega] ORTEGA, J. M.; Matrix Analysis: A Second Course. Plenum Press, New York, 1987.
- [Parlett] PARLETT, B. N.; The Symmetric Eigenvalue Problem. Classics in Applied Mathematics 20, SIAM, 1997.

RELATÓRIOS TÉCNICOS

DEPARTAMENTO DE CIÊNCIA DA COMPUTAÇÃO

Instituto de Matemática e Estatística da USP

A listagem contendo os relatórios técnicos anteriores a 1997 poderá ser consultada ou solicitada à Secretaria do Departamento, pessoalmente, por carta ou e-mail (mac@ime.usp.br).

Flávio Soares Corrêa da Silva e Daniela Vasconcelos Carbogim
A TWO-SORTED INTERPRETATION FOR ANNOTATED LOGIC
RT-MAC-9801, fevereiro de 1998, 17 pp.

Flávio Soares Corrêa da Silva, Wamberto Weber Vasconcelos, Jaume Agustí, David Robertson e Ana Cristina V. de Melo.
WHY ONTOLOGIES ARE NOT ENOUGH FOR KNOWLEDGE SHARING
RT-MAC-9802, outubro de 1998, 15 pp.

J. C.de Pina e J. Soares
ON THE INTEGER CONE OF THE BASES OF A MATROID
RT-MAC-9803, novembro de 1998, 16 pp.

K. Okuda and S.W.Song
REVISITING HAMILTONIAN DECOMPOSITION OF THE HYPERCUBE
RT-MAC-9804, dezembro 1998, 17pp.

Markus Endler
AGENTES MÓVEIS: UM TUTORIAL
RT-MAC-9805, dezembro 1998, 19pp.

Carlos Alberto de Bragança Pereira, Fabio Nakano e Julio Michael Stern
A DYNAMIC SOFTWARE CERTIFICATION AND VERIFICATION PROCEDURE
RT-MAC-9901, março 1999, 21pp.

Carlos E. Ferreira e Dilma M. Silva
BCC DA USP: UM NOVO CURSO PARA OS DESAFIOS DO NOVO MILÊNIO
RT-MAC-9902, abril 1999, 12pp.

Ronaldo Fumio Hashimoto and Junior Barrera

A SIMPLE ALGORITHM FOR DECOMPOSING CONVEX STRUCTURING ELEMENTS

RT-MAC-9903, abril 1999, 24 pp.

Jorge Euler, Maria do Carmo Noronha e Dilma Menezes da Silva

ESTUDO DE CASO: DESEMPENHO DEFICIENTE DO SISTEMA OPERACIONAL LINUX PARA CARGA MISTA DE APLICAÇÕES.

RT-MAC-9904, maio 1999, 27 pp.

Carlos Humes Junior e Paulo José da Silva e Silva

AN INEXACT CLASSICAL PROXIMAL POINT ALGORITHM VIEWED AS DESCENT METHOD IN THE OPTIMIZATION CASE

RT-MAC-9905, maio 1999, pp.

Carlos Humes Junior and Paulo José da Silva e Silva

STRICT CONVEX REGULARIZATIONS, PROXIMAL POINTS AND AUGMENTED LAGRANGIANS

RT-MAC-9906, maio 1999, 21 pp.

Ronaldo Fumio Hashimoto, Junior Barrera, Carlos Eduardo Ferreira

A COMBINATORIAL OPTIMIZATION TECHNIQUE FOR THE SEQUENTIAL DECOMPOSITION OF EROSIONS AND DILATIONS

RT-MAC-9907, maio 1999, 30 pp.

Carlos Humes Junior and Marcelo Queiroz

ON THE PROJECTED PAIRWISE MULTICOMMODITY FLOW POLYHEDRON

RT-MAC-9908, maio 1999, 18 pp.

Carlos Humes Junior and Marcelo Queiroz

TWO HEURISTICS FOR THE CONTINUOUS CAPACITY AND FLOW ASSIGNMENT GLOBAL OPTIMIZATION

RT-MAC-9909, maio 1999, 32 pp.

Carlos Humes Junior and Paulo José da Silva e Silva

AN INEXACT CLASSICAL PROXIMAL POINT ALGORITHM VIEWED AS A DESCENT METHOD IN THE OPTIMIZATION CASE

RT-MAC-9910, julho 1999, 13 pp.

Markus Endler and Dilma M. Silva and Kunio Okuda

A RELIABLE CONNECTIONLESS PROTOCOL FOR MOBILE CLIENTS

RT-MAC-9911, setembro 1999, 17 pp.

David Robertson, Fávio S. Corrêa da Silva, Jaume Agustí and Wamberto W. Vasconcelos
A LIGHTWEIGHT CAPABILITY COMMUNICATION MECHANISM
RT-MAC-9912, novembro 1999, 14 pp.

Flávio S. Corrêa da Silva, Jaume Agustí, Roberto Cássio de Araújo and Ana Cristina V. de Melo
KNOWLEDGE SHARING BETWEEN A PROBABILISTIC LOGIC AND BAYESIAN BELIEF NETWORKS
RT-MAC-9913, novembro 1999, 13 pp.

Ronaldo F. Hashimoto, Junior Barrera and Edward R. Dougherty
FINDING SOLUTIONS FOR THE DILATION FACTORIZATION EQUATION
RT-MAC-9914, novembro 1999, 20 pp.

Marcelo Finger and Wamberto Vasconcelos
SHARING RESOURCE-SENSITIVE KNOWLEDGE USING COMBINATOR LOGICS
RT- MAC-2000-01, março 2000, 13pp.

Marcos Alves e Markus Endler
PARTICIONAMENTO TRANSPARENTE DE AMBIENTES VIRTUAIS DISTRIBUÍDOS
RT- MAC-2000-02, abril 2000, 21pp.

Paulo Silva, Marcelo Queiroz and Carlos Humes Junior
A NOTE ON "STABILITY OF CLEARING OPEN LOOP POLICIES IN MANUFACTURING SYSTEMS"
RT- MAC-2000-03, abril 2000, 12 pp.

Carlos Alberto de Bragança Pereira and Julio Michael Stern
FULL BAYESIAN SIGNIFICANCE TEST: THE BEHRENS-FISHER AND COEFFICIENTS OF VARIATION PROBLEMS
RT-MAC-2000-04, agosto 2000, 20 pp.

Telba Zalkind Irony, Marcelo Lauretto, Carlos Alberto de Bragança Pereira and Julio Michael Stern
A WEIBULL WEAROUT TEST: FULL BAYESIAN APPROACH
RT-MAC-2000-05, agosto 2000, 18 pp.

Carlos Alberto de Bragança Pereira and Julio Michael Stern
INTRINSIC REGULARIZATION IN MODEL SELECTION USING THE FULL BAYESIAN SIGNIFICANCE TEST
RT-MAC-2000-06, outubro 2000, 18 pp.

Douglas Moreto and Markus Endler
EVALUATING COMPOSITE EVENTS USING SHARED TREES
RT-MAC-2001-01, janeiro 2001, 26 pp.

Vera Nagamura and Markus Endler
COORDINATING MOBILE AGENTS THROUGH THE BROADCAST CHANNEL
RT-MAC-2001-02, janeiro 2001, 21 pp.

Júlio Michael Stern
THE FULLY BAYESIAN SIGNIFICANCE TEST FOR THE COVARIANCE PROBLEM
RT-MAC-2001-03, fevereiro 2001, 15 pp.

Marcelo Finger and Renata Wassermann
TABLEAUX FOR APPROXIMATE REASONING
RT- MAC-2001-04, março 2001, 22 pp.

Julio Michael Stern
*FULL BAYESIAN SIGNIFICANCE TESTS FOR MULTIVARIATE NORMAL
STRUCTURE MODELS*
RT-MAC-2001-05, junho 2001, 20 pp.

Paulo Sérgio Naddeo Dias Lopes and Hernán Astudillo
VIEWPOINTS IN REQUIREMENTS ENGINEERING
RT-MAC-2001-06, julho 2001, 19 pp.

Fabio Kon
O SOFTWARE ABERTO E A QUESTÃO SOCIAL
RT- MAC-2001-07, setembro 2001, 15 pp.

Isabel Cristina Italiano, João Eduardo Ferreira and Osvaldo Kotaro Takai
ASPECTOS CONCEITUAIS EM DATA WAREHOUSE
RT – MAC-2001-08, setembro 2001, 65 pp.

Marcelo Queiroz, Carlos Humes Junior and Joaquim Júdice
ON FINDING GLOBAL OPTIMA FOR THE HINGE FITTING PROBLEM
RT – MAC –2001-09, novembro 2001, 39 pp.

Marcelo Queiroz, Joaquim Júdice and Carlos Humes Junior
THE SYMMETRIC EIGENVALUE COMPLEMENTARITY PROBLEM
RT – MAC –2001-10, novembro 2001, 33 pp.