

UNIVERSIDADE DE SÃO PAULO
Instituto de Ciências Matemáticas e de Computação
ISSN 0103-2577

**On Pseudo Almost Periodic Solutions to Some Neutral
Functional-Differential Equations**

**Eduardo Hernández M.
Toka Diagana**

Nº 237

NOTAS

Série Matemática



São Carlos – SP
Ago./2005

SYSNO.	1455856
DATA	/ /
ICMC - SBAB	

Uma Nota sobre Soluções Pseudo Quase Periódicas de Equações Funcionais do Tipo Neutro.

Eduardo Hernández & Toka Diagana

Resumo

O objetivo deste artigo é estabelecer a existência e unicidade de soluções pseudo quase periódicas para a equação abstrata do tipo neutro

$$\begin{aligned}\frac{d}{dt}D(u_t) &= AD(u_t) + g(t, u_t), \\ u_0 &= \varphi \in C([-p, 0], X),\end{aligned}$$

onde A é gerador infinitesimal de um C_0 -semigrupo de operadores lineares limitados definido sobre um espaço de Banach X ; $D\psi = \psi(0) - f(t, \psi)$ e f, g são funções apropriadas.

On Pseudo Almost Periodic Solutions to Some Neutral Functional-Differential Equations

Eduardo Hernández M and Toka Diagana

AMS-Subject Classification: 35R10; 34K40; 47D06.

Keywords: Partial functional differential equations, pseudo almost periodic function.

Abstract

This paper discusses the existence and uniqueness of pseudo almost periodic solutions to a class of partial neutral functional-differential equations. Under some suitable assumptions, existence and uniqueness results are obtained. An example is given to illustrate our abstract results.

1 Introduction

The main goal of this paper is establish the existence and uniqueness of a *pseudo almost periodic* solution to the abstract neutral Cauchy problem

$$\frac{d}{dt}D(u_t) = AD(u_t) + g(t, u_t), \quad (1)$$

$$u_0 = \varphi \in C([-p, 0], \mathbb{X}), \quad (2)$$

where A is the infinitesimal generator of an uniformly exponentially stable semigroup of linear operators $(T(t))_{t \geq 0}$ on a Banach space \mathbb{X} , $Du = u(0) - f(t, u)$, the history $u_t : [-p, 0] \mapsto \mathbb{X}$ defined by $u_t(\theta) = u(t + \theta)$ belongs to some phase space \mathcal{B} defined axiomatically, and f, g are some appropriate functions.

The existence of almost automorphic, almost periodic, asymptotically almost periodic, and pseudo almost periodic solutions is one of most attractive topics in the qualitative theory of differential equations due to their significance and applications in physical sciences.

The concept of the pseudo almost periodicity (p.a.p. for short) which is the central issue in this paper was first initiated by C. Y. Zhang in the earlier nineties [23, 24]—and it is a natural generalization of the well-known (Bochner) almost periodicity (a.p.). Thus this new concept is welcome to implement another existing generalization of the (Bochner) almost periodicity, the notion of asymptotically almost periodicity (a.a.p.) due to Fréchet, see, e.g., [5], [8], and [16]. For more on these and related issues, see, e.g., [1], [2], [3], [4], [6], [7], [14], [23] and the references therein.

Some contributions related to pseudo almost periodic solutions to abstract differential equations and partial differential equations have recently been made in [12, 13, 7, 6, 4, 14].

Existence results concerning almost periodic and asymptotically almost periodic solutions to ordinary neutral differential equations and abstract partial neutral differential equations have recently been established in [15, 18, 11]. However, the existence of pseudo almost periodic to functional-differential equations with delay, especially, abstract partial neutral differential equations is an untreated topic and this is the main motivation of the present paper.

This paper is organized as follows. In Section 2 we introduce some notations, definitions and properties related to the theory of almost periodic and pseudo almost periodic functions. Furthermore, we establish some preliminary results on the composition of pseudo almost periodic functions of class p . Finally, in the Section 4, we establish existence and uniqueness of pseudo almost periodic solutions to the above-mentioned problem. We next illustrate the previous existence result by an example.

2 Preliminaries

In this section we introduce the required background and some preliminaries that we need in the sequel. In this paper, $A : D(A) \subset \mathbb{X} \rightarrow \mathbb{X}$ denotes the infinitesimal generator of an uniformly asymptotically stable semigroup of linear operators $(T(t))_{t \geq 0}$ on a Banach space $(\mathbb{X}, \|\cdot\|)$, and M, w are some positive constants such that $\|T(t)\| \leq Me^{-wt}$, $t \geq 0$.

To deal with pseudo almost periodic solutions we will need to introduce some classical and new concepts. In what follows, $(\mathbb{Z}, \|\cdot\|_{\mathbb{Z}})$ and $(\mathbb{W}, \|\cdot\|_{\mathbb{W}})$ stand for Banach spaces. In addition to that $C(\mathbb{R}, \mathbb{Z})$ and $BC(\mathbb{R}, \mathbb{Z})$ denote the collection of continuous functions, and the Banach space of bounded continuous functions from \mathbb{R} into \mathbb{Z} equipped with the sup norm defined by $\|u\|_{\infty} := \sup_{t \in \mathbb{R}} \|u(t)\|$, respectively. Similar definitions apply for both $C(\mathbb{R} \times \mathbb{Z}, \mathbb{W})$ and $BC(\mathbb{R} \times \mathbb{Z}, \mathbb{W})$. The notation $B_r(x, \mathbb{Z})$ stands for the open ball centered at $x \in \mathbb{Z}$ with radius $r > 0$.

Definition 1 *A function $f \in C(\mathbb{R}, \mathbb{Z})$ is almost periodic (a.p.) if for each $\varepsilon > 0$ there exists a relatively dense subset of \mathbb{R} denoted by $\mathcal{H}(\varepsilon, f, \mathbb{Z})$ (i.e., there exists $\delta > 0$ such that $[a, a + \delta] \cap \mathcal{H}(\varepsilon, f, \mathbb{Z}) \neq \{\emptyset\}$ for each $a \in \mathbb{R}$) such that $\|f(t + \tau) - f(t)\|_{\mathbb{Z}} < \varepsilon$ for each $t \in \mathbb{R}$ and each $\tau \in \mathcal{H}(\varepsilon, f, \mathbb{Z})$. The collection of those functions will be denoted by $AP(\mathbb{Z})$.*

We need the next Lemma in the sequel.

Lemma 1 [22, p. 25] *A function $f \in C(\mathbb{R}, \mathbb{Z})$ is almost periodic if and only if the set of functions $\{\sigma_{\tau}f : \tau \in \mathbb{R}\}$, where $(\sigma_{\tau}f)(t) = f(t + \tau)$, is relatively compact in $C(\mathbb{R}, \mathbb{Z})$.*

Similarly, a function $F \in C(\mathbb{R} \times \mathbb{Z}, \mathbb{W})$ is called almost periodic in $t \in \mathbb{R}$ uniformly in $z \in \mathbb{Z}$ if for each $\varepsilon > 0$ and for all compact $K \subset \mathbb{Z}$, if there exists a relatively dense subset of \mathbb{R} denoted by $\mathcal{H}(\varepsilon, F, K)$ such that $\|F(t + \tau, z) - F(t, z)\|_{\mathbb{W}} < \varepsilon$ for all $t \in \mathbb{R}$, $z \in K$, and $\tau \in \mathcal{H}(\varepsilon, F, K)$. The collection of those functions will be denoted $AP(\mathbb{Z}, \mathbb{W})$.

The notation $PAP_0(\mathbb{Z})$ stands for the space of functions

$$PAP_0(\mathbb{Z}) := \left\{ u \in BC(\mathbb{R}, \mathbb{Z}) : \lim_{r \rightarrow \infty} \frac{1}{2r} \int_{-r}^r \|u(t)\|_{\mathbb{Z}} dt = 0 \right\}.$$

To study the issues related to delay we need to introduce the new space of functions defined for each $p \in \mathbb{R}$ by

$$PAP_0(\mathbb{Z}, p) := \left\{ u \in BC(\mathbb{R}, \mathbb{Z}) : \lim_{r \rightarrow \infty} \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \|u(\theta)\|_{\mathbb{Z}} \right) dt = 0 \right\}.$$

In addition to the above-mentioned spaces, the present setting requires the introduction of the following new function spaces

$$\begin{aligned} PAP_0(\mathbb{Z}, \mathbb{W}) &= \left\{ u \in BC(\mathbb{R} \times \mathbb{Z}, \mathbb{W}) : \lim_{r \rightarrow \infty} \frac{1}{2r} \int_{-r}^r \|u(t, z)\|_{\mathbb{W}} dt = 0 \right\}, \text{ and} \\ PAP_0(\mathbb{Z}, \mathbb{W}, p) &:= \left\{ u \in BC(\mathbb{R} \times \mathbb{Z}, \mathbb{W}) : \lim_{r \rightarrow \infty} \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \|u(\theta, z)\|_{\mathbb{W}} \right) dt = 0 \right\}, \end{aligned}$$

where in both cases the limit (as $r \mapsto \infty$) is uniform in $z \in \mathbb{Z}$.

In view of the above, it is clear that $PAP_0(\mathbb{Z}, p)$ and $PAP_0(\mathbb{Z}, \mathbb{W}, p)$ are continuously embedded in $PAP(\mathbb{Z})$ and $PAP_0(\mathbb{Z}, \mathbb{W})$, respectively. Furthermore, it is not hard to see that $PAP_0(\mathbb{Z}, p)$ and $PAP_0(\mathbb{W}, \mathbb{Z}, p)$ are closed in $PAP_0(\mathbb{Z})$ and $PAP_0(\mathbb{W}, \mathbb{Z})$, respectively. Consequently, using [14, Lemma 1.2], one obtains the following:

Lemma 2 *The spaces $PAP_0(\mathbb{Z}, p)$ and $PAP_0(\mathbb{W}, \mathbb{Z}, p)$ endowed with the uniform convergence topology are Banach spaces.*

Definition 2 *A function $f \in BC(\mathbb{R}, \mathbb{Z})$ is called pseudo almost periodic (p.a.p.) if $f = g + \varphi$, where $g \in AP(\mathbb{Z})$ and $\varphi \in PAP_0(\mathbb{Z})$. The class of those functions will be denoted by $PAP(\mathbb{Z})$.*

Definition 3 *A function $F \in BC(\mathbb{R} \times \mathbb{Z}, \mathbb{W})$ is called uniformly pseudo almost periodic (u.p.a.p.) if $F = G + \Phi$, where $G \in AP(\mathbb{Z}, \mathbb{W})$ and $\Phi \in PAP_0(\mathbb{Z}, \mathbb{W})$. The class of those functions will be denoted by $UPAP(\mathbb{Z}, \mathbb{W})$.*

We need to introduce two new notions of pseudo almost periodicity that we will use in the sequel.

Definition 4 *A function $F \in BC(\mathbb{R}, \mathbb{Z})$ is called pseudo almost periodic of class p (p.a.p.p.) if $F = G + \varphi$, where $G \in AP(\mathbb{Z})$ and $\varphi \in PAP_0(\mathbb{Z}, p)$. The class of those functions will be denoted by $PAP(\mathbb{Z}, p)$.*

Definition 5 *A function $F \in BC(\mathbb{R} \times \mathbb{Z}, \mathbb{W})$ is called uniformly pseudo almost periodic of class p (u.p.a.p.p.) if $F = G + \varphi$, where $G \in AP(\mathbb{R} \times \mathbb{Z}, \mathbb{W})$ and $\varphi \in PAP_0(\mathbb{Z}, \mathbb{W}, p)$. The class of those functions will be denoted by $UPAP(\mathbb{Z}, \mathbb{W}, p)$.*

3 Preliminary Results

To establish our main result on the existence of pseudo almost periodic solutions we need to prove some preliminaries results related to the composition of pseudo almost periodic functions of class p . Basically, those results are inspired from ideas and estimates given in [14]. Thus, for the sake of clarity, proofs of those results will be given.

Theorem 1 *Let $F \in PAP(\mathbb{Z}, \mathbb{W})$ and let $h \in PAP(\mathbb{W}, p)$. Assume that there exists a function $L_F : \mathbb{R} \mapsto [0, \infty)$ satisfying*

$$\|F(t, z_1) - F(t, z_2)\|_{\mathbb{W}} \leq L_F(t) \|z_1 - z_2\|_{\mathbb{Z}}, \quad \forall t \in \mathbb{R}, \forall z_1, z_2 \in \mathbb{Z}. \quad (3)$$

If

$$\limsup_{r \rightarrow \infty} \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} L_F(\theta) \right) dt < \infty, \quad \text{and} \quad (4)$$

$$\lim_{r \rightarrow \infty} \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} L_F(\theta) \right) \xi(t) dt = 0 \quad (5)$$

for each $\xi \in PAP(\mathbb{R})$, then the function $t \mapsto F(t, h(t))$ belongs to $PAP(\mathbb{W}, p)$.

Proof: Assume that $F = F_1 + \varphi$, $h = h_1 + h_2$, where $F_1 \in AP(\mathbb{Z}, \mathbb{W})$, $\varphi \in PAP(\mathbb{Z}, \mathbb{W}, p)$, $h_1 \in AP(\mathbb{Z})$ and $h_2 \in AP(\mathbb{Z}, p)$. Consider the decomposition

$$F(t, h(t)) = F_1(t, h_1(t)) + [F(t, h(t)) - F(t, h_1(t))] + \varphi(t, h_1(t)).$$

Since $F_1(\cdot, h_1(\cdot)) \in AP(\mathbb{W})$, it remains to prove that both $[F(\cdot, h(\cdot)) - F(\cdot, h_1(\cdot))]$ and $\varphi(\cdot, h_1(\cdot))$ belong to $PAP(\mathbb{W}, p)$. Using (3) above it follows that

$$\begin{aligned} \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \|F(\theta, h(\theta)) - F(\theta, h_1(\theta))\| \right) dt \\ \leq \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} L_F(\theta) \|h_2(\theta)\| \right) dt \\ \leq \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} L_F(\theta) \right) \cdot \left(\sup_{\theta \in [t-p, t]} \|h_2(\theta)\| \right) dt. \end{aligned}$$

which implies that $[F(\cdot, h(\cdot)) - F(\cdot, h_1(\cdot))] \in PAP(\mathbb{W}, p)$, by (5).

Since $h_1(\mathbb{R})$ is relatively compact in \mathbb{Z} and F_1 is uniformly continuous on sets of the form $\mathbb{R} \times K$ with $K \subset \mathbb{Z}$ being compact, for $\epsilon > 0$ there exists $\delta \in (0, \epsilon)$ such that

$$\|F_1(t, z) - F_1(t, \bar{z})\| \leq \epsilon, \quad z, \bar{z} \in h_1(\mathbb{R})$$

with $\|z - \bar{z}\| < \delta$.

Now, fix $z_1, \dots, z_n \in h_1(\mathbb{R})$ such that $h_1(\mathbb{R}) \subset \bigcup_{i=1}^n B_\delta(z_i, \mathbb{Z})$. Obviously, the sets $E_i = h_1^{-1}(B_\delta(z_i))$ form an open covering of \mathbb{R} , and therefore using the sets $B_1 = E_1$, $B_2 = E_2 \setminus E_1$ and $B_i = E_i \setminus \bigcup_{j=1}^{i-1} E_j$ one obtains a covering of \mathbb{R} by disjoint open sets.

For $t \in B_i$, $h_1(t) \in B_\delta(z_i)$

$$\begin{aligned} \|\varphi(t, h_1(t))\| &\leq \|F(t, h_1(t)) - F(t, z_i)\| + \|-F_1(t, h_1(t)) + F_1(t, z_i)\| + \|\varphi(t, z_i)\| \\ &\leq L_F(t) \|h_1(t) - z_i\| + \epsilon + \|\varphi(t, z_i)\| \\ &\leq L_F(t)\epsilon + \epsilon + \|\varphi(t, z_i)\|. \end{aligned}$$

Using the previous inequality it follows that

$$\begin{aligned} &\frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \|\varphi(t, h_1(t))\| \right) dt \\ &\leq \frac{1}{2r} \sum_{i=1}^n \int_{B_i \cap [-r, r]} \left(\sup_{\theta \in [t-p, t]} \|\varphi(\theta, h_1(\theta))\| \right) dt \\ &\leq \frac{1}{2r} \sum_{i=1}^n \int_{B_i \cap [-r, r]} \left(\sup_{j=1, \dots, n} \left[\sup_{\theta \in [t-p, t] \cap B_j} \|\varphi(\theta, h_1(\theta))\| \right] \right) dt \\ &\leq \frac{1}{2r} \sum_{i=1}^n \int_{B_i \cap [-r, r]} \left(\sup_{j=1, \dots, n} \left[\sup_{\theta \in [t-p, t] \cap B_j} \|F(\theta, h_1(\theta)) - F(\theta, z_j)\| \right] \right) dt \\ &\quad + \frac{1}{2r} \sum_{i=1}^n \int_{B_i \cap [-r, r]} \left(\sup_{j=1, \dots, n} \left[\sup_{\theta \in [t-p, t] \cap B_j} \|F_1(\theta, h_1(\theta)) - F_1(\theta, z_j)\| \right] \right) dt \\ &\quad + \frac{1}{2r} \sum_{i=1}^n \int_{B_i \cap [-r, r]} \left(\sup_{j=1, \dots, n} \left[\sup_{\theta \in [t-p, t] \cap B_j} \|\varphi(\theta, z_j)\| \right] \right) dt \\ &\leq \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} L_F(\theta)\epsilon + \epsilon \right) dt + \sum_{i=1}^n \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \|\varphi(\theta, z_j)\| \right) dt \end{aligned}$$

which clearly shows that $\varphi(\cdot, h_1(\cdot)) \in PAP(\mathbb{W}, p)$ and completes the proof. ■

Remark 1 Note that assumptions (4) and (5) are verified by many functions. Examples include constants functions, functions in $PAP(\mathbb{R}, p)$, and functions of $L^1(\mathbb{R})$ which are decreasing on $[0, \infty)$ and nondecreasing on $[-\infty, 0)$ among others.

To study the system (1) we need the following result:

Theorem 2 If $u \in PAP(\mathbb{Z}, p)$, then $t \rightarrow u_t$ belongs to $PAP(C([-p, 0], \mathbb{Z}), p)$.

Proof: Assume that $u = h + g$ where $h \in AP(\mathbb{Z})$ and $g \in PAP(\mathbb{Z}, p)$. Clearly, $u_t = h_t + g_t$ and from Lemma 1 we infer that h_t is almost periodic. On the other hand, for $r > 0$ we

see that

$$\begin{aligned}
& \frac{1}{2r} \int_{-r}^r \left[\sup_{\theta \in [t-p, t]} \left(\sup_{\xi \in [-p, 0]} \|g(\theta + \xi)\| \right) \right] dt \\
& \leq \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-2p, t]} \|g(\theta)\| \right) dt \\
& \leq \frac{1}{2r} \int_{-r-p}^{r-p} \left(\sup_{\theta \in [t-p, t]} \|g(\theta)\| + \sup_{\theta \in [t, t+p]} \|g(\theta)\| \right) dt \\
& \leq \frac{1}{2r} \int_{-r-p}^{r-p} \left(\sup_{\theta \in [t-p, t]} \|g(\theta)\| \right) dt + \frac{1}{2r} \int_{-r-p}^{r-p} \left(\sup_{\theta \in [t, t+p]} \|g(\theta)\| \right) dt \\
& \leq \frac{r+p}{r} \frac{1}{2(r+p)} \int_{-r-p}^{r+p} \left(\sup_{\theta \in [t-p, t]} \|g(\theta)\| \right) dt + \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \|g(\theta)\| \right) dt
\end{aligned}$$

which enables to complete the proof. ■

Our existence result requires the next theorem.

Theorem 3 Let $u \in PAP_0(\mathbb{Z}, p)$. If v is the function defined by

$$v(t) := \int_{-\infty}^t T(t-s)u(s)ds, \quad \forall t \in \mathbb{R},$$

then $v \in PAP_0(\mathbb{Z}, p)$.

Proof: For $r > 0$ we get

$$\begin{aligned}
& \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \int_{-\infty}^{\theta} \|T(\theta-s)\| \|u(s)\| ds \right) dt \\
& \leq \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} \int_{-\infty}^{\theta} e^{-w(\theta-s)} \|u(s)\| ds \right) dt \\
& \leq \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} e^{wp} \int_{-\infty}^{\theta} e^{-w(t-s)} \|u(s)\| ds \right) dt \\
& \leq \frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} e^{wp} \int_{-\infty}^t e^{-w(t-s)} \|u(s)\| ds \right) dt \\
& \leq \frac{e^{wp}}{2r} \int_{-r}^r \int_{-\infty}^t e^{-w(t-s)} \|u(s)\| ds dt \\
& \leq \frac{e^{wp}}{2r} \int_{-\infty}^{-r} \int_{-r}^r e^{-w(t-s)} \|u(s)\| dt ds + \frac{e^{wp}}{2r} \int_{-r}^r \int_s^r e^{-w(t-s)} \|u(s)\| dt ds \\
& \leq \frac{e^{wp} \|u\|_{\infty}}{2rw} \int_{-\infty}^{-r} e^{w(s+r)} ds + \frac{e^{wp}}{2rw} \int_{-r}^r \|u(s)\| dt.
\end{aligned}$$

Consequently,

$$\frac{1}{2r} \int_{-r}^r \left(\sup_{\theta \in [t-p, t]} v(\theta) \right) dt \leq \frac{e^{wp} \|u\|_{\infty}}{2rw^2} + \frac{e^{wp}}{2r} \int_{-r}^r \|u(s)\| dt$$

which converges to zero as $r \rightarrow \infty$. The proof is now complete. ■

4 Existence results

In this section we study the existence of pseudo almost periodic solution for the neutral system. The next definition of a mild solution is inspired by semigroup theory.

Definition 6 A continuous function $u : [\sigma, \sigma + a) \rightarrow \mathbb{X}$, $a > 0$, is a mild solution of the neutral system (1) on $[\sigma, \sigma + a)$, if $u_{\sigma} = \varphi$ and

$$u(t) = T(t - \sigma)(\varphi(0) + f(\sigma, \varphi)) + f(t, u_t) + \int_{\sigma}^t T(t - s)g(s, u_s)ds, \quad t \in [\sigma, \sigma + a).$$

Since the semigroup is uniformly exponentially stable, the next concept of pseudo almost periodic mild solution is clear.

Definition 7 A function $u \in BC(\mathbb{R}, \mathbb{X})$ is a mild pseudo almost periodic solution of neutral system (1), if for each $t \in \mathbb{R}$ and

$$u(t) = f(t, u_t) + \int_{-\infty}^t T(t - s)g(s, u_s)ds, \quad t \in \mathbb{R}.$$

To discuss the existence of pseudo almost periodic solutions we introduce the next conditions on the functions f, g .

Theorem 4 The functions $f, g : \mathbb{R} \times \mathcal{B} \rightarrow \mathbb{X}$ are continuous and there exist a constant $L_f > 0$ and a continuous function $L_g : \mathbb{R} \rightarrow [0, \infty)$ such that

$$\begin{aligned} \|f(t, \psi_1) - f(t, \psi_2)\| &\leq L_f \|\psi_1 - \psi_2\|_{\mathcal{B}}, \\ \|g(t, \psi_1) - g(t, \psi_2)\| &\leq L_g(t) \|\psi_1 - \psi_2\|_{\mathcal{B}}, \end{aligned}$$

for all $t \in \mathbb{R}$ and every $\psi_i \in \mathcal{B}$ ($i = 1, 2$). If

$$\Theta = L_f + M \sup_{t \in \mathbb{R}} \int_{-\infty}^t e^{-w(t-s)} L_g(s) ds < 1, \quad (6)$$

then there exists a unique pseudo almost periodic mild solution to (1).

Proof: Let $\Gamma : PAP(\mathbb{X}, p) \rightarrow C(\mathbb{R}, \mathbb{X})$ be the operator defined by

$$\Gamma u(t) := -f(t, u_t) + \int_{-\infty}^t T(t - s)g(s, u_s)ds, \quad t \in \mathbb{R}.$$

Clearly, Γu is well-defined and continuous. Moreover, from Theorems 1, 2 and 3 it easily follows that $\Gamma u \in PAP(\mathbb{X}, p)$ whenever $u \in AP(\mathbb{X}, p)$. On the other hand, for $u, v \in PAP(\mathbb{X}, p)$ we get

$$\begin{aligned} \|\Gamma u(t) - \Gamma v(t)\| &\leq L_f \|u_t - v_t\|_{\mathcal{B}} + M \int_{-\infty}^t e^{-w(t-s)} L_g(s) \|u_s - v_s\|_{\mathcal{B}} ds \\ &\leq \left(L_f + M \sup_{t \in \mathbb{R}} \int_{-\infty}^t e^{-w(t-s)} L_g(s) ds \right) \|u - v\|_{\infty} \\ &\leq \Theta \|u - v\|_{\infty} \end{aligned}$$

which prove that Γ is a strict contraction. Consequently, by the Banach fixed-point principle there exists a unique mild solution to (1), which obviously is pseudo almost periodic. ■

5 Example

This section is devoted to a concrete example related to the existence of pseudo almost periodic solutions to the neutral system (1)-(2).

In what follows, we set $\mathbb{X} = (L^2[0, \pi], \|\cdot\|_2)$ and define A the linear operator

$$D(A) := \{f \in \mathbb{X} : f'' \in L^2[0, \pi], f(0) = f(\pi) = 0\}, \quad Af := f'', \quad \forall f \in D(A).$$

It is well known that A is the infinitesimal generator of an analytic semigroup $(T(t))_{t \geq 0}$ on \mathbb{X} . Moreover, A has a discrete spectrum, with eigenvalues of the form $-n^2, n \in \mathbb{N}$, whose corresponding (normalized) eigenfunctions are given by $z_n(\xi) := \sqrt{\frac{2}{\pi}} \sin(n\xi)$.

The corresponding semigroup to A is defined by $T(t)f = \sum_{n=1}^{\infty} e^{-n^2 t} \langle f, z_n \rangle z_n$ for each $f \in \mathbb{X}$, with the estimate $\|T(t)\| \leq e^{-t}$ for every $t \geq 0$.

Consider the first-order neutral system

$$\begin{aligned} \frac{\partial}{\partial t} \left[u(t, \xi) + \int_{-p}^0 b(s) u(t+s, \xi) ds \right] &= \frac{\partial^2}{\partial \xi^2} \left[u(t, \xi) + \int_{-p}^0 b(s) u(t+s, \xi) ds \right] \\ &\quad + a_0(\xi) u(t, \xi) + \int_{-p}^0 a(s) u(t+s, \xi) ds, \end{aligned} \quad (7)$$

$$u(t, 0) = u(t, \pi) = 0, \quad (8)$$

where $(t, \xi) \in \mathbb{R} \times [0, \pi]$ and a, b, a_0, a_1 are continuous functions.

By defining the substituting operators

$$\begin{aligned} f(t, \psi)(\xi) &:= \int_{-p}^0 b(s) \psi(s, \xi) ds \\ g(t, \psi)(\xi) &:= a_0(\xi) \psi(0, \xi) + \int_{-p}^0 a(s) \psi(s, \xi) ds, \end{aligned}$$

the system (7)-(8) can be rewritten as a system of the form (1)-(2). Moreover, it is clear that f, g are bounded linear operators and that

$$\begin{aligned}\|f(t, \psi)\| &\leq \left[\sqrt{\int_{-p}^0 b^2(s) ds} \right] \cdot \|\psi\|_{\mathcal{B}}, \\ \|g(t, \psi)\| &\leq \left(\|a_0\|_{\infty} + \sqrt{\int_{-p}^0 a^2(s) ds} \right) \cdot \|\psi\|_{\mathcal{B}},\end{aligned}$$

for all $t \in \mathbb{R}$ and $\psi_i \in \mathcal{B} (i = 1, 2)$.

The next result is a straightforward consequence of Theorem 4.

Proposition 1 *Under the previous assumptions, the neutral system (7)-(8) has a unique pseudo almost periodic mild solution whenever*

$$\sqrt{\int_{-p}^0 b^2(s) ds} + \left(\|a_0\|_{\infty} + \sqrt{\int_{-p}^0 a^2(s) ds} \right) < 1.$$

References

- [1] M. ADIMY AND K. EZZINBI, Strict Solutions of Nonlinear Hyperbolic Neutral Differential Equations. *Appl. Math. Lett.* **12** (1999), no. 1, 107–112.
- [2] E. AIT DADS AND K. EZZINBI, Existence of Pseudo Almost Periodic Solution for Some Abstract Semilinear Functional-Differential Equation. *Dynam. Systems Appl.* **11** (2002), no. 4, 493–498.
- [3] E. AIT DADS AND K. EZZINBI, Pseudo Almost Periodic Solutions of Some Delay Differential Equations. *J. Math. Anal. Appl.* **201** (1996), no. 3, 840–850.
- [4] B. AMIR AND L. MANIAR, Composition of Pseudo Almost Periodic Functions and Cauchy Problems with Operator of Nondense Domain. *Ann. Math. Blaise Pascal* **6** (1999), no. 1, 1–11.
- [5] C. CORDUNEANU, *Almost Periodic Functions*, Second Edition, Chelsea, New York (1989).
- [6] C. CUEVAS AND M. PINTO, Existence and Uniqueness of Pseudo Almost Periodic Solutions of Semilinear Cauchy Problems with non Dense Domain. *Nonlinear Anal.* **45** (2001), no. 1, Ser. A: Theory Methods, 73–83.
- [7] T. DIAGANA, Pseudo Almost Periodic Solutions to Some Differential Equations. *Nonlinear Anal.* **60** (2005), no. 7, 1277–1286.
- [8] A. M. FINK, *Almost Periodic Differential Equations*, Lecture Notes in Mathematics, Vol. **377**, Springer-Verlag, New York-Berlin, 1974.

- [9] J. K. HALE AND S. M. VERDUYN LUNEL, *Introduction to Functional-Differential Equations*. Applied Mathematical Sciences, **99**. Springer-Verlag, New York, 1993.
- [10] J. K. HALE, Partial Neutral Functional-Differential Equations. *Rev. Roumaine Math. Pures Appl.* **39** (1994), no. 4, 339-344.
- [11] E. M. HERNÁNDEZ AND M. L. PELICER, Asymptotically Almost Periodic and Almost Periodic Solutions for Partial Neutral Differential Equations, *Applied Mathematics Letters*, 2005 (in press).
- [12] H. R. HENRÍQUEZ AND C. H. VÁSQUEZ, Almost Periodic Solutions of Abstract Retarded Functional-Differential Equations with Unbounded Delay. *Acta Appl. Math.* **57** (1999), no. 2, 105-132.
- [13] Y. HINO AND S. MURAKAMI, Limiting Equations and Some Stability Properties for Asymptotically Almost Periodic Functional-Differential Equations with Infinite Delay. *Tôhoku Math. J. (2)* **54** (2002), no. 2, 239-257.
- [14] H.-X. LI, F.-L. HUANG, AND J.-Y. LI, Composition of Pseudo Almost Periodic Functions and Semilinear Differential Equations. *J. Math. Anal. Appl.* **255** (2001), no. 2, 436-446.
- [15] N. MINH MAN AND N. VAN MINH, On the Existence of Quasi-Periodic and Almost Periodic Solutions of Neutral Functional-Differential Equations. *Commun. Pure Appl. Anal.* **3** (2004), no. 2, 291-300.
- [16] G. M. N'GUÉRÉKATA, *Topics in Almost Automorphy*, Springer-Verlag, New York, 2005.
- [17] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer-Verlag, New York, (1983).
- [18] R. YUAN, Existence of Almost Periodic Solutions of Neutral Functional-Differential Equations via Liapunov-Razumikhin Function. *Z. Angew. Math. Phys.* **49** (1998), no. 1, 113-136.
- [19] J. WU AND H. XIA, Rotating waves in Neutral Partial Functional-Differential Equations. *J. Dynam. Differential Equations* **11** (1999), no. 2, 209-238.
- [20] J. WU AND H. XIA, Self-Sustained Oscillations in a Ring Array of Coupled Lossless Transmission Lines. *J. Differential Equations* **124** (1996), no. 1, 247-278.
- [21] J. WU, *Theory and Applications of Partial Functional-Differential Equations*. Applied Mathematical Sciences **119** Springer-Verlag, New York, 1996.
- [22] S. ZAIDMAN, *Almost-periodic functions in abstract spaces*. Res. Notes in Math. **126**. Pitman (Advanced Publishing Program), Boston, MA, 1985.

- [23] C. Y. ZHANG, Pseudo Almost Periodic Solutions of Some Differential Equations, *J. Math. Anal. Appl.* **151** (1994), 62-76.
- [24] C. Y. ZHANG, Pseudo Almost Periodic Solutions of Some Differential Equations II, *J. Math. Anal. Appl.* **192** (1995), 543-561.

Eduardo Hernández M.

Departamento de Matemática
Instituto de Ciências Matemáticas e da Computação
Universidade de São Paulo - Campus de São Carlos
Caixa Postal 668
13560-970 São Carlos, SP. Brazil
E-mail : lalohm@icmc.sc.usp.br

Toka Diagana

Department of Mathematics,
Howard University
2441 Sixth Street N.W.,
Washington, D.C. 20059, USA.
E-mail : tdiagana@howard.edu

NOTAS DO ICMC

SÉRIE MATEMÁTICA

- 236/2005 HERNÁNDEZ, E. M.; DIAGANA, T. – On pseudo almost periodic solutions to some neutral functional-differential equations.
- 235/2005 HERNÁNDEZ, E. M.; DIAGANA, T. – Existence and uniqueness of pseudo almost periodic solutions to some abstract partial neutral functional-differential equations.
- 234/2005 HERNÁNDEZ, E. Existence results for a impulsive abstract partial differential equation with state-dependent delay.
- 233/2005 KUSHNER, L. – Quasi-homogeneous maps in two variables.
- 232/2005 GIMENES, L.P.; FEDERSON, M. - Oscillation by impulses for a second order delay differential equations.
- 231/2005 BIRBRAIR, L.; COSTA, J.; FERNANDES, A.; RUAS, M. – K-Bi-Lipschitz equivalence of real function-germs.
- 230/2005 COSTA, J.C.F.; SITTA, A.M. – Path formulation for Z_2 O Z_2 -equivariant bifurcation problems.
- 229/2005 FERNANDES, A.; SOARES, C.H.; SANTOS, R. – Topological triviality of family of functions and sets.
- 228/2005 BÁGIO, D.; DIAS, I.; PAQUES, A.. – On self-dual normal bases.
- 227/2005 ALVAREZ, S.; BIRBRAIR, L.; COSTA, J.; FERNANDES, A. – Topological K-equivalence of analytic function-germs.