

RT-MAT 2003-25

**HOMOGENEITY RANK OF REAL
REPRESENTATIONS OF COMPACT LIE GROUPS**

**CLAUDIO GORODSKI AND
FABIO PODESTÀ**

Outubro 2003

Esta é uma publicação preliminar (“preprint”).

HOMOGENEITY RANK OF REAL REPRESENTATIONS OF COMPACT LIE GROUPS

CLAUDIO GORODSKI AND FABIO PODESTÀ

ABSTRACT. The main result of this paper is the classification of the real irreducible representations of compact Lie groups with vanishing homogeneity rank.

1. INTRODUCTION

Let a compact Lie group G act smoothly on a smooth manifold M . The codimension of the principal orbits in M is called the cohomogeneity $\text{cohom}(G, M)$ of the action. Püttmann, starting from an inequality for the dimension of the fixed point set of a maximal torus in G due to Bredon ([Bre72], p. 194), introduced in [Pue02] the *homogeneity rank* of (G, M) as the integer

$$\begin{aligned}\text{homrk}(G, M) &= \text{rk } G - \text{rk } G_{\text{princ}} - \text{cohom}(G, M) \\ &= \text{rk } G - \text{rk } G_{\text{princ}} + (\dim G - \dim G_{\text{princ}}) - \dim M,\end{aligned}$$

where G_{princ} is a principal isotropy subgroup of the action and, for a compact Lie group K , $\text{rk } K$ denotes its rank, namely the dimension of a maximal torus. We will see in the next section that orbit-equivalent actions have the same homogeneity rank.

This invariant, although not with this name, had already been considered by Huckleberry and Wurzbacher who proved that a Hamiltonian action of a compact Lie group on a symplectic manifold has vanishing homogeneity rank if and only if the principal orbits are coisotropic with respect to the invariant symplectic form (see [HW90], Theorem 3, p. 267 for this result and other characterizations of this property). If $\rho : G \rightarrow \text{U}(V)$ is a complex representation where V is a complex vector space endowed with an invariant symplectic structure, then the G -action is automatically Hamiltonian, and it has vanishing homogeneity rank if and only if every principal orbit is coisotropic; this condition can be proved to be equivalent to the fact that a Borel subgroup of the complexified group G^c has an open orbit in V , and also to the fact that the naturally induced representation of G on the ring of regular functions $\mathbb{C}[V]$ splits into the sum of mutually inequivalent irreducible representations (see e.g. [Kra85], p. 199). Complex representations with these equivalent properties are called *coisotropic* or *multiplicity-free*; Kac [Kac80] classified the irreducible multiplicity-free representations and, later, Benson and Ratcliff [BR96] and, independently, Leahy [Lea98] classified the reducible ones.

In this paper we consider the case of an irreducible representation $\rho : G \rightarrow \text{O}(V)$ of a compact Lie group G on a real vector space V with vanishing homogeneity rank. Since representations admitting an invariant complex structure have null homogeneity rank if and only if they are multiplicity-free, we will deal only with irreducible representations of real type, also called absolutely irreducible, namely those which admit no invariant complex structure. Our main result is the following theorem.

Date: October 9, 2003.

Theorem 1. *An absolutely irreducible representation ρ of a compact connected Lie group G has vanishing homogeneity rank if and only if it is either orbit-equivalent to the isotropy representation of a non-Hermitian symmetric space of inner type or it is one of the following representations:*

G	ρ	d	c
$\mathrm{Sp}(1) \times \mathrm{Sp}(n)$, $n \geq 2$	(standard) ³ $\otimes_{\mathbf{H}}$ (standard)	$8n$	3
$\mathrm{SO}(4) \times \mathrm{Spin}(7)$	(standard) $\otimes_{\mathbf{R}}$ (spin)	32	5
$\mathrm{Sp}(1) \times \mathrm{Spin}(11)$	(standard) $\otimes_{\mathbf{H}}$ (spin)	64	6

where d denotes the dimension of the representation space and c denotes its cohomogeneity.

2. PRELIMINARIES

Let (G, V) be an absolutely irreducible representation of a compact Lie group G on a real vector space V . It is shown in Corollary 1.2 in [Pue02] that the homogeneity rank of a linear representation is non positive. In this regard, the representations with vanishing homogeneity rank are precisely those with maximal homogeneity rank. The following monotonicity property that is stated on p. 375 in [Pue02] and is valid for smooth actions on smooth manifolds will be the basis of the method of our classification. Since there is no proof in [Pue02], we include one for the sake of completeness.

Proposition 2. *Let (G, M) be a smooth action. If G' be a closed subgroup of G , then $\mathrm{homrk}(G', M) \leq \mathrm{homrk}(G, M)$.*

Proof. We first prove the statement in the case in which M is G -homogeneous, i. e. we prove that given a homogeneous space $M = G/H$, where G is a compact Lie group and H is a closed subgroup, for every closed subgroup G' of G we have

$$\mathrm{homrk}(G', G/H) \leq \mathrm{rk} G - \mathrm{rk} H.$$

We prove this by induction on the dimension of the manifold, the initial case $\dim M = 1$ being clear. Fix the point $o = [H] \in G/H$, a maximal torus T_H of H , and a maximal torus T of G containing T_H . Since conjugation of G' by elements of G does not affect the homogeneity rank, we can assume that a maximal torus T' of G' sits inside T . Then we have

$$\mathrm{rk} G' - \mathrm{rk} G'_o \leq \dim T' - \dim(T' \cap G'_o) = \dim T' \cdot o \leq \dim T \cdot o,$$

where G'_o denotes the isotropy subgroup of G' at o . Therefore

$$\mathrm{rk} G' - \mathrm{rk} G'_o \leq \mathrm{rk} G - \mathrm{rk} H.$$

We now consider the slice representation of G'_o on the normal space W to the orbit $G' \cdot o$; we can assume that the dimension k of W is at least 2, since otherwise G'_o contains a principal isotropy subgroup of $(G, G/H)$ as a subgroup of finite index and the claim follows immediately. Denote by S the unit sphere in W with respect to a G'_o -invariant inner product in W and apply the induction hypothesis. Since G'_o is a closed subgroup of $\mathrm{SO}(k)$, we have

$$\begin{aligned} \mathrm{homrk}(G'_o, S) &= \mathrm{rk} G'_o - \mathrm{rk} G'_{\mathrm{princ}} - \mathrm{cohom}(G', M) + 1 \\ &\leq \mathrm{homrk}(\mathrm{SO}(k), S) \\ &= \frac{1 + (-1)^k}{2}, \end{aligned}$$

where G'_{princ} denotes a principal isotropy subgroup of G' on M . It then follows that

$$\text{homrk}(G', M) \leq \text{rk } G' - \text{rk } G'_o + \frac{-1 + (-1)^k}{2} \leq \text{rk } G - \text{rk } H,$$

and our claim is proved.

In the general case, we fix a G -regular point $p \in M$ and observe that a point $q \in G \cdot p$ is principal for the G' -action on $G \cdot p$ if and only if it is principal for the G' -action in M ; this means that $\text{cohom}(G', M) = \text{cohom}(G, M) + \text{cohom}(G', G \cdot p)$. We know from the previous case that

$$\text{homrk}(G', G \cdot p) \leq \text{homrk}(G, G \cdot p),$$

and now our claim follows by subtracting $\text{cohom}(G, M)$ from both members of the above inequality. \square

Corollary 3. *Let (G, V) be a representation of a compact Lie group G on a real vector space V . If (G, V) is not of vanishing homogeneity rank, then the action of a closed subgroup of G on V is never of vanishing homogeneity rank.*

The preceding corollary indicates a strategy to classify representations with vanishing homogeneity rank. First we observe that the standard representation of $\text{SO}(n)$ on \mathbf{R}^n is of vanishing homogeneity rank if and only if n is even. Then we need to decide which of the maximal subgroups of $\text{SO}(n)$, where n is even, act absolutely irreducibly on \mathbf{R}^n with vanishing homogeneity rank. For each example that we encounter, we examine which of its maximal subgroups still act absolutely irreducibly on \mathbf{R}^n with vanishing homogeneity rank, and so on. The process will eventually yield all the closed subgroups of $\text{SO}(n)$ that act absolutely irreducibly on \mathbf{R}^n with vanishing homogeneity rank. The effectiveness of this strategy is elucidated by the following well known result of Dynkin [Dyn52a].

- Theorem 4** (Dynkin). (1) *Let G be a maximal connected subgroup of $\text{SO}(n)$. Then G is conjugate in $\text{O}(n)$ to one of the following:*
- (a) $\text{SO}(k) \times \text{SO}(n - k)$, where $1 \leq k \leq n - 1$;
 - (b) $\rho(\text{SO}(p) \times \text{SO}(q))$, where $pq = n$ and $3 \leq p \leq q$, and ρ is the real tensor product of the vector representations;
 - (c) $\text{U}(k)$, where $2k = n$;
 - (d) $\rho(\text{Sp}(p) \times \text{Sp}(q))$, where $4pq = n \neq 4$, and ρ is the quaternionic tensor product of the vector representations;
 - (e) $\rho(G_1)$, where G_1 is simple and ρ is a real form of a complex irreducible representation of degree n of real type.
- (2) *Let G be a maximal connected subgroup of $\text{SU}(n)$. Then G is conjugate to one of the following:*
- (a) $\text{SO}(n)$;
 - (b) $\text{Sp}(k)$, where $2k = n$;
 - (c) $\text{S}(\text{U}(k) \times \text{U}(n - k))$, where $1 \leq k \leq n - 1$;
 - (d) $\rho(\text{SU}(p) \times \text{SU}(q))$, where $pq = n$ and $p \geq 3$ and $q \geq 2$, and ρ is the complex tensor product of the vector representations;
 - (e) $\rho(G_1)$, where G_1 is simple and ρ is a complex irreducible representation of degree n of complex type.
- (3) *Let G be a maximal connected subgroup of $\text{Sp}(n)$. Then G is conjugate to one of the following:*

- (a) $U(n)$;
- (b) $Sp(k) \times Sp(n-k)$, where $1 \leq k \leq n-1$;
- (c) $\rho(SO(p) \times Sp(q))$, where $pq = n$ and $p \geq 3$ and $q \geq 1$, and ρ is the real tensor product of the vector representations;
- (d) $\rho(G_1)$, where G_1 is simple and ρ is a complex irreducible representation of degree $2n$ of quaternionic type.

Recall that a symmetric space of compact type $X = L/G$ is said to be of *inner type* if $\text{rk } L = \text{rk } G$; otherwise, X is said to be of *outer type* (compare Theorem 8.6.7 on p. 255 in [Wol84]). Moreover, the isotropy representation of X is absolutely irreducible if and only if X is non-Hermitian. The following lemma implies that the isotropy representations of symmetric spaces of semisimple type that have vanishing homogeneity rank are precisely those coming from non-Hermitian symmetric spaces of inner type.

Lemma 5. *Let (G, V) be the isotropy representation of a symmetric space of compact type $X = L/G$. Then $\text{homrk}(G, V) = 0$ if and only if $\text{rk } G = \text{rk } L$.*

Proof. Let $\mathfrak{l} = \mathfrak{g} + V$ be the Cartan decomposition of X with respect to the involution, where \mathfrak{l} and \mathfrak{g} respectively denote the Lie algebras of L and G . Let $\mathfrak{a} \subset V$ be a maximal Abelian subspace. By the structural theory of symmetric spaces, it is known that the dimension of \mathfrak{a} is equal to the cohomogeneity of (G, V) , and that the centralizer \mathfrak{m} of \mathfrak{a} in \mathfrak{g} is the Lie algebra of a principal isotropy subgroup of (G, V) . It follows that $\text{homrk}(G, V) = \text{rk } \mathfrak{g} - \text{rk } \mathfrak{m} - \dim \mathfrak{a}$. Let \mathfrak{t} be a Cartan subalgebra of \mathfrak{m} . Then it is easily seen that $\mathfrak{t} + \mathfrak{a}$ is a Cartan subalgebra of \mathfrak{l} . Now $\text{rk } \mathfrak{m} = \dim \mathfrak{t}$, $\text{rk } \mathfrak{l} = \dim \mathfrak{t} + \dim \mathfrak{a}$, and hence $\text{homrk}(G, V) = \text{rk } \mathfrak{g} - \text{rk } \mathfrak{l} = \text{rk } G - \text{rk } L$ which proves our thesis. \square

We will also use Theorem 1.3 of [Pue02] which, for convenience of the reader, we restate here.

Theorem 6 (Püttmann). *Let (G, M) be an isometric action of the compact Lie group G on a Riemannian manifold M . Then, for any $x \in M$, we have*

$$\dim \nu_x(G \cdot x)^{G_x} \leq \text{cohom}(G, M) - (\text{rk } G_x - \text{rk } G_{\text{princ}}),$$

where $\nu_x(G \cdot x)$ denotes the normal space to the orbit $G \cdot x$ at x , $\nu_x(G \cdot x)^{G_x}$ denotes the fixed point subspace of G_x in $\nu_x(G \cdot x)$, and G_{princ} is a principal isotropy subgroup of (G, M) .

The following proposition implies that orbit-equivalent actions have the same homogeneity rank.

Proposition 7. *Let (G, M) be a smooth action. If G' is a closed subgroup of G , and G and G' have the same orbits in M , then $\text{homrk}(G', M) = \text{homrk}(G, M)$.*

Proof. It is clearly enough to prove that if G' and G act transitively on the same manifold M , then $\text{homrk}(G', M) = \text{homrk}(G, M)$. If we represent $M = G/H = G'/H'$ for suitable closed subgroups $H \subset G$ and $H' \subset G'$, then we claim that

$$\text{rk } G - \text{rk } H = \text{rk } G' - \text{rk } H'.$$

This follows from the fact that, given a homogeneous space $M = G/H$ with G compact, the number $\chi_{\pi}(M) := \text{rk } H - \text{rk } G$ is a homotopy invariant of M (see [Oni94], p. 207) \square

Finally, we state the following direct consequences of the definition of homogeneity rank, which we shall repeatedly use in our arguments.

Remark 8. Let (G, M) be a smooth action of a compact Lie group G on a smooth manifold M . Then:

- (a) If $\text{homrk}(G, M) = 0$, then $\dim M \leq \dim G + \text{rk } G$.
- (b) If G' is a connected closed subgroup of G having the same homogeneity rank, then $\text{rk } G' \geq \text{rk } G - \text{rk } G_{\text{princ}}$. Moreover if G_{princ} is finite, then $G = G'$ (indeed, G and G' are orbit-equivalent both with finite principal isotropy, hence G and G' have the same Lie algebra).

3. THE CLASSIFICATION

In this section, we apply the strategy discussed in the previous section to classify absolutely irreducible representations with vanishing homogeneity rank. It is enough to consider orthogonal representations of even degree $2n$. According to Theorem 4, the maximal connected subgroups of $\text{SO}(2n)$ acting absolutely irreducibly on $V = \mathbf{R}^{2n}$ are: $\rho(\text{SO}(p) \times \text{SO}(q))$, where $pq = 2n$ and $3 \leq p \leq q$, and ρ is the real tensor product of the vector representations; $\rho(\text{Sp}(p) \times \text{Sp}(q))$, where $4pq = 2n \neq 4$, and ρ is the quaternionic tensor product of the vector representations; and $\rho(G_1)$, where G_1 is simple and ρ is a real form of a complex irreducible representation of degree $2n$ of real type.

3.1. The case of $\rho(\text{SO}(p) \times \text{SO}(q))$ and its maximal subgroups. Here $pq = 2n$ and $3 \leq p \leq q$. We have that ρ is the isotropy representation of the symmetric space $\text{SO}(p+q)/\text{SO}(p) \times \text{SO}(q)$, and $\text{rk } \text{SO}(p+q) = \text{rk } \text{SO}(p) + \text{rk } \text{SO}(q)$ because not both of p, q are odd. It follows from Corollary 3 that this is an example.

Next we must investigate maximal connected subgroups G of $\rho(\text{SO}(p) \times \text{SO}(q))$. We shall consider separately three cases which cover all the possibilities.

3.1.1. $G = \rho(G_1)$, where $G_1 = K \times \text{SO}(q)$, and $K \subset \text{SO}(p)$ is a maximal connected subgroup. Set $\hat{G}_1 = \text{SO}(p) \times \text{SO}(q)$, $\hat{G} = \rho(\hat{G}_1)$. There is a \hat{G}_1 -regular point $x \in V$ whose connected principal isotropy subgroup is given by $\hat{G}_{1x} = \text{SO}(q-p) \subset \text{SO}(q)$. The isotropy subgroup of G_1 at x is the intersection $\hat{G}_{1x} \cap G_1$, and its connected component is $\text{SO}(q-p)$. If G has vanishing homogeneity rank on V , then Theorem 6 applied to x gives

$$\begin{aligned} \dim \nu_x(Gx)^{G_x} &\leq \text{cohom}(G, V) - (\text{rk } G_x - \text{rk } G_{\text{princ}}) \\ &= \text{rk } G - \text{rk } G_x \\ &= \text{rk } K + \text{rk } \text{SO}(q) - \text{rk } \text{SO}(q-p) \\ &= \text{rk } K + \text{rk } \text{SO}(p), \end{aligned}$$

where we have used that not both of p, q are odd. Note that $\dim \nu_x(\hat{G}x) = p$ and $\dim \nu_x(Gx) = p + \dim \text{SO}(p) - \dim K$. It is clear that

$$\nu_x(Gx)^{G_x} \supset \nu_x(\hat{G}x),$$

since $\nu_x(Gx) \supset \nu_x(\hat{G}x)$, $G_x \subset \hat{G}_x$ and x is \hat{G} -regular. It follows that

$$\dim \nu_x(Gx)^{G_x} \geq p.$$

Combining with the above we get that

$$\text{rk } K + \text{rk } \text{SO}(p) \geq p \geq 2\text{rk } \text{SO}(p),$$

and therefore

$$\text{rk } K = \text{rk } \text{SO}(p).$$

By the classification of maximal subgroups of maximal rank of $\mathrm{SO}(p)$, (see, for example, section 8.10 in [Wol84]), since K is irreducible and of real type on \mathbf{R}^p , we must have $K = \mathrm{SO}(p)$.

3.1.2. $G = \rho(G_1)$, where $G_1 = \mathrm{SO}(p) \times K$, and $K \subset \mathrm{SO}(q)$ is a maximal connected subgroup. According to Theorem 4, we need to consider three cases.

(a) $K = \mu(K_1)$, where K_1 is simple and μ is an absolutely irreducible representation of degree q . Of course we need only to consider representations μ such that $\mu(K_1)$ is a proper subgroup of $\mathrm{SO}(q)$.

We may assume $p < q$. Remark 8(a) gives that $p^2 < pq \leq \frac{p(p-1)}{2} + [\frac{p}{2}] + r$, where $r = \dim K_1 + \mathrm{rk} K_1$, and $[x]$ denotes the greatest integer contained not exceeding x . This implies that

$$(1) \quad p^2 < 2r,$$

and that

$$(2) \quad p^2 - 2qp + 2r \geq 0.$$

Equation (2) and $p \geq 3$ then imply that

$$(3) \quad q \leq \frac{r}{3} + \frac{3}{2}.$$

Let s be the minimal degree of an absolutely irreducible representation of K and such that its image is not the full $\mathrm{SO}(s)$. Then

$$(4) \quad s > \frac{r}{3} + \frac{3}{2}$$

is a sufficient condition for (G, V) not to be of vanishing homogeneity rank. We next run through the possibilities for K_1 .

- $K_1 = \mathrm{SU}(m)$, where $m \geq 2$. Here $r = m^2 + m - 2$. If $m = 2$, then $s = 5$, and (4) holds. If $m \geq 3$, then $s = m^2 - 1$ (realized by the adjoint representation), and (4) holds.
- $K_1 = \mathrm{Sp}(m)$, where $m \geq 2$. Here $r = 2m^2 + 2m$. If $m = 2$, then $s = 10$ (realized by the adjoint representation) and (3) holds. If $m \geq 3$ then $s = 2m^2 - m - 1$ (realized by the second fundamental representation) and (3) holds.
- $K_1 = \mathrm{Spin}(m)$, where $m \geq 7$. Here $r = \frac{m(m-1)}{2} + [\frac{m}{2}]$. All irreducible representations of real type violate (3), except possibly the (half-)spin representations of B_{4k-1} , B_{4k} , D_{4k} . These have respectively $q = 2^{4k-1}$, 2^{4k} , 2^{4k-1} . Condition (3) is respectively

$$3 \cdot 2^{4k} \leq 64k^2 - 16k + 9,$$

$$3 \cdot 2^{4k+1} \leq 64k^2 + 16k + 9,$$

$$3 \cdot 2^{4k} \leq 64k^2 + 9.$$

The only cases that survive are B_3 and D_4 . In the case of B_3 we have $q = 8$ and $r = 24$. Then (1) implies that $p = 3, 4, 5, 6$. Next we use (2) to get rid of $p = 5, 6$. We end up with $p = 3$ and $p = 4$, and this gives the admissible cases $(\mathrm{SO}(3) \times \mathrm{Spin}(7), \mathbf{R}^3 \otimes \mathbf{R}^8)$ and $(\mathrm{SO}(4) \times \mathrm{Spin}(7), \mathbf{R}^4 \otimes \mathbf{R}^8)$, but note that the first one of these is orbit-equivalent to $(\mathrm{SO}(3) \times \mathrm{SO}(8), \mathbf{R}^3 \otimes \mathbf{R}^8)$. In the case of D_4 , we have that $\mu(\mathrm{Spin}(8)) = \mathrm{SO}(8)$, and we rule this out.

- K_1 is an exceptional group. Here (4) holds in each case, so there are no examples, see the table below.

K_1	r	s
G_2	16	7
F_4	56	26
E_6	84	78
E_7	140	133
E_8	256	248

(b) $K = \mu(\mathrm{SO}(k) \times \mathrm{SO}(l))$, where $3 \leq k \leq l$ and $q = kl$, and μ is the real tensor product of the vector representations. Here $r = \frac{k^2+12}{2} + \underbrace{\left[\frac{k}{2} \right] - \frac{k}{2} + \left[\frac{l}{2} \right] - \frac{l}{2}}_{=\theta}$. Note that $-1 \leq \theta \leq 0$.

Then (3) is $k^2 + l^2 - 6kl + 2\theta + 9 \geq 0$. Set $m = l - k \geq 0$. Then $m^2 - 4km - 4k^2 + 9 + 2\theta \geq 0$. This implies that

$$(5) \quad 2k + \sqrt{8k^2 - 9 - 2\theta} \leq m.$$

If $pk \leq l$, since the action of $\mathrm{SO}(p) \times \mu(\mathrm{SO}(k) \times \mathrm{SO}(l)) \subset \mathrm{SO}(p) \times \mathrm{SO}(q)$ on $\mathbf{R}^p \otimes \mathbf{R}^q$ is the same thing as the action of $\mu'(\mathrm{SO}(p) \times \mathrm{SO}(k)) \times \mathrm{SO}(l) \subset \mathrm{SO}(pk) \times \mathrm{SO}(l)$ on $\mathbf{R}^{pk} \otimes \mathbf{R}^l$, where μ' is the real tensor product of the vector representations, this case has already been considered in section 3.1.1. So now we assume that

$$(6) \quad pk > l.$$

Note that $q > \sqrt{2r}$. This implies via (2) that $3 \leq p \leq q - \sqrt{q^2 - 2r}$. Combining this with (6) we have $l < kq - k\sqrt{q^2 - 2r}$ and then $l^2(1 - 2k^2) + 2k^2r > 0$. Substituting the value of r we get $k \leq l < k\sqrt{\frac{k^2+2\theta}{k^2-1}}$. We deduce that $\theta = 0$ and

$$(7) \quad 0 \leq m < k \left(\frac{k}{\sqrt{k^2-1}} - 1 \right).$$

Now (5) and (7) combined imply that $3k + \sqrt{8k^2 - 9} < \frac{k^2}{\sqrt{k^2-1}}$, which is impossible for $k \geq 3$.

(c) $K = \mu(\mathrm{Sp}(k) \times \mathrm{Sp}(l))$, where $q = 4kl \neq 4$, and μ is the quaternionic tensor product of the vector representations. We postpone this case to section 3.4.

3.1.3. $G = \{(x, \sigma(x)) : x \in \mathrm{SO}(p)\}$, where $p = q$ and σ is an automorphism of $\mathrm{SO}(p)$. Here Remark 8(a) immediately implies that (G, V) cannot have vanishing homogeneity rank.

3.2. The case of $\rho(\mathrm{Sp}(p) \times \mathrm{Sp}(q))$ and its maximal subgroups. Here $4pq = 2n \neq 4$ and $p \leq q$. We have that ρ is the isotropy representation of the symmetric space $\mathrm{Sp}(p+q)/\mathrm{Sp}(p) \times \mathrm{Sp}(q)$, and $\mathrm{rk} \, \mathrm{Sp}(p+q) = \mathrm{rk} \, \mathrm{Sp}(p) + \mathrm{rk} \, \mathrm{Sp}(q)$. It follows from Corollary 3 that this is an example.

Next we must investigate maximal connected subgroups G of $\rho(\mathrm{Sp}(p) \times \mathrm{Sp}(q))$. We shall consider three cases separately which cover all the possibilities.

3.2.1. $G = \rho(G_1)$, where $G_1 = K \times \mathrm{Sp}(q)$, and $K \subset \mathrm{Sp}(p)$ is a maximal connected subgroup. Set $\hat{G}_1 = \mathrm{Sp}(p) \times \mathrm{Sp}(q)$. There is a \hat{G}_1 -regular point $x \in V$ whose principal isotropy subgroup is given by $\hat{G}_{1x} = \mathrm{Sp}(1)^p \times \mathrm{Sp}(q-p)$. Let \mathcal{O} be the orbit of \hat{G}_1 through x . A point $y \in \mathcal{O}$ is G_1 -regular in \mathcal{O} if and only if it is G_1 -regular in V ; moreover the isotropy subgroup G_{1y} is given by the intersection of G_1 with a suitable conjugate of \hat{G}_{1x} in \hat{G}_1 . This means that a principal isotropy subgroup of G_1 contains a subgroup isomorphic to $\mathrm{Sp}(q-p)$. If (G, V) has vanishing homogeneity rank, it then follows that

$$\dim V = 4pq \leq \dim K + \mathrm{rk} K + 2q^2 + 2q - (2(q-p)^2 + 2(q-p)),$$

hence

$$(8) \quad 2p^2 - 2p \leq \dim K + \mathrm{rk} K.$$

According to Theorem 4, there are two cases to be considered. But if K is of the form $\mu(\mathrm{SO}(k) \times \mathrm{Sp}(l))$, where μ is the real tensor product of the vector representations, we refer to section 3.4. So we can assume that K is of the form $\mu(K_1)$, where K_1 is simple and μ is a complex irreducible representation of degree $2p$ of quaternionic type. Let s be half the minimal degree of a complex irreducible representation of quaternionic type of K_1 and such that its image is not the full $\mathrm{Sp}(s)$. The list of the values of s for each compact simple group K_1 is given by the following table (groups not appearing in the table do not admit quaternionic representations):

K_1	s
$\mathrm{SU}(4a+2)$, $a \geq 1$	$\frac{1}{2}(4a+2)$
$\mathrm{Spin}(8a+3)$, $a \geq 1$	2^{4a}
$\mathrm{Spin}(8a+4)$, $a \geq 1$	2^{4a}
$\mathrm{Spin}(8a+5)$, $a \geq 1$	2^{4a+1}
$\mathrm{Sp}(a)$, $a \geq 3$	$\frac{a}{3}(2a^2 - 3a - 2)$
$\mathrm{Sp}(2)$	8
$\mathrm{Sp}(1)$	2
E_7	28

It is now easy to see that (8) implies $K_1 = \mathrm{Sp}(1)$ and $p = 2$, and then the only admissible case is $G_1 = \mathrm{Sp}(1) \times \mathrm{Sp}(q)$, where $\mathrm{Sp}(1) \subset \mathrm{Sp}(2)$ via the irreducible representation of degree 4.

3.2.2. $G = \rho(G_1)$, where $G_1 = \mathrm{Sp}(p) \times K$, and $K \subset \mathrm{Sp}(q)$ is a maximal connected subgroup.

According to Theorem 4, we need to consider two cases.

(a) $K = \mu(K_1)$, where K_1 is simple and μ is a complex irreducible representation of degree $2q$ of quaternionic type. We may assume $p < q$. Remark 8(a) gives that $4p^2 < 4pq \leq 2p^2 + 2p + r$, where $r = \dim K_1 + \mathrm{rk} K_1$ (note that $r \geq 4$). This implies that

$$p < \frac{1 + \sqrt{1 + 2r}}{2} \quad \text{and} \quad 2p^2 + 2p(1 - 2q) + r \geq 0.$$

From this we get that

$$(9) \quad q \leq \frac{r}{4} + 1,$$

and

$$(10) \quad \text{if } q \geq \frac{1+\sqrt{1+2r}}{2}, \text{ then } p \leq q - \frac{1}{2} - \frac{1}{2}\sqrt{(2q-1)^2 - 2r}.$$

Running through the compact simple groups K_1 that admit quaternionic representations (see table in section 3.2.1) and using (9) and (10), we get the following admissible cases: $K_1 = \mathrm{Sp}(1)$, $p = 1$, $q = 2$; $K_1 = \mathrm{Sp}(3)$, $p = 1$, $q = 7$; $K_1 = \mathrm{Spin}(11)$, $p = 1$, $q = 16$; $K_1 = \mathrm{Spin}(12)$, $p = 1$, $q = 16$; $K_1 = \mathrm{SU}(6)$, $p = 1$, $q = 10$; $K_1 = \mathrm{E}_7$, $p = 1$, $q = 28$. All cases but that of $K_1 = \mathrm{Spin}(11)$ come from isotropy representations of symmetric spaces. (b) $K = \mu(\mathrm{SO}(k) \times \mathrm{Sp}(l))$, where $q = kl$, and μ is the real tensor product of the vector representations. We postpone this case to section 3.4.

3.2.3. $G = \{(x, \sigma(x)) : x \in \mathrm{Sp}(p)\}$, where $p = q$ and σ is an automorphism of $\mathrm{Sp}(p)$. Here Remark 8(a) immediately implies that (G, V) can have vanishing homogeneity rank only if $p = 1$, so this case is out.

3.3. The case of $\rho(G_1)$. Here G_1 is a compact simple Lie group and ρ is an absolutely irreducible representation of G_1 of degree $2n$. Remark 8(a) says that $2n \leq \dim G_1 + \mathrm{rk} G_1$. In particular, this implies that $2 \dim G_1 \geq 2n - 2$, so we can use Lemma 2.6 in [Kol02] to deduce that (G, V) is orbit equivalent to the isotropy representation of a symmetric space.

3.4. The case of $\rho(\mathrm{SO}(m) \times \mathrm{Sp}(p) \times \mathrm{Sp}(q))$, where ρ is the real and quaternionic tensor products of the vector representations. Here $2n = 4mpq$, $m \geq 3$ and $p \leq q$. By direct computation or using Theorem 1.1 in [HH70], we see that:

- (i) if $m \geq 4pq + 2$, then the connected principal isotropy is given by $\mathrm{SO}(m - 4pq)$;
- (ii) if $q \geq mp + 1$, then the connected principal isotropy is given by $\mathrm{Sp}(q - mp)$;
- (iii) in all other cases the connected principal isotropy is trivial.

In case (i) the condition of vanishing homogeneity rank reads

$$4p^2q^2 = p^2 + q^2 + p + q \leq 2p^2 + 2q^2,$$

and this implies $p = q = 1$. In case (ii) we have

$$4p^2(1 - m^2) + 4p(1 + m) + m^2 - m + 2 \left[\frac{m}{2} \right] = 0.$$

If $m = 2l$, then we have $p^2(1 - 4l^2) + p(1 + 2l) + l^2 = 0$, which implies that $1 + 2l$ divides l^2 , impossible. If $m = 2l + 1$, then we have $(4p^2 - 1)l = 2p$, which is impossible. In case (iii), we have the equation

$$8mpq = m^2 - m + 2 \left[\frac{m}{2} \right] + 4p^2 + 4p + 4q^2 + 4q.$$

If $m = 2l$, this reads

$$(11) \quad l^2 - 4pql + p^2 + q^2 + p + q = 0,$$

subject to the constraints

$$\frac{q}{2p} \leq l \leq 2pq, \quad p \leq q, \quad l \geq 2,$$

while if $m = 2l + 1$, we have

$$(12) \quad l^2 - (4pq - 1)l + p^2 + q^2 - 2pq + p + q = 0,$$

subject to the constraints

$$\frac{q}{2p} - \frac{1}{2} \leq l \leq 2pq, \quad p \leq q, \quad l \geq 1.$$

Consider first equation (11). It can be solved in l to yield $l = 2pq \pm \sqrt{\Delta}$, where $\Delta = 4p^2q^2 - p^2 - q^2 - p - q$. If $l = 2pq + \sqrt{\Delta}$, using the fact that $l \leq 2pq$ we have $\Delta = 0$ and then $l = 2pq \leq p + q$, which gives $p = q = 1$, and then $l = 2$, $m = 4$. If $l = 2pq - \sqrt{\Delta}$, then $\frac{q}{2p} \leq l$ implies that

$$q^2(4p^2 - 1) - 4p^2q - 4p^2(p^2 + p) \leq 0,$$

and therefore

$$p \leq q \leq \frac{2p^2 + 2p\sqrt{4p^3 + 4p^2 - 1}}{4p^2 - 1} < p + 2.$$

It then follows that we only need to consider the possibilities $q = p$ and $q = p + 1$. If $q = p$, then we have

$$2 \leq l = 2p^2 - \sqrt{4p^4 - 2p^2 - 2p} \leq 2,$$

so that $l = 2$, $p = q = 1$. If $q = p + 1$, then

$$2 \leq l = 2p^2 + 2p - \sqrt{4p^4 + 8p^3 + 2p^2 - 4p - 2} < 2,$$

which is impossible.

Next we consider equation (12). Here it is useful to note that Remark 8(b) applied to $G = \mathrm{SO}(m) \times \mathrm{Sp}(p) \times \mathrm{Sp}(q)$ viewed as a subgroup of $\mathrm{SO}(m) \times \mathrm{SO}(4pq)$ gives the extra condition

$$(13) \quad \left[\frac{m}{2} \right] \leq p + q.$$

Equation (12) can be solved in l to yield $l = \frac{4pq-1 \pm \sqrt{\Delta_1}}{2}$, where $\Delta_1 = 16p^2q^2 - 4p^2 - 4p - 4q^2 - 4q + 1$. If $l = \frac{4pq-1 + \sqrt{\Delta_1}}{2}$, then (13) implies that

$$2pq \leq l \leq p + q \leq 2q,$$

which gives $p = q = 1$, $\Delta_1 = 1$, $l = 2$ and $m = 5$. If $l = \frac{4pq-1 - \sqrt{\Delta_1}}{2}$, then the inequality $\frac{q}{2p} - \frac{1}{2} \leq l$ implies that

$$q^2(4p^2 - 1) - 4p^2q - 4p^4 - 4p^3 + p^2 \leq 0,$$

which in turn implies

$$p \leq q \leq \frac{2p^2 + p\sqrt{16p^4 + 16p^3 - 4p^2 - 4p + 1}}{4p^2 - 1} < p + 2.$$

Therefore we need only to consider the cases $q = p$ and $q = p + 1$. If $q = p$, then

$$l = \frac{4p^2 - 1 - \sqrt{16p^4 - 8p^2 - 8p + 1}}{2} < 2.$$

This gives $l = 1$, $m = 3$, $p = q = 1$. If $q = p + 1$, then

$$l = 2p^2 + 2p - \frac{1}{2} - \frac{1}{2}\sqrt{16p^4 + 32p^3 + 8p^2 - 16p - 7} < 1,$$

which is impossible.

3.5. The examples and their subgroups. In this section we show that all candidates $G \subset \mathrm{SO}(2n)$ found in the previous sections are actually examples of groups acting absolutely irreducibly on $V = \mathbf{R}^{2n}$ with vanishing homogeneity rank, and these groups do not admit subgroups with the same property. This will complete the proof of Theorem 1.

We first examine the representations that are not orbit-equivalent to isotropy representations of non-Hermitian symmetric spaces of inner type. We have three candidates:

- (1) $G = \mathrm{Sp}(1) \times \mathrm{Sp}(q)$ ($q \geq 2$) acting on $V = S^3(\mathbf{C}^2) \otimes_{\mathbf{H}} \mathbf{C}^{2q} \cong \mathbf{R}^{8q}$;
- (2) $G = \mathrm{SO}(4) \times \mathrm{Spin}(7)$ acting on $V = \mathbf{R}^4 \otimes \mathbf{R}^8 \cong \mathbf{R}^{32}$, where $\mathrm{Spin}(7)$ acts on \mathbf{R}^8 via the spin representation;
- (3) $G = \mathrm{Sp}(1) \times \mathrm{Spin}(11)$ acting on $V = \mathbf{C}^2 \otimes_{\mathbf{H}} \mathbf{C}^{32} \cong \mathbf{R}^{64}$, where $\mathrm{Spin}(11)$ acts on \mathbf{C}^{32} via the spin representation.

We now show that in each case the representation has vanishing homogeneity rank. Indeed, in case 1 we have that a connected principal isotropy is given by $\mathrm{Sp}(q-2)$ (see [GT00], Proposition 7.12), therefore the cohomogeneity is three and the homogeneity rank vanishes. In case 2, a connected isotropy subgroup is trivial. This can be seen by selecting a pure tensor $v \otimes w$ with $v \in \mathbf{R}^4$ and $w \in \mathbf{R}^8$ and computing the connected isotropy, which is $\mathrm{SO}(3) \times G_2$; then the slice representation is given by $\mathbf{R} \oplus \mathbf{R}^3 \otimes \mathbf{R}^7$; starting again with this new representation, we eventually come up with a trivial isotropy. Therefore the cohomogeneity is five and the homogeneity rank vanishes. In case 3, we also have trivial connected principal isotropy and vanishing homogeneity rank. Indeed, if $v \in \mathbf{C}^{32}$ is a highest weight vector for the spin representation of $\mathrm{Spin}(11)$, then the subgroup $H \subset \mathrm{Spin}(11)$ defined by $H = \{g \in \mathrm{Spin}(11) : g \cdot v \in \mathbf{C}^* \cdot v\}$ is given by $\mathrm{U}(5)$. Now if $p : \mathrm{Sp}(1) \times \mathrm{Spin}(11) \rightarrow \mathrm{Spin}(11)$ is the projection, then

$$p((\mathrm{Sp}(1) \times \mathrm{Spin}(11))_v) = \{g \in \mathrm{Spin}(11) : g \cdot v \in \mathrm{Sp}(1) \cdot v\} \supset H.$$

Since H is maximal in $\mathrm{Spin}(11)$, we get that $(\mathrm{Sp}(1) \times \mathrm{Spin}(11))_v$ is given by $\mathbf{T}^1 \cdot \mathrm{SU}(5)$, where \mathbf{T}^1 sits diagonally in the product of a suitable maximal torus in $\mathrm{Sp}(1)$ and the center of H . From this we see that the slice representation at v is given by $\mathbf{R} \oplus \mathbf{C}^5 \oplus \Lambda^2 \mathbf{C}^5$ and the connected principal isotropy is trivial. The cohomogeneity is six and the homogeneity rank vanishes.

We now examine subgroups of the previous examples. In case 1, a maximal subgroup of G leaving no complex structure on V invariant is of the form $G' = \mathrm{Sp}(1) \times K$, where $K \subset \mathrm{Sp}(q)$ is maximal. Since $\mathrm{Sp}(2) \times K$ does not have vanishing homogeneity rank on V by the results of section 3.2.2, and $G' \subset \mathrm{Sp}(2) \times K$, we have that G' does not have vanishing homogeneity rank on V . In cases 2 and 3, G admits no proper subgroups acting with vanishing homogeneity rank because the connected principal isotropy is trivial and then we may apply Remark 8(b).

We finally consider the representations (G, V) that are orbit-equivalent to isotropy representations of non-Hermitian symmetric spaces of inner type, and we classify the subgroups $G' \subset G$ which still act absolutely irreducibly on V with vanishing homogeneity rank. In the following table we list the representations ρ which need to be examined; we denote by c the cohomogeneity of ρ , by d the dimension of V , and by $[[W]]$ a real form of the G -module W .

Case	G	ρ	c	d	$\dim G_{\text{princ}}$
1	$\text{Sp}(1) \cdot \text{SU}(6)$	$\mathbb{C}^2 \otimes_{\mathbb{H}} \Lambda^3 \mathbb{C}^6$	4	40	2
2	$\text{Sp}(1) \cdot \text{Spin}(12)$	$\mathbb{C}^2 \otimes_{\mathbb{H}} \text{(half-spin)}$	4	64	9
3	$\text{Sp}(1) \cdot E_7$	$\mathbb{C}^2 \otimes_{\mathbb{H}} \mathbb{C}^{56}$	4	112	28
4	$\text{Sp}(1) \cdot \text{Sp}(3)$	$\mathbb{C}^2 \otimes_{\mathbb{H}} \mathbb{C}^{14}$	4	28	0
5	$\text{SO}(4)$	$S^3(\mathbb{C}^2) \otimes_{\mathbb{H}} \mathbb{C}^2$	2	8	0
6	$\text{Spin}(16)$	half-spin	8	128	0
7	$\text{SU}(8)$	$[[\Lambda^4 \mathbb{C}^8]]$	7	70	0
8	$\text{SO}(3) \times \text{Spin}(7)$	$\mathbb{R}^3 \otimes \mathbb{R}^8$	3	24	3
9	$\text{Spin}(7)$	spin	1	8	14
10	$\text{Spin}(9)$	spin	1	16	21

Cases 4 through 7 can be dealt with using Remark 8(b). We consider case 1. If G'' is a maximal subgroup of G , we may assume that G'' is of the form $G'' = \text{Sp}(1) \cdot G'''$, where G''' is maximal in $\text{SU}(6)$, since G'' does not leave any complex structure invariant. Now Remark 8(a) implies that $\dim G''' + \text{rk } G''' \geq 36$, and $\text{rk } G''' \leq 5$ implies $\dim G''' \geq 31$, so that $\dim \text{SU}(6)/G''' \leq 4$. If G''' is a proper subgroup of $\text{SU}(6)$, then the left action of $\text{SU}(6)$ on $\text{SU}(6)/G'''$ is almost effective because $\text{SU}(6)$ is simple. Therefore $\dim \text{SU}(6)$ is less than the dimension of the isometry group of $\text{SU}(6)/G'''$, which is at most 10, but this is a contradiction. Hence $G''' = \text{SU}(6)$.

In case 2, again we can assume that $G'' = \text{Sp}(1) \cdot G'''$, where G''' is maximal in $\text{Spin}(12)$. We have $\dim G''' + \text{rk } G''' \geq 60$; since G'' is supposed to act absolutely irreducibly on V , its rank is not maximal by a theorem of Dynkin (see Theorem 7.1, p. 158 in [Dyn52b]), and therefore $\dim G''' \geq 55 = \dim \text{Spin}(11)$. It is known that a subgroup of $\text{Spin}(n)$ of dimension greater or equal to $\dim \text{Spin}(n-1)$ is conjugate to the standard $\text{Spin}(n-1) \subset \text{Spin}(n)$ if $n \neq 4, 8$ (see e.g. [Kob72], p. 49). So, $G''' = \text{Sp}(1) \cdot \text{Spin}(11)$, which is indeed an example with trivial connected principal isotropy by the discussion above.

In case 3, using the same argument as in case 2, we see that $G'' = \text{Sp}(1) \cdot G'''$, where $\dim G''' \geq 102$. An inspection of the list of all maximal subalgebras of E_7 (see Table 12, p. 150 and Theorem 14.1, p. 231 in [Dyn52b]) shows that there is no such proper subgroup.

In case 8, a maximal subgroup G'' acting absolutely irreducibly on V must be of the form $G'' = \text{SO}(3) \times K$, where $K \subset \text{Spin}(7)$ is maximal; arguing as above, we see that $\dim K \geq 18$, so that $\dim(\text{Spin}(7)/K) \leq 3$ and this is impossible, because $\text{Spin}(7)$ is simple.

In case 9, let $K \subset \text{Spin}(7)$ be a maximal subgroup acting absolutely irreducibly on \mathbb{R}^8 . Since K cannot have maximal rank as above, and using Theorem 4, we see that K must be simple of rank at most two and it must admit an irreducible representation of degree 7 and of real type. Moreover, by Remark 8(a), we have $\dim K \geq 6$, hence $\text{rk } K = 2$, and a direct inspection of all such simple groups shows that none of them but G_2 admits an irreducible representation of degree 7. But G_2 does not admit an irreducible representation of degree 8.

In case 10, we consider a maximal subgroup K of $\text{Spin}(9)$ acting absolutely irreducibly on \mathbb{R}^{16} . This means that $\text{rk } K \leq 3$ and $\dim K \geq 13$. Looking at the list of all maximal subgroups of $\text{Spin}(9)$, we see that we can suppose K to be simple and to act irreducibly on \mathbb{R}^9 , via the embedding $K \subset \text{Spin}(9)$. Therefore K must be one of G_2 , $\text{SU}(4)$, $\text{Spin}(7)$ or

$\mathrm{Sp}(3)$, but we immediately see that none of these groups admits an irreducible representation of degree 9 and of real type.

This finishes the proof of Theorem 1.

REFERENCES

- [BR96] C. Benson and G. Ratcliff, *A classification of multiplicity-free representations*, J. Algebra **181** (1996), 152–186.
- [Bre72] G. Bredon, *Introduction to compact transformation groups*, Academic Press, 1972.
- [Dyn52a] E. B. Dynkin, *The maximal subgroups of the classical groups*, Amer. Math. Soc. Trans. **6** (1952), 245–378.
- [Dyn52b] E. B. Dynkin, *Semisimple subalgebras of the semisimple Lie algebras*, Amer. Math. Soc. Trans. **6** (1952), 111–244.
- [GT00] C. Gorodski and G. Thorbergsson, *Representations of compact Lie groups and the osculating spaces of their orbits*, Preprint, University of Cologne, (also E-print math. DG/0203196), 2000.
- [HH70] W.-C. Hsiang and W.-Y. Hsiang, *Differentiable actions of compact connected classical groups, II*, Ann. of Math. (2) **92** (1970), 189–223.
- [HW90] A.T. Huckleberry and T. Wurzbacher, *Multiplicity-free complex manifolds*, Math. Ann. **286** (1990), 261–280.
- [Kac80] V. Kac, *Some remarks on nilpotent orbits*, J. Algebra **64** (1980), 190–213.
- [Kob72] S. Kobayashi, *Transformation groups in differential geometry*, Ergeb. der Math. und ihrer Grenzgeb., Springer-Verlag, 1972.
- [Kol02] A. Kollross, *A classification of hyperpolar and cohomogeneity one actions*, Trans. Amer. Math. Soc. **354** (2002), 571–612.
- [Kra85] H. Kraft, *Geometrische Methoden in der Invariantentheorie*, Aspekte der Math., Vieweg, 1985.
- [Lea98] A. Leahy, *A classification of multiplicity-free representations*, J. Lie Theory **8** (1998), 367–391.
- [Oni94] A. L. Onishchik, *Topology of transitive transformation groups*, Johann Ambrosius Barth, 1994.
- [Pue02] T. Püttmann, *Homogeneity rank and atoms of actions*, Ann. Global Anal. Geom. **22** (2002), 375–399.
- [Wol84] J.A. Wolf, *Spaces of constant curvature*, 5th ed., Publish or Perish, Houston, 1984.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO, 1010,
SÃO PAULO, SP 05508-900, BRASIL

E-mail address: gorodski@ime.usp.br

DIPARTIMENTO DI MATEMATICA E APPL. PER L'ARCHITETTURA, PIAZZA GHIBERTI 27, 50100 FLORENCE, ITALY

E-mail address: Podesta@math.unifi.it

TRABALHOS DO DEPARTAMENTO DE MATEMÁTICA

TÍTULOS PUBLICADOS

- 2002-01 COELHO, F.U. and LANZILOTA, M.A. On non-semiregular components containing paths from injective to projective modules. 13p.
- 2002-02 COELHO, F.U., LANZILOTTA, M.A. and SAVIOLI, A.M.P.D. On the Hochschild cohomology of algebras with small homological dimensions. 11p.
- 2002-03 COELHO, F.U., HAPPEL, D. and UNGER, L. Tilting up algebras of small homological dimensions. 20p.
- 2002-04 SHESTAKOV, I.P. and UMIRBAEV, U.U. Possion brackets and two-generated subalgebras of rings of polynomials. 19p.
- 2002-05 SHESTAKOV, I.P. and UMIRBAEV, U.U. The tame and the wild automorphisms of polynomial rings in three variables. 34p.
- 2002-06 ALENCAR, R. and LOURENÇO, M.L. On the Gelbaum-de Lamadrid's result. 16p.
- 2002-07 GRISHKOV, A. Lie algebras with triality. 28p.
- 2002-08 GRISHKOV, A.N. and GUERREIRO, M. Simple classical Lie algebras in characteristic 2 and their gradations, I. 21p.
- 2002-09 MELO, S.T., NEST, R. and SCHROHE, E. K-Theory of Boutet de Monvel's algebra. 8p.
- 2002-10 POJIDAEV, A.P. Enveloping algebras of Filippov algebras. 17p.
- 2002-11 GORODSKI, C. and THORBERGSSON, G. The classification of taut irreducible representations. 47p.
- 2002-12 BORRELLI, V. and GORODSKI, C. Minimal Legendrian submanifolds of S^{2n+1} and absolutely area-minimizing cones. 13p.
- 2002-13 CHALOM, G. and TREPODE, S. Representation type of one point extensions of quasitilted algebras. 16p.
- 2002-14 GORODSKI, C. and THORBERGSSON, G. Variationally complete actions on compact symmetric spaces. 8p.
- 2002-15 GRISHKOV, A.N. and GUERREIRO, M. Simple classical Lie algebras in characteristic 2 and their gradations, II. 15p.
- 2002-16 PEREIRA, Antônio Luiz and PEREIRA, Marcone Corrêa. A Generic Property for the Eigenfunctions of the Laplacian. 28p.

- 2002-17 GALINDO, P., LOURENÇO, M.L. and MORAES, L.A. Polynomials generated by linear operators. 10p.
- 2002-18 GRISHKOV, A. and SIDKI, S. Representing idempotents as a sum of two nilpotents of degree four. 9p.
- 2002-19 ASSEM, I. and COELHO, F.U. Two-sided gluings of tilted algebras. 27p.
- 2002-20 ASSEM, I. and COELHO, F.U. Endomorphism rings of projectives over Laura algebras. 10p.
- 2002-21 CONDORI, L.O. and LOURENÇO, M.L. Continuous homomorphisms between topological algebras of holomorphic germs. 11p.
- 2002-22 MONTES, R.R. and VERDERESI, J.A. A new characterization of the Clifford torus. 5p.
- 2002-23 COELHO, F.U., DE LA PEÑA, J.A. and TREPODE, S. On minimal non-tilted algebras. 27p.
- 2002-24 GRISHKOV, A.N. and ZAVARNITSINE, A.V. Lagrange's theorem for Moufang Loops. 21p
- 2002-25 GORODSKI, C., OLMOS, C. and TOJEIRO, R. Copolarity of isometric actions. 23p.
- 2002-26 MARTIN, Paulo A. The Galois group of $x^n - x^{n-1} - \dots - x - 1$. 18p.
- 2002-27 DOKUCHAEV, M.A. and MILIES, C.P. Isomorphisms of partial group rings. 12p.
- 2002-28 FUTORNY, V. and OVSIENKO, S. An analogue of Kostant theorem for special PBW algebras. 14p.
- 2002-29 CHERNOUSOVA, Zh. T., DOKUCHAEV, M.A., KHIBINA, M.A., KIRICHENKO, V.V., MIROSHNICHENKO, S.G. and ZHURAVLEV, V.N. Tiled orders over discrete valuation rings, finite Markov chains and partially ordered sets. I. 36p.
- 2002-30 FERNÁNDEZ, J.C.G. On commutative power-associative nilalgebras. 10p.
- 2002-31 FERNÁNDEZ, J.C.G. Superalgebras and identities. 11p.
- 2002-32 GRICHKOV, A.N. , GIULIANI, M.L.M. and ZAVARNITSINE, A.V. The maximal subloops of the simple Moufang loop of order 1080. 10p.
- 2002-33 ZAVARNITSINE, A.V. Recognition of the simple groups $L_3(q)$ by element orders. 20p.
- 2002-34 ZUBKOV, A.N. and MARKO, F. When a Schur superalgebra is cellular? 13p.
- 2002-35 COELHO, F.U. and SAVIOLI, A.M.P.D. On shod extensions of algebras. 11p.

- 2003-01 COELHO, F.U. and LANZILOTTA, M.A. Weakly shod algebras. 28p.
- 2003-02 GREEN, E.L., MARCOS, E. and ZHANG, P. Koszul modules and modules with linear presentations. 26p.
- 2003-03 KOSZMIDER, P. Banach spaces of continuous functions with few operators. 31p.
- 2003-04 GORODSKI, C. Polar actions on compact symmetric spaces which admit a totally geodesic principal orbit. 11p.
- 2003-05 PEREIRA, A.L. Generic Hyperbolicity for the equilibria of the one-dimensional parabolic equation $u_t = (a(x)u_x)_x + f(u)$. 19p.
- 2003-06 COELHO, F.U. and PLATZECK, M.I. On the representation dimension of some classes of algebras. 16p.
- 2003-07 CHERNOUSOVA, Zh. T., DOKUCHAEV, M.A., KHIBINA, M.A., KIRICHENKO, V.V., MIROSHNICHENKO, S.G., ZHURAVLEV, V.N. Tiled orders over discrete valuation rings, finite Markov chains and partially ordered sets. II. 43p.
- 2003-08 ARAGONA, J., FERNANDEZ, R. and JURIAANS, S.O. A Discontinuous Colombeau Differential Calculus. 20p.
- 2003-09 OLIVEIRA, L.A.F., PEREIRA, A.L. and PEREIRA, M.C. Continuity of attractors for a reaction-diffusion problem with respect to variation of the domain. 22p.
- 2003-10 CHALOM, G., MARCOS, E., OLIVEIRA, P. Gröbner basis in algebras extended by loops. 10p.
- 2003-11 ASSEM, I., CASTONGUAY, D., MARCOS, E.N. and TREPODE, S. Quotients of incidence algebras and the Euler characteristic. 19p.
- 2003-12 KOSZMIDER, P. A space $C(K)$ where all non-trivial complemented subspaces have big densities. 17p.
- 2003-13 ZAVARNITSINE, A.V. Weights of the irreducible $SL_3(q)$ -modules in defining characteristic. 12p.
- 2003-14 MARCOS, E. N. and MARTÍNEZ-VILLA, R. The odd part of a N -Koszul algebra. 7p.
- 2003-15 FERREIRA, V.O., MURAKAMI, L.S.I. and PAQUES, A. A Hopf-Galois correspondence for free algebras. 12p.
- 2003-16 KOSZMIDER, P. On decompositions of Banach spaces of continuous functions on Mrówka's spaces. 10p.
- 2003-17 GREEN, E.L., MARCOS, E.N., MARTÍNEZ-VILLA, R. and ZHANG, P. D-Koszul Algebras. 26p.
- 2003-18 TAPIA, G. A. and BARBANTI, L. Um esquema de aproximação para equações de evolução. 20p.

- 2003-19 ASPERTI, A. C. and VILHENA, J. A. Björling problem for maximal surfaces in the Lorentz-Minkowski 4-dimensional space. 18p.
- 2003-20 GOODAIRE, E. G. and MILIES, C. P. Symmetric units in alternative loop rings. 9p.
- 2003-21 ALVARES, E. R. and COELHO, F. U. On translation quivers with weak sections. 10p.
- 2003-22 ALVARES, E.R. and COELHO, F.U. Embeddings of non-semiregular translation quivers in quivers of type $Z\Delta$. 23p.
- 2003-23 BALCERZAK, M., BARTOSZEWCZ, A. and KOSZMIDER, P. On Marczewski-Burstin representable algebras. 6p.
- 2003-24 DOKUCHAEV, M. and ZHUKAVETS, N. On finite degree partial representations of groups. 24p.
- 2003-25 GORODSKI, C. and PODESTÀ, F. Homogeneity rank of real representations of compact Lie groups. 13p.

Nota: Os títulos publicados nos Relatórios Técnicos dos anos de 1980 a 2001 estão à disposição no Departamento de Matemática do IME-USP.
Cidade Universitária “Armando de Salles Oliveira”
Rua do Matão, 1010 - Cidade Universitária
Caixa Postal 66281 - CEP 05315-970