

VOLUME 61(1)

2012

ISSN 0031-0174

The PALAEOBOTANIST

Established 1952

SPECIAL ISSUE

Indo-Brazilian Symposium on Glimpses of Gondwana Research



Birbal Sahni Institute of Palaeobotany
Lucknow, India

EDITORIAL BOARD

Chief Editor

NARESH C. MEHROTRA

Editors

R.P. TIWARI, India

RAJNI TEWARI, India

M. SHANMUKHAPPA, India

Assistant Editor

RATAN KAR, India

EDITORIAL ADVISORY COMMITTEE

DAVID J. BATTEN, UK

S. N. BHALLA, India

DAVID DILCHER, USA

MARTIN J. HEAD, Canada

ROBERTO IANNUZZI, Brazil

V.A. KRASSILOV, Israel

CHENG-SEN LI, P.R. China

SERGE NAUGOLYNKH, Russia

RAJIV NIGAM, India

MARY E. BERNARDES-DE-OLIVEIRA, Brazil

JAMES B. RIDING, UK

GAR W. ROTHWELL, USA

VLADIMIR SERGEEV, Russia

ROBERT A. SPICER, UK

The Palaeobotanist : ISSN 0031-0174. Published in March and September by the Birbal Sahni Institute of Palaeobotany, 53 University Road, Lucknow 226 007 (U.P.), India.

© 2012, Birbal Sahni Institute of Palaeobotany. The appearance of the code at the top of the first page of a paper in this Journal indicates the owner's copyright consent that copies of the research papers may be made for personal or internal use. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purpose, for creating new collective work, for resale or for copying or distributing outside India.

Manuscripts for publication and correspondence on editorial matters should be addressed to the *Chief Editor*, *The Palaeobotanist, Birbal Sahni Institute of Palaeobotany, 53 University Road, Lucknow 226 007, India*. Email: director@bsip.res.in

All business communications should be addressed to the *Registrar, Birbal Sahni Institute of Palaeobotany, 53 University Road, Lucknow 226 007, India*. Email: registrar@bsip.res.in

Phone : 0091-522-2742997

Fax : 0091-522-2740485 / 2740098

Email : palaeobotanist@gmail.com

Web-site : www.bsip.res.in

The views expressed in the papers included in this issue are those of the authors only.

Price per volume : ₹1600.00 (Indian subscribers); US\$ 120.00 (Foreign subscribers, inclusive of air mail postage).

Issued : March 2012

Printed at : Army Printing Press, 33 Nehru Road, Sadar, Lucknow 226 002, India

Palaeobotanical evidence of wildfire in the Upper Permian of India: Macroscopic charcoal remains from the Raniganj Formation, Damodar Basin

ANDRÉ JASPER^{1*}, MARGOT GUERRA-SOMMER², DIETER UHL³,
 MARY E.C. BERNARDES-DE-OLIVEIRA⁴, AMIT K. GHOSH⁵, RAJNI TEWARI⁵
 AND MARIELA INÉS SECCHI¹

¹Centro Universitário UNIVATES, Lajeado, Rio Grande do Sul, Brazil.

²Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil.

³Senckenberg Forschungsinstitut und Naturmuseum, Frankfurt am Main, Germany and Senckenberg Centre for Human Evolution and Palaeoenviroment, Tübingen University, Tübingen, Germany.

⁴Universidade Guarulhos, Guarulhos, São Paulo, Brazil and Universidade de São Paulo, São Paulo, Capital, Brazil.

⁵Birbal Sahni Institute of Palaeobotany, 53 University Road, Lucknow 226 007, India.

*Corresponding author: ajasper@univates.br

(Received 11 November, 2011; revised version accepted 9 January, 2012)

ABSTRACT

Jasper A, Guerra-Sommer M, Uhl D, Bernardes-De-Oliveira MEC, Ghosh AK, Tewari R & Secchi MI 2012. Palaeobotanical evidence of wildfire in the Upper Permian of India: Macroscopic charcoal remains from the Raniganj Formation, Damodar Basin. The Palaeobotanist 61(1): 75-82.

Macroscopic fossil charcoal has been discovered in the carbonaceous shales associated with Seam-VI of Raniganj Formation, Upper Permian, Damodar Basin, India. A pycnoxylic gymnosperm wood is described and confirms the occurrence of palaeo-wildfire in this area during the Late Permian. The integration of the data presented in the current study with previously published data for the Raniganj Formation, principally related to the occurrence of (pyrogenic) inertinites within coal layers, demonstrates that palaeo-wildfires were common events during the deposition of the preserved material. In addition, the presence of charcoal in Permian sediments associated with coal levels at different Gondwana localities demonstrates that wildfires have been relatively common events across the continent during this period.

Key-words—Charcoal, Gymnosperm woods, Palaeo-wildfires, Upper Permian, Raniganj Formation, Raniganj Coalfield, Damodar Basin.

भारत के ऊपरी पर्मियन में वन अग्नि के पुरावानस्पतिक साक्ष्य: रानीगंज शैलसमूह दामोदर द्वोणी से प्राप्त स्थूल लकड़ी के कोयले के अवशेष

एंड्रे जैसपर, मार्गोट गुएरा-सॉमर, डाइटर उह्ल, मेरी ई सी बर्नार्डीज-डी-ऑलीवीरा, अमित के थोष, रजनी तिवारी एवं मेरिएला आइन्स सेक्ची

सारांश

रानीगंज शैलसमूह ऊपरी पर्मियन, दामोदर द्वोणी, भारत की संस्तर-पट्टम कार्बनमय जीवाश्म शेल सहयोगी मैंग्नूल जीवाश्म चारकोल अन्वेषित किया गया है। वनदारुक अनावृतवीजी काष्ठ वर्गित की गई है तथा अंतिम पर्मियन के द्वीरान इस क्षेत्र में पुरा-दावार्नन की घटना की पुरी करती है। रानीगंज शैलसमूह हेतु पृथ्वी में प्रकाशित आंकड़े के साथ भीजुदा अध्ययन में प्रस्तुत आंकड़े का एकाकरण, कोयला परतों के साथ (अग्निजनिक) इनटीनाइटों की सेट्ड्यांतिक रूप से प्राप्ति

प्रदर्शित करती है कि परिवर्तित पश्चात के निकेपण के दौरान पुरा-दावानि आम घटनाएँ थीं। इसके अलावा, विभिन्न गोडवाना क्षेत्रों में कोयला स्तरों के सहयोगी पर्मियन अवसादों में धारकोल की मीजूदगी प्रदर्शित करती है कि इस अवधि में समूचे महाद्वीप में दावानि सापेक्षतया आम घटनाएँ रही हैं।

सकेत-शब्द—लकड़ी का कोयला, अनावृतवीजी काष्ठ, पुरा-दावानि, ऊपरी पर्मियन, रानीगंज शैलसमृद्ध, रानीगंज कोयलाक्षेत्र, इमोदर द्वारा।

Evidências Paleobotânicas de Paleoincêndios no Permiano Superior da Índia: Registro de Charcoal Macroscópico da Formação Raniganj, Bacia Damodar Valley

RESUMO

Charcoal macroscópico fóssil foi descoberto em níveis ricos em matéria orgânica associados à Sequência VI da Formação Raniganj, Permiano Superior, Bacia Damodar Valley, Índia. Um lenho picnoxílico gimnospérmico é descrito e confirma a ocorrência de paleoincêndios vegetacionais na área durante o Neopermiano. A integração dos dados aqui apresentados com aqueles já publicados acerca da Formação Raniganj, principalmente aqueles relacionados à ocorrência de inertinitas (de origem pirogênica) em níveis de carvão, demonstraram que paleoincêndios foram eventos comuns durante a deposição dos níveis estudados. Além disso, a presença de charcoal em sedimentos Permianos associados a níveis de carvão em diferentes localidades do Gondwana confirma que este tipo de evento foi relativamente comum no continente durante esse período.

Palavras-chave—Charcoal, Lenhos Gimnospérmicos, Paleoincêndios vegetacionais, Permiano Superior, Formação Raniganj, Mina Raniganj, Bacia Damodar Valley

INTRODUCTION

Fire plays an important role as a major source of disturbance in many modern ecosystems (Bowman *et al.*, 2009; Flannigan *et al.*, 2009) and it is expected that the occurrence of fire in many areas worldwide may change/increase drastically with changing climate (Flannigan *et al.*, 2009; Westerling *et al.*, 2011). Thus, it is of great interest for the understanding of the interactions between climate, fire-ecology and vegetation to study such interactions under past climate change scenarios. As such palaeobotanical studies can act as long-term experiments on time-scales not available to neo-ecologists.

A period that is of special interest for the understanding of these interactions is the Permian, as this is the only period during the history of the Earth which experienced a long term global climate change from an icehouse into a greenhouse climate after the conquest of the continents by land-plants (Gastaldo *et al.*, 1996).

About a decade ago there were only a few substantiated records of Permian macroscopic fossil charcoal (Scott, 2000). However, a number of studies has subsequently demonstrated the almost ubiquitous presence of macroscopic fossil charcoal in many Permian deposits from the Northern Hemisphere; i.e. North America (DiMichele *et al.*, 2004), Europe (Rößler, 2001; Uhl & Kerp, 2003; Noll *et al.*, 2003; Uhl *et al.*, 2004, 2008; Šimůnek & Martínek, 2009) and China (Wang & Chen, 2001). In contrast, for large parts of Gondwana the record of Permian macroscopic fossil charcoal is still scarce to non-existent.

Despite numerous studies on Gondwana inertinites of assumed problematic origin (cf. Scott, 2000), the first unequivocal record of charcoal as a direct palaeobotanical evidence of palaeowildfires on Gondwana was published by Glasspool (2000) based on material from the Late Permian of the Sydney Basin, Australia. Subsequent studies also confirmed the presence of charcoal in Permian sediments from South Africa (Glasspool, 2003), Jordan (Uhl *et al.*, 2007) and Brazil (Jasper *et al.*, 2008, 2011a, b). Mishra *et al.* (1990) and Navale and Saxena (1989) described high inertinite levels in Permian coals of India which are probably the result of fires. However, remains of macroscopic fossil charcoal have not been reported so far.

Jasper *et al.* (2008) demonstrated that the charcoalified remains discovered in Early Permian sediments of the Quitéria outcrop were related to basic types of gymnosperm wood and fragments of lycopsids. These authors also inferred that potential sources of ignition for the wildfires in the studied area could have been the volcanic activities in nearby areas.

In the Faxinal Coalfield, charcoal remains have also been discovered in a tonstein layer, originating from volcanic ashfall tuffs, interbedded within a coal layer (Jasper *et al.*, 2011a, b).




Fig. 1—Stratigraphy of Permian Gondwana in Damodar Basin (after Mukhopadhyay *et al.*, 2010).




Fig. 2—Geological map of a part of the Raniganj Coalfield in the Damodar Basin, West Bengal, India showing the study area and sample location (modified after Murthy *et al.*, 2010).

Considering that the charcoal remains are preserved in this layer, the connection with the volcanic activity seems very likely.

In this context additional analyses are necessary to evaluate the palaeo-wildfire coverage on Gondwana during the Permian, the discovery of the first macroscopic fossil charcoal remains from India is very significant. In this way, this study helps to fill one of the remaining geographical gaps for the Late Palaeozoic Gondwana, describing material coming from the Upper Permian of Raniganj Formation (Lopingian), Raniganj Coalfield, Damodar Basin.

GEOLOGICAL AND STRATIGRAPHIC SETTING

Stratigraphically the Permian Gondwana of Damodar Basin includes Talchir, Karharbari, Barakar, Barren Measures and Raniganj formations (Fig. 1). Lithologically the Raniganj Formation is characterized by the association of coal, organic rich carbonaceous shale, siltstone and sandstone. However, the coal seams are usually thinner and less frequent. The sandstones vary from fine to coarse grained and the fine grained sandstones are usually micaceous. Predominantly, the sandstones are arkose to subarkose and occasionally calcareous. The carbonaceous shales are black in colour while

the siltstones are grey to dark grey in colour. The maximum thickness of the Raniganj Formation in the Raniganj Coalfield is about 900 m and there are 10 coal seams (R – I to R – X) in this formation. The Raniganj Formation overlies the sandstones of the Barren Measures Formation and underlies the claystones of the Panchet Formation.

According to Murthy *et al.* (2010), the Raniganj Coalfield (23°03'–23°51' N and 86°42'–87°28' E), explored as an open pit mine, corresponds to the oriental segment of the basin, represented by an elliptical deposit with approximately 3,000 km². The abundance of organic rich shales in the Raniganj Formation shows levels of palaeosoils and stems in growing position. Hence, Ghosh (2002) inferred a hot humid palaeoenvironment, which is confirmed by the high percentage of *Densipollenites* pollen grains (Tewari & Tripathi, 1992). According to Mukhopadhyay *et al.* (2010) the coaly sediments were deposited during a regressive phase and represent an anoxic floodplain enabling the deposition of important peat areas which have produced economically viable coal layers at the Raniganj, Jharia and Singrauli areas.

Associated with the coal layers occurs a typical *Glossoptris* flora which reached its climax during the deposition of the Raniganj Formation and which was the main organic matter source for the Permian coals in this area. The




Fig. 3—Litholog of the studied section showing the location of yielding samples.




Fig. 4—Field photograph showing the location of study area.

charcoal remains studied here were collected in organic rich shale from the top of the Raniganj Formation, associated with well-preserved *Glossopteris* leaves, indicating that palaeo-wildfires affected the Upper Permian environments in this region.

MATERIAL AND METHODS

During field work of the first Indo-Brazilian Workshop (in November 2010), samples with well-preserved *Glossopteris* leaves were collected from the Open Cast Mine of Sonepur Bazaar area of the Raniganj Coalfield (Fig. 2). The material for the present study comes from the shale associated with the Seam-VI of the Raniganj Formation (Figs 3 & 4). Some of these samples also include fragments of macroscopic fossil charcoal (stored in the Setor de Botânica e Paleobotânica do MCN/UNIVATES under acronym PbU, specimens 815 to 820).

The material which shows characteristics of macroscopic charcoal (black colour and streak, silky lustre, *sensu* Jones & Chaloner, 1991; Scott, 1989, 2000, 2010), was mechanically extracted from the collected hand specimens with the aid of

preparation needles and tweezers under a binocular stereomicroscope in the laboratory. Due to the very fragile nature of these specimens, they could not be cleaned with water or any acids to remove adhering mineral remains or non-charred humic material.

The fragments were mounted on standard stubs with LeitC (Plano), and subsequently examined with the aid of a JEOL JSM 6490 LV Scanning Electron Microscope (SEM) at the Senckenberg Forschungsinstitut und Naturmuseum Frankfurt.

RESULTS

General characters of the charcoal remains:

The charcoal fragments measure between $0.3 \times 0.1 \times 0.2$ and $0.6 \times 0.2 \times 0.4$ cm and have no abraded edges. The SEM analyses, showing well-preserved anatomical details and homogenised cell-walls (Pl. 1.1), allowed to confirm that the samples were charred (*sensu* Jones & Chaloner, 1991; Scott, 1989, 2000, 2010).

However, the woody tissues, which were probably compressed during diagenesis, have been shattered into more or less small pieces (Pl. 1.2, 3). Due to this excessive mechanical damage it was not possible to observe complete tissue characteristics, thus hampering a specific taxonomic/systematic affiliation.

Although the anatomical structures of charred soft wood are generally well preserved, charcoalification can induce changes of certain anatomical features, like the appearance of pits and especially cross-field pits (Jones & Chaloner, 1991; Gerards *et al.*, 2007). In the studied material, closing walls of pits, which are easily subjected to such changes, could not be observed. The description of the pitting is based on their apertures.

Anatomical characters of the charcoal remains:

Pycnoxylic secondary wood, in radial view tracheids is $25-30 \mu\text{m}$ in width. They exhibit uniseriate (Pl. 1.4-1.6) or biseriate bordered pitting (Pl. 1.7). Pits are circular to oval ($3-7 \mu\text{m}$ in diameter) with circular to oval apertures (Pl. 1.5). When biseriate they are arranged sub-oppositely to alternate (Pl. 1.7). Rays are rare and composed of parenchymatous cells, $14-19 \mu\text{m}$ in width and $18-31 \mu\text{m}$ in height (Pl. 1.8). Rays are uniseriate and 5-7 cells high. Cross-field pits, leaf traces or growth rings are not visible.

DISCUSSION

The fragmentary nature of the charcoal remains, does not allow for establishing specific taxonomical affinities. The bordered pitting and presence of uniseriate rays only permitted

to infer a gymnospermous affinity of the material. In addition, the absence of abraded edges allows the inference of an autochthonous/parautochthonous origin for the material without any considerable transport prior to burial.

Considering the association of the remains to *Glossopteris* leaves and previous studies about the Gondwanan macroscopic charcoal (Jasper *et al.* 2008; 2011a, b), the correlation with gymnosperms can be considered as an unsurprising result. This follows the tendency observed in other Permian coal bearing strata of the Gondwana, in which the fire reached the mire and the surrounding vegetation, dominated by the so called *Glossopteris* flora.

On the other hand, the additional confirmation of the occurrence of palaeo-wildfires associated with coal bearing strata of the Permian Gondwana helps to refine scenarios for such events during the period. So far such scenarios are mainly based on data from mesoscopic charcoal, from the Sydney Basin, Australia (Glasspool, 2000), and from macroscopic charcoal, from the Paraná Basin, Brazil (Jasper *et al.*, 2011b).

Despite the high concentration of inertinites in the Gondwana coals some authors prefer different explanations for the origin of these macerals in the Late Palaeozoic Gondwana (White, 1925; Jurasky, 1929; Schopf, 1975; Taylor *et al.*, 1988). However, an increasing number of localities containing macroscopic charcoal remains, like the material presented here, have recently been found in different associations all over Gondwana, which are linked to inertinite bearing coal layers or in clastic levels directly above such coals.

Recent studies (Scott & Glasspool, 2006, 2007; Scott, 2010; Hudspith *et al.*, 2011) reinforced that inertinites observed in Permian coals around the world, even in high abundance, were quite likely generated by palaeo-wildfires. The abundance of structured inertinites in Permian coals from the Sattupalli Coalfield, Godavari Graben, India, has recently been connected to such fire events (Singh *et al.*, 2011).

The autochthonous/parautochthonous origin of the charcoals studied here is a strong evidence to the fact that palaeo-wildfires in Gondwana frequently reached coal forming environments during the Permian.

CONCLUSIONS

From the evidence presented here, it is possible to draw the following conclusions:

1. The charcoalified plant remains from the Raniganj Formation testify to the occurrence of wildfire in the Damodar Basin, India during the Permian.
2. The preserved charred plant fragments support the hypothesis of autochthonous/parautochthonous origin for the material.
3. The charred wood remains are related to gymnosperms.




PLATE 1

4. The frequent occurrence of palaeo-wildfires in Gondwana during the Late Palaeozoic is reinforced.

Acknowledgements—We thank Claudia Franz, Senckenberg Forschungsinstitut and Naturmuseum Frankfurt, for technical assistance with SEM facilities. A. Jasper acknowledges financial support by FAPERGS (Project 11/1307-7) and CNPq (Projects 301671/2009-5, 401771/2010-5 and 474857/2011-5). M. Guerra-Sommer acknowledges financial support by CNPq. M.E.C. Bernardes-de-Oliveira and R. Tewari are thankful to CNPq (Brazil) and to the Department of Science and Technology (India) for the financial grants to the Indo-Brazil S & T Programme of International Cooperation (DST/INT/India/RPO-24/2007 and Brazil/CNPq Project 490829/2007). M.E.C. Bernardes-de-Oliveira is thankful to CNPq (Project 304265/2009-8); M.I. Secchi acknowledges financial grant of CAPES (master scholarship 23038.009744/2010-48), FAPERGS and CNPq. R. Tewari and A.K. Ghosh are also thankful to the Director, BSIP for his kind permission to publish this collaborative work. The authors also thank Dr Madhav Kumar, one of the referees of *The Palaeobotanist* for the relevant contributions to the manuscript.

REFERENCES

Bowman DMJS, Balch JK, Artaxo P, Bond WJ, Carlson JM, Cochrane MA, D'Antonio CM, Defries RS, Doyle JC, Harrison SP, Johnston FH, Keeley JE, Krawchuk MA, Kull CA, Marston JB, Moritz MA, Prentice IC, Roos CL, Scott AC, Swetnam TW, Van Der Werf GR & Pyne SJ 2009. Fire in the Earth System. *Science* 324: 481-484.

DiMichele WA, Hook RW, Nelson WJ & Chaney DS 2004. An unusual Middle Permian flora from the Blaine Formation (Pease River Group: Leonardian-Guadalupian Series) of King County, West Texas. *Journal of Paleontology* 78: 765-782.

Flannigan MD, Krawchuk MA, Groot WJ de, Wotton BM & Gowman LM 2009. Implications of changing climate for global wildland fire. *International Journal of Wildland Fire* 18: 483-507.

Gastaldo RA, DiMichele WA & Pfefferkorn HW 1996. Out of the Icethouse into the Greenhouse: a Late Paleozoic Analog for modern global vegetational change. *GSA Today* 6: 2-7.

Gerards T, Damblon F, Wauthoz B & Gerrienne P 2007. Comparison of cross-field pitting in fresh, dried and charcoalized softwoods. *IAWA Journal* 28: 49-60.

Ghosh SC 2002. The Raniganj Coal Basin: an example of an Indian Gondwana Rift. *Sedimentary Geology* 147: 155-176.

Glasspool IJ 2000. A major fire event recorded in the mesofossils and petrology of the Late Permian, Lower Whybrow coal seam, Sydney Basin, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* 164: 373-396.

Glasspool IJ 2003. Hypautochthonous-allochthonous coal deposition in the Permian, South African, Witbank Basin No. 2 seam; a combined approach using sedimentology, coal petrology and palaeontology. *International Journal of Geology* 53: 81-135.

Hudspith V, Scott AC, Collinson ME, Pronina N & Beeley T 2011. Evaluating the extent to which wildfire history can be interpreted from inertinite distribution in coal pillars: an example from the Late Permian, Kuznetsk Basin, Russia. *International Journal of Coal Geology*. <http://dx.doi.org/10.1016/j.coal.2011.07.009>.

Jasper A, Uhl D, Guerra-Sommer M & Mosbrugger V 2008. Palaeobotanical evidence of wildfires in the Late Palaeozoic of South America – Early Permian, Rio Bonito Formation, Paraná Basin, Rio Grande do Sul, Brazil. *Journal of South American Earth Science* 26: 435-444.

Jasper A, Uhl D, Guerra-Sommer M, Hamad AMBA & Machado NTG 2011a. Charcoal remains from a tonstein layer in the Faxinal Coalfield, Lower Permian, southern Paraná Basin, Brazil. *Anais da Academia Brasileira de Ciências* 83: 471-48.

Jasper A, Uhl D, Guerra-Sommer M, Bernardes-de-Oliveira MEC & Machado NTG 2011b. Upper Paleozoic charcoal remains from South America: multiple evidences of fire events in the coal bearing strata of the Paraná Basin, Brazil. *Palaeogeography, Palaeoclimatology, Palaeoecology* 306: 205-218.

Jones TP & Chaloner WG 1991. Fossil charcoal, its recognition and palaeoatmospheric significance. *Palaeogeography, Palaeoclimatology, Palaeoecology* 97: 39-50.

Jurasky KA 1929. Neue Untersuchungen und Gedanken zur Entstehung fossiler Holzkohle. In: Stutzer O (Editor)—*Fusit-Vorkommen, Entstehung und praktische Bedeutung der Faserkohle*. Schriften aus dem Gebiet der Brennstoff-Geologie, Verlag von Ferdinand Enke, Stuttgart, Heft, 2: 23-41.

Misra HK, Chandra TK & Verma RP 1990. Petrology of some Permian coals of India. *International Journal of Coal Geology* 16: 47-71.

Mukhopadhyay G, Mukhopadhyay SK, Roychowdhury M & Parui PK 2010. Stratigraphic correlation between different Gondwana basins of India. *Journal of Geological Society of India* 76: 251-266.

Murthy S, Chakraborti B & Roy MD 2010. Palynodating of subsurface sediments, Raniganj Coalfield, Damodar Basin, West Bengal. *Journal of Earth System Science* 119: 701-710.

Navale GKB & Saxena R 1989. An appraisal of coal petrographic facies in Lower Gondwana (Permian) coal seams of India. *International Journal of Coal Geology* 12: 553-558.

Noll R, Uhl D & Lausberg S 2003. Brandstrukturen an Kieselhölzern der Donnersberg Formation (Oberes Rotliegen, Unterperm) des Saar-Nahe Beckens (SW-Deutschland). *Veröffentlichungen Museum für Naturkunde Chemnitz* 26: 63-72.

Rößler R (Ed.) 2001. Der Versteinerte Wald von Chemnitz. *Museum für Naturkunde Chemnitz*. 252 pp.

Schoop JM 1975. Modes of fossil preservation. *Reviews of Palaeobotany & Palynology* 20: 27-53.

Scott AC 1989. Observations on the nature and origin of fusain. *International Journal of Coal Geology* 12: 443-475.



PLATE 1

SEM images of the charcoal samples of the Raniganj Coalfield.

1. Tracheids presenting homogenized cell walls (arrow).
2. Tracheids shattered into more or less small pieces.
3. Detail of 2 exhibiting shattered characteristic of the charcoal remains.
4. Tracheids of Raniganj Coalfield wood remains exhibiting uniserial bordered pitting, with presence of circular to oval pits tracheids.
5. Tracheids of Raniganj Coalfield wood remains exhibiting uniserial bordered pitting.
6. Tracheids of Raniganj Coalfield wood remains exhibiting uniserial bordered pitting.
7. Tracheids of Raniganj Coalfield wood remains with biserial bordered pitting sub-oppositely arranged.
8. Detail of Raniganj Coalfield wood remains presenting uniserial ray (arrow).

Scott AC 2000. The pre-quaternary history of fire. *Palaeogeography, Palaeoclimatology, Palaeoecology* 164: 281-329.

Scott AC 2010. Charcoal recognition, taphonomy and uses in palaeoenvironmental analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 291: 11-39.

Scott AC & Glasspool I 2006. The diversification of Paleozoic fire systems and fluctuation in atmospheric oxygen concentration. *PNAS* 103: 10861-10865.

Scott AC & Glasspool I 2007. Observations and experiments on the origin and formation of inertinite group macerals. *International Journal of Coal Geology* 70: 53-66.

Šimůnek Z & Martinek K 2009. A study of Late Carboniferous and Early Permian plant assemblages from the Boskovice Basin, Czech Republic. *Review of Palaeobotany & Palynology* 155: 275-307.

Singh PK, Prachiti PK, Kalpana MS, Manikyamba C, Singh MP, Lakshminarayana G, Singh AK & Naik AS 2011. Petrographic characteristics and carbon isotopic composition of Permian coal: Implications on depositional environment of Sattupalli Coalfield, Godavari Valley, India. *International Journal of Coal Geology*. <http://dx.doi.org/10.1016/j.coal.2011.10.002>.

Taylor GH, Liu S & Diesel CFK 1988. The cold climate origin of inertinite-rich Gondwana coals. *International Journal of Coal Geology* 11: 1-22.

Tiwari RS & Tripathi A 1992. Marker assemblage zones of spore and pollen species through Gondwana Palaeozoic and Mesozoic sequence in India. *Palaeobotanist* 40: 194-236.

Uhl D, Abu Hamad AMB, Kerp H & Bandel K 2007. Evidence for palaeo-wildfire in the Late Permian palaeotropics – charcoalified wood from the Um Irna Formation of Jordan. *Review of Palaeobotany & Palynology* 144: 221-230.

Uhl D, Jasper A, Abu Hamad AMB & Montenari M 2008. Permian and Triassic wildfires and atmospheric oxygen levels. *Proceedings from the WSEAS International Conference on Environmental and Geological Science and Engineering*, 1, pp. 179-188.

Uhl D & Kerp H 2003. Wildfires in the Late Palaeozoic of Central Europe – The Zechstein (Upper Permian) of NW Hesse (Germany). *Palaeogeography, Palaeoclimatology, Palaeoecology* 199: 1-15.

Uhl D, Lausberg S, Noll R & Stäpf KRG 2004. Wildfires in the Late Palaeozoic of Central Europe – an overview of the Rotliegend (Upper Carboniferous-Lower Permian) of the Saar-Nahe Basin (SW-Germany). *Palaeogeography, Palaeoclimatology, Palaeoecology* 207: 23-35.

Wang Z & Chen A 2001. Traces of arborescent lycopods and dieback of the forest vegetation in relation to terminal Permian mass extinction in North China. *Review of Palaeobotany & Palynology* 117: 217-243.

Westerling AL, Turner MG, Smithwick EAH, Romme WH & Ryan MG 2011. Continued warming could transform Greater Yellowstone fire regimes by mid-21st century. <http://dx.doi.org/10.1073/pnas.1110199108>.

White D 1925. Environmental conditions of deposition of coal. *American Institute of Mining, Metallurgy and Engineering Transaction* 71: 3-34.