

Molecular detection of *Rickettsia* genus in chigger mites (Trombidiformes: Trombiculidae) collected on small mammals in southeastern Brazilian

Detecção molecular do gênero *Rickettsia* em ácaros trombiculídeos (Trombidiformes: Trombiculidae) coletados em pequenos mamíferos do sudeste brasileiro

Fernando de Castro Jacinavicius^{1,2} ; Ricardo Bassini-Silva² ; Sebastián Muñoz-Leal² ; Cal Welbourn³ ; Ronald Ochoa⁴ ; Marcelo Bahia Labruna² ; Darci Moraes Barros-Battesti^{2,5*} 

¹ Laboratório Especial de Coleções Zoológicas, Instituto Butantan, São Paulo, SP, Brasil

² Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia – FMVZ, Universidade de São Paulo – USP, São Paulo, SP, Brasil

³ Florida State Collection of Arthropods, Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

⁴ Systematic Entomology Laboratory, United States Department of Agriculture, Agricultural Research Service, Beltsville, MD, USA

⁵ Departamento de Patologia Veterinária, Faculdade de Ciências Agrárias e Veterinárias – FCAV, Universidade Estadual Paulista “Júlio de Mesquita Filho” – UNESP, Jaboticabal, SP, Brasil

Received April 23, 2019

Accepted August 08, 2019

Abstract

Chiggers are ectoparasites of vertebrates and may cause trombiculiasis or transmit pathogens to their hosts. Specimens collected from rodents and marsupials were morphologically identified as *Herpetacarus hertigi*, *Eutrombicula tinami*, *Kymoceta* sp., *Quadrasetta brasiliensis*, *Quadrasetta falconensis*, *Quadrasetta flochi*, *Quadrasetta mackenziei*, *Quadrasetta pazca*, *Quadrasetta trapezoides*, *Quadrasetta* sp., *Serratacarus* sp., and *Trombewingia bakeri*. These mites were submitted individually to molecular analyses for the detection of bacteria of the genus *Coxiella*, *Hepatozoon* and *Rickettsia*. Samples were positive to *Rickettsia* only. Obtained sequences for the *gltA* (350 pb) and *ompA* (488 pb) genes were identical to “*Candidatus Rickettsia colombianensi*”, a species previously detected in ticks. In addition, molecular identification of mites based on 18S rDNA sequences are provided for *H. hertigi*, *Kymoceta* sp., *Q. brasiliensis*, *Q. pazca*, *Q. trapezoides*, *Quadrasetta* sp., and *T. bakeri* for the first time. This is the first report of the detection of a *Rickettsia* sp. in chigger mites collected on rodents in Brazil.

Keywords: Chiggers, “*Candidatus Rickettsia colombianensi*”, ectoparasites, Rodentia, Didelphimorphia, Brazil.

Resumo

Os trombiculídeos são ectoparasitas de vertebrados e podem causar trombiculíase ou transmitir patógenos ao hospedeiro. Exemplares coletados em roedores e marsupiais foram identificados morfologicamente como *Herpetacarus hertigi*, *Eutrombicula tinami*, *Kymoceta* sp., *Quadrasetta brasiliensis*, *Quadrasetta falconensis*, *Quadrasetta flochi*, *Quadrasetta mackenziei*, *Quadrasetta pazca*, *Quadrasetta trapezoides*, *Quadrasetta* sp., *Serratacarus* sp. e *Trombewingia bakeri*. Estes ácaros foram submetidos individualmente à análise molecular para detecção de bactérias dos gêneros *Coxiella*, *Hepatozoon* e *Rickettsia*. Amostras foram positivas somente para *Rickettsia*. Sequências obtidas para os genes *gltA* (350 pb) e *ompA* (488 pb) foram idênticas à “*Candidatus Rickettsia colombianensi*”, uma espécie anteriormente detectada em carrapatos. Além disso, foram fornecidas sequências de DNA 18S para identificação molecular de *H. hertigi*, *Kymoceta* sp., *Q. brasiliensis*, *Q. pazca*, *Q. trapezoides*, *Quadrasetta* sp. e *T. bakeri*. Este é o primeiro registro da detecção de *Rickettsia* em ácaros trombiculídeos coletados em roedores do Brasil.

Palavras-chave: Trombiculídeos, “*Candidatus Rickettsia colombianensi*”, ectoparasitas, Rodentia, Didelphimorphia, Brasil.

*Corresponding author: Darci Moraes Barros-Battesti. Departamento de Patologia Veterinária, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista “Júlio de Mesquita Filho” – UNESP, Via de Acesso Professor Paulo Donato Castelan, s/n, Vila Industrial, CEP 14884-900, Jaboticabal, SP, Brasil. e-mail: barros.battesti@gmail.com



Introduction

Larval stages of trombiculid mites (Trombidiformes: Trombiculidae), also known as chiggers, typically parasitize terrestrial vertebrates and during feeding they inject digestive enzymes into the skin of their hosts. If infected, chiggers may transmit pathogens and/or cause severe skin reactions to the host, a disease commonly known as trombiculiasis (SANTIBÁÑEZ et al., 2015). In the Asia-Pacific region, chiggers are known as tsutsugamushi (“tsutsuga” = disease, “mushi” = bug), and are recognized vectors of *Orientia tsutsugamushi*, the agent of an acute febrile disease in humans, commonly known as Scrub typhus (RAPMUND et al., 1969; TAKAHASHI et al., 2004; KELLY et al., 2009; PHASOMKUSOLSIL et al., 2009).

Pathogenic microorganisms other than *O. tsutsugamushi* also can be found in trombiculid mites. Frank (1977) detected the presence of the protozoan *Hepatozoon erhardovae* in chiggers of the species *Hirsutiella zachvatkini* (Schluger 1948) in Austria, and *Coxiella burnetii*, the causative agent of Q-fever, was detected once in African chiggers (DANIEL, 1961; KEPKA, 1965). Furthermore, Blanc et al. (1952) successfully infected *Neotrombicula autumnalis* (Shaw 1790) with *C. burnetii* under laboratory conditions, and Le Gac et al. (1953) reported a Q-fever case in a laboratory technician who was handling chiggers from an endemic area in Equatorial Africa. On the other hand, bacteria of genus *Rickettsia* were detected in chiggers from Ukraine (VYSOTSKAYA & SCHLUGER, 1953), Korea (CHOI et al., 2007) and China (HUANG et al., 2017). In particular, *Rickettsia monacensis* and *Rickettsia helvetica* were detected in mites from Slovakia (MIŤKOVÁ et al., 2015).

In Brazil, Fonseca (1932) emphasized the importance of mites as potential vectors in the epidemiological cycle of rickettsial diseases. Recently, Bassini-Silva et al. (2018a) detected a *Rickettsia felis*-like agent in the trombiculid *Blankaartia sinnamaryi* (Floch

& Fauran, 1956) parasitizing birds, which corresponded to the first report of a rickettsial agent in Brazilian mites.

Here, we are contributing with the knowledge on molecular studies of chiggers parasitizing small mammals. In addition, we report the molecular detection of a *Rickettsia* sp. in three species of chiggers collected from three hosts.

Materials and Methods

Molecular analyses

Chiggers collected in São Paulo state, preserved in ethyl alcohol PA 100%, stored in a -80°C freezer, housed in the Acari Collection of the Instituto Butantan (IBSP) of São Paulo City, Brazil, were individually submitted to DNA extraction using the Guanidine Isothiocyanate (GT) lysis protocol following Chomczynski (1993). Each mite was placed into a plastic microtube and punctured in the idiosomal region with a sterile needle (1.20 * 40 - 18G). After DNA extraction, exoskeletons of the specimens were recovered and slide-mounted using Hoyer’s medium in order to perform morphological identifications according to Walter & Krantz (2009).

Conventional PCRs targeting a partial fragment of the mite 18S ribosomal gene and a section of the mite mitochondrial cytochrome oxidase I (COI) gene were initially performed using thermal conditions and employed primers listed in Table 1. All reactions included a positive (DNA extracted from *B. sinnamaryi*) and a negative (DNA-free ultrapure water) control. As 18S rRNA and COI gene-PCRs were used as endogenous controls, negative samples were excluded from further analyses.

Positive samples were then screened for agents in the genera *Coxiella*, *Hepatozoon* and *Rickettsia* through conventional and heminested PCR protocols. All the reactions included a negative

Table 1. Oligonucleotide primers used for the amplification of the genes 18S rRNA and COI (endogenous controls), *gltA* and *ompA* (*Rickettsia*), 16S rRNA (*Coxiella*) and 18S rRNA (*Hepatozoon*).

Organisms	Gene	Sequence 5'-3'	Size (pb)	Reference
<i>Rickettsia</i> spp.	<i>gltA</i>	CS-78 (GCAAGTATCGGTGAGGATGTAAT) CS-323 (GCTTCCTTAAAATTCAATAATCAGGAT)	401	Labruna et al. (2004)
<i>Rickettsia</i> spp. (SPF)	<i>ompA</i> (1st round)	Rr190.70 (ATGGCGAATATTTCTCCAAAA) Rr190.701R2 (GTTCCGTTAATGGCAGCATCT)	632	Regnery et al. (1991) and Roux et al. (1996)
<i>Rickettsia</i> spp. (SPF)	<i>ompA</i> (hemi-nested)	Rr190.70 (ATGGCGAATATTTCTCCAAAA) Rr190.602 (AGTGCAGCATTCGCTCCCCCT)	532	Regnery et al. (1991)
<i>Coxiella</i> spp.	16S	16SrRNA F (GGGGAAGAAAGTCTCAAGGGTAATACCTT) 16SrRNA R (TGCATCGAATTAAACCACATGCTCCACCGC)	532	Almeida et al. (2012)
<i>Hepatozoon</i> spp.	18S	HEP142-169-F (GCTTGAAACACTCTARTTTTCTCAAAG) HEP743-718-R (ACAATAAAGTAAAAACAYTTCAAAG)	574	Almeida et al. (2013)
Mite	COI	COI 772 (TGATTTTTTGGTCACCCAGAAG) COI 773 (TACAGCTCCTATAGATAAAAC)	408	Navajas et al. (1994) adapted by Soller et al. (2001)
Mite	COI	bcdF01 CATTTCCHACTAAYCATAARGATATTGG bcdR04 TATAAACYTCDGGATGNCCAAAAA	560-680	Dabert et al. (2008, 2010)
Mite	COI	LCO1490 GGTCACAAATCATAAAGATATTGG LCO2198 TAAACTTCAGGGTGACCAAAAAATCA	710	Folmer et al. (1994)
Mite	18S	18S-1F (ATATTGGAGGGCAAGTCTGG) 18S-1R (TGGCATCGTTTATGGTTAG)	500	Otto & Wilson (2001)

SPF: Spotted Fever Group rickettsiae.

control (DNA-free Milli-Q water) and an appropriate positive control, which consisted of DNA of *Coxiella* sp., extracted from infected *Ornithodoros* ticks; *Hepatozoon canis*, extracted from blood of an infected dog; and *Rickettsia vini* strain Breclav, extracted from an infected Vero cell culture. Reactions yielding amplicons of the expected size were treated with ExoSAP-IT (USB Corporation®, OH) following the manufacturer instructions (3 µl of ExoSAP with 7.5 µl of the amplified DNA). Sanger sequencing of the amplicons was performed at the “Centro de pesquisa sobre Genoma Humano e Células Tronco do Instituto de Biociências da USP”. Obtained sequences were assembled and trimmed with Geneious R9 (KEARSE et al., 2012), and then submitted to BLASTn analyses (NCBI, 2019) in order to infer closest similarities with other homologous sequences (ALTSCHUL et al., 1990).

Morphological tools

The slide-mounted voucher specimens were identified to the genus level using the key by Brennan & Goff (1977), and to the species level based on the original descriptions cited in the Brazilian checklist of chiggers (JACINAVICIUS et al., 2018b). Additional comparisons were made with type series and other specimens deposited in the mite collection of the United State National Museum (USNM), currently housed at the Systematic Entomology Laboratory, Beltsville, Maryland, USA (BARC-USDA-ARS).

Results

The 317 chiggers examined at the IBSP were collected from nine cricetid rodents (*Akodon montensis*, *Akodon* sp., *Delomys sublineatus*, *Euryoryzomys russatus*, *Hylaeamys megacephalus*, *Necomys lasiurus*, *Necomys squamipes*, *Oligoryzomys* sp. and *Thaptomys nigrita*), one echimyid rodent (*Thrichomys fosteri*), and four marsupials (*Didelphis aurita*, *Gracilinanus agilis*, *Monodelphis americana*, *Monodelphis domestica* and *Thylamys macrurus*). After DNA extraction, mites were slide mounted and morphologically identified generating the following vouchers: *Herpetacarus hertigi* (Brennan, 1970) (22), *Eutrombicula tinami* (Oudemans, 1910) (14), *Kymocta* sp. (7), *Quadrasetta brasiliensis* Goff and Gettinger, 1989 (175), *Quadrasetta falconensis* Goff and Brennan, 1977 (6), *Quadrasetta flochi* (Brennan and Jones, 1960) (5), *Quadrasetta mackenziei* (Yunker and Brennan, 1964) (4), *Quadrasetta pazca* (Brennan and Jones, 1964) (54), *Quadrasetta trapezoides* (Brennan and Jones, 1964) (11), *Quadrasetta* sp. (9), *Serratacarus* sp. (1) and *Trombewingia bakeri* (Fonseca, 1955) (9).

A total of 20 samples yielded expected size amplicons for the mite 18S rRNA gene. Obtained sequences corresponded to a unique haplotype for each of the following seven chigger species. The GenBank accession numbers are: one *H. hertigi* (MG817637), two *Kymocta* sp. (MG817642), four *Q. brasiliensis* (MG817643), two *Q. pazca* (MG817644), seven *Q. trapezoides* (MG817645), one *Quadrasetta* sp. (KY934461), and three *T. bakeri* (MG817646). One amplified sample of *Q. pazca* did not generate a good quality sequence and was not deposited in GenBank. Attempts to amplify fragments of the COI gene from all samples were unsuccessful. After BLASTn analyses, consensus sequences of *H. hertigi*, *Kymocta*

sp., *Q. brasiliensis*, *Q. pazca*, *Q. trapezoides*, *Quadrasetta* sp., and *T. bakeri* were 98.63% (431/437-pb), 99.51% (408/410-pb), 99.50% (397/399-pb), 99.75% (404/405-pb), 98.77% (400/405-pb), 98.39% (428/435-pb), and 99.50% (397/399-pb) identical with a homologous sequence of *Eutrombicula splendens* (Ewing, 1913) (KY922159).

All 20 samples positive for 18S rDNA PCR assays were tested for *Coxiella*, *Hepatozoon* and *Rickettsia*. A total of 13 samples yielded rickettsial *gltA* amplicons, but only six were successfully sequenced. In addition, one sample also yielded rickettsial *ompA* amplicons. PCR targeting 16S rDNA gene of *Coxiella* sp. and 18S rDNA gene of *Hepatozoon* sp. were negative. Sequences of *Rickettsia* were submitted to BLASTn analysis. Six identical *gltA* sequences were 100% (350/350 bp) identical to “*Candidatus Rickettsia colombianensi*” (MG970682). These sequences came from one larva of *H. hertigi*, three larvae of *Q. trapezoides* and two of *T. bakeri*. Besides that, one of the successfully-sequenced *ompA* samples was 100% (488/488 bp) identical to “*Ca. Rickettsia colombianensi*” (MG970683) as well. Sequences of *Rickettsia* generated in the current study were deposited in GenBank under the accession numbers MG906649, MG906650, MG906652, MG906653, MG906654, MG906656.

Overall, one species of *H. hertigi* collected in Fontes do Ipiranga State Park (PEFI), São Paulo municipality, parasitizing an *Oligoryzomys* sp.; three specimens of *Q. trapezoides* from Morro Grande, Cotia municipality, collected on *Necomys squamipes*; and two specimens of *T. bakeri* collected in Campos do Jordão municipality, on *Akodon montensis*, were positive for “*Ca. Rickettsia colombianensi*”.

Discussion

The successful amplification rates for the mite 18S rRNA gene upon individual chiggers, was 13.63% (3/22) for *H. hertigi*, 28.57% (2/7) for *Kymocta* sp., 2.28% (4/175) for *Q. brasiliensis*, 5.55% (3/54) for *Q. pazca*, 63.64% (1/11) for *Q. trapezoides*, 11.11% (1/9) for *Quadrasetta* sp., and 33.33% (3/9) for *T. bakeri*. Amplification of this same locus was unsuccessful for *E. tinami*, *Q. falconensis*, *Q. flochi*, *Q. mackenziei* and *Serratacarus* sp. We were successful in amplifying only 7.26% (20/317) of all individually tested species, while Bassini-Silva et al. (2018a) obtained a 72.5% (29/40) of success in amplifying the same gene for *B. sinnamaryi*. Park et al. (2015) successfully amplified the 18S rRNA gene in 50% (38/76) for *Helenicula miyagawai* (Sasa, Kumada and Miura, 1951) mites, and in 87.5% (7/8) for *Leptotrombidium scutellare* (Nagayo, Miyagawa, Mitamura, Tamiya and Tenjin, 1921).

Of the 73 chiggers species reported from Brazil, 18S rDNA sequences were previously available for only three species: *B. sinnamaryi* (MG783391) (BASSINI-SILVA et al., 2018a), *Eutrombicula daemonei* Bassini-Silva and Jacinavicius, 2018 (MG707783, MG70778) (BASSINI-SILVA et al., 2018b), and *Q. brasiliensis* (MF113413, MF113412, KY934462, KY934463, KY934464) (JACINAVICIUS et al., 2018a). In the present study, we provide new 18S rDNA sequences for *H. hertigi*, *Kymocta* sp., *Q. pazca*, *Q. trapezoides*, and *T. bakeri*. According to Hillis & Dixon (1991) and Cruickshank (2002), the 18S rRNA gene

has a slow rate of evolution and it is suitable to infer family and subfamily level phylogenies. As the availability of sequences of this locus for chiggers from South America is still scarce, the creation of a South American chigger gene bank may afford the inclusion of these taxa in future phylogenetic studies dealing with the relationships between mites in higher groupings.

The mitochondrial COI gene has a faster rate of sequence divergence if compared with nuclear ribosomal genes (OTTO & WILSON, 2001). Kampen et al. (2004), Moniuszko et al. (2015) and Kumlert et al. (2018) successfully amplified this gene for some chigger species of the genera *Ascoschoengastia*, *Blankaartia*, *Hirsutella*, *Leptotrombidium*, *Neotrombicula*, *Schoengastia*, *Schoutedenichia*, and *Walchia*, from Europe and Asia. Although in the current study we employed this same pairs of primers, none of our samples yielded amplicons, which is in the line with previous studies in Brazil (BASSINI-SILVA et al., 2018a; JACINAVICIUS et al., 2018a). We highlight then the need for testing new sets or designing new primers that will successfully amplify this gene for South American chiggers.

Rickettsial DNA identical to two "*Ca. Rickettsia colombianensi*" genes was detected in three chigger species for the first time, all from the state of São Paulo. "*Ca. Rickettsia colombianensi*" was originally characterized from the tick *Amblyomma dissimile* Koch, 1844 in Colombia (MIRANDA et al., 2012). In Brazil, this species of *Rickettsia* sp. was recently detected in *A. dissimile* from Amazon biome of Amapá state (LUZ et al., 2018), and the current finding extends its occurrence to further Acari representatives, in particular, to trombiculid mites in the Atlantic rainforest biome of the São Paulo state. "*Candidatus R. colombianensi*" belongs to the spotted fever group of rickettsiae, and its isolation in Vero cells has demonstrated a marked cytopathic effect (MIRANDA et al., 2019). Although infection in cattle and iguanas has been suggested in Colombia (MIRANDA et al., 2012), the role of "*Ca. Rickettsia colombianensi*" as a pathogenic agent is still obscure.

The single record of *Rickettsia* in chiggers from Brazil was documented by Bassini-Silva et al. (2018a), and correspond to a *Rickettsia felis*-like agent detected in *B. sinnamaryi*, collected on *Tachyphonus coronatus* (Aves: Passeriformes) from the state of Minas Gerais. However, *Rickettsia* spp. have been reported from various mites collected on birds and mammals (VYSOTSKAYA & SCHLUGER, 1953; HASE et al., 1978; CHOI et al., 2007; TAKAHASHI et al., 2004; MIŤKOVÁ et al., 2015; HUANG et al., 2017; BASSINI-SILVA et al., 2018a), which suggest that our finding is not an isolated case. On the other hand, we had no success in detecting the presence of *Coxiella* sp. and *Hepatozoon* sp. in the analyzed chiggers.

This is the first report of *Rickettsia* detection in chiggers collected on rodents in Brazil. As studies on chiggers as possible vectors of *Rickettsia* are scarce in the Neotropical region, more prospections are needed to determine if these mites are, in fact, involved in the epidemiological cycles of rickettsial diseases.

Acknowledgements

To Gabrielle Ribeiro de Andrade and Maria Cristina Ferreira do Rosário for technical contributions (CNPq no. 377343/2015-3 and 377342/2015-7, respectively). To Debra D. Creel for the technical

support. To Melissa K. Miller, entomologist, Public Health Command - Atlantic, Fort Meade, Maryland instead of entomologist U.S. Army Forces, for her valuable comments on this manuscript. This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq no. 454907/2014-1 and 440639/2015-8 to DMB-B, 377976/2014-8 to FCJ), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP no. 2010/51875-9 to DMB-B and 2017/01416-7 to RB-S). Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA; USDA is an equal opportunity provider and employer. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

References

- Almeida AP, Marcili A, Leite RC, Nieri-Bastos FA, Domingues LN, Martins JR, et al. *Coxiella* symbiont in the tick *Ornithodoros rostratus* (Acari: argasidae). *Ticks Tick Borne Dis* 2012; 3(4): 203-206. <http://dx.doi.org/10.1016/j.ttbdis.2012.02.003>. PMID:22480930.
- Almeida AP, Souza TD, Marcili A, Labruna MB. Novel *Ehrlichia* and *Hepatozoon* agents infecting the crab-eating fox (*Cerdocyon thous*) in southeastern Brazil. *J Med Entomol* 2013; 50(3): 640-646. <http://dx.doi.org/10.1603/ME12272>. PMID:23802461.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1990; 215(3): 403-410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2). PMID:2231712.
- Bassini-Silva R, Jacinavicius FC, Maturano R, Muñoz-Leal S, Ochoa R, Bauchan G, et al. *Blankaartia sinnamaryi* (Trombidiformes: Trombiculidae) parasitizing birds in southeastern Brazil, with notes on *Rickettsia* detection. *Rev Bras Parasitol Vet* 2018a; 27(3): 354-362. <http://dx.doi.org/10.1590/s1984-296120180057>. PMID:30184002.
- Bassini-Silva R, Jacinavicius FC, Muñoz-Leal S, Maturano R, Welbourn WC, Ochoa R, et al. A new species of the genus *Eutrombicula* Ewing, 1938 (Trombidiformes: Trombiculidae) and new records for the species *Eutrombicula batatas* (Linnaeus, 1758) in Brazil. *Acarologia* 2018b; 58(4): 976-986. <http://dx.doi.org/10.24349/acarologia/20184304>.
- Blanc G, Joyeux C, Bruneau J. Observations sur les larves de *Trombicula autumnalis* (SHAW) dans le centre de la France. Recherches sur leur rôle possible dans la transmission de la maladie de Derrick-Burnet (Q-Fever). *Arch Inst Pasteur Maroc* 1952; 4: 314-325.
- Brennan JM, Goff ML. Keys to the genera of chiggers of the Western Hemisphere (Acarina: trombiculidae). *J Parasitol* 1977; 63(3): 554-566. <http://dx.doi.org/10.2307/3280021>. PMID:68115.
- Choi YJ, Lee EM, Park JM, Lee KM, Han SH, Kim JK, et al. Molecular detection of various rickettsiae in mites (Acari: Trombiculidae) in southern Jeolla Province, Korea. *Microbiol Immunol* 2007; 51(3): 307-312. <http://dx.doi.org/10.1111/j.1348-0421.2007.tb03912.x>. PMID:17380050.
- Chomczynski P. A reagent for the single-step simultaneous isolation of RNA, DNA and proteins from cell and tissue samples. *Biotechniques* 1993; 15(3): 532-534, 536-537. PMID:7692896.
- Cruickshank RH. Molecular markers for the phylogenetics of mites and ticks. *Syst Appl Acarol* 2002; 7(0): 3-14. <http://dx.doi.org/10.11158/saa.7.1.1>.

- Dabert J, Ehrnsberger R, Dabert M. *Glaucales tytonis* sp. n. (Analoidea, Xolalgidae) from the barn owl *Tyto alba* (Strigiformes, Tytonidae): compiling morphology with DNA barcode data for taxon description in mites (Acari). *Zootaxa* 2008; 1719: 41-52.
- Dabert M, Witalinski W, Kazmierski A, Olszanowski Z, Dabert J. Molecular phylogeny of acariform mites (Acari, Arachnida): strong conflict between phylogenetic signal and long-branch attraction artifacts. *Mol Phylogenet Evol* 2010; 56(1): 221-241. <http://dx.doi.org/10.1016/j.ympev.2009.12.020>. PMID:20060051.
- Daniel M. The bionomics and developmental cycle of some chiggers (Acariformes, Trombiculidae) in the Slovak Carpathians. *Cs Parasitol* 1961; 8: 31-118.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 1994; 3(5): 294-299. PMID:7881515.
- Fonseca F, Notas de Acaerologia I. Papel dos acarídeos do gênero *Trombicula* na transmissão das Rickettsias patogênicas e aplicação dessa hipótese à *Rickettsia brasiliensis* Montero, 1931. *Mem Inst Butantan* 1932; 3: 127-133.
- Frank C. Ein Beitrag zur Biologie von *Hepatozoon erhardovae* Krampitz 1964 in Rötelmäusen aus der Südweststeiermark und des Neusiedlerseegbietes (Burgenland). *Parasitol Res* 1977; 53(3): 251-254. <http://dx.doi.org/10.1007/BF00389940>.
- Hase T, Roberts LW, Hildebrandt PK, Cavanaugh DC. Stylostome formation by *Leptotrombidium* mites (Acari: trombiculidae). *J Parasitol* 1978; 64(4): 712-718. <http://dx.doi.org/10.2307/3279967>. PMID:98623.
- Hillis DM, Dixon MT. Ribosomal DNA: molecular evolution and phylogenetic inference. *Q Rev Biol* 1991; 66(4): 411-453. <http://dx.doi.org/10.1086/417338>. PMID:1784710.
- Huang Y, Zhao L, Zhang Z, Liu M, Xue Z, Ma D, et al. Detection of a novel *Rickettsia* from *Leptotrombidium scutellare* Mites (Acari: Trombiculidae) from Shandong of China. *J Med Entomol* 2017; 54(3): 544-549. <http://dx.doi.org/10.1093/jme/tjw234>. PMID:28399204.
- Jacinavicius FC, Bassini-Silva R, Mendoza-Roldan JA, Muñoz-Leal S, Hingst-Zaher E, Ochoa R, et al. A contribution to the knowledge of *Quadrasetta brasiliensis* Goff and Gettinger, 1989 (Trombidiformes: Trombiculidae), with description of the deutonymph instar. *Acarologia* 2018a; 58(2): 442-456. <http://dx.doi.org/10.24349/acarologia/20184252>.
- Jacinavicius FC, Bassini-Silva R, Mendoza-Roldan JA, Pepato AR, Ochoa R, Welbourn C, et al. A checklist of chiggers from Brazil, including new records (Acari: Trombidiformes: Trombiculidae and Leeuwenhoekiidae). *ZooKeys* 2018b; 743(1): 1-41. <http://dx.doi.org/10.3897/zookeys.743.22675>. PMID:29670435.
- Kampen H, Scholer A, Metzen M, Oehme R, Hartelt K, Kimmig P, et al. *Neotrombicula autumnalis* (Acari, Trombiculidae) as a vector for *Borrelia burgdorferi* sensu lato? *Exp Appl Acarol* 2004; 33(1-2): 93-102. <http://dx.doi.org/10.1023/B:APPA.0000029975.92510.90>. PMID:15285141.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 2012; 28(12): 1647-1649. <http://dx.doi.org/10.1093/bioinformatics/bts199>. PMID:22543367.
- Kelly DJ, Fuerst PA, Ching WM, Richards AL. Scrub typhus: the geographic distribution of phenotypic and genotypic variants of *Orientia tsutsugamushi*. *Clin Infect Dis* 2009; 48(Suppl.3): S203-S230. <http://dx.doi.org/10.1086/596576>. PMID:19220144.
- Kepka O. Merkblätter über angewandte Parasitenkunde und Schädlingsbekämpfung, Merkblatt Nr. 12: Die Herbstmilbe (*Neotrombicula autumnalis*). *Angew Parasitol Beilage* 1965; 6(4): 1-13.
- Kumlert R, Chaisiri K, Anantatat T, Stekolnikov AA, Morand S, Prasartvit A, et al. Autofluorescence microscopy for paired-matched morphological and molecular identification of individual chigger mites (Acari: Trombiculidae), the vectors of scrub typhus. *PLoS One* 2018; 13(3): e0193163. <http://dx.doi.org/10.1371/journal.pone.0193163>. PMID:29494599.
- Labruna MB, Whitworth T, Horta MC, Bouyer DH, McBride JW, Pinter A, et al. *Rickettsia* species infecting *Amblyomma cooperi* ticks from an area in the state of São Paulo, Brazil, where Brazilian spotted fever is endemic. *J Clin Microbiol* 2004; 42(1): 90-98. <http://dx.doi.org/10.1128/JCM.42.1.90-98.2004>. PMID:14715737.
- Le Gac P, Giroud P, Andre M, Roger F. Deux modes de contamination professionnelle par la fièvre Q. *Bull Soc Pathol Exot* 1953; 46: 659-662.
- Luz HR, Silva-Santos E, Costa-Campos CE, Acosta I, Martins TF, Muñoz-Leal S, et al. Detection of *Rickettsia* spp. in ticks parasitizing toads (*Rhinella marina*) in the northern Brazilian Amazon. *Exp Appl Acarol* 2018; 75(3): 309-318. <http://dx.doi.org/10.1007/s10493-018-0270-y>. PMID:29846852.
- Miranda J, Mattar S, Puerta-González A, Muskus C, Oteo JA. Genome sequence of “*Candidatus Rickettsia colombianensi*”, a novel tick-associated bacterium distributed in Colombia. *Microbiol Resour Announc* 2019; 8(14): e01433-18. <https://dx.doi.org/10.1128%2FMRA.01433-18>.
- Miranda J, Portillo A, Oteo JA, Mattar S. *Rickettsia* sp. Strain Colombianensi (Rickettsiales: Rickettsiaceae): a new proposed *Rickettsia* detected in *Amblyomma dissimile* (Acari: Ixodidae) from Iguanas and free-living larvae ticks from vegetation. *J Med Entomol* 2012; 49(4): 960-965. <http://dx.doi.org/10.1603/ME11195>. PMID:22897060.
- Mitková K, Berthová L, Kalúz S, Kazimírová M, Burdová L, Kocianová E. First detections of *Rickettsia helvetica* and *R. monacensis* in ectoparasitic mites (Laelapidae and Trombiculidae) infesting rodents in south-western Slovakia. *Parasitol Res* 2015; 114(7): 2465-2472. <http://dx.doi.org/10.1007/s00436-015-4443-x>. PMID:25843569.
- Moniuszko H, Zaleśny G, Małol J. Host-associated differences in morphometric traits of parasitic larvae *Hirsutiella zachvatkini* (Actinotrichida: trombiculidae). *Exp Appl Acarol* 2015; 67(1): 123-133. <http://dx.doi.org/10.1007/s10493-015-9925-0>. PMID:26002309.
- National Center for Biotechnology Information – NCBI. Blastn [online]. 2019 [cited 2019 Mar 10]. Available from: www.ncbi.nlm.nih.gov/blast
- Navajas M, Gutierrez J, Bonato O, Bolland HR, Mapangou-Divassa S. Intraspecific diversity of the cassava green mite *Mononychellus progressivus* (Acari: Tetranychidae) using comparisons of mitochondrial and nuclear ribosomal DNA sequences and cross-breeding. *Exp Appl Acarol* 1994; 18(6): 351-360. <http://dx.doi.org/10.1007/BF00116316>. PMID:7628252.
- Otto JC, Wilson KJ. Assessment of the usefulness of ribosomal 18S and mitochondrial COI sequences in Prostigmata phylogeny. In: Proctor HC, Norton RA, Colloff MJ. *Acarology: Proceedings of the 10th International Congress*; 2011; Collingwood, Australia. Melbourne: CSIRO Publishing; 2001. p. 100-109.
- Park SW, Ha NY, Ryu B, Bang JH, Song H, Kim Y, et al. Urbanization of scrub typhus disease in South Korea. *PLoS Negl Trop Dis* 2015; 9(5): e0003814. <http://dx.doi.org/10.1371/journal.pntd.0003814>. PMID:26000454.

- Phasomkusolsil S, Tanskul P, Ratanatham S, Watcharapichat P, Phulsuksombati D, Frances SP, et al. Transstadial and transovarial transmission of *Orientia tsutsugamushi* in *Leptotrombidium imphalum* and *Leptotrombidium chiangraiensis* (Acari: trombiculidae). *J Med Entomol* 2009; 46(6): 1442-1445. <http://dx.doi.org/10.1603/033.046.0628>. PMID:19960694.
- Rapmund G, Upham RW Jr, Kundin WD, Manikumar C, Chan TC. Transovarial development of scrub typhus rickettsiae in a colony of vector mites. *Trans R Soc Trop Med Hyg* 1969; 63(2): 251-258. [http://dx.doi.org/10.1016/0035-9203\(69\)90155-2](http://dx.doi.org/10.1016/0035-9203(69)90155-2). PMID:4978468.
- Regnery RL, Spruill CL, Plikaytis BD. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol* 1991; 173(5): 1576-1589. <http://dx.doi.org/10.1128/jb.173.5.1576-1589.1991>. PMID:1671856.
- Roux V, Fournier PE, Raoult D. Differentiation of spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein rOmpA. *J Clin Microbiol* 1996; 34(9): 2058-2065. PMID:8862558.
- Santibáñez PA, Palomar M, Portillo A, Santibáñez S, Oteo JA. The role of chiggers as human pathogens. In: Samie A. *An overview of tropical diseases*. London: Intech; 2015. p. 173-202. <http://dx.doi.org/10.5772/61978>.
- Schluger EG. New red mites (Acari, Trombidiidae) of the fauna of the USSR. *Entomol Obozr* 1948; 30: 157-164.
- Soller R, Wohltmann A, Witte H, Blohm D. Phylogenetic relationships within terrestrial mites (Acari: Prostigmata, Parasitengona) inferred from comparative DNA sequence analysis of the mitochondrial cytochrome oxidase subunit I gene. *Mol Phylogenet Evol* 2001; 18(1): 47-53. <http://dx.doi.org/10.1006/mpev.2000.0855>. PMID:11161741.
- Takahashi M, Misumi H, Urakami H, Nakajima S, Furui S, Yamamoto S, et al. Mite vectors (Acari: Trombiculidae) of scrub typhus in a new endemic area in northern Kyoto, Japan. *J Med Entomol* 2004; 41(1): 107-114. <http://dx.doi.org/10.1603/0022-2585-41.1.107>. PMID:14989353.
- Vysotskaya SO, Schluger EG. Licinki krasnotelok-parazity gryzunov Leningradskoj oblasti. *Parazitologicheskii Sbornik* 1953; 15: 345-352.
- Walter DE, Krantz GW. Collecting, rearing, and preparing specimens. In: Krantz GW, Walter DE. *A manual of acarology*. Lubbock: Texas Tech University Press; 2009. p. 83-96.