

A Systematic Mapping Study of Emotional Response Evaluation Instruments

Suzane Santos dos Santos and Kamila Rios da Hora
Rodrigues^[0000-0001-9176-8709]

University of São Paulo, São Carlos, São Paulo, Brazil
suzanesantos@usp.br, kamila.rios@icmc.usp.br

Abstract. In the context of the study of emotions, the limits that define it can be so confusing that everything can be easily characterized as an emotion. Experts are not unanimous about what is an emotion and what is not, and if this is a challenge for experts, for other professionals it is an even bigger barrier. For example, analyzing users' emotional aspects when interacting with interactive computational solutions is very difficult for Computer professionals, in several situations, this evaluation is the responsibility of the specialist professionals in the studied domain. In an attempt to alleviate this problem, this systematic mapping study seeks to identify different instruments for evaluating emotional responses from different contexts to help professionals of many domains choose suitable artifacts for their evaluations. We identified 32 studies that describe 18 different instruments, these instruments are mainly from the field of Psychology and aimed at adults.

Keywords: Computational Systems Evaluation · Emotional Aspects · Emotional Response · Systematic Mapping.

1 Introduction

There are different ways to evaluate a product, whether it is a physical good, a service provided, or a computational solution. In order to conduct a practical assessment, it is crucial to know how and when to use the different types of assessment available in the literature. Human-Computer Interaction (HCI) is a research area of Computer Science in which evaluation has high relevance. During the evaluation stage, problems in the interface and user interaction, not noticed in the design and development stages, are identified and corrected. Thus, after a systematic and careful evaluation, the user has the chance to receive a safer, more effective product that, above all, does not harm their experience while using the product.

In the context of evaluation and HCI, another concept intrinsically associated with usability is the concept of User eXperience (UX). According to Nilsen [35], the user experience encompasses all aspects of the end user's interaction with its services and products. More specifically, UX is related to how people feel about a product and their pleasure and satisfaction when using it.

User emotion, in turn, is no longer just related to unexpected system response or frustration with an incomprehensible error message. It is now understood that a wide range of emotions plays an essential role in all tasks performed on the computer. When interacting with computer systems, users' emotions are a fundamental aspect to help understand the user experience [4, 46].

This recent change in user emotion concerning interactive systems has raised the need to understand better what emotion is and how it influences the user during interaction. However, even though the term emotion is used very often, and several studies in the literature address this issue, there is no consensus on the concept, which is controversial even for specialists in the field [45].

Given that HCI is the intersection between Psychology and Social Sciences on the one hand and the combination of Computer Science and technology on the other, it is crucial to understand how different areas of knowledge understand and assess individuals' emotional responses. Hence, we conducted a Systematic Mapping (SM) study whose objective is to identify instruments for evaluating emotional responses and find instruments from other fields that can be systematized and incorporated into the area of Computing or other areas.

This paper is divided as follows: Section 2 describes the theoretical foundation, Section 3 describes the protocol for planning, conducting and reporting the systematic mapping. Section 4 contains the synthesis of the results obtained through the mapping. In section 5 we make the final considerations and our conclusions on the subject.

2 Theoretical Foundation

In this section, we present a summary of the study of emotions. In the literature, there are several definitions of emotion. According to Young [59], emotion is an acute disorder of the individual, of psychological origin, involving behavior, conscious experience, and visceral functioning. For Ekman [12], emotion refers to the process by which an elicitor is assessed automatically or in an extended way. An affect program may or may not be triggered, organized responses may occur, although more or less managed by attempts to control emotional behavior.

According to Izard [21], emotion is a complex concept with neurophysiological, neuromuscular, and phenomenological aspects. At the neurophysiological level, emotion is defined primarily in terms of patterns of electrochemical activity in the nervous system. At the neuromuscular level, emotion is primarily a facial activity, and facial patterns and secondary is a bodily response. At the phenomenological level, emotion is essentially a motivating experience or experience that has immediate meaning and importance. These definitions were found in the work proposed by Kleinginna and Kleinginna [25], in which the authors compiled a compilation of 92 definitions and nine skeptical statements from a variety of sources in the emotion literature.

For Coan and Allen [9], emotion is too broad a class of events to be a single scientific category. As psychologists use the term, it includes the euphoria of winning an Olympic gold medal, a brief startle with an unexpected noise, a

deep, unrelenting pain, the fleeting pleasant sensations of a warm breeze. While it can also mean cardiovascular changes in response to the display of a movie, stalking and murder of an innocent victim, lifelong love for a child, feeling excited about no known reason, and interest in a newsletter.

The boundaries of emotion can be so confusing that everything can easily be characterized as emotion. Experts are not unanimous about what is an emotion and what is not. All the different types of events included in this term are essential, some of the vital importance. Nevertheless, it is increasingly evident that not all events can be explained in the same way. No description and evaluation framework can do justice to this heterogeneous class of events without differentiating one type of event from another [9].

2.1 Emotion Evaluation

The definition of Scherer [44] states that: “Emotion is defined as an episode of synchronized and interrelated changes in the states of all or most of the five subsystems of the organism in response to the evaluation of a stimulus event external or internal as relevant to the main concerns of the organism”. Therefore, we adopted this definition in this study. The rationale behind this choice lies in the fact that this definition is one of the most comprehensive. The five components are:

1. Cognitive evaluations, which have the function of evaluating objects and events;
2. Behavioral trends (action trends), responsible for preparing and directing activities;
3. Motor expressions (facial and vocal expressions), which communicate reactions and behavioral intentions;
4. Physiological reactions (physical symptoms), responsible for regulating the body;
5. Subjective feelings (conscious experience), which monitors the organism’s internal state and interaction with the environment.

This study focuses on the subjective feelings component. This means that we only considered self-report instruments for the systematic mapping study. In the field of Psychology, a self-report is any test, measure, or survey that is based on an individual’s own account of their symptoms, behaviors, beliefs, or feelings. Examples that are widely used are interviews (structured or not) and questionnaires, which are usually applied using paper and pencil, or online.

3 Methodology

This section outlines the protocol used to carry out this study. The protocol consists of five activities: defining the research questions, the search process, research strategy (inclusion/exclusion criteria), data extraction strategy, and synthesis of the extracted data.

3.1 Research Question

The present systematic mapping study addresses one leading research question, which we named RQ: “What self-report instruments are used to evaluate individuals’ emotional responses?”.

By answering this research question, we can discover which self-report instruments are used to assess individuals’ emotional responses. Since this is a broad question and is not limited to a specific area, the results may include well-known and widely used instruments, as well as innovative instruments that may be discovered and disseminated.

3.2 Search Process

The search process aimed at identifying studies that will answer our research question. In order to achieve this goal, we create a search string for the search process, gathering the most relevant terms related to the search question and combining them by logical operators. To obtain relevant and valuable results for this study, some iterations were carried out until reaching the terms that composed the following string used: (**”emotion evaluation” OR ”emotional evaluation” OR ”emotional response evaluation” OR ”evaluation of emotion”**).

The procedure consisted of an automated search into well-known digital libraries in both Computing and Health areas. The electronic search was performed on:

- ACM Digital Library¹;
- IEEE Xplore²;
- PubMed digital libraries³;
- Scielo⁴;
- Scopus⁵;
- Virtual Health Library⁶;
- Web of Science⁷.

3.3 Search Strategy

Inclusion (IC) and exclusion (EC) criteria were defined for the studies returned by the search string, as shown in Table 1.

The selection process followed six steps:

1. Execution of the search on the bases previously chosen;

¹ <https://dl.acm.org/>

² <https://ieeexplore.ieee.org>

³ <https://pubmed.ncbi.nlm.nih.gov/>

⁴ <https://www.scielo.br/>

⁵ <https://www.scopus.com/>

⁶ <https://bvshalud.org/en/>

⁷ <https://www.webofscience.com/>

Table 1. Selection Criteria.

Criteria	Code	Description
Inclusion	IC.1	Study that explicitly addresses the evaluation of users' emotional responses.
Exclusion	EC.1	Study that does not address the evaluation of users' emotional response.
	EC.2	Study that presents instruments that are not self-report.
	EC.3	Study written in languages other than English.

2. Removal of duplicates studies;
3. Selection through title and abstract;
4. Application of selection criteria in the studies selected in step 3, in the full text;
5. Application of quality criteria in the final set of selected studies;
6. And finally, data extraction.

Quality criteria (see Table 2) were adopted to ensure that the selected studies were relevant to answering the research question raised. The possible answers to the questions were “yes”, “partially”, or “no”, quantified with the values “1”, “0.5”, and “0”, respectively. For the paper to be considered sufficient quality to have its data extracted for the research, it was necessary to reach a minimum score of 3.5 points. The studies that did not reach the minimum score were eliminated. The quality criteria applied to the studies are described in Table 2.

Table 2. Quality Criteria

Quality Criteria	Question
QC.1	Does the study define who the target audience is?
QC.2	Does the study describe which emotions it assesses?
QC.3	Is the assessment procedure replicable?
QC.4	Can the instrument be digitized?

The database search returned a total of 1410 studies, of which 736 were duplicates. Thus, only 674 went through the first iteration of the inclusion and exclusion criteria. At this stage, based on reading the title and abstract, we selected 70 studies. In the second iteration of the selection criteria, we read the full text of the remaining studies and applied the quality criteria, the final set of studies consisted of 52 studies. The identified works are described in Table 3.

Table 3. Set of studies identified.

Id	Authors and Year	Source	Instrument
1	Müller <i>et al.</i> (2021) [34]	Scopus	University of California, Los Angeles Loneliness Scale (UCLA-3)
2	Bojan <i>et al.</i> (2021) [6]	Scopus	Positive and Negative Affect Schedule (PANAS)
3	Önder (2020) [60]	Scopus	The Oxford Happiness Questionnaire (OHP)
4	Oh <i>et al.</i> (2019) [36]	Scopus	The Profile of Mood States (POMS) and Semantic Differential (SD)
6	Racine (2017) [41]	Scopus	Self-Assessment Manikin (SAM)
7	Park <i>et al.</i> (2017) [39]	Scopus	A modified semantic differential (SD) and Profile of Mood States (POMS)
8	Carmel <i>et al.</i> (2017) [8]	Scopus	State Trait Anxiety Inventor (STAI)
9	Miśkiewicz <i>et al.</i> (2016) [33]	Scopus	Positive and Negative Affect Schedule (PANAS)
10	Balconi <i>et al.</i> (2016) [3]	Scopus	Self-Assessment Manikin (SAM)
11	Maffei <i>et al.</i> (2015) [30]	Web of Science	Self-Assessment Manikin (SAM)
12	Balconi <i>et al.</i> (2015) [2]	Scopus	An adapted version of SAM
13	Ermes <i>et al.</i> (2014) [14]	Scopus	PAD Semantic Scale and EmoCards
14	Melnyk <i>et al.</i> (2013) [32]	Scopus	Beck Youth Inventory (2nd Edition: BYI II)
15	Xavier and Neris (2012) [57]	Scopus	Self-Assessment Manikin (SAM)
16	Somaini <i>et al.</i> (2011) [48]	Scopus	State-Trait Anxiety Inventory Y-1 (STAI) Self-Assessment Manikin (SAM)
17	Vuoskoski and Eerola (2011) [55]	Scopus	POMS-A
18	Emery and Hess (2008) [13]	Scopus	Positive and Negative Affect Schedule (PANAS)
19	Gaina <i>et al.</i> (2004) [16]	Scopus	Profile of Mood States (POMS)
20	Shibata <i>et al.</i> (1993) [47]	Scopus	Semantic Differential (SD)
21	Park and Chong (2019) [38]	Web of Science	Music emotion assessment tool (MEAT)
22	Aguilar <i>et al.</i> (2008) [1]	Web Of Science	Self-Assessment Manikin (SAM)
23	Wood and Moreau (2006) [56]	Web Of Science	Modification of Differential Emotions Scale (DES-II)
24	Jayanthi <i>et al.</i> (2018) [23]	Web Of Science	Rosenberg Self-esteem Scale (RSE)
25	Gozansky <i>et al.</i> (2021) [18]	PubMed	The Revised University of California, Los Angeles (R-UCLA) Loneliness Scale and The Short Depression, Anxiety and Stress Scale (DASS-21)
26	Igasaki <i>et al.</i> (2020) [20]	PubMed	Profile of Mood States 2nd Edition (POMS 2)
27	Philpott <i>et al.</i> (2016) [40]	PubMed	Hospital Anxiety and Depression Scale (HADS)
28	St. Jacques <i>et al.</i> (2015) [50]	PubMed	Self-Assessment Manikin (SAM) and Positive and Negative Affect Schedule (PANAS)
29	Tempesta <i>et al.</i> (2010) [53]	PubMed	Self-Assessment Manikin (SAM)
30	Gil <i>et al.</i> (2011) [17]	PubMed	State Trait Anxiety Inventor (STAI) and Edinburgh Postnatal Depression Scale (EPDS)
31	Martinez <i>et al.</i> (2018) [31]	Virtual Health Library	Geriatric Depression Scale (GDS)
32	Firoozi <i>et al.</i> (2013) [15]	Virtual Health Library	Beck Depression Inventory (BDI)

3.4 Data Analysis

The data extraction process was carried out systematically, throughout a form for recording the information necessary for answer the research question, containing the following fields:

1. Study identifier (ID);
2. Title;
3. Authors;
4. Year;
5. Search base;
6. Evaluation instrument;
7. Instrument origin field;
8. Emotions evaluated by the instrument;
9. Target audience;
10. and Evaluation procedure.

In Table 3 we show the ID, authors, year, source and the evaluation instrument, and Table 4, in turn, shows the instruments used in the studies described in Table 3. In total, we identified 19 assessment instruments, some instruments, such as the POMS, have variations and only one version appears in the table.

4 Results

The instruments described in the tables were divided into four categories: screening instruments, non-verbal instruments, instruments based on rating scales, and instruments based on the semantic differential.

4.1 Screening Instruments

A screening test is done to detect potential health disorders or diseases in people who do not have any disease symptoms. The goal is early detection and lifestyle changes or surveillance to reduce the risk of disease or detect it early enough to treat it most effectively. Brief psychological measures can be used to screen individuals for a range of mental health conditions. Screening measures are often questionnaires completed by clients. Screening tends to be quick to administer, but results are only indicative: if a positive result is found on a screening test, then the screening test can be followed up by a more definitive test [54]. The ID's of instruments that fall into this category are: 14, 23, 25, 27, 30, 31 and 32.

The following is a brief description of the instruments in this category:

- Beck Youth Inventory (BYI II): this instrument uses five self-report inventories to assess symptoms of depression, anxiety, anger, disruptive behavior, and self-concept in children and adolescents [51];
- Differential Emotion Sacale (DES-II): the DES is a standardized instrument that reliably divides the individual's description of emotion experience into validated, discrete categories of emotion [22];

Table 4. Instruments identified.

Instrument	Origin field	Target audience	Emotional Aspects involved
UCLA-3	Psychology	Teens and adults	Subjective feelings of loneliness
PANAS	Psychology	Adults	Positive and negative affect
OHQ	Psychology	No restrictions described	Happiness
POMS	Psychology	Adults	Tension-anxiety, depression-dejection, anger-hostility, fatigue, confusion and vigor
SD	Psychology	No restrictions described	Connotative meaning of objects, events, and concepts
SAM	Psychology	No restrictions described	Valence/Pleasure, arousal and dominance/control
STAI	Psychology	Adults	State anxiety and trait anxiety
PAD Semantic Scale	Psychology	No restrictions described	Pleasure, arousal and dominance
EmoCards	Psychology	No restrictions described	Excited (neutral or pleasant), average pleasant, calm (pleasant, neutral or unpleasant), average unpleasant and excited unpleasant
BYI	Psychology	Children and adolescents	Depression, anxiety, anger, disruptive behavior, and self-concept
MEAT	Psychology of Music	Adults	Happiness, sadness, anger, and fear
DES-II	Psychology	No restrictions described	Seven positives and seven negatives emotions variables
RES	Psychology	Teens	Both positive and negative feelings about the self
DASS-II	Psychology	Adults	The negative emotional states of depression, anxiety, and stress/tension
HADS	Psychology	No restrictions described	States of depression and anxiety in the setting of an hospital medical outpatient clinic
EPDS	Psychology	Cisgender women	Postpartum depression symptoms on the third day after childbirth
GDS	Geriatric Psychiatry	Older adults	Symptoms of depression
BDI	Psychology	13 years old and above	Key symptoms of depression including mood, pessimism, sense of failure, etc.

- The Short Depression, Anxiety and Stress Scale (DASS-21): The Depression, Anxiety and Stress Scale - 21 Items (DASS-21) is a set of three self-report scales designed to measure the emotional states of depression, anxiety and stress. Each of the three DASS-21 scales contains 7 items, divided into sub-scales with similar content [29];
- Hospital Anxiety and Depression Scale (HADS): This instrument measure anxiety and depression in a general medical population of patients. HADS focuses on non-physical symptoms so that it can be used to diagnose depression in people with significant physical ill-health [52];
- Edinburgh Postnatal Depression Scale (EPDS): The 10-question Edinburgh Postnatal Depression Scale (EPDS) is a valuable and efficient way of identifying patients at risk for “perinatal” depression. The EPDS is easy to administer and has proven to be an effective screening tool. This instrument can only be applied by a specialist [10];
- Geriatric Depression Scale (GDS): GDS is a self-report measure of depression in older adults. Users respond in a “Yes/No” format. This form can be completed in approximately 5 to 7 minutes, making it ideal for people who are easily fatigued or are limited in their ability to concentrate for longer periods of time. GDS is a scale widely used and it is an instrument that non-specialists can administer [58];
- Beck Depression Inventory (BDI): The Beck Depression Inventory (BDI) is a 21-item, self-report rating inventory that measures characteristic attitudes and symptoms of depression. The BDI has been developed in different forms, including several computerized forms [5].

Screening tests for emotional disorders are usually administered by trained professionals. Systematizing them would be possible with the help of domain experts.

4.2 Non-verbal Instruments

Non-verbal instruments have no age restriction. They can be applied to children, the elderly, people with communication difficulties, and low education. The Self-Assessment Manikin (SAM) (ID’s: 6, 10, 11, 12, 15, 16, 22 and 28) is a non-verbal instrument that is also based on a 9-point Likert Scale.

The Self-Assessment Manikin (see Fig. 1) is an image-based questionnaire developed by Bradley and Lang [7] to measure emotional response. The questionnaire, widely used in evaluations by Computing professionals, was designed to measure three characteristics of an emotional response (pleasure, arousal and dominance), identified as central to emotion in research conducted by Lang et al. [27]. SAM can be considered free of language; that is, any individual, of any schooling, can answer it.

EmoCards (2) is an instrument made up of eight cards and is manually administered. The Emocard was inspired by the model of [43] and has eight emotions, each of these emotions is represented by a male and female face, totaling 16 cards, as shown in Fig. 2.




Fig. 1. SAM (Extracted from Bradley and Lang [7]).




Fig. 2. EmoCards (Extracted from Russell [43]).

4.3 Instruments based on Rating Scales

One of the most common rating scales is the Likert scale. The original Likert scale is a set of statements offered for a real or hypothetical situation under study. Participants are asked to show their level of agreement (from strongly disagree to agree strongly) with the given statement (items) on a metric scale. Here all the statements in combination reveal the specific dimension of the attitude towards the issue, hence, necessarily inter-linked with each other [24].

UCLA (1, 25) POMS (4, 7, 17, 19, 26), PANAS (2, 9, 18, 28), STAI (8, 16, 30), and OHQ (3), are instruments based on a Likert scale, as follows they will be briefly described.

- University of California, Los Angeles Loneliness Scale (UCLA): The UCLA Loneliness Scale is a commonly used measure of loneliness. It was originally released in 1978 as a 20-item scale. It has since been revised several times and shorter versions have been introduced for situations where 20 questions are too many, such as phone surveys [42];
- The Profile of Mood States (POMS): POMS questionnaires contain a series of descriptive words/statements that describe feelings people have. Subjects self report on each of these areas using a 5-point Likert scale. There are several versions of the POMS questionnaire. Currently, the most commonly used is the POMS 2, which is available for adults aged 18 years and older (POMS 2–A) and another for adolescents 13 to 17 years of age (POMS 2–Y). Both POMS 2 instruments are available as full-length (65 items) and short versions (35 items) [26];
- : Positive and Negative Affect Schedule (PANAS): PANAS is a self-report questionnaire consisting of two 10-item scales to measure positive and negative affect. Each item is rated on a 5-point scale from 1 to 5 [11];
- State-Trait Anxiety Inventory (STAI): The State-Trait Anxiety Inventory (STAI) is a commonly used measure of trait and state anxiety. It can be used in clinical settings to diagnose anxiety and to distinguish it from depressive syndromes. Form Y, its most popular version, has 20 items for assessing trait anxiety and 20 for state anxiety. All items are rated on a 4-point scale [49];
- The Oxford Happiness Questionnaire (OHP): The Oxford Happiness Questionnaire (OHQ) is a widely-used scale for assessment of personal happiness. Each item of questionnaire each presented as a single statement can be endorsed on a uniform six-point Likert scale [19].

4.4 Instruments based on the Semantic Differential

Developed by Osgood, Suci and Tannenbaum [37], the Semantic Differential usually takes a bipolar adjective scale of 5 or 7 points. This form usually differs according to the number of points on the scale, the degree, and marking of these points. The authors created this method when they realized the need to assess the affectivity and qualities of a concept, as well as ways to quantify the effective meaning of attitudes, opinions, perceptions, social image, personality,

preferences, and interests of people or patients with content related to their health, treatment, and illness, which are not directly measurable. The works whose ID's are 4, 7 and 20 use instruments based on the semantic differential [28].

The next section describes the potential threats related to the systematic mapping conducted

4.5 Threats to Validity

This subsection aims at presenting the most common threats to validity of this research. Such threats are described as follows:

- Study inclusion/exclusion bias: If inclusion/exclusion criteria are conflicting, or very generic ones;
- Construction of the search string: Problems with string construction can result in the search returning a large number of studies (including many irrelevant ones) or missing some relevant studies;
- Data extraction bias: The data extraction phase can be hampered by the use of “open questions” on the variables collected, whose treatment is not explicitly discussed in the protocol;
- Researcher bias: Finally, this threat refers to potential bias the authors of studies may have, while interpreting or synthesizing the extracted results.

5 Final Remarks

This study describes a systematic mapping of emotion evaluation instruments. Its contributions are the protocol planning and the mapping results. For each study selected, we extracted and summarized their information. The self-report instruments found in this study are mainly from the Psychology field and are aimed at adults. Most instruments are administered manually. Some instruments are already used in the Computing area.

The mapping also answered our research question and brought us several self-report instruments used in different domains to assess different emotions. Our objective is to offer a framework composed of a system with several of these systematized instruments (with the support of domain professionals), so that professionals can carry out their assessments and obtain data and results in real time.

We identified 32 papers that describe 18 different instruments. For each of these techniques we extracted and summarized their information. The self-report instruments that are used to assess the emotional responses of users listed in this study are mainly from the field of Psychology and aimed at adults. Most instruments are administered manually. Some instruments are already used in the Computing area, such as the SAM, the Semantic Differential and EmoCards.

We believe that this study is relevant for our field because computer professionals develop applications for several other areas, a frequent example of the tools focused on the Health area. In many cases, the application developer does

not have feedback from the end-user or the specialist, and this occurs because, in most situations, the evaluation is in charge of the domain specialists themselves. Therefore, it is of utmost importance that the professionals who work to create them have the necessary tools to evaluate them. One way to achieve this goal is to analyze users' emotional responses to these interactive systems. The identification of instruments that assess users' emotional responses in different areas is, therefore, essential so that new instruments can be disseminated.

6 Acknowledgments

The authors would like to thank the CAPES, Brazilian agency, for their financial support.

References

1. Aguilar de Arcos, F., Verdejo-García, A., Ceverino, A., Montañez-Pareja, M., López-Juárez, E., Sánchez-Barrera, M., López-Jiménez, Á., Pérez-García, M.: Dysregulation of emotional response in current and abstinent heroin users: negative heightening and positive blunting. *Psychopharmacology* **198**(2), 159–166 (2008)
2. Balconi, M., Cotelli, M., Brambilla, M., Manenti, R., Cosseddu, M., Premi, E., Gasparotti, R., Zanetti, O., Padovani, A., Borroni, B.: Understanding emotions in frontotemporal dementia: the explicit and implicit emotional cue mismatch. *Journal of Alzheimer's Disease* **46**(1), 211–225 (2015)
3. Balconi, M., Pala, F., Manenti, R., Brambilla, M., Cobelli, C., Rosini, S., Benussi, A., Padovani, A., Borroni, B., Cotelli, M.: Facial feedback and autonomic responsiveness reflect impaired emotional processing in parkinson's disease. *Scientific Reports* **6**, 31453 (08 2016). <https://doi.org/10.1038/srep31453>
4. Barbosa, S., Silva, B.: *Interação Humano-Computador*. Elsevier Brasil (2010), <https://books.google.com.br/books?id=qk0skwr.cewC>
5. Beck, A.T., Ward, C.H., Mendelson, M., Mock, J., Erbaugh, J.: An inventory for measuring depression. *Archives of general psychiatry* **4**(6), 561–571 (1961)
6. Bojan, K., Stavropoulos, T., Lazarou, I., Nikolopoulos, S., Kompatsiaris, I., Tsolaki, M., Mukaetova-Ladinska, E., Christogianni, A.: The effects of playing the cosma cognitive games in dementia. *International Journal of Serious Games* **8** (03 2021). <https://doi.org/10.17083/ijsg.v8i1.412>
7. Bradley, M.M., Lang, P.J.: Measuring emotion: the self-assessment manikin and the semantic differential. *Journal of behavior therapy and experimental psychiatry* **25**(1), 49–59 (1994)
8. Carmel, S., King, D., O'Rourke, N., Bachner, Y.: Subjective well-being: gender differences in holocaust survivors-specific and cross-national effects. *Aging and Mental Health* **21**, 668–675 (04 2017). <https://doi.org/10.1080/13607863.2016.1148660>
9. Coan, J.A., Allen, J.J.: *Handbook of emotion elicitation and assessment*. Oxford university press (2007)
10. Cox, J.L., Holden, J.M., Sagovsky, R.: Detection of postnatal depression: development of the 10-item edinburgh postnatal depression scale. *The British journal of psychiatry* **150**(6), 782–786 (1987)

11. Crawford, J.R., Henry, J.D.: The positive and negative affect schedule (panas): Construct validity, measurement properties and normative data in a large non-clinical sample. *British Journal of Clinical Psychology* **43**(3), 245–265 (2004). <https://doi.org/https://doi.org/10.1348/0144665031752934>, <https://bpspsychub.onlinelibrary.wiley.com/doi/abs/10.1348/0144665031752934>
12. Ekman, P.: Biological and cultural contributions to body and facial movement. *The anthropology of the body* (1977)
13. Emery, L., Hess, T.M.: Viewing instructions impact emotional memory differently in older and young adults. *Psychology and Aging* **23**(1), 2 (2008)
14. Ermes, V., Janß, A., Radermacher, K., Röcker, C.: Analyzing the benefits of integrative multi-dimensional assessments of usability features in interaction-centered user studies. In: *Proceedings of the 8th International Conference on Pervasive Computing Technologies for Healthcare*. p. 227–230. *PervasiveHealth '14*, ICST (Institute for Computer Sciences, Social-Informatics and Telecommunications Engineering), Brussels, BEL (2014). <https://doi.org/10.4108/icst.pervasivehealth.2014.255142>, <https://doi.org/10.4108/icst.pervasivehealth.2014.255142>
15. Firoozi, M., Besharat, M.A., Boogar, E.: Emotional regulation and adjustment to childhood cancer: Role of the biological, psychological and social regulators on pediatric oncology adjustment. *Iranian journal of cancer prevention* **6**, 65–72 (03 2013)
16. Gaina, A., Zhang, M., Sekine, M., Hamanishi, S., Wang, H., Nasermodeli, A., Kagamimori, S.: Improvement of daytime rapid eye movement parameters following a hot bath in night-shift workers. *Sleep and Biological Rhythms* **2**, 144–149 (2004)
17. Gil, S., Teissèdre, F., Chambres, P.: The evaluation of emotional facial expressions in early postpartum depression mood: A difference between adult and baby faces? *Psychiatry research* **186**, 281–6 (04 2011). <https://doi.org/10.1016/j.psychres.2010.06.015>
18. Gozansky, E., Moscona, G., Okon-Singer, H.: Identifying variables that predict depression following the general lockdown during the covid-19 pandemic. *Frontiers in Psychology* **12** (05 2021). <https://doi.org/10.3389/fpsyg.2021.680768>
19. Hills, P., Argyle, M.: The oxford happiness questionnaire: a compact scale for the measurement of psychological well-being. *Personality and Individual Differences* **33**(7), 1073–1082 (2002). [https://doi.org/https://doi.org/10.1016/S0191-8869\(01\)00213-6](https://doi.org/https://doi.org/10.1016/S0191-8869(01)00213-6), <https://www.sciencedirect.com/science/article/pii/S0191886901002136>
20. Igasaki, T., Hiramatsu, S., Yanagihara, D., Baba, Y.: Emotion evaluation during working on a puzzle by spatiotemporal pattern of band power of electroencephalogram. vol. 2020, pp. 1043–1046 (07 2020). <https://doi.org/10.1109/EMBC44109.2020.9176312>
21. Izard, C.E.: *The face of emotion*. (1971)
22. Izard, C.E.: *Patterns of emotions: A new analysis of anxiety and depression*. Academic Press (2013)
23. Jayanthi, M., Kumar, R., Swathi, S.: Investigation on association of self-esteem and students' performance in academics. *International Journal of Grid and Utility Computing* **9**, 211 (01 2018). <https://doi.org/10.1504/IJGUC.2018.093976>
24. Joshi, A., Kale, S., Chandel, S., Pal, D.K.: Likert scale: Explored and explained. *British Journal of Applied Science & Technology* **7**(4), 396 (2015)
25. Kleinginna, P.R., Kleinginna, A.M.: A categorized list of emotion definitions, with suggestions for a consensual definition. *Motivation and emotion* **5**(4), 345–379 (1981)

26. Lane, A., Terry, P., Fogarty, G.: Construct validity of the profile of mood states (01 2007)
27. Lang, P.J., Greenwald, M.K., Bradley, M.M., Hamm, A.O.: Looking at pictures: Affective, facial, visceral, and behavioral reactions. *Psychophysiology* **30**(3), 261–273 (1993)
28. Lopes, J.d.L., Nogueira-Martins, L.A., Andrade, A.L.d., Barros, A.L.B.L.d.: Escala de diferencial semântico para avaliação da percepção de pacientes hospitalizados frente ao banho. *Acta Paulista de Enfermagem* **24**, 815 – 820 (00 2011), http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-21002011000600015nrm=iso
29. Lovibond, P.F., Lovibond, S.H.: The structure of negative emotional states: Comparison of the depression anxiety stress scales (dass) with the beck depression and anxiety inventories. *Behaviour research and therapy* **33**(3), 335–343 (1995)
30. Maffei, A., Vencato, V., Angrilli, A.: Sex differences in emotional evaluation of film clips: Interaction with five high arousal emotional categories. *PLOS ONE* **10**(12), 1–13 (12 2016). <https://doi.org/10.1371/journal.pone.0145562>, <https://doi.org/10.1371/journal.pone.0145562>
31. Martinez, M., Multani, N., Anor, C., Misquitta, K., Tang-Wai, D., Keren, R., Fox, S., Lang, A., Marras, C., Tartaglia, M.: Emotion detection deficits and decreased empathy in patients with alzheimer’s disease and parkinson’s disease affect caregiver mood and burden. *Frontiers in Aging Neuroscience* **10** (04 2018). <https://doi.org/10.3389/fnagi.2018.00120>
32. Melnyk, B., Kelly, S., Lusk, P.: Outcomes and feasibility of a manualized cognitive-behavioral skills building intervention: Group cope for depressed and anxious adolescents in school settings. *Journal of child and adolescent psychiatric nursing : official publication of the Association of Child and Adolescent Psychiatric Nurses, Inc* **27** (10 2013). <https://doi.org/10.1111/jcap.12058>
33. Miśkiewicz, H., Antoszewski, B., Iljin, A.: Personality traits and decision on breast reconstruction in women after mastectomy. *Polish Journal of Surgery* **88**, 209–214 (2016)
34. Müller, F., Röhr, S., Reininghaus, U., Riedel-Heller, S.G.: Social isolation and loneliness during covid-19 lockdown: Associations with depressive symptoms in the german old-age population. *International Journal of Environmental Research and Public Health* **18**(7) (2021). <https://doi.org/10.3390/ijerph18073615>, <https://www.mdpi.com/1660-4601/18/7/3615>
35. Norman, D., Nielsen, J.: The definition of user experience (ux). <https://www.nngroup.com/articles/definition-user-experience/> (2006), acesso em: 02 Fev. 2021
36. Oh, Y.A., Kim, S.O., Park, S.A.: Real foliage plants as visual stimuli to improve concentration and attention in elementary students. *International Journal of Environmental Research and Public Health* **16**, 796 (03 2019). <https://doi.org/10.3390/ijerph16050796>
37. Osgood, C.E., Suci, G.J., Tannenbaum, P.H.: The measurement of meaning. No. 47, University of Illinois press (1957)
38. Park, H.Y., Chong, H.J.: A comparative study of the perception of music emotion between adults with and without visual impairment. *Psychology of Music* **47**(2), 225–240 (2019). <https://doi.org/10.1177/0305735617745148>, <https://doi.org/10.1177/0305735617745148>
39. Park, S.A., Song, C., Oh, Y.A., Miyazaki, Y., Son, K.C.: Comparison of physiological and psychological relaxation using measurements of heart rate variability, prefrontal

- cortex activity, and subjective indexes after completing tasks with and without foliage plants. *International Journal of Environmental Research and Public Health* **14**, 1087 (09 2017). <https://doi.org/10.3390/ijerph14091087>
40. Philpott, A.L., Andrews, S.C., Staios, M., Churchyard, A., Fisher, F.: Emotion evaluation and social inference impairments in huntington's disease. *Journal of Huntington's disease* **5**(2), 175–183 (2016)
 41. Racine, S.: Emotional ratings of high- and low-calorie food are differentially associated with cognitive restraint and dietary restriction. *Appetite* **121** (11 2017). <https://doi.org/10.1016/j.appet.2017.11.104>
 42. Russell, D., Peplau, L., Cutrona, C.: The revised ucla loneliness scale: Concurrent and discriminate validity evidence. *Journal of Personality and Social Psychology* **39**, 472–480. (10 1980). <https://doi.org/10.1037/0022-3514.39.3.472>
 43. Russell, J.: A circumplex model of affect. *Journal of Personality and Social Psychology* **39**, 1161–1178 (12 1980). <https://doi.org/10.1037/h0077714>
 44. Scherer, K.R.: Appraisal considered as a process of multilevel sequential checking. *Appraisal processes in emotion: Theory, methods, research* **92**(120), 57 (2001)
 45. Scherer, K.R.: What are emotions? and how can they be measured? *Social Science Information* **44**(4), 695–729 (2005). <https://doi.org/10.1177/0539018405058216>
 46. Sears, A., Jacko, J.A.: *Human-computer interaction fundamentals*. CRC press (2009)
 47. Shibata, S., Ohba, K., Inooka, N.: Emotional evaluation of human arm motion models. *Proceedings of 1993 2nd IEEE International Workshop on Robot and Human Communication* pp. 346–351 (1993)
 48. Somaini, L., Manfredini, M., Amore, M., Zaimovic, A., Raggi, M., Leonardi, C., Gerra, M., Donnini, C., Gerra, G.: Psychobiological responses to unpleasant emotions in cannabis users. *European archives of psychiatry and clinical neuroscience* **262**, 47–57 (07 2011). <https://doi.org/10.1007/s00406-011-0223-5>
 49. Spielberger, C.D., Gonzalez-Reigosa, F., Martinez-Urrutia, A., Natalicio, L.F., Natalicio, D.S.: The state-trait anxiety inventory. *Revista Interamericana de Psicologia/Interamerican Journal of Psychology* **5**(3 & 4) (1971)
 50. St. Jacques, P.L., Grady, C., Davidson, P.S., Chow, T.W.: Emotional evaluation and memory in behavioral variant frontotemporal dementia. *Neurocase* **21**(4), 429–437 (2015)
 51. Steer, R.A., Kumar, G., Beck, J.S., Beck, A.T.: Evidence for the construct validities of the beck youth inventories with child psychiatric outpatients. *Psychological reports* **89**(3), 559–565 (2001)
 52. Stern, A.F.: The Hospital Anxiety and Depression Scale. *Occupational Medicine* **64**(5), 393–394 (06 2014). <https://doi.org/10.1093/occmed/kqu024>, <https://doi.org/10.1093/occmed/kqu024>
 53. Tempesta, D., Couyoumdjian, A., Curcio, G., Moroni, F., Marzano, C., De Gennaro, L., Ferrara, M.: Lack of sleep affects the evaluation of emotional stimuli. *Brain research bulletin* **82**, 104–8 (04 2010). <https://doi.org/10.1016/j.brainresbull.2010.01.014>
 54. Tools, P.: Psychological assessment tools for mental health. <https://www.psychologytools.com/download-scales-and-measures/> (2021), acesso em: 02 Fev. 2021
 55. Vuoskoski, J.K., Eerola, T.: The role of mood and personality in the perception of emotions represented by music. *Cortex* **47**(9), 1099–1106 (2011). <https://doi.org/https://doi.org/10.1016/j.cortex.2011.04.011>, <https://www.sciencedirect.com/science/article/pii/S0010945211001067>
 56. Wood, S., Moreau, C.: From fear to loathing? how emotion influence the evaluation and early use of innovations. *Journal of Marketing - J MARKETING* **70**, 44–57 (07 2006). <https://doi.org/10.1509/jmkg.70.3.44>

57. Xavier, R., Neris, V.: A hybrid evaluation approach for the emotional state of information systems users. *ICEIS 2012 - Proceedings of the 14th International Conference on Enterprise Information Systems* **3**, 45–53 (01 2012)
58. Yesavage, J.A., Brink, T.L., Rose, T.L., Lum, O., Huang, V., Adey, M., Leirer, V.O.: Development and validation of a geriatric depression screening scale: a preliminary report. *Journal of psychiatric research* **17**(1), 37–49 (1982)
59. Young, P.T.: Emotion in man and animal; its nature and relation to attitude and motive. (1943)
60. Önder, I.: Association of happiness with morningness - eveningness preference, sleep-related variables and academic performance in university students. *Biological Rhythm Research* (11 2020). <https://doi.org/10.1080/09291016.2020.1848266>