

THE PROPERTY (D) AND THE ALMOST LIMITED COMPLETELY CONTINUOUS OPERATORS

M. L. LOURENÇO and V. C. C. MIRANDA*

Department of Mathematics, Institute of Mathematics and Statistics, University of São Paulo,
R. do Matão, 1010 – Butantã, 05508-090 São Paulo, Brazil
e-mails: lourencomarylilian@gmail.com, colferaiv@gmail.com

(Received August 31, 2021; revised April 14, 2022; accepted May 23, 2022)

Abstract. In this paper, we study some geometric properties in Banach lattices and the class of almost limited completely continuous operators. For example, we study Banach lattices with property (d) and we give a new characterization of this property in terms of the solid hull of almost limited sets.

1. Introduction

Throughout this paper, X and Y will denote real Banach spaces, and E and F will denote Banach lattices. We denote by B_X the closed unit ball of X . In a Banach lattice the additional lattice structure provides a large number of tools that are not available in more general Banach spaces. This fact facilitates the study of geometric properties of Banach lattices, as well as of the properties of operators acting between lattices. A number of questions arise naturally about the relationship between the properties of the operators and the lattice structure. Among these questions, one of the most frequent one is the so-called majorization problem, in which we study which properties S inherits from the operator T for given positive operators $S \leq T$ between Banach lattices E and F . These theories have won notoriety in many articles that have been developed over the last 20 years. Our interest is to study some geometric properties of Banach spaces in the context of Banach lattices and their consequences in operator theory. To state our results, we need to fix some notation and recall some definitions. We denote by E^+ the set of all positive elements in E . For a set $A \subset E$,

* Corresponding author.

Supported by a CAPES PhD's scholarship (88882.377946/2019-01).

Key words and phrases: almost limited set, almost limited completely continuous operator, Banach lattice, property (d), strong Gelfand–Phillips property.

Mathematics Subject Classification: 46B42, 47B65.

$A^+ = A \cap E^+$. Recall that $A \subset E$ is said to be *solid* if $|y| \leq |x|$ with $x \in A$ implies that $y \in A$. In addition, if F is a solid subspace of E , we say that F is an *ideal*. The *solid hull of A* , denoted by $\text{sol}(A)$, (resp. the *ideal generated by A* , E_A), is the smallest solid set that contains A (resp., the smallest ideal that contains A). A *band* B in E is an ideal such that $\sup(A) \in B$ for every subset $A \subset B$ which has supremum in E . A norm $\|\cdot\|$ of a Banach lattice E is called *order continuous* if for each net (x_α) such that $x_\alpha \downarrow 0$ in E imply that $\|x_\alpha\| \rightarrow 0$, where the notation $x_\alpha \downarrow 0$ means that the net (x_α) is decreasing and $\inf\{x_\alpha : \alpha\} = 0$. There are some equivalent statements of this property which can be found in [13, Section 2.4].

A Banach lattice E has *property (d)* if $|f_n| \xrightarrow{\omega^*} 0$ for every disjoint weak* null sequence $(f_n) \subset E'$. This property was studied by Wnuk in [20], however, this terminology was introduced later in [8]. In [6], Chen, Chen and Ji show that if E is a σ -Dedekind complete Banach lattice, then E has property (d). In [20] Wnuk showed that ℓ_∞/c_0 has property (d) but it is not σ -Dedekind complete. It is also clear that if the lattice operations in E' are weak* sequentially continuous, E has property (d). The reciprocal does not hold since ℓ_∞ has property (d), but the lattice operations of ℓ'_∞ are not weak* sequentially continuous. Indeed, by Josefson–Nussenzweig theorem there exists weak* null sequence $(f_n) \subset \ell'_\infty$ with $\|f_n\| = 1$ for all n , however $(|f_n|)$ is not weak* null, because for $e = (1, 1, 1, \dots)$, we have that $|f_n|(e) = \sup_{x \in [-e, e]} |f_n(x)| = \|f_n\| = 1$ for all n .

We recall a few geometric definitions in Banach spaces and its lattice versions. A subset $A \subset X$ is said to be *limited* if every weak* null sequence in X' converges uniformly to zero on A . We say that X has the *Dunford–Pettis* property* (DP*) if every relatively weakly compact set is limited. If every limited set of X is relatively compact it is said that X has the *Gelfand–Phillips property* (GP), or equivalently every limited weakly null sequence is norm null. Separable, reflexive and Schur Banach spaces have the GP property. The study and consequences of limited sets, DP* property and GP property in Banach spaces can be found, for example in [4, 5, 7]. We observe that a σ -Dedekind complete Banach lattice has the GP property if and only if it has order-continuous norm (see [19, Theorem 4.5]).

Following the idea and technique of Bouras [3] who considered the disjoint versions of the geometric properties of Banach spaces for lattices, Chen, Chen and Ji introduced the concept of almost limited sets in a Banach lattice [6]. A bounded set $A \subset E$ is an *almost limited set* if every disjoint weak* null sequence in E' converges uniformly to zero on A . We say that E has the *weak Dunford–Pettis* property* (wDP*) if every relatively weakly compact set is almost limited and E has the *strong Gelfand–Phillips property* (sGP) if every almost limited set is relatively compact. It is known that c_0 and c are Banach lattices with the sGP (see [2, p. 17]) without the wDP* property (see [6, Remark 3.4] and [2, Example 2.10], respectively) and the

Banach lattices ℓ_∞ and $L_1[0, 1]$ have the wDP* (see [6, Remark 3.4]) and do not have the sGP property (see [2, p. 19] for the fact that $L_1[0, 1]$ does not enjoy the sGP property).

The aim of this paper is to study property (d). In particular, we improve a result given in [11], about the solid hull of an almost limited set, and we also improve [2, Theorem 3.10] concerning operators that maps almost limited sets onto almost limited sets. As a consequence we give a new characterization of this property in terms of the solid hull of almost limited sets. We also apply these results to establish some properties of the almost limited completely continuous operators between Banach lattices, including a majority problem concerning this class of operators.

We refer the reader to [1,13] for Banach lattices theory and positive operators.

2. Banach lattices with property (d)

The property (d) and the almost limited sets are deeply connected. In [11], the authors proved that for σ -Dedekind complete Banach lattices, the solid hull of an almost limited set is also an almost limited set. In the next Lemma, we improve such result by showing that it holds for Banach lattices with property (d).

LEMMA 2.1. *Let E be a Banach lattice with property (d). If $A \subset E$ is almost limited, then $\text{sol}(A)$ also is almost limited.*

PROOF. Assume that $\text{sol}(A)$ is not almost limited. We are going to prove that A cannot be almost limited. Since $\text{sol}(A)$ is not almost limited, there exist a disjoint weak* null sequence $(f_n) \subset E'$ and a sequence $(x_n) \subset \text{sol}(A)$ such that $|f_n(x_n)| \geq \varepsilon$ for all n . In particular, for each n , there exist $y_n \in A$ such that $|x_n| \leq |y_n|$. On the other hand, it follows from [1, Theorem 1.23] that for each n , there exist $g_n \in E'$ satisfying $|g_n| \leq |f_n|$ and $|f_n|(|y_n|) = g_n(y_n)$. Now, as (f_n) is a disjoint weak* null sequence in E' and E has property (d), $|f_n| \xrightarrow{\omega^*} 0$. This implies that $g_n \xrightarrow{\omega^*} 0$. However,

$$\varepsilon \leq |f_n(x_n)| \leq |f_n|(|x_n|) \leq |f_n|(|y_n|) = g_n(y_n).$$

So, A cannot be almost limited, against the assumption. \square

Lemma 2.1 is not true for general Banach lattices. In fact, the singleton $\{e\}$ in c , where c denotes the Banach lattice of all real convergent sequences, is almost limited. However, its solid hull $\text{sol}(\{e\}) = [-e, e] = B_c$ is not almost limited. It is known that every order interval in a Banach lattice E is almost limited if and only if E has property (d) (see [12, Proposition 2.3]). Using this fact and Lemma 2.1, we have the following result.

PROPOSITION 2.2. *For a Banach lattice E , the following assertions are equivalent:*

1. *E has property (d).*
2. *Every order interval is almost limited.*
3. *The solid hull of every almost limited set also is almost limited.*

In [2], the authors gave an example where bounded linear operators do not map almost limited sets onto almost limited sets. The next result improves [2, Theorem 3.10], where F is σ -Dedekind complete.

THEOREM 2.3. *Let E and F be Banach lattices. Let $T: E \rightarrow F$ be an order bounded operator and let $A \subset E$ be an almost limited solid set. If F has property (d), then $T(A)$ is an almost limited subset of F .*

PROOF. By [12, Theorem 2.7], it suffices to prove that $f_n(Tx_n) \rightarrow 0$ for each disjoint sequence $(x_n) \subset A^+$ and each disjoint sequence $f_n \xrightarrow{\omega^*} 0$ in $(F')^+$. Considering the sequences $(x_n) \subset E$ and $(T'f_n) \subset E'$, there exists a disjoint sequence $(g_n) \subset E'$ satisfying $|g_n| \leq |T'f_n|$ and $g_n(x_n) = T'f_n(x_n)$ for all n (see [1, p. 77]). We claim that $g_n \xrightarrow{\omega^*} 0$. Indeed, if $x \in E^+$, since T is order bounded, there exists a $y \in F^+$ with $T[-x, x] \leq [-y, y]$. Hence

$$\begin{aligned} |T'f_n|(x) &= \sup\{|T'f_n(u)| : |u| \leq x\} \\ &\leq \sup\{|f_n(v)| : |v| \leq y\} = |f_n|(y) = f_n(y) \rightarrow 0. \end{aligned}$$

Now, since $x \in E^+$, it follows that $|g_n(x)| \leq |g_n|(|x|) = |T'f_n|(|x|) \rightarrow 0$, which yields the claim. As $A \subset E$ is almost limited, $(x_n) \subset A$ and $(g_n) \subset E'$ is a disjoint weak* null sequence, $g_n(x_n) \rightarrow 0$. Hence $T'f_n(x_n) \rightarrow 0$. \square

Now we give more examples of Banach lattices satisfying property (d). Let (E_n) be a family of Riesz spaces, the direct sum $\bigoplus_{n \in \mathbb{N}} E_n$ is a Riesz space with the following partial order relation $(x_n) \leq (y_n) \Leftrightarrow x_n \leq y_n$ for all $n \in \mathbb{N}$. Further, $(x_n) \vee (y_n) = (x_n \vee y_n)$ and $(x_n) \wedge (y_n) = (x_n \wedge y_n)$. If, in addition, each E_n is a Banach lattice, we can consider the following Banach lattices:

$$\left(\bigoplus_{n \in \mathbb{N}} E_n \right)_0 = \left\{ (x_n)_n : x_n \in E_n, \lim_n \|x_n\| = 0 \right\}$$

and

$$\left(\bigoplus_{n \in \mathbb{N}} E_n \right)_p = \left\{ (x_n)_n : x_n \in E_n, \sum_n \|x_n\|^p < \infty \right\},$$

where $1 \leq p < \infty$. In $(\bigoplus_{n \in \mathbb{N}} E_n)_0$ we consider the supremum norm, while in $(\bigoplus_{n \in \mathbb{N}} E_n)_p$ the p -norm. Moreover, $(\bigoplus_{n \in \mathbb{N}} E_n)'_0 = (\bigoplus_{n \in \mathbb{N}} E_n)_1$ and

$(\bigoplus_{n \in \mathbb{N}} E_n)'_p = (\bigoplus_{n \in \mathbb{N}} E_n)_q$ where $1/p + 1/q = 1$ (see [1, Theorem 4.6]). It is natural to ask under which conditions the above Banach lattices have property (d).

PROPOSITION 2.4. *Let (E_n) be a family of Banach lattices. Then $E = (\bigoplus_{n \in \mathbb{N}} E_n)_0$ has property (d) if, and only if E_n has property (d) for all n .*

PROOF. Suppose that all E_n have property (d). Let $(f_k) \subset E' = (\bigoplus_{n \in \mathbb{N}} E'_n)_1$ be a disjoint weak* null sequence. In particular, if $f_k = (f_k^n)_n$ with $f_k^n \in E'_n$ for all $k \in \mathbb{N}$ and $n \in \mathbb{N}$, we have that $(f_k^n)_k$ is disjoint in E'_n , because if $j \neq k$, $0 = f_k \wedge f_j = (f_k^n \wedge f_j^n)_n$, and so $f_k^n \wedge f_j^n = 0$ for all $n \in \mathbb{N}$ and for all $k \neq j$. In addition, since the linear embedding of E_n into E is continuous, we have that $f_k^n \xrightarrow{\omega^*} 0$, when $k \rightarrow \infty$, in E'_n . This implies that $|f_k^n| \xrightarrow{\omega^*} 0$ as $k \rightarrow \infty$ for all n .

Let $x = (x_n) \in E$. We claim that $|f_k|(x) \rightarrow 0$. First since $f_k \xrightarrow{\omega^*} 0$ in E' , there exists $M > 0$ such that

$$\| |f_k^n| \| = \| f_k^n \| \leq \sum_n \| f_k^n \| = \| |f_k| \|_1 \leq M \quad \text{for all } n, k \in \mathbb{N}.$$

Finally, let $\varepsilon > 0$. As $\|x_n\| \rightarrow 0$, there exists $n_0 \in \mathbb{N}$ such that $\|x_n\| < \varepsilon/(2M)$ for all $n > n_0$. On the other hand, as $|f_k^n|(x) \rightarrow 0$ when $k \rightarrow \infty$ for all $n \in \mathbb{N}$, for each $n \in \mathbb{N}$, there exists $k_n \in \mathbb{N}$ such that

$$| |f_k^n|(x) | < \varepsilon/(2n_0) \quad \text{for all } k \geq k_n.$$

Let $k_0 = \max\{k_1, \dots, k_{n_0}\}$. If $k \geq k_0$,

$$\begin{aligned} | |f_k|(x) | &\leq \sum_{n=1}^{n_0} | |f_k^n|(x_n) | + \sum_{n=n_0+1}^{\infty} | |f_k^n|(x_n) | \\ &\leq \sum_{n=1}^{n_0} | |f_k^n|(x_n) | + \sum_{n=n_0+1}^{\infty} \| f_k^n \| \| x_n \| < \sum_{n=1}^{n_0} \varepsilon/(2n_0) + \| |f_k| \|_1 \varepsilon/(2M) < \varepsilon. \end{aligned}$$

This means that $|f_k| \xrightarrow{\omega^*} 0$ as desired.

Now, we assume that E_n does not have property (d) for some n . Then, there exists a disjoint weak* null sequence $(f_n^k)_k \subset E'_n$ such that $(|f_n^k|)_k$ is not weak* null. Without loss of generality, assume that there exist $\varepsilon > 0$ and an element $x_n \geq 0$ in E_n such that $|f_n^k|(x_n) \geq \varepsilon$ for all k . Consider $f_k = (0, \dots, 0, f_n^k, 0, \dots) \in E'$ for all $k \in \mathbb{N}$. We have that

$$|f_k| \wedge |f_j| = (0, \dots, 0, |f_n^k| \wedge |f_j^k|, 0, \dots) = 0,$$

which implies that (f_k) is disjoint in E' . Further, if $x = (x_i) \in E$, $f_k(x) = f_n^k(x_n) \rightarrow 0$ as $k \rightarrow \infty$. Hence $(f_k) \subset E'$ is weak* null. However,

$$|f_k|(0, \dots, 0, x_n, 0, \dots) = |f_n^k|(x_n) \geq \varepsilon$$

for all $k \in \mathbb{N}$. Thus E lacks property (d). \square

Using the same arguments of Proposition 2.4 and doing the necessary adaptations we can prove the next result.

PROPOSITION 2.5. *Let (E_n) be a family of Banach lattices and let $1 \leq p < \infty$. Then $E = (\bigoplus_{n \in \mathbb{N}} E_n)_p$ has property (d) if and only if E_n has property (d) for all n .*

Let $E_n = E$ for each n . Then $(\bigoplus_{n \in \mathbb{N}} E_n)_0$ and $(\bigoplus_{n \in \mathbb{N}} E_n)_p$ are denoted by $c_0(E)$ and $\ell_p(E)$, respectively, for all $1 \leq p < \infty$. As a consequence of Proposition 2.4 and Proposition 2.5, we get the following result.

COROLLARY 2.6. *For a Banach lattice E , the following are equivalent:*

1. E has property (d).
2. $c_0(E)$ has property (d).
3. $\ell_p(E)$ has property (d) for some $1 \leq p < \infty$.
4. $\ell_p(E)$ has property (d) for all $1 \leq p < \infty$.

We observe that every bounded linear operator $T: E \rightarrow c_0$ is uniquely determined by a weak* null sequence $(x'_n) \subset E'$ such that $T(x) = (x'_n(x))$ for all $x \in E$. When this sequence is disjoint, we say that T is a *disjoint operator*.

The next Theorem characterizes property (d) with the regularity of a disjoint linear operator on E . We recall that a linear operator between Banach lattices is said to be *regular* if it can be written as difference of two positive operators, or equivalently, if it is dominated by a positive operator.

THEOREM 2.7. *A Banach lattice E has property (d) if and only if every disjoint linear operator on E is regular.*

PROOF. Assume that E has property (d) and let $T: E \rightarrow c_0$ be a disjoint linear operator. Then there is a disjoint sequence $(x'_n) \subset E'$ with $T(x) = (x'_n(x))$ for all n . Since E has property (d), we can consider the positive operator $S: E \rightarrow c_0$ defined by $S(x) = (|x'_n|(x))$ for all $x \in E$. It follows that $S \geq T$, thus T is regular.

Let $(x'_n) \subset E'$ be a disjoint weak* null sequence and let $T: E \rightarrow c_0$ be the disjoint operator such that $T(x) = (x'_n(x))$. By hypothesis, T is regular. So we can find a positive linear operator $S \geq T$. Let $(y'_n) \subset (E')^+$ be the weak* null sequence such that $S(x) = (y'_n(x))$ for all x . In particular, if $x \in E^+$, then $x'_n(x) \leq y'_n(x)$ for all n . Thus $x'_n \leq y'_n$ and $(x'_n)^+ \leq y'_n$. Consequently,

$(x'_n)^+ \xrightarrow{\omega^*} 0$ in E' . Hence $(x'_n)^- = (x'_n)^+ - x'_n \xrightarrow{\omega^*} 0$ in E' . Therefore, $|x'_n| \xrightarrow{\omega^*} 0$. \square

We remark that property (d) is not preserved by closed sublattices. For instance, $E = \ell_\infty$ has property (d), however $F = c$ is a closed sublattice of E without property (d). However, we have that property (d) is preserved by projection bands. Recall that a band B in E is said to be a *projection band* if there exists a linear projection $P: E \rightarrow E$ satisfying $P(E) = B$ and $0 \leq Px \leq x$ for all $x \in E^+$. Such projection is called a *band projection* (see [13, Definition 1.2.1]).

PROPOSITION 2.8. *Let E be Banach lattice with property (d). If F is a projection band in E , then F has property (d).*

PROOF. Let $P: E \rightarrow E$ denote the band projection associated to the projection band F . In particular, $0 \leq Px \leq x$ holds for all $x \in E^+$. We claim that P is interval preserving, i.e. $[0, Px] = P([0, x])$ for all $x \in E^+$. Indeed, if $y \in [0, Px]$, i.e. $0 \leq y \leq Px$, we get that $y \in F$, because $Px \in F$ and F is an ideal in E , and so $Py = y$. Since $Px \leq x$, we have that $0 \leq Py = y \leq Px \leq x$, which yields that $y \in P([0, x])$. Thus $[0, Px] \subset P([0, x])$. On the other hand, if $y \in P([0, x])$, there exists $z \in [0, x]$ such that $y = Pz$, which implies that $0 \leq y = Pz \leq Px$, hence $y \in [0, Px]$. Then P is interval preserving. Consequently, P' is a Riesz homomorphism (see [13, Theorem 1.4.19]). In order to prove that F has property (d), let (y'_n) be a disjoint weak* null sequence in F' . As P' is a Riesz homomorphism, we get that $(P'y'_n)$ is a disjoint weak* null sequence in E' , and so $|P'y'_n| \xrightarrow{\omega} 0$ in E' . Finally, given $x \in F^+$, we have that

$$\begin{aligned} |y'_n|(x) &= \sup\{|y'_n(y)| : |y| \leq x, y \in F\} \\ &= \sup\{|y'_n \circ P(y)| : |y| \leq x, y \in F\} \\ &\leq \sup\{|P'y'_n(y)| : |y| \leq x, y \in E\} = |P'y'_n|(x) \rightarrow 0. \end{aligned}$$

Then $|y'_n| \xrightarrow{\omega^*} 0$ in F' . \square

As a consequence of Proposition 2.8, we have the following example:

EXAMPLE 2.9. Let $E = (\bigoplus_{n \in \mathbb{N}} \ell_2^n)_0$. It follows from a commentary after [14, Lemma 10] that $E^\perp = \{f \in E''' : f(x) = 0, \forall x \in E\}$ is a projection band in E''' . Thus, by Proposition 2.8, E^\perp has property (d).

Let E be a Banach lattice and let F be a closed ideal in E . The quotient space E/F is a Banach lattice endowed with the partial order $[x] \leq [y]$ if and only if there exist $x_1 \in [x]$ and $y_1 \in [y]$ such that $x_1 \leq y_1$ (see [1, p. 100]). Recall that $(E/F)'$ is isometric isomorphic to the annihilator F^\perp (see [9,

Proposition 2.6]). As a consequence, we get the next theorem which is a generalization of Proposition 2.8.

THEOREM 2.10. *Let E be a Banach lattice and let F be a closed ideal in E . If E has property (d), then E/F has property (d).*

PROOF. Let $\pi: E \rightarrow E/F$ be the canonical projection and consider its dual operator $\pi': (E/F)' \rightarrow F^\perp$, which is an isometric isomorphism. We show that π' is a Riesz homomorphism, i.e. $|\pi'f| = \pi'|f|$ for all $f \in (E/F)'$. Indeed, given $f \in (E/F)'$ and $x \in E^+$, it follows from [1, Theorem 1.18] that

$$\begin{aligned} |\pi'f|(x) &= \sup\{|\pi'f(y)| : |y| \leq x\} \\ &= \sup\{|f([y])| : |[y]| \leq [x]\} = |f|([x]) = (\pi'|f|)(x). \end{aligned}$$

If $(f_n) \subset (E/F)'$ be a disjoint weak* null sequence, as π' is a Riesz homomorphism, then $(\pi'f_n)$ is a disjoint sequence in F^\perp and so in E' . If $x \in E$, we have that $\pi'f_n(x) = f_n([x]) \rightarrow 0$. Thus $(\pi'f_n)$ is a disjoint weak* null sequence in E' . As E has property (d), $\pi'|f_n| = |\pi'f_n| \xrightarrow{\omega^*} 0$ in E' . If $[x] \in E/F$, we have that $|f_n|([x]) = \pi'|f_n|(x) \rightarrow 0$. Hence $|f_n| \xrightarrow{\omega^*} 0$ and consequently E/F has property (d). \square

It follows from Theorem 2.10 that ℓ_∞/c_0 has property (d), while it is not σ -Dedekind complete (see [20, Remark 1.5]). This shows that property (d) is, in a sense, a more robust concept than σ -Dedekind completeness.

We conclude this section with the following question:

QUESTION. *Let E be a Banach lattice with property (d). Has $\ell_\infty(E)$ property (d)?*

3. The class of the almost limited completely continuous operators

The study of certain geometric properties in Banach spaces is largely connected with observing the behavior of bounded linear operators. Whenever a new class or a new property appears in Banach spaces, it is natural to study the relationship of such property in the context of operators. The objective here is to study some operator classes from properties in Banach lattices, more specifically, we will study the class of the almost limited completely continuous operators which is connected to the sGP property in Banach lattices.

Let us recall some necessary definitions. We recall that $T: X \rightarrow Y$ is a *completely continuous* linear operator if $Tx_n \rightarrow 0$ for every weakly null sequence $(x_n) \subset X$. Since the limited and weakly convergent sequences play an important role in the study of Banach spaces with the GP property, the

authors in [15] introduced and studied the class of limited completely continuous operators. Following [15], a linear operator $T: X \rightarrow Y$ is *limited completely continuous* (lcc) if $Tx_n \rightarrow 0$ for every limited weakly null sequence $(x_n) \subset X$. The class of completely continuous operators and the class of limited completely continuous operators will be denoted by $Cc(X; Y)$ and $Lcc(X; Y)$, respectively.

As the lattice structure is, in general, distinct of the Banach space, in the setting of Banach lattices is natural study known properties in the class of Banach spaces under a Banach lattice point of view. By considering almost limited sequences instead of limited sequences, the authors in [2] introduced the class of alcc operators and studied its relation with the sGP property. Following [2], a bounded linear operator $T: E \rightarrow Y$ is said to be an *almost limited completely continuous* (alcc operator) if $Tx_n \rightarrow 0$ in Y for every almost limited weakly null sequence $(x_n) \subset E$. This class is denoted by $L^a cc(E; Y)$. Moreover, the authors in [2] showed that if E has the sGP property (resp. E has the wDP* property), then $L^a cc(E; Y) = L(E; Y)$ (resp. $L^a cc(E; Y) = Cc(E; Y)$) for every Banach space Y (see Theorem 3.4 and page 22 in [2], respectively).

In the Banach space theory, it is usual to connect classes of operators with classes of sets. In [16], the authors introduced the class of the L-limited sets and studied its relations with the class of lcc operators, a subset A of a dual space X' is a *L-limited set* if every limited weakly null sequence (x_n) in X converges uniformly to zero on A .

In the same way the authors in [2] introduced the alcc operators in order to establish a “lattice” version of the lcc operators by using almost limited sequences instead of limited sequences, we are going to consider almost limited sequences in the definition above in the place of limited sequences. Following this idea, we introduce the class of strong L-limited sets.

DEFINITION 3.1. A subset A of a dual space E' is called an *strong L-limited set* if every almost limited weakly null sequence (x_n) in E converges uniformly to zero on A .

It is clear that every strong L-limited subset of E' is L-limited. We will see after Proposition 3.4 that there exist L-limited sets which are not strong L-limited.

EXAMPLE 3.2. Let $E = C(K)$ where K is a compact metric space. It follows from [2, Theorem 2.12] that every almost limited weakly null sequence in E is norm null. Consequently, $B_{E'}$ is a strong L-limited set.

REMARK 3.3. Let E be a Banach lattice and let Y be a Banach space. A bounded linear operator $T: E \rightarrow Y$ is alcc if and only if $T'(B_{Y'})$ is a strong L-limited set in E' .

In [2] the authors showed that if E' has a weak unit or if E has order continuous norm, then E has the sGP if and only if every almost limited

weakly null sequence is norm null. From this fact, the following result is immediate.

PROPOSITION 3.4. *Assume that E' has a weak unit or E has order continuous norm. Then E has the sGP property if and only if $B_{E'}$ is a strong L-limited set.*

As a consequence of Proposition 3.4, we have that $B_{L_p[0,1]}$ with $1 < p \leq \infty$ is not strong L-limited, even though they are L-limited (see [16, Theorem 2.3]). Moreover, since c_0 and ℓ_p with $1 \leq p < \infty$ have order continuous norm and the sGP property (see [2, p. 17]), Proposition 3.4 yields that B_{ℓ_p} with $1 \leq p \leq \infty$ is a strong L-limited set.

We observe that $B_{L_\infty[0,1]} = \text{sol}(\{\chi_{[0,1]}\})$. Thus, the solid hull of a strong L-limited set is not necessarily strong L-limited. In the next Theorem, we give conditions on E so that the solid hull of a strong L-limited also is strong L-limited.

THEOREM 3.5. *Let E be a Banach lattice with order continuous norm. If the lattice operations in E are sequentially weakly continuous, then the solid hull of a strong L-limited subset of E' also is strong L-limited.*

PROOF. Let $A \subset E'$. Assume that $\text{sol}(A)$ is not strong L-limited. Then there exists an almost limited weakly null sequence $(x_n) \subset E$ such that $\sup_{x' \in \text{sol}(A)} |x'(x_n)| \geq \varepsilon$ for all n and for some $\varepsilon > 0$. For each n , there exists $x'_n \in \text{sol}(A)$ such that $|x'_n(x_n)| \geq \varepsilon$. In particular, for each n , we can find $y'_n \in A$ such that $|x'_n| \leq |y'_n|$. On the other hand, by [1, Theorem 1.23], there exists $(z''_n) \subset E''$ such that $|z''_n| \leq J_E(|x_n|)$ in E'' and $z''_n(y'_n) = J_E(|x_n|)(y'_n)$ for every n . Since E has order continuous norm, E is an ideal of E'' (see [1, Theorem 4.9]). Then for each n , there exists $z_n \in E$ such that $z''_n = J_E(z_n)$. Thus

$$\varepsilon \leq |x'_n(x_n)| \leq |x'_n|(|x_n|) \leq |y'_n|(|x_n|) = z''_n(y'_n) = y'_n(z_n)$$

holds for all n . Since $|z_n| \leq |x_n|$ for all n and $|x_n| \xrightarrow{\omega} 0$ in E , by assumption we have that $z_n \xrightarrow{\omega} 0$ in E . As E has order continuous norm, it has property (d), by Lemma 2.1, (z_n) is almost limited. Hence A cannot be strong L-limited, a contradiction. \square

It is known that every weakly compact operator is lcc (see [15, Corollary 2.5]). As a consequence, every relatively weakly compact set in the dual of a Banach space is L-limited (see [16]). We observe that, if E is a non discrete reflexive Banach lattice (e.g. $L_p[0,1]$ with $1 < p < \infty$), then E does not have the sGP property by [2, Theorem 2.8]. On the other hand, as E has order continuous norm, we have that I_E cannot be alcc. As a consequence, $B_{E'}$ is a relatively weakly compact subset of E' which is not strong L-limited. In Proposition 3.6 and Proposition 3.7, we establish connections

between strong L-limited sets, relatively weakly compact sets, alcc operators and weakly compact operators.

PROPOSITION 3.6. *Let E be a Banach lattice. The following assertions are equivalent:*

1. *Every relatively weakly compact set in E' is strong L-limited.*
2. *For every Banach space Y , every weakly compact operator $T: E \rightarrow Y$ is alcc.*

PROOF. (1) \Rightarrow (2). Let $T: E \rightarrow Y$ be a weakly compact operator. So $T'(B_{Y'})$ is relatively weakly compact in E' and, as consequence, a strong L-limited set. Therefore T is an alcc operator.

(2) \Rightarrow (1). Assume that there exists a relatively weakly compact set $A \subset E'$ that is not strong L-limited. Without loss of generality, we can assume that there exists $\varepsilon > 0$, $(x'_n) \subset A$ and an almost limited weakly null sequence such that $|x'_n(x_n)| \geq \varepsilon$ for all n . Since A is relatively weakly compact, there exists a subsequence (x'_{n_k}) which converges weakly to $x' \in E'$. Define the weakly compact operator $T: E \rightarrow c_0$ by $T(x) = ((x'_{n_k} - x')(x))_n$ (see [1, Theorem 5.26]). On the other hand, let $k_0 \in \mathbb{N}$ such that $|x'(x_{n_k})| < \varepsilon/2$ for all $k \geq k_0$. If $k \geq k_0$,

$$\|Tx_{n_k}\| \geq |x'_{n_k}(x_{n_k}) - x'(x_{n_k})| \geq \varepsilon/2.$$

Thus T is not an alcc operator. \square

Using the same arguments of [16, Theorem 2.8] and doing the necessary adaptations, we can prove the following Proposition.

PROPOSITION 3.7. *Let E be a Banach lattice. The following assertions are equivalent:*

1. *Every strong L-limited set in E' is relatively weakly compact.*
2. *For every Banach space Y , every alcc operator $T: E \rightarrow Y$ is weakly compact.*

From Propositions 3.6 and 3.7, we get the next corollary.

COROLLARY 3.8. *Let E be a Banach lattice. The class of strong L-limited sets coincides with the class of relatively weakly compact sets in E' if and only if $L_{cc}^a(E; Y) = W(E; Y)$, for every Banach space Y .*

Recall that a linear operator $T: E \rightarrow F$ is said to be order bounded if it takes order bounded sets from E onto order bounded sets in F . In the next result, we prove that every order bounded linear operator from a Banach lattice with property (d) into a Banach lattice with the sGP property is alcc.

PROPOSITION 3.9. *Let E and F be Banach lattices with property (d). Let $T: E \rightarrow F$ be an order bounded linear operator. If F has the sGP property, then T is an alcc operator.*

PROOF. Let $(x_n) \subset E$ be an almost limited weakly null sequence and let $A = \text{sol}((x_n)_n)$. By Lemma 2.1 and Theorem 2.3, we have that $T(A)$ is almost limited. As $(Tx_n) \subset T(A)$, (Tx_n) is an almost limited sequence in F . Then $Tx_n \xrightarrow{\omega} 0$ in F , and so $Tx_n \rightarrow 0$ in Y since F has the sGP property. \square

Now, we give some remarks concerning Proposition 3.9.

REMARKS 3.10. 1. It is known that for every $1 \leq p \leq \infty$, the Banach lattice $E = L_p[0, 1]$ has property (d). Therefore, if F is any Banach lattice with both property (d) and the sGP property, we get from Proposition 3.9 that $L^r(E; F) \subset L^a cc(E; F)$.

2. Let E be a Banach lattice that contains a lattice copy of ℓ_1 . Then there exists a positive projection $P: E \rightarrow \ell_1$ (see [13, Proposition 2.3.11]). As a consequence of Proposition 3.9, we get that P is an alcc operator. In particular, Banach lattices with the positive Schur property contains lattice copy of ℓ_1 (see [18, p. 19]). Besides, we get from Remark 3.3 that $P'(B_{\ell_\infty})$ is a strong L-limited set in E' .

3. There is a Banach lattice F with property (d) which does not have the sGP property, such that, for any Banach lattice E , every order bounded operator $T: E \rightarrow F$ is alcc. Indeed, let K be a compact metric space and assume that K is σ -Stonian (i.e. the closure of every open F_σ set is open). It follows from [2, Theorem 2.12] that $F = C(K)$ does not have the sGP property. On the other hand, [13, Proposition 2.1.4] yields that $F = C(K)$ is σ -Dedekind complete, so it has property (d). Given a Banach lattice E , we claim that every bounded operator $T: E \rightarrow F$ is completely continuous, hence alcc. Indeed, if $x_n \xrightarrow{\omega} 0$ in E , we have that $Tx_n \xrightarrow{\omega} 0$ in F . In particular, (Tx_n) is a bounded sequence. As $C(K)$ has a unit, (Tx_n) is contained in an order interval. Since $C(K)$ has property (d), it follows from Proposition 2.2 that order intervals are almost limited sets. Consequently, (Tx_n) is an almost limited weakly null sequence. [2, Theorem 2.12] yields that $\|Tx_n\| \rightarrow 0$.

The next example shows that the assumption of T being order bounded linear operator in Proposition 3.9 is essential.

EXAMPLE 3.11. Let (r_k) denote the Rademacher functions and let $T: L_1[0, 1] \rightarrow c_0$ be the operator given by

$$T(f) = \left(\int_0^1 f(t) r_k(t) dt \right)_k.$$

We observe that T is well-defined since $r_n \xrightarrow{\omega} 0$ in $L_1[0, 1]$ and that T is not an alcc operator, once the sequence (r_n) is almost limited weakly null

in $L_1[0, 1]$, but

$$\|T(r_n)\|_\infty = \sup_{k \in \mathbb{N}} \left| \int_0^1 r_n(t) r_k(t) dt \right| \geq 1 \quad \text{for all } n \in \mathbb{N}.$$

We claim that T is not an order bounded operator. Indeed, as c_0 is Dedekind complete, it suffices to check that T is not regular. Suppose that T is regular. So we assume that there exists $S \geq 0$ with $S \geq T$. The components of S lead to a weak* null positive sequence $(\varphi_n) \subset (L_1[0, 1])'$ such that $S(f) = (\varphi_n(f))_n$. Since $(L_1[0, 1])' = L_\infty[0, 1]$, there exists $(f_n) \subset (L_\infty[0, 1])^+$ with $\varphi_n(f) = \int_0^1 f_n(t) f(t) dt$. Since $T \leq S$, we get that $r_n \leq f_n$ in $L_\infty[0, 1]$ which means that $f_n \geq r_n^+$. But

$$\varphi_n(1) = \int_0^1 f_n(t) dt \geq \int_0^1 r_n^+(t) dt = 1/2$$

for all n , a contradiction.

In the class of the positive linear operators in Banach lattices, we have a dominated type problem. For instance, let $S, T: E \rightarrow F$ be positive operators such that $S \leq T$. The question is, if T has some property, does S also has it? Here we are interested in the class of alcc operators and we see that in general the answer is negative, as the next example shows.

EXAMPLE 3.12. Let $R_1: c_0 \rightarrow \ell_\infty$, $S_1, S_2: L_1[0, 1] \rightarrow \ell_\infty$ be given by

$$R_1(\alpha_i)_i = (\alpha_i)_i,$$

$$S_2(f) = \left(\int_0^1 f dx, \int_0^1 f dx, \dots \right) \quad \text{and} \quad S_1(f) = \left(\int_0^1 f r_i^+(x) dx \right)_i.$$

Define $S, T: c_0 \oplus L_1[0, 1] \rightarrow \ell_\infty$ by

$$S((\alpha_i), f) = R_1((\alpha_i)) + S_1(f) \quad \text{and} \quad T((\alpha_i), f) = R_1((\alpha_i)) + S_2(f).$$

Note that $0 \leq S \leq T$.

We claim that T is an alcc operator. If $(x_n, f_n) \subset c_0 \oplus L_1[0, 1]$ is an almost limited weakly null sequence, then $(x_n) \subset c_0$ and $(f_n) \subset L_1[0, 1]$ are both almost limited weakly null sequences. As c_0 has the sGP property, we have that $\|x_n\|_\infty \rightarrow 0$. So,

$$\|R_1(x_n)\| = \|x_n\|_\infty \rightarrow 0.$$

On the other hand, as $f_n \xrightarrow{\omega} 0$ in $L_1[0, 1]$, it follows that $\int_0^1 f_n \rightarrow 0$, what implies that

$$\|S_2(f_n)\|_\infty = \sup_i \left| \int_0^1 f_n \right| = \left| \int_0^1 f_n \right| \rightarrow 0.$$

Therefore

$$T(x_n, f_n) = R_1(x_n) + S_2(f_n) \rightarrow 0.$$

It is easy to verify that S is not an alcc operator. In fact, $(r_n) \subset L_1[0, 1]$ is an almost limited weakly null sequence. However, $\|S_1(r_n)\|_\infty \geq 1/2$ for all n . Hence $(0, r_n)$ is an almost limited weakly null sequence in $c_0 \oplus L_1[0, 1]$ such that $S(0, r_n) = S_1(r_n)$ that is not norm null.

Also note that R_1 is not completely continuous, since $e_n \xrightarrow{\omega} 0$ in c_0 and $\|e_n\| = 1$. Hence neither S nor T is completely continuous.

We observe that the order bounded linear operator $S_1 : L_1[0, 1] \rightarrow \ell_\infty$ given in Example 3.12 is a positive non alcc operator. This shows that the assumption of F having the sGP property in Proposition 3.9 is essential.

The following result was inspired in [1, Theorem 5.89].

THEOREM 3.13. *Let E and F be Banach lattices and let $S, T : E \rightarrow F$ be positive operators such that $S \leq T$ and T is an alcc operator. If E has property (d) and the lattice operations in E are weakly sequentially continuous, then S is an alcc operator.*

PROOF. Let (x_n) be an almost limited weakly null sequence. By Lemma 2.1, $\text{sol}((x_n)_n)$ is an almost limited set in E . In particular, $(|x_n|)$ is an almost limited sequence. On the other hand, since the lattice operations in E are weakly sequentially continuous, $|x_n| \rightarrow 0$. As T is alcc, $T|x_n| \rightarrow 0$. Thus $Sx_n \leq S|x_n| \leq T|x_n| \rightarrow 0$. \square

The following theorem was inspired by Kalton–Saab theorem's ([1, Theorem 5.90]).

THEOREM 3.14. *Let E and F be Banach lattices and let $S, T : E \rightarrow F$ be positive operators such that $S \leq T$ and T is an alcc operator. If E is σ -Dedekind complete and F has order continuous norm, then S is an alcc operator.*

PROOF. Let $(x_n) \subset E$ be an almost limited weakly null sequence. We want to prove that $Sx_n \rightarrow 0$ in F . Let $A = \text{sol}((x_n)_n)$. We observe that for every disjoint sequence $(y_n) \subset A$, $\|Ty_n\| \rightarrow 0$. [Indeed, by [1, Theorem 3.34], $y_n \xrightarrow{\omega} 0$ in E . However, since A is almost limited (Lemma 2.1), we get that (y_n) is an almost limited weakly null sequence and the statement holds since T is alcc.] Therefore, given $\varepsilon > 0$, by [1, Theorem 4.36], we have that there exists $0 \leq u \in E_A$ such that

$$(1) \quad \|T((|x_n| - u^+))\| < \varepsilon/6 \quad \text{for all } n \in \mathbb{N}.$$

Now, since E is σ -Dedekind complete and F has order continuous norm, by Theorem 4.87 of [1], there exist positive operators $M_1, \dots, M_k : E \rightarrow E$

and $P_1, \dots, P_k: F \rightarrow F$ such that $0 \leq \sum_{i=1}^k P_i T M_i \leq T$ on E and

$$(2) \quad \left\| \left| S - \sum_{i=1}^k P_i T M_i \right| (u) \right\| < \varepsilon/3.$$

We claim that $T M_i(x_n) \rightarrow 0$, where $n \rightarrow \infty$ for all $i = 1, \dots, k$. Since T is an alcc operator, it suffices to prove that $(M_i(x_n))$ is an almost limited weakly null sequence in E . As E has property (d), $A \subset E$ is a solid almost limited set, $M_i: E \rightarrow E$ is a positive operator and $M_i(x_n) \xrightarrow{\omega} 0$ in E . It follows from Theorem 2.3 that $M_i(A)$ is an almost limited subset of E . In particular, $(M_i(x_n))_n$ is an almost limited weakly null sequence for each $i = 1, \dots, k$ as we stated. Consequently,

$$\left\| \sum_{i=1}^k P_i T M_i(x_n) \right\| \leq \sum_{i=1}^k \|P_i\| \|T M_i(x_n)\| \rightarrow 0.$$

Thus, there exists $n_0 \in \mathbb{N}$ such that

$$(3) \quad \left\| \sum_{i=1}^k P_i T M_i(x_n) \right\| < \varepsilon/3 \quad \text{for all } n \geq n_0.$$

Finally, by using the same argument used in the proof of Kalton–Saab's theorem ([1, Theorem 5.90]), we get that $\|Sx_n\| < \varepsilon$ for all $n \geq n_0$. Here, we give a short version of the proof. Since $0 \leq S \leq T$ and $0 \leq \sum_{i=1}^k P_i T M_i \leq T$, we have that

$$\begin{aligned} & \left| S(x_n) - \sum_{i=1}^k P_i T M_i(x_n) \right| \\ & \leq \left| S - \sum_{i=1}^k P_i T M_i \right| (|x_n| - u)^+ + \left| S - \sum_{i=1}^k P_i T M_i \right| (u) \\ & \leq 2T(|x_n| - u)^+ + \left| S - \sum_{i=1}^k P_i T M_i \right| (u). \end{aligned}$$

Now, from (1) and (2),

$$\left\| S(x_n) - \sum_{i=1}^k P_i T M_i \right\| \leq 2\varepsilon/3.$$

Finally, $\|Sx_n\| < \varepsilon$ for all $n \geq n_0$ holds from above inequality and (3). \square

Next, we will study when an alcc operator is completely continuous or compact. Since every compact operator is completely continuous, it follows that $K(E; Y) \subset Cc(E; Y) \subset L^a cc(E; Y)$. If every alcc operator is a compact operator, we have that $K(E; Y) = Cc(E; Y) = L^a cc(E; Y)$. So $Cc(E; Y) = L^a cc(E; Y)$ must be a necessary condition for $K(E; Y) = L^a cc(E; Y)$. In the next proposition we give conditions so that these spaces are distinct.

PROPOSITION 3.15. *Let E be an infinite dimensional reflexive lattice with the sGP property. Then there exists a Banach space Z such that $K(E; Z) \neq L^a cc(E; Z)$.*

PROOF. By above commentary, E cannot have the wDP* property and hence it neither can have the DP* property. By Theorem 2.8 of [15], there exists a Banach space Z such that $Cc(E; Z) \subsetneq Lcc(E; Z)$. On the other hand, as E has the sGP property, then $L^a cc(E; Y) = Lcc(E; Y) = L(E; Y)$ for all Banach space Y . It follows that $K(E; Z) = Cc(E; Z) \neq L^a cc(E; Z)$. \square

It is known that if E has the wDP* property, then every alcc operator on E is completely continuous (see [2, p. 22]). Using Proposition 3.9, we will show that if E has the property (d) and if every alcc operator on E is completely continuous, then E has the wDP* property.

PROPOSITION 3.16. *Let E be Banach lattice with property (d). If $Cc(E; F) = L^a cc(E; F)$ for all Banach lattices F , then E has the wDP* property.*

PROOF. Assume that E has property (d) and suppose by contradiction that E does not have the wDP* property. By [12, Theorem 3.1], there exist $(x_n) \subset E^+$ a weakly null sequence and $(f_n) \subset (E')^+$ a weak* null sequence such that $f_n(x_n) \geq \varepsilon$ for all n and some $\varepsilon > 0$. Consider the positive linear operator $T: E \rightarrow c_0$ defined by $T(x) = (f_n(x))_n$ for all $x \in E$, by Proposition 3.9 we have that T is an alcc operator. But T is not completely continuous, once $\|Tx_n\| \geq |f_n(x_n)| \geq \varepsilon$ for all n . \square

[10, Corollary 5] shows that every completely continuous linear operator on a Banach space X is compact if and only if X does not contain an isomorphic copy of ℓ_1 . In particular, if E is a Banach lattice that does not contain copy of ℓ_1 and if E has the wDP* property, then every alcc operator on E must be compact.

Recall that a linear operator $T: E \rightarrow Y$ is called *AM-compact* if it maps order intervals in E to relatively compact sets in Y . In the next proposition, we study a relation between the class of alcc operators and the class of AM-compact operators.

PROPOSITION 3.17. *Let E and F be two Banach lattices. If E has property (d), then every alcc operator $T: E \rightarrow F$ is AM-compact.*

PROOF. Let $T: E \rightarrow F$ be an alc operator and let $A \subset E$ be an order interval. It follows from Proposition 2.2 that A is an almost limited set, and so $T(A)$ is a relatively compact subset of F (see [2, Theorem 3.2]). Thus T is an AM-compact operator. \square

Acknowledgement. The authors are thankful to the referee for careful reading and considered suggestions leading to a better paper.

References

- [1] C. Aliprantis and O. Burkinshaw, *Positive Operators*, Springer (Dordrecht, 2006).
- [2] H. Ardakani, S. M. S. M. Mosadegh, M. Moshtaghioun and M. Salimi, The strong Gelfand–Phillips property in Banach lattices, *Banach J. Math. Anal.*, **10** (2016), 15–26.
- [3] K. Bouras, Almost Dunford–Pettis sets in Banach lattices, *Rend. Circ. Mat. Palermo*, **62** (2013), 227–236.
- [4] J. Bourgain and J. Diestel, Limited operators and strict cosingularity, *Math. Nachr.*, **119** (1984), 55–58.
- [5] H. Carrión, P. Galindo, M. L. Lourenço, A stronger Dunford–Pettis property, *Studia Math.*, **184** (2008), 205–2016.
- [6] J. X. Chen, Z. L. Chen and G. X. Ji, Almost limited sets in Banach lattices, *J. Math. Anal. Appl.*, **412** (2014), 547–553.
- [7] L. Drewnowski, On Banach spaces with the Gelfand–Phillips property, *Math. Z.*, **193** (1986), 405–411.
- [8] A. Elbour, Some characterizations of almost limited operators, *Positivity*, **21** (2017), 865–874.
- [9] M. Fabian, P. Habala, P. Hajék, V. Montesinos and V. Zizler, *Banach Space Theory. The Basis for Linear and Nonlinear Analysis*, Springer (2011).
- [10] J. M. Gutiérrez, Weakly continuous functions on Banach spaces not containing ℓ_1 , *Proc. Amer. Math. Soc.*, **119** (1993), 147–152.
- [11] J. H'michane, N. Hafidi and L. Zraoula, On the class of disjoint limited completely continuous operators, *Positivity*, **22** (2018), 1419–1431.
- [12] N. Machrafi, A. Elbour and M. Moussa, Some characterizations of almost limited sets and applications, arXiv:1312.2770v2 (2014).
- [13] P. Meyer-Nieberg, *Banach Lattices*, Springer-Verlag (1991).
- [14] M. L. Lourenço and V. C. C. Miranda, A note on the Banach lattice $c_0(\ell_n^2)$ and its dual, *Carpathian Math. Publ.* (2023), to appear.
- [15] M. Salimi and M. Moshtaghioun, The Gelfand–Phillips property in closed subspaces of some operator spaces, *Banach J. Math. Anal.*, **5** (2011), 84–92.
- [16] M. Salimi and M. Moshtaghioun, A new class of Banach spaces and its relation with some geometric properties of Banach spaces, *Abstr. Appl. Anal.* (2012), 1085–1092.
- [17] C. Stegall, Duals of certain spaces with the Dunford–Pettis property, *Notices Amer. Math. Soc.*, **19** (1972), 799.
- [18] W. Wnuk, Banach lattices with properties of the Schur type: a survey, *Conf. Sem. Mat. Univ. Bari*, **249** (1993), 1–25.
- [19] W. Wnuk, *Banach Lattices with Order Continuous Norms*, Polish Scientific Publishers PWN (Warsaw, 1999).
- [20] W. Wnuk, On the dual positive Schur property in Banach lattices, *Positivity*, **17** (2013), 759–773.