

# A history of over 40 years of potentially pathogenic free-living amoeba studies in Brazil – a systematic review

Natália Karla Bellini<sup>1/+</sup>, Otavio Henrique Thiemann<sup>2,3</sup>, María Reyes-Batlle<sup>4</sup>, Jacob Lorenzo-Morales<sup>4,5</sup>, Adriana Oliveira Costa<sup>1/+</sup>

<sup>1</sup>Universidade Federal de Minas Gerais, Faculdade de Farmácia, Departamento de Análises Clínicas e Toxicológicas, Belo Horizonte, MG, Brasil

<sup>2</sup>Universidade de São Paulo, Instituto de Física de São Carlos, São Carlos, SP, Brasil

<sup>3</sup>Universidade Federal de São Carlos, Departamento de Genética e Evolução, São Carlos, SP, Brasil

<sup>4</sup>Universidad de La Laguna, Instituto Universitario de Enfermedades Tropicales y Salud Pública de Canarias, Departamento de Obstetricia, Ginecología, Pediatría, Medicina Preventiva y Salud Pública, Toxicología, Medicina Legal y Forense y Parasitología, Red de Investigación Cooperativa en Enfermedades Tropicales, Tenerife, Islas Canarias, Spain

<sup>5</sup>Instituto de Salud Carlos III, Consorcio Centro de Investigación Biomédica en Red MP de Enfermedades Infecciosas, Madrid, Spain

Free-living amoeba (FLA) group includes the potentially pathogenic genera *Acanthamoeba*, *Naegleria*, *Balamuthia*, *Sappinia*, and *Vermamoeba*, causative agents of human infections (encephalitis, keratitis, and disseminated diseases). In Brazil, the first report on pathogenic FLA was published in the 70s and showed meningoencephalitis caused by *Naegleria* spp. FLA studies are emerging, but no literature review is available to investigate this trend in Brazil critically. Thus, the present work aims to integrate and discuss these data. Scopus, PubMed, and Web of Science were searched, retrieving studies from 1974 to 2020. The screening process resulted in 178 papers, which were clustered into core and auxiliary classes and sorted into five categories: wet-bench studies, dry-bench studies, clinical reports, environmental identifications, and literature reviews. The papers dating from the last ten years account for 75% (134/178) of the total publications, indicating the FLA topic has gained Brazilian interest. Moreover, 81% (144/178) address *Acanthamoeba*-related matter, revealing this genus as the most prevalent in all categories. Brazil's Southeast, South, and Midwest geographic regions accounted for 96% (171/178) of the publications studied in the present work. To the best of our knowledge, this review is the pioneer in summarising the FLA research history in Brazil.

Key words: free-living amoeba - Brazil - literature review - *Acanthamoeba* - *Naegleria* - *Balamuthia*

Free-living amoeba (FLA) is a protozoan group including Excavata and Amoebozoa lineages capable of living free in the environment and alternatively proliferating within a host, therefore named amphizoic amoebas.<sup>(1)</sup> *Acanthamoeba* spp., *Naegleria fowleri*, *Sappinia pedata*, *Balamuthia mandrillaris*, and more recently *Vermamoeba vermiformis*, include pathogenic amoebae causative agents of human infections.<sup>(1,2,3,4)</sup>

FLAs have been ubiquitously isolated around the globe from both natural and artificial environments.<sup>(5,6)</sup> These organisms share common aspects in their life cycle as the existence of a trophozoite stage able to feed, divide, and move by pseudopodium projection and constrictions of the cytoplasm. Another stage is the cyst, which can

withstand adverse conditions as food scarcity, unbalance in pH, salt concentration, and temperature.<sup>(6)</sup> *Naegleria* spp. possess an additional flagellate transient form that can escape from harsh environmental conditions.<sup>(1,7)</sup>

FLAs have attracted the interest of human health agencies due to their involvement as opportunistic and non-opportunistic infections that affect the central nervous system (CNS), the cornea, and other organs. CNS infections can be classified either as Primary Amoebic Meningoencephalitis (PAM), an acute infection, caused by *N. fowleri*<sup>(8)</sup> or Granulomatous Amoebic Encephalitis (GAE), a subacute to chronic illness caused by *Acanthamoeba* spp. and *B. mandrillaris*.<sup>(2,9)</sup> *Acanthamoeba* spp. and *B. mandrillaris* have also been identified in skin lesions and disseminated infections that, similarly to GAE, affect predominantly debilitated or immunocompromised patients.<sup>(10)</sup> FLAs can also affect the cornea, causing a progressive, sight-threatening infection termed amoebic or *Acanthamoeba* keratitis (AK), which has *Acanthamoeba* spp. as the main etiological agent.<sup>(2,11,12)</sup> *S. pedata* accounts for a single case of encephalitis in humans,<sup>(13,14)</sup> while *V. vermiformis* has been mainly reported in keratitis cases, most of them in co-infection with *Acanthamoeba*.<sup>(4)</sup> Despite the widespread recognition of FLAs as pathogens, infections still result in several deaths or events of visual impairment. These outcomes are mainly associated with the lack of fast and reliable diagnostic methods and effective treatments.<sup>(15,16)</sup> Besides their importance as human pathogens, FLAs have been investigated in relation to their ecological interaction with the aquatic and soil microbial community.

doi: 10.1590/0074-02760210373

Financial support: OHT had financial support from Biota-Fapesp Program (Process # 2018/20693-4); JLM had support from Red de Investigación Cooperativa en Enfermedades Tropicales - RICET, Spain (project # RD16/0027/0001, Programme Redes Temáticas de Investigación Cooperativa, FIS); Consorcio Centro de Investigación Biomédica en Red - CIBER de Enfermedades Infecciosas, Instituto de Salud Carlos III, Spain (Project # CB21/13/00100); Ministerio de Sanidad, Gobierno de España; and Cabildo de Tenerife, Tenerife innova, Marco Estratégico de Desarrollo Insular-MEDI, Fondo de Desarrollo de Canarias -FDCAN (Project # 21/0587).

+ Corresponding authors: nkbellini@gmail.com / adrincosta@gmail.com

https://orcid.org/0000-0003-2408-3210

https://orcid.org/0000-0002-1793-3013

Received 22 November 2021

Accepted 28 March 2022



Previous analyses have shown FLA harboring intracellular organisms from the ambient, proposing the amoeba as a vehicle to bacteria, fungi, protozoan, and viruses, the so-called amoeba resistant microorganisms (ARM).<sup>(17,18)</sup> Through lysing cells, ARMs can disseminate in the environment or a host. Therefore, FLA is referred to as Trojan horses for the microbial community,<sup>(17,19)</sup> emphasising the significance of investigating its ecological importance throughout the world.<sup>(18)</sup>

Water-related outbreaks caused by protozoans in Latin America were revised in a survey that pointed out the Brazilian prominence accounting for 30% of the reports (20/66). Comparing the causative agents of these outbreaks, *Acanthamoeba* and *Cyclospora* shared the fourth position in the ranking.<sup>(20)</sup> Environmental isolation of FLA in the Brazilian territory relies on recent reports that call attention to the country's potential on harboring FLAs. Efforts to address gaps in FLA knowledge include developing more reliable diagnosis,<sup>(21,22)</sup> investigation of factors related to pathogenicity,<sup>(23,24)</sup> and development of therapeutic approaches.<sup>(25,26)</sup> Those are just a portion of the literature produced in Brazil on the FLA field, and controversially, there is no literature review devoted to summing up and debating these data critically. Thus, the following central question guided the present literature review: What is the Brazilian research contribution to the free-living amoeba literature until 2020?

*Identification, systematic documentation, and screening of the data* - The data analysis followed the general workflow as suggested by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline.<sup>(27)</sup> Fourteen keywords were defined [Supplementary data (Table II)] and used as queries in the databases Scopus, Web of Science, and PubMed/Medline, all accessed through the CAPES News Portal ([www.periodicos.capes.gov.br](http://www.periodicos.capes.gov.br)) between March 22 to 24, 2021. We have defined 13 keywords (#1 to #13) and a geographical region of interest (#14) to select topics of interest [Supplementary data (Table II)].

The search entry varied according to the database, as described in the Supplementary data (Table II), providing an initial dataset composed of 1512 articles (Fig. 1). We managed the literature dataset using Mendeley version 1.17.10<sup>(28)</sup> and State of the Art through Systematic Review (StArt) version 3.4 beta.<sup>(29)</sup> The StArt tool allowed to identify and subtract duplicate references, corresponding to literature overlaps intra-database and inter-databases. We designed inclusion and exclusion criteria [Supplementary data (Table II)] to accept and reject papers in the first round of data examination. The keywords listed in Supplementary data (Table II) were contrasted against the title, affiliation, and abstract sections of the 475 records selected for screening. We have withdrawn 277 of them (Fig. 1 - Screening step). Next, we examined the main text of the remaining paper according to Supplementary data (Table II - III) criteria. Study limitations included: records available in the gray literature (e.g., google scholar) are absent; the period cutoff excluding the literature made public in 2021; dissertations, letters, and protocols are also absent in this review. Following litera-

ture recommendations concerned with diminishing study bias, no language restrictions were applied.<sup>(30)</sup> The screening and application of inclusion/exclusion criteria resulted in 178 articles gathering the Brazilian research contribution to the free-living amoeba literature (Fig. 1).

Data analysis considered (1) the geographic scope and (2) research domains of the Brazilian literature on FLA topics. A primary analysis exposed a need for two clustering levels concerning the research domain. Based on the central research question addressed in each study we divided the first level into core and auxiliary classes. We settled as *core* the investigations with FLAs as the major topic addressed in all sections of the paper. We sorted some studies including FLAs in any part of the main text, but as a complement to other topics, as *auxiliary*. The second level of clustering consisted of five research domains in which both core/auxiliary papers were categorised: *dry-bench*, *wet-bench*, *clinical*, *environmental* and *review*.

Publications categorised as wet-bench included *in vitro* or *in vivo* procedures conducted with isolates or cultures without a primary intention of prospecting the environment (environmental) or describing cases (clinical). Dry-bench grouped the bioinformatics-based research whose experimental approach is *in silico*. And review account to literature reviews. We assessed the main text of each paper, and we added the additional two layers of clustering to the data extraction forms. Two researchers have conducted the eligibility checking and data extraction processes to reinforce decisions taken during both steps. We have solved in consensus any uncertainty related to studies inclusion, exclusion, and clustering.




Fig. 1: Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow chart describing the protocol employed in the systematic review and the number of citations (n) retrieved in each step.

The first and second levels of clustering were displayed in the number or percentage of papers and sorted according to date of publication, federative unit, and FLA species. Occasionally, papers encompassing multiple federative units' affiliation were scored numerous times and obtained percentages exceeding 100%. However, regardless of the number of authors from the same federative unit, we gave the "1" score.

*FLA research in Brazil had an expansion in the last ten years, and the Southern, South, and Midwest regions lead the contribution* - The FLA topic is an emerging field in Brazil, in which 134 papers from a total of 178 papers (~ 75%) were published in the last ten years of the 1974 to 2020 period (Fig. 2A). The remaining 44 papers (~ 25%) are dated in a time window of 37 years from this period. The data indicate that after 2008, FLA knowledge increased progressively.

Papers classified as the core class accounted for 120 articles predominantly originating from Brazilian scientific institutions in the Southeast (SP, MG, ES, RJ), South (SC, PR, RS), and the Midwest (GO, DF, MS) regions, named here with the acronym **SSM** (Fig. 2B). Although the Southeast region leads the ranking, with ~ 64% of the total reports, the state of Rio Grande do Sul (RS) represents the federative unit with more contributions. The map has also shown studies from the North (PA) and Northeast (PB, and SE) regions with lower frequencies. Overall, it

depicts half of the country (13 out of 26 federative unities) contributing on FLA topics sorted as core (Fig. 2B).

A total of 58 articles were classified in the auxiliary class (Fig. 2C). Most of them were originated in universities in the Southeast region, resembling the tendency perceived for the core class of papers (Fig. 2B). Altogether, FLA studies in Brazil have a predominant origin in centres from the SSM region (96% - 171/178). Factors explaining this regional concentration are the heterogeneous funding distributions<sup>(31)</sup> and spatial distributions of universities through the country.<sup>(32)</sup> Regardless of this trend, the overall landscape of FLA research in Brazil has highlighted half of the federative unities being committed to expanding the knowledge on FLA, therefore reinforcing the country's relevance as a thriving contributor to the field.

*Wet-bench research predominates in Brazilian FLA literature* - Five research domains composed the second level of clustering in the core and auxiliary classes of papers, as shown in Fig. 3. The research domain classified as wet-bench represents 58 out of 120 core class papers (Fig. 3A), mainly with studies addressing the biochemical characterisation of FLA strains isolated by the authors and commercial lineages (e.g., ATCC strains). Regarding the territorial distribution, wet-bench investigations reproduced the prevalence of the SSM region, in which Rio Grande do Sul (RS) and São Paulo (SP) occupied the first




Fig 2: progression and distribution of free-living amoeba (FLA) studies in Brazil. A - Bar graph with the number of core and auxiliary classes of publications per year representing the first level of clustering. Geographical distribution of core (B) and auxiliary (C) classes of papers according to federative units declared in the affiliations. A single paper with multiple federative unities affiliation was scored once per federative unit so that the total percentage resulted higher than 100% in both B and C panels.

and second positions in the ranking with 40.7% and 22% of the published papers, respectively (Fig. 3B).

The higher occurrence of clinical assigned reports is on papers describing keratitis (18/29), followed by encephalitis (7/29). Fewer reporting cases of disseminated diseases (1/29), skin infections (1/29), besides the presence of FLA in urine and feces (2/29), were reported. Summarising the Brazilian publications on FLA distribution, the environmental category identified 25 out of 120 papers in the core class.

The remaining 58 papers sorted in the auxiliary class included 60% (n = 34) of wet-bench research [Supplementary data (Figure)], reinforcing the predominance of this research domain in FLA national studies. Review, clinical and environmental tags account for 13, 6, and 5 papers, respectively, out of the 58 in this class, while no study was classified as dry-bench. Once again, the Southeast region leads the number of studies, covering topics ranging from waterborne protozoa to more specific issues as virus replication.

*Acanthamoeba leads the ranking of FLA studies in Brazil* - The *Acanthamoeba* genus was predominant in 81% (144/178) of papers [Supplementary data (Table I)], a tendency depicted in clinical, environmental, wet-bench and review categories of both the core and auxiliary classes (Fig. 4). For the dry-bench category, both *Acanthamoeba* and *Naegleria* shared the same number of publications (one paper each) in the core class.

To a lesser extent, some papers addressed two FLA genera in a single research, as revealed by combinations of *Vermamoeba* and *Acanthamoeba* in the clinical category, as well as *Naegleria* and *Acanthamoeba* in the environmental category, both within the core class (Fig. 4). Multiple emphases occurred in papers discussing more than two FLA genera in a single paper, for example, *Naegleria*, *Acanthamoeba*, and *Balamuthia* identified in clinical publications of the auxiliary class (Fig. 4). Regardless of the single or multiple FLA genus in the same article, those classified in the environmental category accounted for *Acanthamoeba*, *Vermamoeba*, *Balamuthia*, and *Naegleria* reports in the Brazilian territory (Fig. 4). *Sappinia* was mentioned only in two papers classified as reviews in the auxiliary class.<sup>(33,34)</sup> The predominance of *Acanthamoeba*-related investigations is an expected finding considering that this species is the most frequent FLA in the environment<sup>(2)</sup> and presents higher viability than *Naegleria* in dust samples.<sup>(35)</sup> In addition, while other pathogenic species of FLAs are involved in rare brain infections, *Acanthamoeba* can cause both cerebral and corneal AK infections, the latter showing an increasing number of cases in recent years.<sup>(2)</sup>

*Diversity of approaches in wet-bench references* - The wet-bench category included approaches that varied from morphological characterisation to the development of diagnostic and therapeutic tools. Biochemical assays aiming to isolate and characterise nuclear and mitochondria




Fig. 3: discriminative free-living amoeba (FLA) literature that has FLAs as the central topic (core) in Brazil according to five categories: clinical, environmental, review, dry, and wet-bench. (A) Stacked bar graph comparing the number of papers published per year per category. (B) Territorial scope depicting the percentage of papers per federative unit.




Fig. 4: the diversity of the free-living amoeba (FLA) genus per category and class of references. From intense to pale shades, the color spectrum, indicates the prevalence of FLA genera per number of papers. A, N, V, B, and S are *Acanthamoeba*, *Naegleria*, *Vermamoeba*, *Balamuthia*, and *Sappinia*, respectively. M: multiple topics (two or more FLA genera included in the main scope).

drial DNA of *Acanthamoeba* were the pioneer reports on FLA in Brazil, being published in 1974 and 1975.<sup>(36,37)</sup> A long gap of reports on wet-bench research occurred until 1992 when a study reported an isoenzymatic and antigenic assessment towards classifying *Acanthamoeba* isolates in groups I, II and III.<sup>(38)</sup> Further, the randomly amplified polymorphic DNA (RAPD) technique was indicated as a suitable strategy to rapid genotyping new isolates.<sup>(39)</sup> More recent studies adopted high throughput sequencing with bioinformatics pipelines to decipher nuclear and mitochondrial DNA of *Acanthamoeba* and *Vermamoeba* strains.<sup>(40)</sup> Proteomics and metabolomics analysis are also represented by studies devoted to characterising *Acanthamoeba*, which provided informative molecular elements with the potential to be investigated as drug targets or as epitopes for antibodies in diagnostic tools.<sup>(41,42,43)</sup> Additional targets for detection purposes are cation:proton antiporter protein family (CPA) and a calreticulin extracellular domain.<sup>(44,45)</sup> The former was identified through the monoclonal antibody mAb3,




Fig. 5: progression of the methods used in Brazil to monitor FLA presence in clinical (A) and environmental (B) samples.

which positively reacted with several *Acanthamoeba* pathogenic strains.<sup>(44)</sup> The latter study described the use of a recombinant polypeptide, produced from a predicted sequence of *A. castellanii* calreticulin, which was recognised by sera from rats with GAE in the ELISA test.<sup>(44)</sup> These contributions meet the urgent demand for faster detection techniques since late diagnosis is a factor contributing for the high mortality or visual sequelae of FLA infections.<sup>(1,2,6,8)</sup> Other *Acanthamoeba* wet-bench research is concerned with culture manipulation, as the proposal of strategies for prolonging culture storage<sup>(46)</sup> and removing contaminants from cultures.<sup>(47)</sup> Additional approaches included investigating the role of phosphate transporters in metabolism,<sup>(48)</sup> description of ultrastructural cyst wall morphology,<sup>(49)</sup> and genomic features.<sup>(50,51)</sup> Once the primary status of FLAs is not parasitic, a relevant issue in the research field is understanding factors associated with the pathogenic behavior. Brazilian studies explored *in vitro* *Acanthamoeba* properties, such as the interaction of trophozoites with the host cell or the extracellular matrix.<sup>(24,52,53,54)</sup> Contact-independent elements were also included, with contributions that described released proteases through the zymographic technique, indicating the serine protease class as the predominant type produced by *Acanthamoeba*.<sup>(55,56,57,58)</sup> Of note, an additional contribution showed for the first time that *Acanthamoeba* produces and releases extracellular vesicles (EVs) that carry proteases and can be/taken up by target cells, causing cytotoxicity.<sup>(59)</sup> Apart

from the entirely *in vitro* investigations aforementioned, a study adopted a rat model to induce systemic and cerebral acanthamoebiasis as a strategy to reactivate the virulence of *Acanthamoeba* cultures.<sup>(60)</sup>

Efforts to develop novel and improve existing therapies to combat FLA infections have been performed in Brazil, including tests with both first-line options and novel compounds. Polyhexamethylene biguanide and chlorhexidine digluconate were evaluated alone or combined against *Acanthamoeba* cysts.<sup>(61)</sup> Those biguanides and diamidines (e.g., propamidine and hexamidine isethionate), are currently the main therapeutic lines for *Acanthamoeba* keratitis.<sup>(62)</sup> As they exert cytotoxicity to the cornea, searching for new amoebicidal products is still necessary. Thirteen studies proposed in Brazil included dose-response assays with natural and synthetic substances in this direction. Examples are bacteriocin-like compounds,<sup>(63)</sup> alpha-helical and beta-hairpin antimicrobial peptides,<sup>(64)</sup> S-nitrosothiols,<sup>(65)</sup> silver nanoparticles,<sup>(66)</sup> plant extract, or essential oils directly assayed.<sup>(67-73)</sup> Two studies have explored alternatives such as photodynamic therapy by using riboflavin and curcuminoids as photosensitisers for treating *Acanthamoeba* keratitis.<sup>(74,75)</sup> The effectiveness of contact lens multipurpose solutions against *Acanthamoeba*, a topic debated by the FLA community,<sup>(76)</sup> has been also investigated in Brazil,<sup>(77,78)</sup> including the suggestion of imidazolium salt, an anti-amoebic substance, as a suitable option for lens storage.<sup>(79)</sup> Alternatively, therapeutic strategies were concerned with molecules essential to cell division, energy production, cell differentiation, or movement. Examples consist of two Brazilian studies testing the role of small interfering RNA (siRNA) carried in liposomes as an anti-amoebic agent by *in vitro* and *in vivo* assays.<sup>(26,80)</sup> Even though mentions of other FLA species were less frequent than *Acanthamoeba*, *B. mandrillaris* is represented in a study describing the amoeba-host cell interaction.<sup>(81)</sup> *Naegleria* was the focus of three reports; in one of them, mass spectrometry (MALDI-TOF MS) was proposed as a tool to discriminate *N. fowleri* from non-pathogenic species (e.g., *N. italica*, *N. jadini*, *N. gruberi*).<sup>(82)</sup> In two papers, proteins of *Naegleria* were characterised by biophysical approaches, providing insights about its metabolism.<sup>(83,84)</sup>

Wet-bench Brazilian research also included ecological information on FLA and amoebic resistant microorganisms (ARMs). Some reports discuss interactions of FLAs with fungi and bacteria, addressing both endosymbionts and pathogens of concern in human health.<sup>(19,85-90)</sup> Another type of ARMs studied in Brazil comprises human adenovirus<sup>(91)</sup> and giant viruses such as the Tupanvirus, isolated from *V. vermiformis* and *A. polyphaga* mimivirus.<sup>(92,93)</sup> We discussed below other reports on giant virus classified in the auxiliary class. Apart from intra-amoebic microorganisms, a wet-bench study investigated the ecological relation between amoeba and mosquitoes, demonstrating for the first time that *A. polyphaga* can infect *Aedes aegypti*.<sup>(94)</sup> Moreover, the role of FLA as vehicles of pathogenic microorganisms debated worldwide<sup>(95,96,97)</sup> and also present in Brazilian surveys<sup>(85,88,91,98)</sup> reinforce the need of monitoring FLA presence in the environment.

*Clinical references and the risks of FLA infections for human health in Brazil* - As shown in Fig. 3, 29 out of 178 references were classified in the clinical category, including mainly *Acanthamoeba* keratitis,<sup>(99-104)</sup> PAM,<sup>(105)</sup> and GAE,<sup>(106)</sup> reports (Table I).

FLA infections reported in Brazil present the occurrence of *Naegleria* spp. in encephalitis cases,<sup>(105,107)</sup> and *Acanthamoeba* spp. as the commonest genus of keratitis,<sup>(108-111)</sup> skin infections, disseminated disease, and even encountered in human feces<sup>(112)</sup> and urine.<sup>(113)</sup> *Vermamoeba* and *Balamuthia* were reported in two records, and no *Sappinia* cases were described (Table I). Of seven encephalitis cases, only one describes the recovery of the patient.<sup>(114)</sup> It is dated from 1978, reporting the recovery of a 14-year-old boy with a habit of bathing in lagoons. Water samples were collected, confirming the presence of *N. fowleri*.<sup>(114)</sup>

*Acanthamoeba* infections, basically keratitis, represent the highest incidence of amoeba contaminations in Brazil. In this group, investigations dated after 2010 described molecular approaches (Table I-D), indicating this technology became more accessible in the laboratory routine.

The widely debated misdiagnosis of FLA infection<sup>(62)</sup> has also been reported in the Brazilian literature, as discussed in a retrospective study of cases of amoebic keratitis, in which the herpes virus was the most frequent initial suspicion.<sup>(115)</sup> The same occurred for amoeba brain infections, in which a fatal case in Brazil was first hypothesised and treated for viral meningitis.<sup>(116)</sup> Despite that, the overall tendency to rapidly and accurately diagnose FLA proposed in Brazilian surveys may decrease the misdiagnosis. On the other hand, monitoring of intra-FLA resistant microorganisms is an issue of concern worldwide<sup>(1,97,117)</sup> that is still absent in the clinical category of Brazilian papers.

*Environmental reports and the assessment of FLA as a water quality marker in Brazil* - *Acanthamoeba*, *Naegleria*, *Vermamoeba*, and *Balamuthia* were detected in Brazilian habitats, including university buildings, hospitals, swimming pools, tap and freshwater, dust, soil samples, and inside insects<sup>(118-131)</sup> (Table II).

The first isolation in the environment is dated from 1986 and pointed out the presence of *N. fowleri* in an artificial lake in Rio de Janeiro city, state of Rio de Janeiro, the Southeast region.<sup>(132)</sup> Dust samples from hospitals and university buildings yielded positive results for *Naegleria* spp. in the municipalities of Presidente Prudente and Santos, respectively, both in São Paulo state, the Southeast region.<sup>(118,133)</sup> Although based on morphological analysis, a study indicated the *N. fowleri* presence by its ability for supporting 43°C and flagellating.<sup>(1,133)</sup>

*Naegleria* and *Vermamoeba* (*Hartmanella*) identifications were less frequent compared to *Acanthamoeba*,<sup>(134,135)</sup> but they were found coexisting in the same habitats (Table II).<sup>(136,137)</sup> Environmental findings of *B. mandrillaris* were reported to be associated with *Acanthamoeba* populations in air conditioning systems of hospitals.<sup>(138)</sup> Across the globe, FLAs have been sampled

TABLE I  
List of cases of free-living amoeba (FLA) infections in Brazil - from 1977 to 2020

| #  | Year | Case / Patient - Outcome <sup>a</sup>       | Identified FLA                                            | Method <sup>b</sup> | Ref   |
|----|------|---------------------------------------------|-----------------------------------------------------------|---------------------|-------|
| 1  | 1977 | Encephalitis / Human - Death                | <i>Acanthamoeba</i> ; <i>Hartmannella</i>                 | M                   | (106) |
| 2  | 1978 | Encephalitis / Human - Survived             | <i>Naegleria</i> spp.                                     | M                   | (114) |
| 3  | 1992 | Encephalitis / Human - Death                | <i>Leptomyxid amoeba</i>                                  | M, I                | (152) |
| 4  | 2007 | Encephalitis / Human - Death                | <i>Balamuthia mandrillaris</i>                            | M, I                | (154) |
| 5  | 2010 | Encephalitis / Human - Death                | <i>Balamuthia mandrillaris</i>                            | M, I, D             | (116) |
| 6  | 2012 | Encephalitis / Cattle - Death               | <i>N. fowleri</i>                                         | M, I                | (105) |
| 7  | 2019 | Encephalitis / Cow - Death                  | <i>N. fowleri</i>                                         | M, I                | (107) |
| 8  | 2000 | Keratitis / Human                           | <i>A. astronyxis</i>                                      | M, D                | (39)  |
| 9  | 2003 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (99)  |
| 10 | 2004 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (100) |
| 11 | 2004 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (153) |
| 12 | 2007 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (22)  |
| 13 | 2008 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (101) |
| 14 | 2009 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (102) |
| 15 | 2011 | Keratitis / Human                           | <i>A. castellanii</i>                                     | M                   | (155) |
| 16 | 2011 | Keratitis / Human                           | <i>Acanthamoeba</i> T4                                    | M, D                | (74)  |
| 17 | 2013 | Keratitis / Human                           | <i>Acanthamoeba</i> T4, T11, T2 and T4/T2; T4/T13; T4/T16 | M, D                | (103) |
| 18 | 2013 | Keratitis / Human                           | <i>A. castellanii</i> , <i>A. rhysodes</i>                | M, I, D             | (61)  |
| 19 | 2017 | Keratitis / Human                           | <i>A. castellanii</i> T4                                  | M, D                | (104) |
| 20 | 2017 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M, D                | (21)  |
| 21 | 2018 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (87)  |
| 22 | 2018 | Keratitis / Human                           | <i>Acanthamoeba</i> spp.                                  | M                   | (115) |
| 23 | 2018 | Keratitis / Human                           | <i>Acanthamoeba</i> T4                                    | M, D                | (108) |
| 24 | 2018 | Keratitis / Human                           | <i>Acanthamoeba</i> spp T4 (co-infection) <sup>c</sup>    | M, D                | (109) |
| 25 | 2018 | Keratitis / Human                           | <i>Acanthamoeba</i> T4                                    | M, D                | (110) |
| 26 | 2014 | Skin infection / dogs - Survived            | <i>Acanthamoeba</i> T3, T4, T5, T16                       | M, D                | (111) |
| 27 | 2015 | Disseminated amoebic disease / dogs - Death | <i>Acanthamoeba</i> spp.                                  | M, I                | (112) |
| 28 | 1985 | Other / human feces                         | <i>Acanthamoeba</i> spp., <i>Hartmannella</i> spp.        | M                   | (151) |
| 29 | 2009 | Other / urine samples                       | <i>Acanthamoeba</i> spp.                                  | M                   | (113) |

<sup>a</sup>: the disease outcome (survival/death) was not applicable for both keratitis and other clinical samples; <sup>b</sup>: M - microscopy (morphology - M); immunohistochemistry (I); DNA based approach (D); <sup>c</sup>: co-infection case (*Acanthamoeba* spp and *Candida albicans*). Ref: references.

in natural habitats related to water and soil<sup>(139,140,141)</sup> besides extreme environments such as treatment water plants, chlorinated swimming pools, hot spring water, and ice samples.<sup>(142,143,144)</sup> Global warming can favor the spreading of thermophilic pathogenic FLA<sup>(145,146)</sup> and intra-FLA pathogenic bacteria.<sup>(147)</sup> In Brazil, a tropical zone directly exposed to global warming effects,<sup>(148)</sup> the presence of potentially pathogenic FLA strains described in water-related samples (Table II) can pose a health concern. Although the country's regulatory legislation of water quality does not address FLA risks for humans,<sup>(149)</sup> the need for research on tools to detect them is identified in the global literature.<sup>(150)</sup>

*Progression of FLA detection and characterisation tools in Brazil* - With the expansion of studies on FLA in the country, methods for detection and characterisation of environmental and clinical samples have improved, with a tendency to adopt molecular approaches over time (Fig. 5).

Microscopy-based methods for detection and characterisation were adopted in all studies (54/54) and in some, (17/54) it was the only option to identify the FLA. This approach predominates until 2005 (Fig. 5). Between 2010 and 2020, 23 out of 34 FLA environmental and clinical investigations included molecular-based tools for FLA detection, in contrast with five "morphological"

TABLE II  
A summary of the environmental isolations of free-living amoeba (FLA) in Brazil, from 1986 to 2020

| Year | Sample                                           | FLA identification                                                                                                                          | Method <sup>a</sup> | Ref   |
|------|--------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|---------------------|-------|
| 1986 | Lake                                             | <i>N. fowleri</i>                                                                                                                           | M,I                 | (132) |
| 2003 | Hospitals                                        | <i>Acanthamoeba</i> , <i>Naegleria</i>                                                                                                      | M                   | (118) |
| 2007 | Public hospitals                                 | <i>Acanthamoeba</i> spp.                                                                                                                    | M                   | (35)  |
| 2008 | Contact lenses cases                             | <i>Acanthamoeba</i> spp.                                                                                                                    | M, D                | (119) |
| 2009 | University buildings                             | <i>Acanthamoeba</i> ; <i>Naegleria</i>                                                                                                      | M, I                | (133) |
| 2009 | Water samples                                    | <i>Acanthamoeba</i> ACC01                                                                                                                   | M,D                 | (135) |
| 2009 | Swimming pools                                   | <i>Acanthamoeba</i> spp.                                                                                                                    | M,D                 | (120) |
| 2010 | Hospital localities                              | <i>Acanthamoeba</i> T4, T5, T3                                                                                                              | M                   | (121) |
| 2010 | Public hospitals                                 | <i>Acanthamoeba</i> spp.                                                                                                                    | M,D                 | (126) |
| 2011 | Tap water                                        | <i>Acanthamoeba</i> T2,T6, T4                                                                                                               | M,D                 | (127) |
| 2012 | Soil and water samples                           | <i>Acanthamoeba</i> T4, T5, T2-T6                                                                                                           | M                   | (128) |
| 2012 | <i>Aedes aegypti</i> larvae                      | <i>Acanthamoeba</i> T4, T3, T5                                                                                                              | M,D                 | (129) |
| 2012 | Soil and water samples                           | <i>Acanthamoeba</i> T7, T8, T9, T17                                                                                                         | M,D                 | (130) |
| 2012 | Public hospitals                                 | <i>A. castellanii</i> , <i>A. polyphaga</i> , <i>A. lenticulata</i>                                                                         | M,D                 | (131) |
| 2013 | Bromeliads leaves                                | <i>Acanthamoeba</i> T4, T2/T6, T16                                                                                                          | M,D                 | (120) |
| 2013 | Dust of domicile                                 | <i>Acanthamoeba</i> spp.                                                                                                                    | M,I                 | (121) |
| 2015 | Air conditioned samples and contact lenses cases | <i>Acanthamoeba</i> T4, T5, T3                                                                                                              | M,D                 | (98)  |
| 2015 | Mineral bottled water                            | <i>Acanthamoeba</i> T5, T4, T11                                                                                                             | M,D                 | (122) |
| 2016 | Hot tubes, swimming pools                        | <i>Acanthamoeba</i> T3, T5, T4, T15                                                                                                         | M,D                 | (123) |
| 2016 | Swimming pools                                   | <i>Acanthamoeba</i> spp.                                                                                                                    | M,D                 | (91)  |
| 2017 | Air conditioned cooling tower samples            | <i>Acanthamoeba</i> spp T4, T5, <i>Vermamoeba vermiformis</i> , <i>Naegleria</i> sp., <i>N. australiensis</i>                               | M,D                 | (136) |
| 2018 | Dust, sewage, soil, biofilm, sea samples         | <i>Acanthamoeba</i> T1, T15 and T18                                                                                                         | M,D                 | (134) |
| 2020 | Air conditioned in hospitals                     | <i>Acanthamoeba</i> T3, T4, <i>Balamuthia mandrillaris</i>                                                                                  | M,D                 | (138) |
| 2020 | River water                                      | <i>N. philippinensis</i> , <i>N. canariensis</i> , <i>N. australiensis</i> , <i>N. gruberi</i> , <i>N. dobsoni</i> <i>Hartmannella</i> spp. | M,D                 | (137) |
| 2020 | Air conditioned in hospitals                     | <i>Acanthamoeba</i> T4, T5, T11.                                                                                                            | M,D                 | (157) |

a: M - microscopy (morphology - M); immunohistochemistry (I); DNA based approach (D). Ref: references.

and six “Morphological & Immunohistochemical” based analyses. Moreover, it is worth noting that DNA-based investigations were more prevalent on environmental (18/25) than clinical (9/29) references (Fig. 5).

Overall, the studies included microscopy-based investigations combined with hematoxylin-eosin (HE),<sup>(151,152)</sup> calcofluor and Giemsa stains.<sup>(153)</sup> Anti-*B. mandrillaris* antisera has been used for immunofluorescence purposes.<sup>(116,154)</sup> Before performing microscope slides, environmental and clinical samples were often cultured in non-nutritive agar plates<sup>(155)</sup> with an additional supply of bacteria such as *Escherichia coli*,<sup>(153)</sup> *Enterobacter*,<sup>(22,153)</sup> *Aerobacter aerogenes*,<sup>(151)</sup> *Micrococcus luteus* and *Pseudomonas aeruginosa*.<sup>(156)</sup> Osmotolerance and thermotolerance assays have been performed in NNA plates for checking pathogenicity.<sup>(124,128,157)</sup> Molecular-based approaches comprised PCR assays in which 18S rDNA regions<sup>(50,158)</sup> or ITS regions<sup>(159)</sup> were used to genotyping FLA isolates, and the

V3 region of 16S rDNA of bacteria was used to detect intra-FLA bacteria.<sup>(98)</sup> Among the wet-bench category of papers, one study has employed a high-throughput sequencing (HTS) methodology to analyse mitochondrial gene sequences elucidative to investigate diversity in Amoebozoa.<sup>(40)</sup>

*Reviews and dry-bench studies in the core class* - A group of literature reviews covered a diversity of FLA-related topics. The first literature review dates from 2000, and it is devoted to reporting predisposing factors commonly associated with amoebic keratitis, including clinical aspects, diagnosis strategies, and therapeutic options.<sup>(160)</sup> Two comprehensive reviews addressed mechanisms developed by intra *Acanthamoeba* microorganisms to enhance their survival and multiplication chances.<sup>(161,162)</sup> Since several ARMs are capable of proliferating in FLAs and in the human host, other reviews discussed the hypothesis that ARMs can use the amoeba

cell to gain virulence before inhabiting humans.<sup>(161)</sup> Subsequent work discussed clinical aspects of *N. fowleri* infection and debated therapeutic challenges when treating PAM patients.<sup>(159)</sup> The latest literature review was published in 2019, describing the role of extracellular vesicles of bacteria, fungi, and protozoans with a particular emphasis on *A. castellanii*.<sup>(163)</sup>

Finally, the dry-bench references gathered relevant information for ARM monitoring. For example, a study accessed genomic signatures shared by *Acanthamoeba* entoorganisms by using oligonucleotide relative frequencies (OnRF) and relative codon usage (RCU) approaches.<sup>(164)</sup> An additional publication described the tridimensional structure of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) of *N. gruberi*.<sup>(165)</sup>

**Auxiliary category** - The auxiliary class grouped 58 references on broad subjects that also embraced FLA topics. It includes virus replication,<sup>(166,167,168,169)</sup> microbial keratitis,<sup>(170,171)</sup> main aspects of central nervous disorders<sup>(172,173,174)</sup> and cutaneous infections,<sup>(175)</sup> reaction of microorganisms against phototherapies,<sup>(176)</sup> among others.

Prospection surveys concerned with waterborne protozoans have shown *Giardia* and *Acanthamoeba* in saline water and oysters samples, highlighting both as a concern for human health.<sup>(177)</sup> Similarly, a foodborne parasite-based examination in salad samples of Brazilian restaurants showed *Acanthamoeba* presence ranking in the first position with 23.5% identifications.<sup>(178)</sup> Although this identification suggests the importance of food as a source of free-living amoeba, as far as we know, there is no evidence of infections elicited by ingesting FLA.

A study on keratitis has indicated the following groups as etiological agents of corneal infections: bacteria (*Staphylococcus*, *Streptococcus*, *Enterobacter*), fungi (*Candida*, *Rhodotorula*), and protozoan groups (*Acanthamoeba*) as possible etiological agents of corneal infections.<sup>(179)</sup> From 2007 to 2014, another study examined 242 patients from a specialised eye hospital whose results indicated bacteria as the first cause of keratitis infection, and the successive positions in the ranking were occupied by fungi and *Acanthamoeba* infections.<sup>(180)</sup>

Concerned with treatments, the auxiliary literature has investigated UV light and natural compounds acting as photosensitisers for treat a diversity of parasitic and microbial infections, including *Acanthamoeba* keratitis.<sup>(181)</sup> A related survey concerned with the ranges of visual acuity recovery has examined the efficacy of Boston keratoprosthesis type I for *Acanthamoeba* keratitis.<sup>(182)</sup> Other contributions are assays to investigate the effectivity of a biosurfactant on *Candida* and *Acanthamoeba*<sup>(183)</sup> and anti-amoebic action of metabolites from *Penicillium*.<sup>(184)</sup>

Sixty-seven percent (39/58) of the auxiliary references describe investigations specific to ARMs. Fungi and FLA interactions addressed *Trichophyton rubrum*,<sup>(90)</sup> *Fusarium*,<sup>(85)</sup> *Sporothrix*,<sup>(185)</sup> *Paracoccidioides* spp.,<sup>(186)</sup> and *Cryptococcus* species,<sup>(183,187,188)</sup> all associated with *Acanthamoeba*. Some of these reports highlight that fungi-FLA interactions in the environment mimic those found between fungi and host cells, possibly contributing to the development of virulence and immune escape strategies.<sup>(187,188,189)</sup>

One study identified *Pseudomonas* in *Acanthamoeba* isolates.<sup>(86)</sup> Another survey adopted a co-culture model between this FLA genus and methicillin-resistant *Staphylococcus aureus*, showing mutual effects on cell growth or differentiation.<sup>(190)</sup> A third report investigated the role of a K(+) transporter gene in the nutrient uptake and replication of *Legionella pneumophila* inside *A. castellanii*.<sup>(191)</sup>

Several reports on ARMs (30/39) have comprised the giant viruses, placing this topic as a highlight in the auxiliary category. Relevant contributions included new virus species and lineages described after prospection in Brazilian environments.<sup>(192-199)</sup> Exploring variable aspects of interaction, a set of reports addressed the morphological description of viral attachment, replication, and release<sup>(200-207)</sup> to the analysis of gene expression regulating the host cell cycle.<sup>(92,93,204,208)</sup> Reviews and editorials<sup>(209,210,211,212)</sup> described methods for the isolation, culture<sup>(213,214,215)</sup> and antiviral biocides assays<sup>(216,217)</sup> also compose the auxiliary list of papers. Additional studies debated virus lateral gene transfer,<sup>(212,218)</sup> or in co-infections with virophages<sup>(219)</sup> that may likely contribute to genetic variation to the amoeba host, and even the opposite situation, in which viruses are capable of integrating amoeba genes into its genome.<sup>(220,221)</sup> Finally, one dry-bench study in the auxiliary class comprises a review that explored the sexual process in Eukaryotes, discussing the expression of meiosis genes in *Acanthamoeba*.<sup>(222)</sup> Altogether, the auxiliary class references added valuable information to the knowledge about FLA and its relationship with other organisms.

So far, no literature review was devoted to critically analysing the Brazilian publications on the FLA topic, although two<sup>(18,223)</sup> out of four literature reviews on *Veramoeba* spp.,<sup>(224)</sup> *Naegleria*,<sup>(15,223)</sup> and *Acanthamoeba*<sup>(18)</sup> mentioned the country, several Brazilian reports are missing in the latest literature reviews.

**Conclusions and directions** - The present work is the first literature review devoted to critically integrating and debating FLA data produced in Brazil. The data described here outlined the main scientific issues related to FLA research in Brazil since the first publication in 1974 and covering the last 46 years. Even though several FLA genera were reported during this study period, most papers were related to *Acanthamoeba*, and the wet-bench investigations predominated.

Regarding the clinical reports, we believe there are many under-reported cases of FLA infections in Brazil, including keratitis and encephalitis cases, as previously suggested by retrospective studies. In addition, the possibility of pathogenic intra-amoebic microorganisms isolated from clinical FLA strains increases the potential of FLA-ARM interaction to be harmful to human health. Due to that, it is urgent to narrow the relations between researchers and clinicians, universities, hospitals, and medical laboratories in the country, aiming to close monitoring cases of FLA diseases, or even bacterial, fungal, and viruses infections likely carried through amoebas.

The Brazilian epidemiologic surveys on FLA in water collections did not consider the limnologic data on the samples. It is advisable for policymakers and

environmental regulatory departments to integrate the efforts towards improving the knowledge on amoeba distribution. In this direction, understanding the correlation between amoeba presence and environmental characteristics may contribute to better evaluating public health risk areas. Finally, national funding opportunities and distribution across Brazil are extremely important to boost the spread of FLA research in the country and contribute to public health awareness.

### ACKNOWLEDGEMENTS

To Marina Elisa de Oliveira for the support provided during the establishment of the protocol in StArt tool.

### AUTHORS' CONTRIBUTION

NKB and AOC - Conceptualisation, validation and investigation; NKB - methodology and writing - original draft preparation; NKB and OHT - Resources; AOC, OHT, MRB and JLM - writing - review & editing, and visualisation; AOC - supervision. All authors have read and agreed to the published version of the manuscript. The authors declare no conflict of interest.

### REFERENCES

- Schuster FL, Visvesvara GS. Free-living amoebae as opportunistic and non-opportunistic pathogens of humans and animals. *Int J Parasitol*. 2004; 34(9): 1001-27.
- Marciano-Cabral F, Cabral G. *Acanthamoeba* spp. as agents of disease in humans. *Clin Microbiol Rev*. 2003; 16(2): 273-307.
- Schuster FL, Dunnebacke TH, Booton GC, Yagi S, Kohlmeier CK, Glaser C, et al. Environmental isolation of *Balamuthia mandril-laris* associated with a case of amebic encephalitis. *J Clin Microbiol*. 2003; 41(7): 3175-80.
- Scheid PL. *Vermamoeba vermiformis* - a free-living amoeba with public health and environmental health significance. *Open Parasitol J*. 2019; 7(1): 40-7.
- Samba-Louaka A, Delafont V, Rodier M-H, Cateau E, Héchar Y. Free-living amoebae and squatters in the wild: ecological and molecular features. *FEMS Microbiol Rev*. 2019; 43(4): 415-34.
- Marciano-Cabral F. Free-living amoebae as agents of human infection. *J Infect Dis*. 2009; 199(8): 1104-6.
- Fritz-Laylin LK, Prochnik SE, Ginger ML, Dacks JB, Carpenter ML, Field MC, et al. The genome of *Naegleria gruberi* illuminates early eukaryotic versatility. *Cell*. 2010; 140(5): 631-42.
- Piñero JE, Chávez-Munguía B, Omaña-molina M, Lorenzo-morales J. *Naegleria fowleri*. *Trends Parasitol*. 2019; 35(10): 848-9.
- Lee JY, Yu IK, Kim SM, Kim JH, Kim HY. Fulminant disseminating fatal granulomatous amebic encephalitis: the first case report in an immunocompetent patient in South Korea. *Yonsei Med J*. 2021; 62(6): 563-7.
- Duggal SD, Rongpharpi SR, Duggal AK, Kumar A, Biswal I. Role of *Acanthamoeba* in granulomatous encephalitis: a review. *J Infect Dis Immune Ther*. 2017; 1(1): 2.
- Soleimani M, Latifi A, Momenai B, Tayebi F, Mohammadi SS, Ghahvehchian H. Management of refractory *Acanthamoeba* keratitis, two cases. *Parasitol Res*. 2021; 120(3): 1121-4.
- Carnt NA, Subedi D, Connor S, Kilvington S. The relationship between environmental sources and the susceptibility of *Acanthamoeba* keratitis in the United Kingdom. *PLoS One*. 2020; 15(3): 1-11.
- Qvarnstrom Y, Da Silva AJ, Schuster FL, Gelman BB, Visvesvara GS. Molecular confirmation of *Sappinia pedata* as a causative agent of amoebic encephalitis. *J Infect Dis*. 2009; 199(8): 1139-42.
- Gelman BB, Popov V, Chaljub G, Nader R, Rauf SJ, Nauta HW, et al. Neuropathological and ultrastructural features of amebic encephalitis caused by *Sappinia diploidea*. *J Neuropathol Exp Neurol*. 2003; 62(10): 990-8.
- Gharpure R, Bliton J, Goodman A, Ali IKM, Yoder J, Cope JR. Epidemiology and clinical characteristics of primary amebic meningoencephalitis caused by *Naegleria fowleri*: a global review. *Clin Infect Dis*. 2020; 73(1): e19-27.
- Król-Turmińska K, Olender A. Human infections caused by free-living amoebae. *Ann Agric Environ Med*. 2017; 24(2): 254-60.
- Scheid P. Relevance of free-living amoebae as hosts for phylogenetically diverse microorganisms. *Parasitol Res*. 2014; 113(7): 2407-14.
- Rayamajhee B, Subedi D, Peguda HK, Willcox MD, Henriquez FL, Carnt N. A systematic review of intracellular microorganisms within *Acanthamoeba* to understand potential impact for infection. *Pathogens*. 2021; 10(2): 1-26.
- Gonçalves DS, Ferreira MS, Gomes KX, Rodríguez-de-La-Noval C, Liedke SC, Costa GCV, et al. Unravelling the interactions of the environmental host *Acanthamoeba castellanii* with fungi through the recognition by mannose - binding proteins. *Cell Microbiol*. 2019; 21(10): e13066.
- Rosado-García FM, Guerrero-Flórez M, Karanis G, Hinojosa MDC, Karanis P. Water-borne protozoa parasites: the Latin American perspective. *Int J Hyg Environ Health*. 2017; 220(5): 783-98.
- Costa AO, Furst C, Rocha LO, Cirelli C, Cardoso CN, Neiva FS, et al. Molecular diagnosis of *Acanthamoeba* keratitis: evaluation in rat model and application in suspected human cases. *Parasitol Res*. 2017; 116(4): 1339-44.
- Barros JN, Mascaro VLD, Lowen M, Martins MC, Foronda A. Diagnosis of *Acanthamoeba* corneal infection by impression cytology: case report. *Arq Bras Oftal*. 2007; 70(2): 343-6.
- Machado ATP, Silva M, Iulek J. Expression, purification, enzymatic characterization and crystallization of glyceraldehyde-3-phosphate dehydrogenase from *Naegleria gruberi*, the first one from phylum Percolozoa. *Protein Expr Purif*. 2016; 127: 125-30.
- da Rocha-Azevedo B, Costa e Silva-Filho F. Biological characterization of a clinical and an environmental isolate of *Acanthamoeba polyphaga*: analysis of relevant parameters to decode pathogenicity. *Arch Microbiol*. 2007; 188(5): 441-9.
- Carrijo-Carvalho LC, Sant'ana VP, Foronda AS, de Freitas D, Ramos F, Carvalho DS. Therapeutic agents and biocides for ocular infections by free-living amoebae of *Acanthamoeba* genus. *Surv Ophthalmol*. 2016; 62(2): 203-18.
- Zorzi GK, Schuh RS, Maschio VJ, Brazil NT, Rott MB, Teixeira HF. Box Behnken design of siRNA-loaded liposomes for the treatment of a murine model of ocular keratitis caused by *Acanthamoeba*. *Colloids Surfaces B Biointerfaces*. 2019; 173: 725-32.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *BMJ*. 2009; 339: b2535.
- Lo Russo G, Spolveri F, Ciancio F, Mori A. Mendeley: an easy way to manage, share, and synchronize papers and citations. *Plast Reconstr Surg*. 2013; 131(6): 946-7.
- Hernandes E, Zamboni A, Fabbri S. Using GQM and TAM to evaluate StArt-a tool that supports Systematic Review. *Clei Electron J*. 2012; 15(1): 13.
- Morrison A, Polisen J, Husereau D, Moulton K, Clark M, Fianer M, et al. The effect of english-language restriction on systematic review-based meta-analyses: a systematic review of empirical studies. *Int J Technol Assess Health Care*. 2012; 28(2): 138-44.

31. McManus C, Baeta Neves AA. Funding research in Brazil. *Scientometrics*. 2021; 126(1): 801-23.
32. Chiarini T, Oliveira VP, Neto FC. Spatial distribution of scientific activities: an exploratory analysis of Brazil, 2000-10. *Sci Public Policy*. 2014; 41(5): 625-40.
33. Sullivan KE, Bassiri H, Bousfiha AA, Costa-Carvalho BT, Freeman AF, Hagin D, et al. Emerging infections and pertinent infections related to travel for patients with primary immunodeficiencies. *J Clin Immunol*. 2017; 37: 650-92.
34. Chimelli L. A morphological approach to the diagnosis of protozoal infections of the central nervous system. *Patholog Res Int*. 2011; 2011: 29085.
35. Carlesso AM, Simonetti AB, Artuso GL, Rott MB. Isolation and identification of potentially pathogenic free-living amoebae in samples from environments in a public hospital in the City of Porto Alegre, Rio Grande do Sul. *Rev Soc Bras Med Trop*. 2007; 40(3): 316-20.
36. Marzzoco A, Colli W. Characterization of the genome of the small free-living amoeba *Acanthamoeba castellanii*. *Biochim Biophys Acta*. 1975; 395(4): 525-34.
37. Marzzoco A, Colli W. Isolation of nuclei and characterization of nuclear DNA of *Acanthamoeba castellanii*. *Biochim Biophys Acta*. 1974; 374(3): 292-303.
38. Moura H, Wallace S, Visvesvara GS. *Acanthamoeba healyi* N. sp. and the isoenzyme and immunoblot profiles of *Acanthamoeba* spp., groups 1 and 3. *J Protozool*. 1992; 39(5): 573-83.
39. Alves JM, Gusmão CX, Teixeira MMG, Freitas D, Foronda AS, Affonso HT. Random amplified polymorphic DNA profiles as a tool for the characterization of Brazilian keratitis isolates of the genus *Acanthamoeba*. *Braz J Med Biol Res*. 2000; 33(1): 19-26.
40. Fuciková K, Lahr DJG. Uncovering cryptic diversity in two amoebozoan species using complete mitochondrial genome sequences. *J Eukaryot Microbiol*. 2016; 63(1): 112-22.
41. Caumo KS, Monteiro KM, Ott TR, Maschio VJ, Wagner G, Ferreira HB, et al. Proteomic profiling of the infective trophozoite stage of *Acanthamoeba polyphaga*. *Acta Trop*. 2014; 140: 166-72.
42. Maschio JV, Virginio VG, Ferreira HB, Rott MB. Comparative proteomic analysis of soluble and surface-enriched proteins from *Acanthamoeba castellanii* trophozoites. *Mol Biochem Parasitol*. 2018; 225: 47-53.
43. Alves DSMM, Alves LM, da Costa TL, de Castro AM, Vinaud MC. Anaerobic metabolism in T4 *Acanthamoeba* genotype. *Curr Microbiol*. 2017; 74(6): 685-90.
44. Sánchez AGC, Virginio VG, Maschio VJ, Ferreira HB, Rott MB. Evaluation of the immunodiagnostic potential of a recombinant surface protein domain from *Acanthamoeba castellanii*. *Parasitology*. 2021; 143(12): 1656-64.
45. Weber-Lima MM, Prado-Costa B, Becker-Finco A, Costa AO, Billilad P, Furst C, et al. *Acanthamoeba* spp. monoclonal antibody against a CPA2 transporter : a promising molecular tool for acanthamoebiasis diagnosis and encystment study. *Parasitology*. 2020; 147(14): 1678-88.
46. Pens CJ, Rott MB. *Acanthamoeba* spp. cysts storage in filter paper. *Parasitol Res*. 2008; 103(5): 1229-30.
47. Alves DSMM, Gurgel-Gonçalves R, Albuquerque P, Cuba-Cuba CA, Muniz-Junqueira MI, Kuckelhaus SAS. A method for microbial decontamination of *Acanthamoeba* cultures using the peritoneal cavity of mice. *Asian Pac J Trop Biomed*. 2015; 5(10): 796-800.
48. Carvalho-Kelly LF, Pralon CF, Rocco-Machado N, Nascimento MT, Carvalho-de-Araújo AD, Meyer-Fernandes JR. *Acanthamoeba castellanii* phosphate transporter (AcPHS) is important to maintain inorganic phosphate influx and is related to trophozoite metabolic processes. *J Bioenerg Biomembr*. 2020; 52(2): 93-102.
49. Lemgruber L, Lupetti P, De Souza W, Vommaro RC, Da Rocha-Azevedo B. The fine structure of the *Acanthamoeba polyphaga* cyst wall. *FEMS Microbiol Lett*. 2010; 305(2): 170-6.
50. Corsaro D, Köhler M, Venditti D, Rott MB, Walochnik J. Recovery of an *Acanthamoeba* strain with two group I introns in the nuclear 18S rRNA gene. *Eur J Protistol*. 2019; 68: 88-98.
51. Corsaro D, Walochnik J, Köhler M, Rott MB. *Acanthamoeba* misidentification and multiple labels: redefining genotypes T16, T19, and T20 and proposal for *Acanthamoeba micheli* sp. nov. (genotype T19). *Parasitol Res*. 2015; 114(7): 2481-90.
52. da Rocha-Azevedo B, Menezes GC, Costa e Silva-Filho F. The interaction between *Acanthamoeba polyphaga* and human osteoblastic cells *in vitro*. *Microb Pathog*. 2006; 40(1): 8-14.
53. da Rocha-Azevedo B, Jamerson M, Cabral GA, Costa e Silva-Filho F, Marciano-Cabral F. *Acanthamoeba* interaction with extracellular matrix glycoproteins: biological and biochemical characterization and role in cytotoxicity and invasiveness. *J Eukaryot Microbiol*. 2009; 56(3): 270-8.
54. Alves DSMM, Moraes AS, Alves LM, Gurgel-Gonçalves R, Lino Jr RS, Cuba-Cuba CA, et al. Experimental infection of T4 *Acanthamoeba* genotype determines the pathogenic potential. *Parasitol Res*. 2016; 115(9): 3435-40.
55. Ferreira GA, Magliano ACM, Pral EMF, Alfieri SC. Elastase secretion in *Acanthamoeba polyphaga*. *Acta Trop*. 2009; 112(2): 156-63.
56. Cirelli C, Mesquita EIS, Chagas IAR, Furst C, Possamai CO, Abrahão JS, et al. Extracellular protease profile of *Acanthamoeba* after prolonged axenic culture and after interaction with MDCK cells. *Parasitol Res*. 2020; 119(2): 659-66.
57. Sant'ana VP, Carrijo-Carvalho LC, Foronda AS, Chudzinski-Tavassi AM, de Freitas D, Ramos F, et al. Cytotoxic activity and degradation patterns of structural proteins by corneal isolates of *Acanthamoeba* spp. *Graefes Arch Clin Exp Ophthalmol*. 2015; 253(1): 65-75.
58. Alfieri SC, Correia CEB, Motegi SA, Pral EMF. Proteinase activities in total extracts and in medium conditioned by *Acanthamoeba polyphaga* trophozoites. *J Parasitol*. 2000; 86(2): 220-7.
59. Gonçalves DS, Ferreira MS, Liedke SC, Gomes KX, de Oliveira GA, Leão PEL, et al. Extracellular vesicles and vesicle-free secretome of the protozoa *Acanthamoeba castellanii* under homeostasis and nutritional stress and their damaging potential to host cells. *Virulence*. 2018; 9(1): 818-36.
60. Veríssimo CM, Maschio VJ, Correa APF, Brandelli A, Rott MB. Infection in a rat model reactivates attenuated virulence after long-term axenic culture of *Acanthamoeba* spp. *Mem Inst Oswaldo Cruz*. 2013; 108(7): 832-5.
61. Mafra CSP, Carrijo-Carvalho LC, Chudzinski-Tavassi AM, Taguchi FMC, Foronda AS, Carvalho FRS, et al. Antimicrobial action of biguanides on the viability of *Acanthamoeba* cysts and assessment of cell toxicity. *Invest Ophthalmol Vis Sci*. 2013; 54(9): 6363-72.
62. Lorenzo-Morales J, Khan NA, Walochnik J. An update on *Acanthamoeba* keratitis: diagnosis, pathogenesis and treatment. *Parasite*. 2015; 22(10): 1-20.
63. Benitez LB, Caumo K, Brandelli A, Rott MB. Bacteriocin-like substance from *Bacillus amyloliquefaciens* shows remarkable inhibition of *Acanthamoeba polyphaga*. *Parasitol Res*. 2011; 108(3): 687-91.
64. Sacramento RS, Martins RM, Miranda A, Dobroff ASS, Daffre S, Foronda AS, et al. Differential effects of alpha-helical and beta-hairpin antimicrobial peptides against *Acanthamoeba castellanii*. *Parasitology*. 2009; 136(8): 813-21.

65. Cariello AJ, de Souza GFP, Foronda AS, Yu MCZ, Hofling-Lima AL, de Oliveira MG. *In vitro* amoebicidal activity of S-nitrosoglutathione and S-nitroso-N-acetylcysteine against trophozoites of *Acanthamoeba castellanii*. J Antimicrob Chemother. 2010; 65(3): 588-91.
66. Borase HP, Patil CD, Sauter IP, Rott MB, Patil SV. Amoebicidal activity of phytosynthesized silver nanoparticles and their *in vitro* cytotoxicity to human cells. FEMS Microbiol Lett. 2013; 345(2): 127-31.
67. Santos IGA, Scher R, Rott MB, Menezes LR, Costa EV, Cavalcanti SCH, et al. Amoebicidal activity of the essential oils of *Lippia* spp. (Verbenaceae) against *Acanthamoeba polyphaga* trophozoites. Parasitol Res. 2016; 115(2): 535-40.
68. Sauter IP, dos Santos JC, Apel MA, Cibulski SP, Roehe PM, von Poser GL, et al. Amebicidal activity and chemical composition of *Pterocaulon polystachyum* (Asteraceae) essential oil. Parasitol Res. 2011; 109(5): 1367-71.
69. Vunda SLL, Sauter IP, Cibulski SP, Roehe PM, Bordignon SAL, Rott MB, et al. Chemical composition and amoebicidal activity of *Croton pallidulus*, *Croton ericoides*, and *Croton isabelli* (Euphorbiaceae) essential oils. Parasitol Res. 2012; 111(3): 961-6.
70. Sauter IP, Rossa GE, Lucas AM, Cibulski SP, Roehe PM, Silva LAA, et al. Chemical composition and amoebicidal activity of *Piper hispidinervum* (Piperaceae) essential oil. Ind Crop Prod. 2012; 40: 292-5.
71. Ródio C, Vianna DR, Kowalski KP, Panatieri LF, von Poser G, Rott MB. *In vitro* evaluation of the ameibicidal activity of *Pterocaulon polystachyum* (Asteraceae) against trophozoites of *Acanthamoeba castellanii*. Parasitol Res. 2008; 104(1): 191-4.
72. Castro LC, Sauter IP, Ethur EM, Kauffmann C, Dall'agnol R, Souza J, et al. *In vitro* effect of *Acanthospermum australe* (Asteraceae) extracts on *Acanthamoeba polyphaga* trophozoites. Rev Bras Plantas Med. 2013; 15(4): 589-94.
73. Panatieri LF, Brazil NT, Faber K, Medeiros-Neves B, von Poser GL, Rott MB, et al. Nanoemulsions containing a coumarin-rich extract from *Pterocaulon balansae* (Asteraceae) for the treatment of ocular *Acanthamoeba* keratitis. AAPS PharmSciTech. 2017; 18(3): 721-8.
74. Kashiwabuchi RT, Carvalho FRS, Khan YA, de Freitas D, Foronda AS, Hirai FE, et al. Assessing efficacy of combined riboflavin and UV-A Light (365 nm) treatment of *Acanthamoeba* trophozoites. Invest Ophthalmol Vis Sci. 2011; 52(13): 9333-8.
75. Corrêa TQ, Geralde MC, Carvalho MT, Bagnato VS, Kurachi C, Souza CWO. Photodynamic inactivation of *Acanthamoeba polyphaga* with curcuminoids: an *in vitro* study. Opt Methods Tumor Treat Detect Mech Tech Photodyn Ther XXV. 2016; 9694: 1-7.
76. Hendiger EB, Padzik M, Sifaoui I, Reyes-Batlle M, López-Arencibia A, Zyskowska D, et al. Silver nanoparticles conjugated with contact lens solutions may reduce the risk of *Acanthamoeba* keratitis. Pathogens. 2021; 10(5): 583.
77. de Aguiar APC, Silveira CO, Winck MAT, Rott MB. Susceptibility of *Acanthamoeba* to multipurpose lens-cleaning solutions. Acta Parasitol. 2013; 58(3): 304-8.
78. Ludwig IH, Meisler DM, Rutherford I, Bican FE, Langston RH, Visvesvara GS. Susceptibility of *Acanthamoeba* to soft contact lens disinfection systems. Invest Ophthalmol Vis Sci. 1986; 27(4): 626-8.
79. Fabres LF, Gonçalves FC, Duarte EOS, Berté FK, da Conceição DKSL, Ferreira LA, et al. *In vitro* amoebicidal activity of imidazolium salts against trophozoites. Acta Parasitol. 2020; 65(2): 317-26.
80. Faber K, Zorzi GK, Brazil NT, Rott MB, Teixeira HF. siRNA-loaded liposomes: inhibition of encystment of *Acanthamoeba* and toxicity on the eye surface. Chem Biol Drug Des. 2017; 90(3): 406-16.
81. da Rocha-Azevedo B, Jamerson M, Cabral GA, Costa e Silva-Filho F, Marciano-Cabral F. The interaction between the amoeba *Balamuthia mandrillaris* and extracellular matrix glycoproteins *in vitro*. Parasitology. 2007; 134(Pt 1): 51-8.
82. Moura H, Izquierdo F, Woolfitt AR, Wagner G, Pinto T, Del Aguila C, et al. Detection of biomarkers of pathogenic *Naegleria fowleri* through mass spectrometry and proteomics. J Eukaryot Microbiol. 2015; 62(1): 12-20.
83. Da Silva MTA, Caldas VEA, Costa FC, Silvestre DAMM, Thiemann OH. Selenocysteine biosynthesis and insertion machinery in *Naegleria gruberi*. Mol Biochem Parasitol. 2013; 188(2): 87-90.
84. Penteado RF, Martini VP, Iulek J. Crystallization and crystallographic analyses of triosephosphate isomerase from *Naegleria gruberi*. Rev Virtual Quim. 2016; 8(6): 1835-41.
85. Nunes TET, Brazil NT, Fuentesfria AM, Rott MB. *Acanthamoeba* and *Fusarium* interactions: a possible problem in keratitis. Acta Trop. 2016; 157: 102-7.
86. Maschio VJ, Corção G, Rott MB. Identification of *Pseudomonas* spp. as amoeba-resistant microorganisms in isolates of *Acanthamoeba*. Rev Inst Med Trop São Paulo. 2015; 57(1): 81-3.
87. Sticca MP, Carrijo-Carvalho LC, Silva IMB, Vieira LA, Souza LB, Belfort Jr R, et al. *Acanthamoeba* keratitis in patients wearing scleral contact lenses. Cont Lens Anterior Eye. 2018; 41(3): 307-10.
88. Medina G, Neves P, Flores-Martin S, Manosalva C, Andaur M, Otth C, et al. Transcriptional analysis of flagellar and putative virulence genes of *Arcobacter butzleri* as an endocytobiont of *Acanthamoeba castellanii*. Arch Microbiol. 2019; 201(8): 1075-83.
89. Medina G, Leýan P, Silva CV, Flores-Martin S, Manosalva C, Fernández H. Intra-amoebic localization of *Arcobacter butzleri* as an endocytobiont of *Acanthamoeba castellanii*. Arch Microbiol. 2019; 201(10): 1447-52.
90. de Faria LV, do Carmo PHF, da Costa MC, Peres NTA, Chagas IAR, Furst C, et al. *Acanthamoeba castellanii* as an alternative interaction model for the dermatophyte *Trichophyton rubrum*. Mycoses. 2020; 63(12): 1331-40.
91. Staggemeier R, Arantes T, Caumo KS. Detection and quantification of human adenovirus genomes in *Acanthamoeba* isolated from swimming pools. An Acad Bras Cienc. 2016; 88: 635-41.
92. Silva LCF, Almeida GMF, Assis FL, Albarnaz JD, Boratto PVM, Dornas FP, et al. Modulation of the expression of mimivirus-encoded translation-related genes in response to nutrient availability during *Acanthamoeba castellanii* infection. Front Microbiol. 2015; 6(6): 539.
93. Boratto P, Albarnaz JD, Almeida GMDF, Botelho L, Fontes ACL, Costa AO, et al. *Acanthamoeba polyphaga* mimivirus prevents amoebal encystment-mediating serine proteinase expression and circumvents cell encystment. J Virol. 2015; 89(5): 2962-5.
94. Rott M, Caumo K, Sauter I, Eckert J, da Rosa L, da Silva O. Susceptibility of *Aedes aegypti* (Diptera: Culicidae) to *Acanthamoeba polyphaga* (Sarcomastigophora: Acanthamoebidae). Parasitol Res. 2010; 107(1): 195-8.
95. Scheid PL. *Amoebophagous fungi* as predators and parasites of potentially pathogenic free-living amoebae. Open Parasitol J. 2018; 6(1): 75-86.
96. Scheid PL, Schwarzenberger R. Free-living amoebae as vectors of cryptosporidia. Parasitol Res. 2011; 109(2): 499-504.
97. Lorenzo-Morales J, Coronado-Álvarez N, Martínez-Carretero E, Maciver SK, Valladares B. Detection of four adenovirus serotypes within water-isolated strains of *Acanthamoeba* in the Canary Islands, Spain. Am J Trop Med Hyg. 2007; 77(4): 753-6.

98. Maschio VJ, Corção G, Bucker F, Caumo K, Rott MB. Identification of *Paenibacillus* as a symbiont in *Acanthamoeba*. *Curr Microbiol.* 2015; 71(3): 415-20.
99. Forseto AS, Nosé W. Diagnosis of *Acanthamoeba* keratitis with confocal microscopy. *Rev Bras de Oftalmol.* 2003; 62: 220-8.
100. Nakano E, Oliveira M, Portellinha W, de Freitas D, Nakano K. Confocal microscopy in early diagnosis of *Acanthamoeba* keratitis. *J Refract Surg.* 2004; 20(5): S737-41.
101. Kashiwabuchi RT, de Freitas D, Alvarenga LS, Vieira L, Contarini P, Sato E, et al. Corneal graft survival after therapeutic keratoplasty for *Acanthamoeba* keratitis. *Acta Ophthalmol.* 2008; 86(6): 666-9.
102. Carvalho FRS, Foronda AS, Mannis MJ, Hofling-Lima AL, Belfort Jr R, de Freitas D. Twenty years of *Acanthamoeba* keratitis. *Cornea.* 2009; 28(5): 516-9.
103. Duarte JL, Furst C, Klisiowicz DR, Klassen G, Costa AO. Morphological, genotypic, and physiological characterization of *Acanthamoeba* isolates from keratitis patients and the domestic environment in Vitoria, Espírito Santo, Brazil. *Exp Parasitol.* 2013; 135(1): 9-14.
104. Sant'ana VP, Foronda AS, de Freitas D, Carrijo-Carvalho LC, Carvalho FRS. Sensitivity of enzymatic toxins from corneal isolate of *Acanthamoeba* protozoan to physicochemical parameters. *Curr Microbiol.* 2017; 74(11): 1316-23.
105. Pimentel LA, Dantas AFM, Uzal F, Riet-Correa F. Meningoencephalitis caused by *Naegleria fowleri* in cattle of northeast Brazil. *Res Vet Sci.* 2012; 93(2): 811-2.
106. Campos R, Gomes MCO, Prigenzi LS, Stecca J. Meningoencefalite por ameba de vida livre apresentação do primeiro caso latino-americano. *Rev Inst Med Trop São Paulo.* 1977; 19(5): 349-51.
107. Henker LC, Cruz RAS, Silva FS, Driemeier D, Sonne L, Uzal FA, et al. Meningoencephalitis due to *Naegleria fowleri* in cattle in southern Brazil. *Rev Bras Parasitol Vet.* 2019; 28(3): 514-7.
108. Fabres LF, Maschio VJ, Santos DL, Kwitko S, Marinho DR, Araújo BS, et al. Virulent T4 *Acanthamoeba* causing keratitis in a patient after swimming while wearing contact lenses in Southern Brazil. *Acta Parasitol.* 2018; 63(2): 428-32.
109. Buchele MLC, Wopereis DB, Casara F, Macedo JP, Rott MB, Monteiro FBF, et al. Contact lens-related polymicrobial keratitis : *Acanthamoeba* spp. genotype T4 and *Candida albicans*. *Parasitol Res.* 2018; 117(11): 3431-6.
110. Alves DSMM, Gonçalves GS, Moraes AS, Alves LM, Carmo Neto JR, Hecht MM, et al. The first *Acanthamoeba* keratitis case in the Midwest region of Brazil: diagnosis, genotyping of the parasite and disease outcome. *Rev Soc Bras Med Trop.* 2018; 51(5): 716-9.
111. Carlesso AM, Mentz MB, Machado MLS, Carvalho A, Nunes TET, Maschio VJ, et al. Characterization of isolates of *Acanthamoeba* from the nasal mucosa and cutaneous lesions of dogs. *Curr Microbiol.* 2014; 68(6): 702-7.
112. Frade MTS, Melo LF, Pessoa CRM, Araújo JL, Figuera RA, Souza AP, et al. Systemic acanthamoebiasis associated with canine distemper in dogs in the semiarid region of Paraíba, Brazil. *Pesq Vet Bras.* 2015; 35(2): 160-4.
113. Santos LC, Oliveira MS, Lobo RD, Higashino HR, Costa SF, Van Der Heijden IM, et al. *Acanthamoeba* spp. in urine of critically ill patients. *Emerg Infect Dis.* 2009; 15(7): 1144-6.
114. Salles-Gomes Jr CE, Barbosa ER, Nóbrega JP, Scaff M, Spina-França A. Primary amebic meningoencephalomyelitis. Report of a case. *Arq Neuropsiquiatr.* 1978; 36(2): 139-42.
115. dos Santos DL, Kwitko S, Marinho DR, Araújo BS, Locatelli CI, Rott MB. *Acanthamoeba* keratitis in Porto Alegre (southern Brazil): 28 cases and risk factors. *Parasitol Res.* 2018; 117(3): 747-50.
116. Silva RA, Araújo SA, Avellar IFDF, Pittella JEH, Oliveira JT, Christo PP. Granulomatous amoebic meningoencephalitis in an immunocompetent patient. *Arch Neurol.* 2010; 67(12): 1516-20.
117. Cateau E, Delafont V, Hechard Y, Rodier MH. Free-living amoebae: What part do they play in healthcare-associated infections? *J Hosp Infect.* 2014; 87(3): 131-40.
118. da Silva MA, da Rosa JA. Isolation of potentially pathogenic free-living amoebas in hospital dust. *Rev Saude Publica.* 2003; 37(2): 242-6.
119. Pens CJ, Costa M, Fadanelli C, Caumo K, Rott MB. *Acanthamoeba* spp. and bacterial contamination in contact lens storage cases and the relationship to user profiles. *Parasitol Res.* 2008; 103(6): 1241-5.
120. Landeli MF, Salton J, Caumo K, Broetto L, Rott MB. Isolation and genotyping of free-living environmental isolates of *Acanthamoeba* spp. from bromeliads in Southern Brazil. *Exp Parasitol.* 2013; 134(3): 290-4.
121. Becker-Finco A, Costa AO, Silva SK, Ramada JS, Furst C, Stinger AE, et al. Physiological, morphological, and immunochemical parameters used for the characterization of clinical and environmental isolates of *Acanthamoeba*. *Parasitology.* 2013; 140(3): 396-405.
122. Maschio VJ, Chies F, Carlesso AM, Carvalho A, Rosa SP, Van Der Sand ST, et al. *Acanthamoeba* T4, T5 and T11 isolated from mineral water bottles in Southern Brazil. *Curr Microbiol.* 2014; 70(1): 6-9.
123. Fabres LF, Santos SPR, Benitez LB, Rott MB. Isolation and identification of *Acanthamoeba* spp. from thermal swimming pools and spas in Southern Brazil. *Acta Parasitol.* 2016; 61(2): 221-7.
124. Caumo K, Frasson AP, Pens CJ, Panatieri LF, Frazzon APG, Rott MB. Potentially pathogenic *Acanthamoeba* in swimming pools: a survey in the southern Brazilian city of Porto Alegre. *Ann Trop Med Parasitol.* 2009; 103(6): 477-85.
125. Carlesso AM, Artuso GL, Caumo K, Rott MB. Potentially pathogenic *Acanthamoeba* isolated from a hospital in Brazil. *Curr Microbiol.* 2010; 60(3): 185-90.
126. Costa AO, Castro EA, Ferreira GA, Furst C, Crozeta MA, Thomas-Soccol V. Characterization of *Acanthamoeba* isolates from dust of a public hospital in Curitiba, Paraná, Brazil. *J Eukaryot Microbiol.* 2010; 57(1): 70-5.
127. Winck MAT, Caumo K, Rott MB. Prevalence of *Acanthamoeba* from tap water in Rio Grande do Sul, Brazil. *Curr Microbiol.* 2011; 63(5): 464-9.
128. Alves DSMM, Moraes AS, Nitz N, Oliveira MGC, Hecht MM, Gurgel-Gonçalves R, et al. Occurrence and characterization of *Acanthamoeba* similar to genotypes T4, T5, and T2 / T6 isolated from environmental sources in Brasília, Federal District, Brazil. *Exp Parasitol.* 2012; 131(2): 239-44.
129. Otta DA, Rott MB, Carlesso AM, Santos O. Prevalence of *Acanthamoeba* spp. (Sarcostomastigophora : Acanthamoebidae) in wild populations of *Aedes aegypti* (Diptera : Culicidae). *Parasitol Res.* 2012; 111: 2017-22.
130. Magliano ACM, Teixeira MMG, Alfieri SC. Revisiting the *Acanthamoeba* species that form star-shaped cysts (genotypes T7, T8, T9, and T17): characterization of seven new Brazilian environmental isolates and phylogenetic inferences. *Parasitology.* 2011; 139(1): 45-52.
131. Zanella J, Costa SOP, Zacaria J, Echeverrigaray S. A rapid and reliable method for the clonal isolation of *Acanthamoeba* from environmental samples. *Brazilian Arch Biol Technol.* 2012; 55(1): 1-6.
132. Salazar HC, Moura H, Fernandez O, Peralta JM. Isolation of *Naegleria fowleri* from a lake in the city of Rio de Janeiro. *Trans R Soc Trop Med Hyg.* 1986; 80(2): 348-9.

133. Teixeira LH, Rocha S, Pinto RMF, Caseiro MM, Costa SOP. Prevalence of potentially pathogenic free-living amoebae from *Acanthamoeba* and *Naegleria* genera in non-hospital, public, internal environments from the city of Santos, Brazil. *Brazilian J Infect Dis*. 2009; 13(6): 395-7.
134. Possamai CO, Loss AC, Costa AO, Falqueto A, Furst C. *Acanthamoeba* of three morphological groups and distinct genotypes exhibit variable and weakly inter-related physiological properties. *Parasitol Res*. 2018; 117(5): 1389-400.
135. Magliano ACM, da Silva FM, Teixeira MMG, Alfieri SC. Genotyping, physiological features and proteolytic activities of a potentially pathogenic *Acanthamoeba* sp. isolated from tap water in Brazil. *Exp Parasitol*. 2009; 123(3): 231-5.
136. Soares SS, Souza TK, Berte FK, Cantarelli VV, Rott MB. Occurrence of infected free-living amoebae in cooling towers of southern Brazil. *Curr Microbiol*. 2017; 74(12): 1461-8.
137. Bellini NK, da Fonseca ALM, Reyes-Batlle M, Lorenzo-Morales J, Rocha O, Thiemann OH. Isolation of *Naegleria* spp. from a Brazilian water source. *Pathogens*. 2020; 9(2): 90.
138. Fonseca JDG, Gómez-Hernández C, Barbosa CG, Rezende-Oliveira K. Identification of T3 and T4 genotypes of *Acanthamoeba* sp. in dust samples isolated from air conditioning equipment of public hospital of Ituiutaba-MG. *Curr Microbiol*. 2020; 77(5): 890-5.
139. Sousa-Ramos D, Reyes-Batlle M, Bellini NK, Rodríguez-Expósito RL, Piñero JE, Lorenzo-Morales J. Free-living amoebae in soil samples from Santiago Island, Cape Verde. *Microorganisms*. 2021; 9(7): 1460.
140. Reyes-Batlle M, Díaz FJ, Sifaoui I, Rodríguez-Expósito R, Rizo-Liendo A, Piñero JE, et al. Free living amoebae isolation in irrigation waters and soils of an insular arid agroecosystem. *Sci Total Environ*. 2021; 753: 141833.
141. Mahmoudi MR, Zebardast N, Masangkay FR, Karanis P. Detection of potentially pathogenic free-living amoebae from the Caspian Sea and hospital ward dust of teaching hospitals in Guilan, Iran. *J Water Health*. 2021; 19(2): 278-87.
142. Le Calvez T, Trouilhé MC, Humeau P, Moletta-Denat M, Frère J, Héchar Y. Detection of free-living amoebae by using multiplex quantitative PCR. *Mol Cell Probes*. 2012; 26(3): 116-20.
143. Schroeder JM, Booton GC, Hay J, Niszl IA, Seal DV, Markus MB, et al. Use of subgenic 18S ribosomal DNA PCR and sequencing for genus and genotype identification of *Acanthamoebae* from humans with keratitis and from sewage sludge. *J Clin Microbiol*. 2001; 39(5): 1903-11.
144. Reyes-Batlle M, Niyati M, Martín-Navarro CM, López-Arencibia A, Valladares B, Martínez-Carretero E, et al. Unusual *Vermaamoeba vermiformis* strain isolated from snow in upon observation of the snow samples cultured in. Mount Teide, Tenerife, Canary Islands, Spain. *Nov Biomed*. 2015; 3(4): 189-92.
145. Nichols G, Lake I, Heavyside C. Climate change and water-related infectious diseases. *Atmosphere (Basel)*. 2018; 9(10): 1-60.
146. Xue J, Lamar FG, Zhang B, Lin S, Lamori JG, Sherchan SP. Quantitative assessment of *Naegleria fowleri* and fecal indicator bacteria in brackish water of lake Pontchartrain, Louisiana. *Sci Total Environ*. 2018; 622-623: 8-16.
147. Marciano-Cabral F, Jamerson M, Kaneshiro ES. Free-living amoebae, *Legionella* and *Mycobacterium* in tap water supplied by a municipal drinking water utility in the USA. *J Water Health*. 2010; 8(1): 71-82.
148. Alves MTR, Machado KB, Ferreira ME, Vieira LCG, Nabout JC. A snapshot of the limnological features in tropical floodplain lakes: the relative influence of climate and land use. *Acta Limnol Bras*. 2019; 31(e10).
149. MMA - Ministério do Meio Ambiente/Conselho Nacional do Meio Ambiente. Resolução CONAMA n 357. Brasil; 2005. [Internet]. 23 June, 2021. [cited 2021 Nov 15]. Available from: <http://www.mma.gov.br/port/conama/legiabre.cfm?codlegi=459>.
150. Plutzer J, Karanis P. Neglected waterborne parasitic protozoa and their detection in water. *Water Res*. 2016; 101: 318-32.
151. de Moura H, Salazar HC, Fernandes O, Lisboa DC, de Carvalho FG. Free-living amoebae in human intestine: Evidence of parasitism. *Rev Inst Med Trop São Paulo*. 1985; 27(3): 150-6.
152. Chimelli L, Hahn MD, Scaravilli F, Wallace S, Visvesvara GS. Granulomatous amoebic encephalitis due to leptomysid amoebae: report of the first Brazilian case. *Trans R Soc Trop Med Hyg*. 1992; 86(6): 635.
153. Ruthes ACC, Wahab S, Wahab N, Moreira H, Moreira L. Conjunctivitis presumably due to *Acanthamoeba*. *Arq Bras Oftal*. 2004; 67(6): 897-900.
154. Silva-Vergara ML, Colombo ERC, Vissotto EDF, Silva ACAL, Chica JEL, Etchebehere RM, et al. Disseminated *Balamuthia mandrillaris* amoeba infection in an AIDS patient from Brazil. *Am J Trop Med Hyg*. 2007; 77(6): 1096-8.
155. Carvalho FRDS, Carrijo-Carvalho LC, Chudzinski-Tavassi AM, Foronda AS, de Freitas D. Serine-like proteolytic enzymes correlated with differential pathogenicity in patients with acute *Acanthamoeba* keratitis. *Clin Microbiol Infect*. 2010; 17(4): 603-9.
156. Moraes J, Alfieri SC. Growth, encystment and survival of *Acanthamoeba castellanii* grazing on different bacteria. *FEMS Microbiol Ecol*. 2008; 66(2): 221-9.
157. Wopereis DB, Bazzo ML, De Macedo JP, Casara F, Golfeto L, Venancio E, et al. Free-living amoebae and its relationship with air quality in hospital environments: isolation and characterization of *Acanthamoeba* spp. from air-conditioning system. *Parasitology*. 2020; 147(7): 789-90.
158. Caumo K, Rott MB. *Acanthamoeba* T3, T4 and T5 in swimming-pool waters from Southern Brazil. *Acta Trop*. 2011; 117(3): 233-5.
159. Bellini NK, Santos TM, da Silva MTA, Thiemann OH. The therapeutic strategies against *Naegleria fowleri*. *Exp Parasitol*. 2018; 187: 1-11.
160. Alvarenga LS, de Freitas D, Hofling-Lima AL. Ceratite por *Acanthamoeba*. *Arq Bras Oftal*. 2000; 63(1): 155-9.
161. Guimaraes AJ, Gomes KX, Cortines JR, Peralta JM, Peralta RHS. *Acanthamoeba* spp. as a universal host for pathogenic microorganisms: one bridge from environment to host virulence. *Microbiol Res*. 2016; 193: 30-8.
162. Silva LKDS, Boratto PVM, La Scola B, Bonjardim CA, Abrahão JS. *Acanthamoeba* and mimivirus interactions: the role of amoebal encystment and the expansion of the 'Cheshire Cat' theory. *Curr Opin Microbiol*. 2016; 31: 9-15.
163. Gonçalves DS, Ferreira MS, Guimarães AJ. Extracellular vesicles from the protozoa *Acanthamoeba castellanii*: their role in pathogenesis, environmental adaptation and potential applications. *Bioengineering*. 2019; 6(1): 12.
164. Serrano-Solis V, Soares PET, de Farias ST. Genomic signatures among *Acanthamoeba* polyphaga entomorphisms unveil evidence of coevolution. *J Mol Evol*. 2019; 87(1): 7-15.
165. Machado ATP, Silva M, Iulek J. Structural studies of glyceraldehyde-3-phosphate dehydrogenase from *Naegleria gruberi*, the first one from phylum. *Biochim Biophys Acta Proteins Proteom*. 2018; 1866: 581-8.
166. Abrahão JS, Dornas FP, Silva LCF, Almeida GM, Boratto PVM, Colson P, et al. *Acanthamoeba polyphaga* mimivirus and other giant viruses: an open field to outstanding discoveries. *Virol J*. 2014; 11: 1-12.

167. Boratto PVM, Dornas FP, Silva LCF, Rodrigues RAL, Oliveira GP, Cortines JR, et al. Analyses of the Kroon virus major capsid gene and its transcript highlight a distinct pattern of gene evolution and splicing among Mimiviruses. *J Virol*. 2018; 92(2): 1-11.
168. Dornas FP, Assis FL, Aherfi S, Arantes T, Abrahão JS, Colson P, et al. A Brazilian Marseillevirus is the founding member of a lineage in family Marseilleviridae. *Viruses*. 2016; 8(76): 1-16.
169. Dornas FP, Khalil JYB, Pagnier I, Raoult D, Abrahão J, La Scola B. Isolation of new Brazilian giant viruses from environmental samples using a panel of protozoa. *Front Microbiol*. 2015; 6: 1-9.
170. Moriyama AS, Hofling-Lima AL. Contact lens-associated microbial keratitis. *Arq Bras Oftalmol*. 2008; 71(7): 32-6.
171. Müller RT, Abedi F, Cruzat A, Witkin D, Baniyadi N, Cavalcanti BM, et al. Degeneration and regeneration of subbasal corneal nerves after infectious keratitis: a longitudinal *in vivo* confocal microscopy study. *Ophthalmology*. 2015; 122(11): 2200-9.
172. Frade MTS, Ferreira JS, Nascimento MJR, Aquino VVF, Macêdo IL, Carneiro RS, et al. Central nervous system disorders diagnosed in dogs. *Pesq Vet Bras*. 2018; 38(5): 935-48.
173. Gasparetto EL, Cabral RF, Hygino LC, Domingues RC. Diffusion imaging in brain infections. *Neuroimaging Clin NA*. 2011; 21(1): 89-113.
174. Chimelli L. Co-infection of HIV and tropical infectious agents that affect the nervous system. *Rev Neurol (Paris)*. 2012; 168(3): 270-82.
175. Kollipara R, Peranteau AJ, Nawas ZY, Tong Y, Woc-colburn L, Yan AC, et al. Emerging infectious diseases with cutaneous manifestations: fungal, helminthic, protozoan and ectoparasitic infections. *J Am Acad Dermatol*. 2016; 75(1): 19-30.
176. Decarli MC, Carvalho MT, Corrêa TQ, Bagnato VS, Souza CWO. Different photoresponses of microorganisms: from bioinhibition to biostimulation. *Curr Microbiol*. 2016; 72(4): 473-81.
177. Leal DAG, Souza DSM, Caumo KS, Fongaro G, Panatieri LF, Durigan M, et al. Genotypic characterization and assessment of infectivity of human waterborne pathogens recovered from oysters and estuarine waters in Brazil. *Water Res*. 2018; 137: 273-80.
178. Perim LV, Custódio NCC, Lima VCV, Igreja JASL, Alves DSMM, Storchilo HR, et al. Occurrence of parasites in salads in restaurants in Aparecida de Goiânia, Goiás, Brazil. *J Trop Pathol*. 2020; 49(3): 207-14.
179. Marujo FI, Hirai FE, Yu MCZ, Hofling-Lima AL, de Freitas D, Sato EH. Distribution of infectious keratitis in a tertiary hospital in Brazil. *Arq Bras Oftalmol*. 2013; 76(6): 370-3.
180. Farias R, Pinho L, Santos R. Epidemiological profile of infectious keratitis. *Rev Bras Oftalmol*. 2017; 76(3): 116-20.
181. Yin R, Dai T, Avci P, Jorge AES, Melo WCMA, Vecchio D, et al. Light based anti-infectives: ultraviolet C irradiation, photodynamic therapy, blue light, and beyond. *Curr Opin Pharmacol*. 2013; 13(5): 731-62.
182. Santos A, Silva LD, Sousa LB, de Freitas D, Oliveira LA. Results with the Boston Type I keratoprosthesis after *Acanthamoeba* keratitis. *Am J Ophthalmol Case Rep*. 2017; 6: 71-3.
183. da Silva A, Nobre Jr H, Sampaio L, Nascimento B, da Silva C, de Andrade Neto JB, et al. Antifungal and antiprotozoal green amino acid-based rhamnolipids: mode of action, antibiofilm efficiency and selective activity against resistant *Candida* spp. strains and *Acanthamoeba castellanii*. *Colloids Surf B Biointerfaces*. 2020; 193: 111148.
184. Lopes FC, Tichota DM, Sauter IP, Meira SMM, Segalin J, Rott MB, et al. Active metabolites produced by *Penicillium chrysogenum* IFL1 growing on agro-industrial residues. *Ann Microbiol*. 2013; 63: 771-8.
185. Tavares PL, Ribeiro AC, Berte FK, Hellwig AHS, Pagani DM, Souza CCT, et al. The interaction between *Sporothrix schenckii* sensu stricto and *Sporothrix brasiliensis* with *Acanthamoeba castellanii*. *Mycoses*. 2020; 63(3): 302-7.
186. Albuquerque P, Nicola AM, Magnabosco DAG, Derengowski LS, Crisóstomo LS, Xavier LCG, et al. A hidden battle in the dirt: soil amoebae interactions with *Paracoccidioides* spp. *PLoS Negl Trop Dis*. 2019; 13(10): e0007742.
187. Derengowski LS, Paes HC, Albuquerque P, Tavares AHFP, Fernandes L, Silva-Pereira I, et al. The transcriptional response of *Cryptococcus neoformans* to ingestion by *Acanthamoeba castellanii* and macrophages provides insights into. *Eukaryot Cell*. 2013; 12(5): 761-74.
188. Rizzo J, Albuquerque PC, Wolf JM, Nascimento R, Pereira MD, Nosanchuk JD, et al. Analysis of multiple components involved in the interaction between *Cryptococcus neoformans* and *Acanthamoeba castellanii*. *Fungal Biol*. 2017; 121(6-7): 602-14.
189. Ribeiro NS, Santos FM, Garcia AWA, Ferrareze PAG, Fabres LF, Schrank A, et al. Modulation of zinc homeostasis in *Acanthamoeba castellanii* as a possible antifungal strategy against *Cryptococcus gattii*. *Front Microbiol*. 2017; 8: 1-11.
190. Souza TK, Soares SS, Benitez LB, Rott MB. Interaction between methicillin-resistant *Staphylococcus aureus* (MRSA) and *Acanthamoeba polyphaga*. *Curr Microbiol*. 2017; 74(5): 541-9.
191. Hori JI, Pereira MSF, Roy CR, Nagai H, Zamboni DS, Celular B, et al. Identification and functional characterization of K(+) transporters encoded by *Legionella pneumophila* kup genes. *Cell Microbiol*. 2013; 15(12): 2006-19.
192. Silva LKDS, Rodrigues RAL, Andrade ACDS, Hikida H, Andreani J, Levasseur A, et al. Isolation and genomic characterization of a new mimivirus of lineage B from a Brazilian river. *Arch Virol*. 2020; 165(4): 853-63.
193. Boratto PVM, Arantes TS, Silva LCF, Assis FL, Kroon EG, La Scola B, et al. Niemeyer virus: a new mimivirus group A isolate harboring a set of duplicated aminoacyl-tRNA synthetase genes. *Front Microbiol*. 2015; 6: 1-11.
194. Boratto PVM, Oliveira GP, Machado TB, Andrade ACSP, Baudoin J-P, Klose T, et al. Yaravirus: a novel 80-nm virus infecting *Acanthamoeba castellanii*. *Proc Natl Acad Sci*. 2020; 117(28): 202001637.
195. Campos RK, Boratto PV, Assis FL, Aguiar ERGR, Silva LCF, Albarnaz JD, et al. Samba virus: a novel mimivirus from a giant rain forest, the Brazilian Amazon. *Virol J*. 2014; 11(1): 1-11.
196. Abrahão J, Silva L, Silva LS, Khalil JYB, Rodrigues R, Arantes T, et al. Tailed giant Tupanvirus possesses the most complete translational apparatus of the known virosphere. *Nat Commun*. 2018; 9(1): 749.
197. Mougari S, Bekliz M, Abrahão J, Di Pinto F, Levasseur A, La Scola B. Guarani Virophage, a new sputnik-like isolate from a Brazilian lake. *Front Microbiol*. 2019; 10: 1003.
198. Rolland C, Andreani J, Louazani AC, Aherfi S, Francis R, Rodrigues R, et al. Discovery and further studies on Giant Viruses at the IHU mediterranean infection that modified the perception of the virosphere. *Viruses*. 2019; 11(4): 312.
199. Abrahão J, Silva L, Oliveira D, Almeida G. Lack of evidence of mimivirus replication in human PBMCs. *Microbes Infect*. 2018; 20(5): 281-3.
200. Souza F, Rodrigues R, Reis E, Lima M, La Scola B, Abrahão J. In-depth analysis of the replication cycle of Orpheovirus. *Virol J*. 2019; 16(1): 1-11.
201. Silva LCF, Araújo R, Rodrigues L, Oliveira GP, Dornas FP, La Scola B, et al. Microscopic analysis of the Tupanvirus cycle in *Vermamoeba vermiformis*. 2019; 10: 1-9.

202. Andrade ACDSP, Rodrigues RAL, Oliveira GP, Andrade KR, Bonjardim CA, La Scola B, et al. Filling gaps about mimivirus entry, uncoating and morphogenesis. *J Virol.* 2017; 22: e01355-17.
203. Schrad JR, Young EJ, Abrahão JS, Cortines JR, Parent KN. Microscopic characterization of the Brazilian giant samba virus. *Viruses.* 2017; 9(30): 1-16.
204. Borges I, Rodrigues RAL, Dornas FP, Almeida G, Aquino I, Bonjardim CA, et al. Trapping the enemy: *Vermamoeba vermiformis* circumvents *Faustovirus mariensis* dissemination by enclosing viral progeny inside cysts. *J Virol.* 2019; 93(14): 1-19.
205. Rodrigues RAL, Silva LKS, Dornas FP, Oliveira DB, Magalhães TFF, Santos DA, et al. *Mimivirus fibrils* are important for viral attachment to the microbial world by a diverse glycoside interaction repertoire. *J Virol.* 2015; 89(23): 11812-9.
206. Andrade ACSP, Boratto PVM, Rodrigues RAL, Bastos TM, Azevedo BL, Dornas FP, et al. New isolates of Pandoraviruses: contribution to the study of replication cycle steps. *J Virol.* 2019; 93(5): 1-12.
207. Silva LCF, Almeida GMF, Oliveira DB, Dornas FP, Campos RK, La Scola B, et al. A resourceful giant: APMV is able to interfere with the human type I interferon system. *Microbes Infect.* 2014; 16(3): 187-95.
208. Rodrigues RAL, Louazani AC, Picorelli A, Oliveira GP, Lobo FP, Colson P, et al. Analysis of a Marseillevirus transcriptome reveals temporal gene expression profile and host transcriptional shift. *Front Microbiol.* 2020; 11: 1-17.
209. Abrahão J, La Scola B. Editorial: large and giant DNA viruses. *Front Microbiol.* 2019; 10: 1-2.
210. Raoult D, Abrahão J. Editorial overview : the megaviromes. *Curr Opin Microbiol.* 2016; 31: 31-3.
211. Rodrigues RAL, Abrahão JS, Drumond BP, Kroon EG. Giants among larges : how gigantism impacts giant virus entry into amoebae. *Curr Opin Microbiol.* 2016; 31(31): 88-93.
212. Oliveira G, La Scola B, Abrahão J. Giant virus vs amoeba: fight for supremacy. *Virol J.* 2019; 16(1): 1-12.
213. Dornas FP, Silva LCF, Almeida GM, Campos RK, Boratto PVM, Franco-Luiz APM, et al. *Acanthamoeba polyphaga* mimivirus stability in environmental and clinical substrates: implications for virus detection and isolation. *PLoS One.* 2014; 9(2): 16-8.
214. Augusto G, Souza P, Queiroz VF, Lima MT, Vinicius E, Reis DS, et al. Virus goes viral: an educational kit for virology classes. *Virol J.* 2020; 17(1): 1-8.
215. Abrahão JS, Boratto P, Dornas FP, Silva LC, Campos RK, Almeida GMF, et al. Growing a giant: evaluation of the virological parameters for mimivirus production. *J Virol Methods.* 2014; 207: 6-11.
216. Boratto PVM, Dornas FP, Andrade KR, Rodrigues R, Peixoto F, Silva LCF, et al. Amoebas as mimivirus bunkers: increased resistance to UV light, heat and chemical biocides when viruses are carried by amoeba hosts. *Arch Virol.* 2014; 159(5): 1039-43.
217. Campos RK, Andrade KR, Ferreira PCP, Bonjardim CA, La Scola B, Kroon EG, et al. Virucidal activity of chemical biocides against mimivirus, a putative pneumonia agent. *J Clin Virol.* 2012; 55(4): 323-8.
218. Borges IA, Assis FL, Silva LKDS, Abrahão J. Rio Negro virophage: sequencing of the near complete genome and transmission electron. *Brazilian J Microbiol.* 2018; 49(Suppl. 1): 260-1.
219. Mougari S, Abrahao J, Oliveira GP, Bou Khalil JY, La Scola B. Role of the R349 gene and its repeats in the MIMIVIRE defense system. *Front Microbiol.* 2019; 10: 1-10.
220. Arantes TS, Rodrigues RAL, Silva LKS, Oliveira GP, de Souza HL, Khalil JYB, et al. The large Marseillevirus explores different entry pathways by forming giant infectious vesicles. *J Virol.* 2016; 90(11): 5246-55.
221. Oliveira GP, Lima MT, Arantes TS, Assis FL, Rodrigues RAL, da Fonseca FG, et al. The investigation of promoter sequences of Marseilleviruses highlights a remarkable abundance of the AAATATTT motif in intergenic regions. *J Virol.* 2017; 91(21): 1-10.
222. Hofstatter PG, Ribeiro GM, Porfírio-Sousa AL, Lahr DJG. The sexual ancestor of all eukaryotes: a defense of the “meiosis toolkit”: a rigorous survey supports the obligate link between meiosis machinery and sexual recombination. *BioEssays.* 2020; 42(9): 1-10.
223. Saberi R, Seifi Z, Dodangeh S, Najafi A, Abdollah Hosseini S, Anvari D, et al. A systematic literature review and meta-analysis on the global prevalence of *Naegleria* spp. in water sources. *Trans-bound Emerg Dis.* 2020; 67: 2389-402.
224. Delafont V, Rodier MH, Maisonneuve E, Cateau E. *Vermamoeba vermiformis*: a free-living amoeba of interest. *Microbial Ecology.* 2018; 76: 991-1001.

TABLE I  
List of references included in the present literature review

| Year        | Title                                                                                                                                                                                        | Category <sup>a</sup> | Target<br>FLA <sup>b</sup> | # <sup>c</sup> |
|-------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|----------------------------|----------------|
| <b>CORE</b> |                                                                                                                                                                                              |                       |                            |                |
| 1974        | Isolation of nuclei and characterization of nuclear DNA of <i>Acanthamoeba castellanii</i>                                                                                                   | W                     | A                          | (1)            |
| 1975        | Characterization of the genome of the small free-living amoeba <i>Acanthamoeba castellanii</i>                                                                                               | W                     | A                          | (2)            |
| 1977        | Primary amoebic meningoencephalitis: first case from Latin-America [Meningoencefalite por ameba de vida livre apresentacao do primeiro caso latino-americano]                                | C                     | N                          | (3)            |
| 1978        | Primary amebic meningoencephalitis - case-report                                                                                                                                             | C                     | N                          | (4)            |
| 1985        | Free-living amoebae in human intestine: evidence of parasitism [Amebas de vida livre no intestino humano evidencias de parasitismo]                                                          | C                     | A, W                       | (5)            |
| 1986        | Isolation of <i>Naegleria fowleri</i> from a lake in the city of Rio de Janeiro, Brazil                                                                                                      | E                     | N                          | (6)            |
| 1988        | <i>Acanthamoeba culbertsoni</i> : susceptibility to soft contact lens disinfection systems                                                                                                   | W                     | A                          | (7)            |
| 1992        | <i>Acanthamoeba healyi</i> N. sp. and the Isoenzyme and Immunoblot Profiles of <i>Acanthamoeba</i> spp., Groups 1 and 3                                                                      | W                     | A                          | (8)            |
| 1992        | Granulomatous amoebic encephalitis due to leptomycid amoebae: Report of the first Brazilian case                                                                                             | C                     | A                          | (9)            |
| 2000        | <i>Acanthamoeba keratitis</i> [Ceratite por <i>Acanthamoeba</i> ]                                                                                                                            | R                     | A                          | (10)           |
| 2000        | Proteinase activities in total extracts and in medium conditioned by <i>Acanthamoeba polyphaga</i> trophozoites                                                                              | W                     | A                          | (11)           |
| 2000        | Random amplified polymorphic DNA profiles as a tool for the characterization of Brazilian keratitis isolates of the genus <i>Acanthamoeba</i>                                                | C                     | A                          | (12)           |
| 2003        | Diagnosis of <i>Acanthamoeba keratitis</i> with confocal microscopy [Microscopia confocal no diagnóstico de ceratite por <i>Acanthamoeba</i> ]                                               | C                     | A                          | (13)           |
| 2003        | Isolation of potentially pathogenic free-living amoebas in hospital dust                                                                                                                     | E                     | A,N                        | (14)           |
| 2004        | Confocal microscopy in early diagnosis of <i>Acanthamoeba keratitis</i>                                                                                                                      | C                     | A                          | (15)           |
| 2004        | Conjunctivitis presumably due to <i>Acanthamoeba</i> [Conjuntivite presumível por <i>Acanthamoeba</i> ]                                                                                      | C                     | A                          | (16)           |
| 2006        | The interaction between <i>Acanthamoeba polyphaga</i> and human osteoblastic cells <i>in vitro</i>                                                                                           | W                     | A                          | (17)           |
| 2006        | The interaction between the amoeba <i>Balamuthia mandrillaris</i> and extracellular matrix glycoproteins <i>in vitro</i>                                                                     | W                     | B                          | (18)           |
| 2007        | Biological characterization of a clinical and an environmental isolate of <i>Acanthamoeba polyphaga</i> : analysis of relevant parameters to decode pathogenicity                            | W                     | A                          | (19)           |
| 2007        | Diagnosis of <i>Acanthamoeba</i> corneal Infection by impression cytology: Case report [Citologia de impressão no diagnóstico de infecção corneana por <i>Acanthamoeba</i> : Relato de caso] | C                     | A                          | (20)           |
| 2007        | Disseminated <i>Balamuthia mandrillaris</i> amoeba infection in an AIDS patient from Brazil                                                                                                  | C                     | B                          | (21)           |
| 2007        | Isolation and identification of potentially pathogenic free-living amoebae in samples from environments in a public hospital in the city of Porto Alegre, Rio Grande do Sul                  | E                     | A                          | (22)           |
| 2008        | <i>Acanthamoeba</i> spp. and bacterial contamination in contact lens storage cases and the relationship to user profiles                                                                     | E                     | A                          | (23)           |
| 2008        | <i>Acanthamoeba</i> spp. cysts storage in filter paper                                                                                                                                       | W                     | A                          | (24)           |
| 2008        | Corneal graft survival after therapeutic keratoplasty for <i>Acanthamoeba keratitis</i>                                                                                                      | C                     | A                          | (25)           |
| 2008        | Growth, encystment and survival of <i>Acanthamoeba castellanii</i> grazing on different bacteria                                                                                             | W                     | A                          | (26)           |
| 2008        | <i>In vitro</i> evaluation of the amebicidal activity of <i>Pterocaulon polystachyum</i> ( <i>Asteraceae</i> ) against trophozoites of <i>Acanthamoeba castellanii</i>                       | W                     | A                          | (27)           |
| 2009        | <i>Acanthamoeba</i> interaction with extracellular matrix glycoproteins: Biological and biochemical characterization and Role in Cytotoxicity and Invasiveness                               | W                     | A                          | (28)           |
| 2009        | <i>Acanthamoeba</i> spp. in urine of critically ill patients                                                                                                                                 | C                     | A                          | (29)           |
| 2009        | Differential effects of alpha-helical and beta-hairpin antimicrobial peptides against <i>Acanthamoeba castellanii</i>                                                                        | W                     | A                          | (30)           |
| 2009        | Elastase secretion in <i>Acanthamoeba polyphaga</i>                                                                                                                                          | W                     | A                          | (31)           |
| 2009        | Genotyping, physiological features and proteolytic activities of a potentially pathogenic <i>Acanthamoeba</i> sp. isolated from tap water in Brazil                                          | E                     | A                          | (32)           |
| 2009        | Potentially pathogenic <i>Acanthamoeba</i> in swimming pools: A survey in the southern Brazilian city of Porto Alegre                                                                        | E                     | A                          | (33)           |
| 2009        | Prevalence of potentially pathogenic free-living amoebae from <i>Acanthamoeba</i> and <i>Naegleria</i> Genera in non-hospital, public, internal environments from the city of Santos, Brazil | E                     | A,N                        | (34)           |
| 2009        | Twenty Years of <i>Acanthamoeba</i> Keratitis                                                                                                                                                | C                     | A                          | (35)           |
| 2010        | Characterization of <i>Acanthamoeba</i> isolates from dust of a public hospital in Curitiba, Paraná, Brazil                                                                                  | E                     | A                          | (36)           |
| 2010        | Granulomatous Amoebic Meningoencephalitis in an Immunocompetent Patient                                                                                                                      | C                     | B                          | (37)           |

| Year | Title                                                                                                                                                                                                                                                                | Category <sup>a</sup> | Target FLA <sup>b</sup> | # <sup>c</sup> |
|------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|-------------------------|----------------|
| 2010 | <i>In vitro</i> amoebicidal activity of S-nitrosoglutathione and S-nitroso-N- acetylcysteine against trophozoites of <i>Acanthamoeba castellanii</i>                                                                                                                 | W                     | A                       | (38)           |
| 2010 | Potentially Pathogenic <i>Acanthamoeba</i> Isolated from a Hospital in Brazil                                                                                                                                                                                        | E                     | A                       | (39)           |
| 2010 | Susceptibility of <i>Aedes aegypti</i> (Diptera: Culicidae) to <i>Acanthamoeba polyphaga</i> (Sarcomastigophora: Acanthamoebidae)                                                                                                                                    | W                     | A                       | (40)           |
| 2010 | The fine structure of the <i>Acanthamoeba polyphaga</i> cyst wall                                                                                                                                                                                                    | W                     | A                       | (41)           |
| 2011 | <i>Acanthamoeba</i> T3, T4 and T5 in swimming-pool waters from Southern Brazil                                                                                                                                                                                       | W                     | A                       | (42)           |
| 2011 | Amoebicidal activity and chemical composition of <i>Pterocaulon polystachyum</i> (Asteraceae) essential oil                                                                                                                                                          | W                     | A                       | (43)           |
| 2011 | Assessing efficacy of combined riboflavin and UV-A light (365 nm) treatment of <i>Acanthamoeba</i> trophozoites                                                                                                                                                      | C                     | A                       | (44)           |
| 2011 | Bacteriocin-like substance from <i>Bacillus amyloliquefaciens</i> shows remarkable inhibition of <i>Acanthamoeba polyphaga</i>                                                                                                                                       | W                     | A                       | (45)           |
| 2011 | Prevalence of <i>Acanthamoeba</i> from tap water in Rio Grande do Sul, Brazil                                                                                                                                                                                        | E                     | A                       | (46)           |
| 2011 | Serine-like proteolytic enzymes correlated with differential pathogenicity in patients with acute <i>Acanthamoeba</i> keratitis                                                                                                                                      | C                     | A                       | (47)           |
| 2012 | A Rapid and Reliable Method for the Clonal Isolation of <i>Acanthamoeba</i> from Environmental Samples                                                                                                                                                               | E                     | A                       | (48)           |
| 2012 | Chemical composition and amoebicidal activity of <i>Croton pallidulus</i> , <i>Croton ericoides</i> , and <i>Croton isabelli</i> (Euphorbiaceae) essential oils                                                                                                      | W                     | A                       | (49)           |
| 2012 | Chemical composition and amoebicidal activity of <i>Piper hispidinervum</i> (Piperaceae) essential oil                                                                                                                                                               | W                     | A                       | (50)           |
| 2012 | Meningoencephalitis caused by <i>Naegleria fowleri</i> in cattle of northeast Brazil                                                                                                                                                                                 | C                     | N                       | (51)           |
| 2012 | Occurrence and characterization of <i>Acanthamoeba</i> similar to genotypes T4, T5, and T2/T6 isolated from environmental sources in Brasília, Federal District, Brazil                                                                                              | E                     | A                       | (52)           |
| 2012 | Prevalence of <i>Acanthamoeba</i> spp. (Sarcomastigophora: Acanthamoebidae) in wild populations of <i>Aedes aegypti</i> (Diptera: Culicidae)                                                                                                                         | E                     | A                       | (53)           |
| 2012 | Revisiting the <i>Acanthamoeba</i> species that form star-shaped cysts (genotypes T7, T8, T9, and T17): Characterization of seven new Brazilian environmental isolates and phylogenetic inferences                                                                   | E                     | A                       | (54)           |
| 2013 | Amoebicidal activity of phytosynthesised silver nanoparticles and their <i>in vitro</i> cytotoxicity to human cells                                                                                                                                                  | W                     | A                       | (55)           |
| 2013 | Antimicrobial action of biguanides on the viability of <i>Acanthamoeba</i> cysts and assessment of cell toxicity                                                                                                                                                     | C                     | A                       | (56)           |
| 2013 | <i>In vitro</i> effect of <i>Acanthospermum australe</i> (Asteraceae) extracts on <i>Acanthamoeba polyphaga</i> trophozoites [Efeito <i>in vitro</i> de extratos de <i>Acanthospermum australe</i> (Asteraceae) sobre trofozoitos de <i>Acanthamoeba polyphaga</i> ] | W                     | A                       | (57)           |
| 2013 | Infection in a rat model reactivates attenuated virulence after long-term axenic culture of <i>Acanthamoeba</i> spp                                                                                                                                                  | W                     | A                       | (58)           |
| 2013 | Isolation and genotyping of free-living environmental isolates of <i>Acanthamoeba</i> spp. from bromeliads in Southern Brazil                                                                                                                                        | E                     | A                       | (59)           |
| 2013 | Morphological, genotypic, and physiological characterization of <i>Acanthamoeba</i> isolates from keratitis patients and the domestic environment in Vitoria, Espírito Santo, Brazil                                                                                 | C                     | A                       | (60)           |
| 2013 | Physiological, morphological, and immunochemical parameters used for the characterization of clinical and environmental isolates of <i>Acanthamoeba</i>                                                                                                              | E                     | A                       | (61)           |
| 2013 | Selenocysteine biosynthesis and insertion machinery in <i>Naegleria gruberi</i>                                                                                                                                                                                      | W                     | N                       | (62)           |
| 2013 | Susceptibility of <i>Acanthamoeba</i> to multipurpose lens-cleaning solutions                                                                                                                                                                                        | W                     | A                       | (63)           |
| 2014 | Characterization of isolates of <i>Acanthamoeba</i> from the nasal mucosa and cutaneous lesions of dogs                                                                                                                                                              | C                     | A                       | (64)           |
| 2014 | Proteomic profiling of the infective trophozoite stage of <i>Acanthamoeba polyphaga</i>                                                                                                                                                                              | W                     | A                       | (65)           |
| 2015 | A method for microbial decontamination of <i>Acanthamoeba</i> cultures using the peritoneal cavity of mice                                                                                                                                                           | W                     | A                       | (66)           |
| 2015 | <i>Acanthamoeba</i> misidentification and multiple labels: redefining genotypes T16, T19, and T20 and proposal for <i>Acanthamoeba micheli</i> sp. nov. (genotype T19)                                                                                               | W                     | A                       | (67)           |
| 2015 | <i>Acanthamoeba polyphaga</i> mimivirus prevents amoebal encystment-mediating serine proteinase expression and circumvents cell encystment                                                                                                                           | W                     | A                       | (68)           |
| 2015 | <i>Acanthamoeba</i> T4, T5 and T11 Isolated From Mineral Water Bottles in Southern Brazil                                                                                                                                                                            | E                     | A                       | (69)           |
| 2015 | Cytotoxic activity and degradation patterns of structural proteins by corneal isolates of <i>Acanthamoeba</i> spp                                                                                                                                                    | W                     | A                       | (70)           |
| 2015 | Identification of <i>Paenibacillus</i> as a Symbiont in <i>Acanthamoeba</i>                                                                                                                                                                                          | E                     | A                       | (71)           |
| 2015 | Identification of pseudomonas spp. as amoeba-resistant microorganisms in isolates of <i>Acanthamoeba</i>                                                                                                                                                             | W                     | A                       | (72)           |
| 2015 | Modulation of the expression of mimivirus-encoded translation-related genes in response to nutrient availability during <i>Acanthamoeba castellanii</i> infection                                                                                                    | W                     | A                       | (73)           |
| 2015 | Systemic acanthamoebiasis associated with canine distemper in dogs in the semiarid region of Paraíba, Brazil                                                                                                                                                         | C                     | A                       | (74)           |
| 2016 | <i>Acanthamoeba</i> and <i>Fusarium</i> interactions: A possible problem in keratitis                                                                                                                                                                                | W                     | A                       | (75)           |
| 2016 | <i>Acanthamoeba</i> and mimivirus interactions: The role of amoebal encystment and the expansion of the ‘Cheshire Cat’ theory                                                                                                                                        | R                     | A                       | (76)           |



| Year | Title                                                                                                                                                                                     | Category <sup>a</sup> | Target<br>FLA <sup>b</sup> | # <sup>c</sup> |
|------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|----------------------------|----------------|
| 2016 | <i>Acanthamoeba</i> spp. as a universal host for pathogenic microorganisms: One bridge from environment to host virulence                                                                 | R                     | A                          | (77)           |
| 2016 | Amebicidal activity of the essential oils of <i>Lippia</i> spp. (Verbenaceae) against <i>Acanthamoeba polyphaga</i> trophozoites                                                          | W                     | A                          | (78)           |
| 2016 | Crystallisation and Crystallographic Analyses of Triose phosphatelsomerase from <i>Naegleria gruberi</i>                                                                                  | W                     | N                          | (79)           |
| 2016 | Detection and quantification of human Adenovirus genomes in <i>Acanthamoeba</i> isolated from swimming pools                                                                              | E                     | A                          | (80)           |
| 2016 | Evaluation of the immunodiagnostic potential of a recombinant surface protein domain from <i>Acanthamoeba castellanii</i>                                                                 | W                     | A                          | (81)           |
| 2016 | Experimental infection of T4 <i>Acanthamoeba</i> genotype determines the pathogenic potential                                                                                             | W                     | A                          | (82)           |
| 2016 | Expression, purification, enzymatic characterization and crystallization of glyceraldehyde-3-phosphate dehydrogenase from <i>Naegleria gruberi</i> , the first one from phylum Percolozoa | W                     | N                          | (83)           |
| 2016 | Isolation and identification of <i>Acanthamoeba</i> spp. from thermal swimming pools and spas in Southern Brazil                                                                          | E                     | A                          | (84)           |
| 2016 | Photodynamic inactivation of <i>Acanthamoeba polyphaga</i> with curcuminoids: an <i>in vitro</i> study                                                                                    | W                     | A                          | (85)           |
| 2016 | Uncovering Cryptic Diversity in Two Amoebozoan Species Using Complete Mitochondrial Genome Sequences                                                                                      | W                     | A, W                       | (86)           |
| 2017 | Anaerobic Metabolism in T4 <i>Acanthamoeba</i> Genotype                                                                                                                                   | W                     | A                          | (87)           |
| 2017 | Interaction Between Methicillin-Resistant <i>Staphylococcus aureus</i> (MRSA) and <i>Acanthamoeba polyphaga</i>                                                                           | W                     | A                          | (88)           |
| 2017 | Modulation of zinc homeostasis in <i>Acanthamoeba castellanii</i> as a possible antifungal strategy against <i>Cryptococcus gattii</i>                                                    | W                     | A                          | (89)           |
| 2017 | Molecular diagnosis of <i>Acanthamoeba</i> keratitis: evaluation in rat model and application in suspected human cases                                                                    | C                     | A                          | (90)           |
| 2017 | Nanoemulsions Containing a Coumarin-Rich Extract from <i>Pterocaulon balansae</i> (Asteraceae) for the Treatment of Ocular <i>Acanthamoeba</i> keratitis                                  | W                     | A                          | (91)           |
| 2017 | Occurrence of Infected Free-Living Amoebae in Cooling Towers of Southern Brazil                                                                                                           | E                     | A, N, W                    | (92)           |
| 2017 | Sensitivity of Enzymatic Toxins from Corneal Isolate of <i>Acanthamoeba</i> Protozoan to Physicochemical Parameters                                                                       | C                     | A                          | (93)           |
| 2017 | siRNA-loaded liposomes: Inhibition of encystment of <i>Acanthamoeba</i> and toxicity on the eye surface                                                                                   | W                     | A                          | (94)           |
| 2017 | Therapeutic agents and biocides for ocular infections by free-living amoebae of <i>Acanthamoeba</i> genus                                                                                 | R                     | A                          | (95)           |
| 2018 | <i>Acanthamoeba</i> keratitis in patients wearing scleral contact lenses                                                                                                                  | C                     | A                          | (96)           |
| 2018 | <i>Acanthamoeba</i> keratitis in Porto Alegre (southern Brazil): 28 cases and risk factors                                                                                                | C                     | A                          | (97)           |
| 2018 | <i>Acanthamoeba</i> of three morphological groups and distinct genotypes exhibit variable and weakly inter-related physiological properties                                               | E                     | A                          | (98)           |
| 2018 | Comparative proteomic analysis of soluble and surface-enriched proteins from <i>Acanthamoeba castellanii</i> trophozoites                                                                 | W                     | A                          | (99)           |
| 2018 | Contact lens-related polymicrobial keratitis: <i>Acanthamoeba</i> spp. genotype T4 and <i>Candida albicans</i>                                                                            | C                     | A                          | (100)          |
| 2018 | Extracellular vesicles and vesicle-free secretome of the protozoa <i>Acanthamoeba castellanii</i> under homeostasis and nutritional stress and their damaging potential to host cells     | W                     | A                          | (101)          |
| 2018 | Structural studies of glyceraldehyde-3-phosphate dehydrogenase from <i>Naegleria gruberi</i> , the first one from phylum Percolozoa                                                       | S                     | N                          | (102)          |
| 2018 | The first <i>Acanthamoeba</i> keratitis case in the midwest region of Brazil: Diagnosis, genotyping of the parasite and disease outcome                                                   | C                     | A                          | (103)          |
| 2018 | The therapeutic strategies against <i>Naegleria fowleri</i>                                                                                                                               | R                     | N                          | (104)          |
| 2018 | Virulent T4 <i>Acanthamoeba</i> causing keratitis in a patient after swimming while wearing contact lenses in Southern Brazil                                                             | C                     | A                          | (105)          |
| 2019 | Box Behnken design of siRNA-loaded liposomes for the treatment of a murine model of ocular keratitis caused by <i>Acanthamoeba</i>                                                        | W                     | A                          | (106)          |
| 2019 | Extracellular vesicles from the protozoa <i>Acanthamoeba castellanii</i> : Their role in pathogenesis, environmental adaptation and potential applications                                | R                     | A                          | (107)          |
| 2019 | Genomic Signatures Among <i>Acanthamoeba polyphaga</i> Entoorganisms Unveil Evidence of Coevolution                                                                                       | S                     | A                          | (108)          |
| 2019 | Intra-amoebic localisation of <i>Arcobacter butzleri</i> as an endocytobiont of <i>Acanthamoeba castellanii</i>                                                                           | W                     | A                          | (109)          |
| 2019 | Meningoencephalitis due to <i>Naegleria fowleri</i> in cattle in southern Brazil [Meningoencefalite por <i>Naegleria fowleri</i> em um bovino no sul do Brasil]                           | C                     | N                          | (110)          |
| 2019 | Recovery of an <i>Acanthamoeba</i> strain with two group I introns in the nuclear 18S rRNA gene                                                                                           | W                     | A                          | (111)          |
| 2019 | Unravelling the interactions of the environmental host <i>Acanthamoeba castellanii</i> with fungi through the recognition by mannose-binding proteins                                     | W                     | A                          | (112)          |

| Year             | Title                                                                                                                                                                       | Category <sup>a</sup> | Target FLA <sup>b</sup> | # <sup>c</sup> |
|------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|-------------------------|----------------|
| 2020             | <i>Acanthamoeba castellanii</i> as an alternative interaction model for the dermatophyte <i>Trichophyton rubrum</i>                                                         | W                     | A                       | (113)          |
| 2020             | <i>Acanthamoeba castellanii</i> phosphate transporter (AcPHS) is important to maintain inorganic phosphate influx and is related to trophozoite metabolic processes         | W                     | A                       | (114)          |
| 2020             | <i>Acanthamoeba</i> spp. monoclonal antibody against a CPA2 transporter: a promising molecular tool for acanthamoebiasis diagnosis and encystment study                     | W                     | A                       | (115)          |
| 2020             | Extracellular protease profile of <i>Acanthamoeba</i> after prolonged axenic culture and after interaction with MDCK cells                                                  | W                     | A                       | (116)          |
| 2020             | Free-living amoebae and its relationship with air quality in hospital environments: Isolation and characterization of <i>Acanthamoeba</i> spp. from air-conditioning system | E                     | A                       | (117)          |
| 2020             | Identification of T3 and T4 Genotypes of <i>Acanthamoeba</i> sp. in Dust Samples Isolated from Air Conditioning Equipment of Public Hospital of Ituiutaba-MG                | E                     | A, B                    | (118)          |
| 2020             | <i>In vitro</i> amoebicidal activity of imidazolium salts against trophozoites                                                                                              | W                     | A                       | (119)          |
| 2020             | Isolation of <i>Naegleria</i> spp. From a Brazilian water source                                                                                                            | E                     | N,W                     | (120)          |
| <b>AUXILIARY</b> |                                                                                                                                                                             |                       |                         |                |
| 2013             | Distribution of infectious keratitis in a tertiary hospital in Brazil                                                                                                       | C                     | A                       | (121)          |
| 2008             | Contact lens-associated microbial keratitis                                                                                                                                 | C                     | A                       | (122)          |
| 2017             | Results with the Boston Type I keratoprosthesis after <i>Acanthamoeba</i> keratitis                                                                                         | C                     | A                       | (123)          |
| 2017             | Epidemiological profile of infectious keratitis                                                                                                                             | C                     | A                       | (124)          |
| 2015             | Degeneration and Regeneration of Sub basal Corneal Nerves after Infectious Keratitis: A Longitudinal <i>in Vivo</i> Confocal Microscopy Study                               | C                     | A                       | (125)          |
| 2018             | Central nervous system disorders diagnosed in dogs                                                                                                                          | C                     | A, N, B                 | (126)          |
| 2013             | Yaravirus: A novel 80-nm virus infecting <i>Acanthamoeba castellanii</i>                                                                                                    | E                     | A                       | (127)          |
| 2020             | Occurrence of parasites in salads in restaurants in Aparecida de Goiânia, Goiás, Brazil                                                                                     | E                     | A                       | (128)          |
| 2018             | Genotypic characterization and assessment of infectivity of human waterborne pathogens recovered from oysters and estuarine waters in Brazil                                | E                     | A                       | (129)          |
| 2020             | Isolation and genomic characterization of a new mimivirus of lineage B from a Brazilian river                                                                               | E                     | A                       | (130)          |
| 2019             | A hidden battle in the dirt: Soil amoebae interactions with <i>Paracoccidioides</i> spp                                                                                     | E                     | A, W                    | (131)          |
| 2016             | Giants among larges: How gigantism impacts giant virus entry into amoebae                                                                                                   | R                     | A                       | (132)          |
| 2011             | Diffusion Imaging in Brain Infections                                                                                                                                       | R                     | A                       | (133)          |
| 2014             | <i>Acanthamoeba polyphaga</i> mimivirus and other giant viruses: An open field to outstanding discoveries                                                                   | R                     | A                       | (134)          |
| 2020             | The Sexual Ancestor of all Eukaryotes: A Defense of the “ Meiosis Toolkit ” A Rigorous Survey Supports the Obligate Link between Meiosis Machinery and Sexual Recombination | R                     | A                       | (135)          |
| 2019             | Discovery and further studies on giant viruses at the IHU mediterranean infection that modified the perception of the virosphere                                            | R                     | A                       | (136)          |
| 2016             | Editorial overview: The megaviromes                                                                                                                                         | R                     | A                       | (137)          |
| 2013             | Light based anti-infectives: Ultraviolet C irradiation, photodynamic therapy, blue light, and beyond                                                                        | R                     | A                       | (138)          |
| 2019             | Emerging infectious diseases with cutaneous manifestations Fungal, helminthic, protozoan and ectoparasitic infections                                                       | R                     | B                       | (139)          |
| 2012             | Co-infection of HIV and tropical infectious agents that affect the nervous system                                                                                           | R                     | A,N,B                   | (140)          |
| 2017             | Emerging Infections and Pertinent Infections Related to Travel for Patients with Primary Immunodeficiencies                                                                 | R                     | A,N,S,B                 | (141)          |
| 2011             | A morphological approach to the diagnosis of protozoal infections of the central nervous system                                                                             | R                     | A,N,S,B                 | (142)          |
| 2019             | Giant virus vs amoeba: Fight for supremacy                                                                                                                                  | R                     | A, W                    | (143)          |
| 2019             | Editorial: Large and giant DNA viruses                                                                                                                                      | R                     | A, W                    | (144)          |
| 2019             | Guarani virophage, a new sputnik-like isolate from a Brazilian lake                                                                                                         | W                     | A                       | (145)          |
| 2019             | Microscopic Analysis of the Tupanvirus Cycle in <i>Vermamoeba vermiformis</i>                                                                                               | W                     | A                       | (146)          |
| 2018             | Lack of evidence of mimivirus replication in human PBMCs                                                                                                                    | W                     | A                       | (147)          |
| 2017             | Filling Knowledge Gaps for Mimivirus Entry, Uncoating, and Morphogenesis                                                                                                    | W                     | A                       | (148)          |
| 2013             | Identification and functional characterization of K <sup>+</sup> transporter encoded by <i>Legionella pneumophila</i> kup genes                                             | W                     | A                       | (149)          |
| 2019             | Transcriptional analysis of flagellar and putative virulence genes of <i>Arcobacter butzleri</i> as an endocytobiont of <i>Acanthamoeba castellanii</i>                     | W                     | A                       | (150)          |
| 2014             | <i>Acanthamoeba polyphaga</i> mimivirus stability in environmental and clinical substrates: Implications for virus detection and isolation                                  | W                     | A                       | (151)          |




| Year | Title                                                                                                                                                                                                             | Category <sup>a</sup> | Target FLA <sup>b</sup> | # <sup>c</sup> |
|------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|-------------------------|----------------|
| 2020 | Antifungal and antiprotozoal green amino acid-based rhamnolipids: Mode of action, antibiofilm efficiency and selective activity against resistant <i>Candida</i> spp. strains and <i>Acanthamoeba castellanii</i> | W                     | A                       | (152)          |
| 2020 | Analysis of a Marseillevirus Transcriptome Reveals Temporal Gene Expression Profile and Host Transcriptional Shift                                                                                                | W                     | A                       | (153)          |
| 2020 | The interaction between <i>Sporothrix schenckii</i> sensu stricto and <i>Sporothrix brasiliensis</i> with <i>Acanthamoeba castellanii</i>                                                                         | W                     | A                       | (154)          |
| 2019 | New Isolates of Pandoraviruses: Contribution to the Study of Replication Cycle Steps                                                                                                                              | W                     | A                       | (155)          |
| 2019 | Role of the R349 gene and its repeats in the mimivire defense system                                                                                                                                              | W                     | A                       | (156)          |
| 2018 | Rio Negro virophage: Sequencing of the near complete genome and transmission electron microscopy of viral factories and particles                                                                                 | W                     | A                       | (157)          |
| 2018 | Analyses of the Kroon virus major capsid gene and its transcript highlight a distinct pattern of gene evolution and splicing among mimiviruses                                                                    | W                     | A                       | (158)          |
| 2017 | The investigation of promoter sequences of marseilleviruses highlights a remarkable abundance of the AAATATTT motif in intergenic regions                                                                         | W                     | A                       | (159)          |
| 2017 | Analysis of multiple components involved in the interaction between <i>Cryptococcus neoformans</i> and <i>Acanthamoeba castellanii</i>                                                                            | W                     | A                       | (160)          |
| 2017 | Microscopic characterization of the Brazilian giant samba virus                                                                                                                                                   | W                     | A                       | (161)          |
| 2016 | The large marseillevirus explores different entry pathways by forming giant infectious vesicles                                                                                                                   | W                     | A                       | (162)          |
| 2016 | Different Photoresponses of Microorganisms: From Bioinhibition to Biostimulation                                                                                                                                  | W                     | A                       | (163)          |
| 2016 | A Brazilian marseillevirus is the founding member of a lineage in family marseilleviridae                                                                                                                         | W                     | A                       | (164)          |
| 2015 | Mimivirus fibrils are important for viral attachment to the microbial world by a diverse glycoside interaction repertoire                                                                                         | W                     | A                       | (165)          |
| 2015 | <i>Niemeyer</i> virus: A new mimivirus group A isolate harboring a set of duplicated aminoacyl-tRNA synthetase genes                                                                                              | W                     | A                       | (166)          |
| 2014 | Samba virus: A novel mimivirus from a giant rain forest, the Brazilian Amazon                                                                                                                                     | W                     | A                       | (167)          |
| 2014 | Amoebas as mimivirus bunkers: Increased resistance to UV light, heat and chemical biocides when viruses are carried by amoeba hosts                                                                               | W                     | A                       | (168)          |
| 2014 | A resourceful giant: APMV is able to interfere with the human type I interferon system                                                                                                                            | W                     | A                       | (169)          |
| 2013 | Active metabolites produced by <i>Penicillium chrysogenum</i> IFL1 growing on agro-industrial residues                                                                                                            | W                     | A                       | (170)          |
| 2013 | The Transcriptional Response of <i>Cryptococcus neoformans</i> to Ingestion by <i>Acanthamoeba castellanii</i> and Macrophages Provides Insights into the Evolutionary Adaptation to the Mammalian Host           | W                     | A                       | (171)          |
| 2012 | Virucidal activity of chemical biocides against mimivirus, a putative pneumonia agent                                                                                                                             | W                     | A                       | (172)          |
| 2019 | In-depth analysis of the replication cycle of Orpheovirus                                                                                                                                                         | W                     | W                       | (173)          |
| 2019 | Trapping the Enemy: <i>Vermamoeba vermiformis</i> Circumvents <i>Faustovirus mariensis</i> Dissemination by Enclosing Viral Progeny inside Cysts                                                                  | W                     | A, W                    | (174)          |
| 2020 | Virus goes viral: An educational kit for virology classes                                                                                                                                                         | W                     | A,W                     | (175)          |
| 2018 | Tailed giant Tupanvirus possesses the most complete translational apparatus of the known virosphere                                                                                                               | W                     | A,W                     | (176)          |
| 2015 | Isolation of new Brazilian giant viruses from environmental samples using a panel of protozoa                                                                                                                     | W                     | A,W                     | (177)          |
| 2014 | Growing a giant: Evaluation of the virological parameters for Mimivirus production                                                                                                                                | W                     | A                       | (178)          |

*a*: category: Wet-bench(W); Dry-bench(S); Review(R); Environmental(E); Clinical(C). *b*: Target FLA: *Acanthamoeba* (A), *Vermamoeba* (V), *Naegleria* (N), *Balamuthia* (B). *c*: the numbers do not correspond to the reference numbers in the original paper.

TABLE II  
Key words (I), acronyms (II) and inclusion/exclusion criteria (III) used to search in the databases

| I. KEY WORDS                                         |                                                                                              |                                                                                                                                                         |
|------------------------------------------------------|----------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| #                                                    | Key word                                                                                     | Synonyms/misspelling                                                                                                                                    |
| 1                                                    | naegleria                                                                                    | OR negleria                                                                                                                                             |
| 2                                                    | free-living amoeba                                                                           | free living amoeba OR freeliving amoeba OR free-living amoebic OR freeliving amoebic OR free-living amebic OR free living amoebic OR free living amebic |
| 3                                                    | amebiasis                                                                                    | OR amoebiasis                                                                                                                                           |
| 4                                                    | amoebic infection                                                                            | OR amebic infection                                                                                                                                     |
| 5                                                    | primary amoebic meningoencephalitis                                                          | primary meningoencephalitis OR primary amebic meningoencephalitis OR amoebic meningoencephalitis OR amebic meningoencephalitis                          |
| 6                                                    | amoebic encephalitis                                                                         | OR amebic encephalitis                                                                                                                                  |
| 7                                                    | granulomatous amoebic encephalitis                                                           | granulomatous encephalitis OR granulomatous amebic encephalitis OR granulomatus amoebic encephalitis                                                    |
| 8                                                    | Acanthamoeba                                                                                 | OR Acanthamoeba OR Acanthameoba OR Acanthamebiasis OR Acanthameba                                                                                       |
| 9                                                    | amoebic keratitis                                                                            | OR amebic keratitis OR amoeba keratitis OR Acanthamoeba keratitis                                                                                       |
| 10                                                   | Sappinia                                                                                     | OR Sappinea OR Sapinia OR sappinia amoebic encephalitis                                                                                                 |
| 11                                                   | Balamuthia                                                                                   | OR Ballamuthia OR Balamuthias OR Balamuthiasis                                                                                                          |
| 12                                                   | Vermamoeba                                                                                   | OR Vermameba                                                                                                                                            |
| 13                                                   | Hartmannella                                                                                 | OR Hartmanella                                                                                                                                          |
| 14                                                   | Brazil                                                                                       | OR brasil                                                                                                                                               |
| II. ACRONYMS                                         |                                                                                              |                                                                                                                                                         |
| Key word #                                           | Web of Science                                                                               | Outcome (# of papers)                                                                                                                                   |
| #1                                                   | TI = (naegleria OR negleria) OR AB = (naegleria OR negleria) OR AK = (naegleria OR negleria) | 1432                                                                                                                                                    |
| #14                                                  | AD =(Brazil OR brasil)                                                                       | 1101096                                                                                                                                                 |
| #1 AND #14                                           | (#1 AND #14)                                                                                 | 14                                                                                                                                                      |
| Key word #                                           | PubMed                                                                                       | Outcome (# of papers)                                                                                                                                   |
| #1                                                   | naegleria[Title/Abstract] OR negleria[Title/Abstract]                                        | 1337                                                                                                                                                    |
| #14                                                  | brazil[Affiliation]                                                                          | 428131                                                                                                                                                  |
| #1 AND #14                                           | naegleria[Title/Abstract] OR negleria[Title/Abstract] AND brazil[Affiliation]                | 11                                                                                                                                                      |
| Key word #                                           | Scopus                                                                                       | Outcome (# of papers)                                                                                                                                   |
| #1                                                   | TITLE-ABS-KEY (naegleria OR negleria)                                                        | 1889                                                                                                                                                    |
| #14                                                  | AFFIL (brazil OR brasil)                                                                     | 1206542                                                                                                                                                 |
| #1 AND #14                                           | TITLE-ABS-KEY (naegleria OR negleria) AND AFFIL (brazil OR brasil)                           | 19                                                                                                                                                      |
| III. CRITERIA                                        |                                                                                              |                                                                                                                                                         |
| Inclusion criteria                                   |                                                                                              | Exclusion criteria                                                                                                                                      |
| 1. To study any pathogenic free-living amoeba genera |                                                                                              | 1. Dissertations and thesis                                                                                                                             |
| 2. Brazilian scope                                   |                                                                                              | 2. Paper published in non-indexed journals                                                                                                              |
| 3. To be authored by Brazilian researchers           |                                                                                              | 3. Not related to free-living amoeba                                                                                                                    |
| 4. Dated from 2020 or earlier                        |                                                                                              | 4. Letter, erratum, book chapter, protocol, conference abstract                                                                                         |
|                                                      |                                                                                              | 5. Affiliations exclusively with foreign institutions (no participation of Brazilian researchers)                                                       |
|                                                      |                                                                                              | 6. Experimental research did not accomplish in Brazil                                                                                                   |
|                                                      |                                                                                              | 7. Dated from 2021                                                                                                                                      |



Discriminative free-living amoeba (FLA) literature in Brazil tagged as AUXILIARY, clustered into five categories: clinical, environmental, review, wet, and dry bench. (A) Stacked bar graph comparing the number of papers published per year per category. (B) Territorial scope depicting the percentage of papers per Federative Unit.

## REFERENCES

- Marzzoco A, Colli W. Isolation of nuclei and characterization of nuclear DNA of *Acanthamoeba castellanii*. *Biochim Biophys Acta*. 1974;374(3):292–303.
- Marzzoco A, Colli W. Characterization of the genome of the small free-living amoeba *Acanthamoeba castellanii*. *Biochim Biophys Acta*. 1975;395(4):525–34.
- Campos R, Gomes MCO, Prigenzi LS, Stecca J. Meningoencefalite por ameba de vida livre apresentação do primeiro caso latino-americano. *Rev Inst Med Trop São Paulo*. 1977;19(5):349–51.
- Salles-Gomes JR CE, Barbosa ER., Nobrega JP, Scaff M., Spina-França A. Primary amebic meningoencephalomyelitis. Report of a case. *Arq Neuropsiquiatr*. 1978;36(2):139–42.
- de Moura H, Salazar HC, Fernandes O, Lisboa DC, de Carvalho FG. Free-living amoebae in human intestine: Evidence of parasitism. *Rev Inst Med Trop São Paulo*. 1985;27(3):150–6.
- Salazar HC, Moura H, Fernandez O, Peralta JM. Isolation of *Naegleria fowleri* from a lake in the city of Rio de Janeiro. *Trans R Soc Trop Med Hyg*. 1986;80(2):348–9.
- de Freitas D, Foronda AS, Sat EH, Júnior RB, Uras R. *Acanthamoeba culbertsoni*: sensibilidade aos diferentes métodos de desinfecção de lentes de contacto hidrofílicas. *Arq Bras Oftal*. 1987;1(3):2–6.
- Moura H, Wallace S, Visvesvara GS. *Acanthamoeba healyi* N. sp. and the isoenzyme and immunoblot profiles of *Acanthamoeba* spp., groups 1 and 3. *J Protozool*. 1992;39(5):573–83.
- Chimelli L, Hahn MD, Scaravilli F, Sara Wallace, Visvesvara GS. Granulomatous amoebic encephalitis due to leptomycid amoebae: report of the first Brazilian case. *Trans R Soc Trop Med Hyg*. 1992;86(6):635.
- Alvarenga LS, de Freitas D, Hofling-Lima AL. Ceratite por *Acanthamoeba*. *Arq Bras Oftal*. 2000;63(1):155–9.
- Alfieri SC, Correia CEB, Motegi SA, Pral EMF. Proteinase activities in total extracts and in medium conditioned by *Acanthamoeba polyphaga* trophozoites. *J Parasitol*. 2000;86(2):220–7.
- Alves JM, Gusmão CX, Teixeira MMG, de Freitas D, Foronda AS, Affonso HT. Random amplified polymorphic DNA profiles as a tool for the characterization of Brazilian keratitis isolates of the genus *Acanthamoeba*. *Braz J Med Biol Res*. 2000;33(1):19–26.
- Forseto AS, Nosé W. Diagnosis of *Acanthamoeba* keratitis with confocal microscopy. Vol. 62, *Rev Bras de Oftalmol*. 2003. p. 220–8.
- da Silva MA, da Rosa JA. Isolation of potentially pathogenic free-living amoebas in hospital dust. *Rev Saude Publica*. 2003;37(2):242–6.
- Nakano E, Oliveira M, Portellinha W, de Freitas D, Nakano K. Confocal microscopy in early diagnosis of *Acanthamoeba* keratitis. *J Refract Surg*. 2004;20(5):S737–741.
- Ruthes ACC, Wahab S, Wahab N, Moreira H, Moreira L. Conjunctivitis presumably due to *Acanthamoeba*. *Arq Bras Oftal*. 2004;67(6):897–900.
- da Rocha-Azevedo B, Menezes GC, Costa e Silva-Filho F. The interaction between *Acanthamoeba polyphaga* and human osteoblastic cells *in vitro*. *Microb Pathog*. 2006;40(1):8–14.
- da Rocha-Azevedo B, Jamerson M, Cabral GA, Costa e Silva-Filho F, Marciano-Cabral F. The interaction between the amoeba *Balamuthia mandrillaris* and extracellular matrix glycoproteins *in vitro*. *Parasitology*. 2007;134(Pt 1):51–8.
- da Rocha-Azevedo B, Costa e Silva-Filho F. Biological characterization of a clinical and an environmental isolate of *Acanthamoeba polyphaga*: analysis of relevant parameters to decode pathogenicity. *Arch Microbiol*. 2007;188(5):441–9.
- Barros JN, Mascaro VLD, Lowen M, Martins MC, Foronda A. Diagnosis of *Acanthamoeba* corneal infection by impression cytology: case report. *Arq Bras Oftal*. 2007;70(2):343–6.
- Silva-vergara ML, Colombo ER da C, Vissotto EDF, Silva ACAL, Chica JEL, Etchebehere RM, et al. Disseminated *Balamuthia mandrillaris* amoeba infection in an AIDS patient from Brazil. *Am J Trop Med Hyg*. 2007;77(6):1096–8.
- Carlesso AM, Simonetti AB, Artuso GL, Rott MB. Isolation and identification of potentially pathogenic free-living amoebae in samples from environments in a public hospital in the City of Porto Alegre, Rio Grande do Sul. *Rev Soc Bras Med Tro*. 2007;40(3):316–20.
- Pens CJ, Costa M da, Fadanelli C, Caumo K, Rott MB. *Acanthamoeba* spp. and bacterial contamination in contact lens storage cases and the relationship to user profiles. *Parasitol Res*. 2008;103(6):1241–5.
- Pens CJ, Rott MB. *Acanthamoeba* spp. cysts storage in filter paper. *Parasitol Res*. 2008;103(5):1229–30.
- Kashiwabuchi RT, de Freitas D, Alvarenga LS, Vieira L, Conatarini P, Sato E, et al. Corneal graft survival after therapeutic keratoplasty for *Acanthamoeba* keratitis. *Acta Ophthalmol*. 2008;86(6):666–9.
- Moraes J, Alfieri SC. Growth, encystment and survival of *Acanthamoeba castellanii* grazing on different bacteria. *FEMS Microbiol Ecol*. 2008;66(2):221–9.
- Ródio C, da Rocha Vianna D, Kowalski KP, Panatieri LF, von Poser G, Rott MB. *In vitro* evaluation of the amebicidal activity of *Pterocaulon polystachyum* (Asteraceae) against trophozoites of *Acanthamoeba castellanii*. *Parasitol Res*. 2008;104(1):191–4.
- da Rocha-Azevedo B, Jamerson M, Cabral GA, Costa e Silva-Filho F, Marciano-Cabral F. *Acanthamoeba* interaction with extracellular matrix glycoproteins: biological and biochemical characterization and role in cytotoxicity and invasiveness. *J Eukaryot Microbiol*. 2009;56(3):270–8.
- Santos LC, Oliveira MS, Lobo RD, Higashino HR, Costa SF, Van Der Heijden IM, et al. *Acanthamoeba* spp. in urine of critically ill patients. *Emerg Infect Dis*. 2009;15(7):1144–6.
- Sacramento RS, Martins RM, Miranda A, Dobroff ASS, Daffre S, Foronda AS, et al. Differential effects of alpha-helical and beta-hairpin antimicrobial peptides against *Acanthamoeba castellanii*. *Parasitology*. 2009;136(8):813–21.
- Ferreira GA, Magliano ACM, Pral EMF, Alfieri SC. Elastase secretion in *Acanthamoeba polyphaga*. *Acta Trop*. 2009;112(2):156–63.
- Magliano ACM, da Silva FM, Teixeira MMG, Alfieri SC. Genotyping, physiological features and proteolytic activities of a potentially pathogenic *Acanthamoeba* sp. isolated from tap water in Brazil. *Exp Parasitol*. 2009;123(3):231–5.
- Caumo K, Frasson AP, Pens CJ, Panatieri LF, Frazzon APG, Rott MB. Potentially pathogenic *Acanthamoeba* in swimming pools: a survey in the southern Brazilian city of Porto Alegre. *Ann Trop Med Parasitol*. 2009;103(6):477–85.
- Teixeira LH, Rocha S, Pinto RMF, Caseiro MM, Costa SOP. Prevalence of potentially pathogenic free-living amoebae from *Acanthamoeba* and *Naegleria* genera in non-hospital, public, internal environments from the city of Santos, Brazil. *Brazilian J Infect Dis*. 2009;13(6):395–7.
- Carvalho FRS, Foronda AS, Mannis MJ, Hofling-Lima AL, Belfort RJr, de Freitas D. Twenty years of *Acanthamoeba* keratitis. *Cornea*. 2009;28(5):516–9.
- Costa AO, Castro EA, Ferreira GA, Furst C, Crozeta MA, Thomas-Soccol V. Characterization of *Acanthamoeba* isolates from dust of a public hospital in Curitiba, Paraná, Brazil. *J Eukaryot Microbiol*. 2010;57(1):70–5.

37. Silva RA, Araújo SDA, Avellar IFDF, Pittella JEH, De Oliveira JT, Christo PP. Granulomatous amoebic meningoencephalitis in an immunocompetent patient. Arch Neurol. 2010;67(12):1516–20.
38. Cariello AJ, Souza GFPe, Foronda AS, Yu MCZ, Hofling-Lima AL, Oliveira MG. *In vitro* amoebicidal activity of S-nitroso-glutathione and S-nitroso-N-acetylcysteine against trophozoites of *Acanthamoeba castellanii*. J Antimicrob Chemother. 2010;65(3):588–91.
39. Carlesso AM, Artuso GL, Caumo K, Rott MB. Potentially pathogenic *Acanthamoeba* isolated from a hospital in Brazil. Curr Microbiol. 2010;60(3):185–90.
40. Rott M, Caumo K, Sauter I, Eckert J, da Rosa L, da Silva O. Susceptibility of *Aedes aegypti* (Diptera: Culicidae) to *Acanthamoeba polyphaga* (Sarcomastigophora: Acanthamoebidae). Parasitol Res. 2010;107(1):195–8.
41. Lemgruber L, Lupetti P, De Souza W, Vommaro RC, da Rocha-Azevedo B. The fine structure of the *Acanthamoeba polyphaga* cyst wall. FEMS Microbiol Lett. 2010;305(2):170–6.
42. Caumo K, Rott MB. *Acanthamoeba* T3, T4 and T5 in swimming-pool waters from Southern Brazil. Acta Trop. 2011;117(3):233–5.
43. Sauter IP, Santos JC, Apel MA, Cibulski SP, Roehe PM, von Poser GL, et al. Amebicidal activity and chemical composition of *Pterocaulon polystachyum* (Asteraceae) essential oil. Parasitol Res. 2011;109(5):1367–71.
44. Kashiwabuchi RT, Carvalho FRS, Khan YA, Freitas D De, Foronda AS, Hirai FE, et al. Assessing efficacy of combined riboflavin and UV-A Light (365 nm) treatment of *Acanthamoeba* trophozoites. Invest Ophthalmol Vis Sci. 2011;52(13):9333–8.
45. Benitez LB, Caumo K, Brandelli A, Rott MB. Bacteriocin-like substance from *Bacillus amyloliquefaciens* shows remarkable inhibition of *Acanthamoeba polyphaga*. Parasitol Res. 2011;108(3):687–91.
46. Winck MAT, Caumo K, Rott MB. Prevalence of *Acanthamoeba* from tap water in Rio Grande do Sul, Brazil. Curr Microbiol. 2011;63(5):464–9.
47. Carvalho FRDS, Carrijo-Carvalho LC, Chudzinski-Tavassi AM, Foronda AS, de Freitas D. Serine-like proteolytic enzymes correlated with differential pathogenicity in patients with acute *Acanthamoeba* keratitis. Clin Microbiol Infect. 2010;17(4):603–9.
48. Zanella J, da Costa SOP, Zacaria J, Echeverrigaray S. A rapid and reliable method for the clonal isolation of *Acanthamoeba* from environmental samples. Brazilian Arch Biol Technol. 2012;55(1):1–6.
49. Vunda SLL, Sauter IP, Cibulski SP, Roehe PM, Bordignon SAL, Rott MB, et al. Chemical composition and amoebicidal activity of *Croton pallidulus*, *Croton ericoides*, and *Croton isabelli* (Euphorbiaceae) essential oils. Parasitol Res. 2012;111(3):961–6.
50. Sauter IP, Rossa GE, Lucas AM, Cibulski SP, Roehe PM, Silva LAA, et al. Chemical composition and amoebicidal activity of *Piper hispidinervum* (Piperaceae) essential oil. Ind Crop Prod. 2012;40:292–5.
51. Pimentel LA, Dantas AFM, Uzal F, Riet-Correa F. Meningoencephalitis caused by *Naegleria fowleri* in cattle of northeast Brazil. Res Vet Sci. 2012;93(2):811–2.
52. Alves DSMM, Moraes AS, Nitz N, Oliveira MGC, Hecht MM, Gurgel-Gonçalves R, et al. Occurrence and characterization of *Acanthamoeba* similar to genotypes T4, T5, and T2/T6 isolated from environmental sources in Brasília, Federal District, Brazil. Exp Parasitol. 2012;131(2):239–44.
53. Otta DA, Rott MB, Carlesso AM, Santos O. Prevalence of *Acanthamoeba* spp. (Sarcomastigophora: Acanthamoebidae) in wild populations of *Aedes aegypti* (Diptera: Culicidae). Parasitol Res. 2012;111(1):2017–22.
54. Magliano ACM, Teixeira MMG, Alfieri SC. Revisiting the *Acanthamoeba* species that form star-shaped cysts (genotypes T7, T8, T9, and T17): characterization of seven new Brazilian environmental isolates and phylogenetic inferences. Parasitology. 2011;139(1):45–52.
55. Borase HP, Patil CD, Sauter IP, Rott MB, Patil S V. Amoebicidal activity of phytosynthesized silver nanoparticles and their *in vitro* cytotoxicity to human cells. FEMS Microbiol Lett. 2013;345(2):127–31.
56. Mafra CSP, Carrijo-Carvalho LC, Chudzinski-tavassi AM, Taguchi FMC, Foronda AS, Carvalho FR de S, et al. Antimicrobial action of biguanides on the viability of *Acanthamoeba* cysts and assessment of cell toxicity. Invest Ophthalmol Vis Sci. 2013;54(9):6363–72.
57. Castro LC, Sauter IP, Ethur EM, Kauffmann C, Dall'agnol R, Souza J, et al. *In vitro* effect of *Acanthospermum australe* (Asteraceae) extracts on *Acanthamoeba polyphaga* trophozoites. Rev Bras Pl Med. 2013;15(4):589–94.
58. Veríssimo CDM, Maschio VJ, Correa APF, Brandelli A, Rott MB. Infection in a rat model reactivates attenuated virulence after long-term axenic culture of *Acanthamoeba* spp. Mem Inst Oswaldo Cruz. 2013;108(7):832–5.
59. Landeli MF, Salton J, Caumo K, Broetto L, Rott MB. Isolation and genotyping of free-living environmental isolates of *Acanthamoeba* spp. from bromeliads in Southern Brazil. Exp Parasitol. 2013;134(3):290–4.
60. Duarte JL, Furst C, Klisiowicz DR, Klassen G, Costa AO. Morphological, genotypic, and physiological characterization of *Acanthamoeba* isolates from keratitis patients and the domestic environment in Vitoria, Espírito Santo, Brazil. Exp Parasitol. 2013;135(1):9–14.
61. Becker-Finco A, Costa AO, Silva SK, Ramada JS, Furst C, Stingher AE, et al. Physiological, morphological, and immunological parameters used for the characterization of clinical and environmental isolates of *Acanthamoeba*. Parasitology. 2013;140(3):396–405.
62. da Silva MTA, Caldas VEA, Costa FC, Silvestre DAMM, Thiemann OH. Selenocysteine biosynthesis and insertion machinery in *Naegleria gruberi*. Mol Biochem Parasitol. 2013;188(2):87–90.
63. de Aguiar APC, de Oliveira Silveira C, Todero Winck MA, Rott MB. Susceptibility of *Acanthamoeba* to multipurpose lens-cleaning solutions. Acta Parasitol. 2013;58(3):304–8.
64. Carlesso AM, Mentz MB, Machado MLS, Carvalho A, Nunes TET, Maschio VJ, et al. Characterization of isolates of *Acanthamoeba* from the nasal mucosa and cutaneous lesions of dogs. Curr Microbiol. 2014;68(6):702–7.
65. Caumo KS, Monteiro KM, Ott TR, Maschio VJ, Wagner G, Ferreira HB, et al. Proteomic profiling of the infective trophozoite stage of *Acanthamoeba polyphaga*. Acta Trop. 2014;140:166–72.
66. Alves DSMM, Gurgel-Gonçalves R, Albuquerque P, Cuba-Cuba CA, Muniz-Junqueira MI, Kuckelhaus SAS. A method for microbial decontamination of *Acanthamoeba* cultures using the peritoneal cavity of mice. Asian Pac J Trop Biomed. 2015;5(10):796–800.
67. Corsaro D, Walochnik J, Köhler M, Rott MB. *Acanthamoeba* misidentification and multiple labels: redefining genotypes T16, T19, and T20 and proposal for *Acanthamoeba micheli* sp. nov. (genotype T19). Parasitol Res. 2015; 114: 2481–2490.
68. Boratto P, Albarnaz JD, Almeida GMDF, Botelho L, Fontes ACL, Costa AO, et al. *Acanthamoeba polyphaga* mimivirus prevents amoebal encystment-mediating serine proteinase expression and circumvents cell encystment. J Virol. 2015;89(5):2962–5.

69. Maschio VJ, Chies F, Carlesso AM, Carvalho A, Rosa SP, Van Der Sand ST, et al. *Acanthamoeba* T4, T5 and T11 isolated from mineral water bottles in Southern Brazil. *Curr Microbiol*. 2014;70(1):6–9.
70. Sant'ana VP, Carrijo-Carvalho LC, Foronda AS, Chudzinski-Tavassi AM, de Freitas D, Ramos F, et al. Cytotoxic activity and degradation patterns of structural proteins by corneal isolates of *Acanthamoeba* spp. *Graefes Arch Clin Exp Ophthalmol*. 2015;253(1):65–75.
71. Maschio VJ, Corção G, Bucker F, Caumo K, Rott MB. Identification of *Paenibacillus* as a symbiont in *Acanthamoeba*. *Curr Microbiol*. 2015;71(3):415–20.
72. Maschio VJ, Corção G, Rott MB. Identification of *Pseudomonas* spp. as amoeba-resistant microorganisms in isolates of *Acanthamoeba*. *Rev Inst Med Trop Sao Palo*. 2015;57(1):81–3.
73. Silva LCF, Almeida GMF, Assis FL, Albarnaz JD, Boratto PVM, Dornas FP, et al. Modulation of the expression of mimivirus-encoded translation-related genes in response to nutrient availability during *Acanthamoeba castellanii* infection. *Front Microbiol*. 2015;6(6):539.
74. Frade MTS, Melo LF, Pessoa CRM, Araújo JL, Figuera RA, Souza AP, et al. Systemic acanthamoebiasis associated with canine distemper in dogs in the semiarid region of Paraíba, Brazil. *Pesq Vet Bras*. 2015;35(2):160–4.
75. Nunes TET, Brazil NT, Fuentefria AM, Rott MB. *Acanthamoeba* and *Fusarium* interactions: a possible problem in keratitis. *Acta Trop*. 2016;157:102–7.
76. Silva LKDS, Boratto PVM, La Scola B, Bonjardim CA, Abrahão JS. *Acanthamoeba* and mimivirus interactions: the role of amoebal encystment and the expansion of the 'Cheshire Cat' theory. *Curr Opin Microbiol*. 2016;31:9–15.
77. Guimaraes AJ, Gomes KX, Cortines JR, Peralta JM, Peralta RHS. *Acanthamoeba* spp. as a universal host for pathogenic microorganisms: one bridge from environment to host virulence. *Microbiol Res*. 2016;193:30–8.
78. Santos IGA, Scher R, Rott MB, Menezes LR, Costa EV, Cavalcanti SCH, et al. Amoebicidal activity of the essential oils of *Lippia* spp. (Verbenaceae) against *Acanthamoeba* polyphaga trophozoites. *Parasitol Res*. 2016;115(2):535–40.
79. Penteado RF, Martini VP, Iulek J. Crystallization and crystallographic analyses of triosephosphate isomerase from *Naegleria gruberi*. *Rev Virtual Quim*. 2016;8(6):1835–41.
80. Staggemeier R, Arantes T, Caumo KS. Detection and quantification of human adenovirus genomes in *Acanthamoeba* isolated from swimming pools. *An Acad Bras Cienc*. 2016;88:635–41.
81. Sánchez AGC, Virgínio VG, Maschio VJ, Ferreira HB, Rott MB. Evaluation of the immunodiagnostic potential of a recombinant surface protein domain from *Acanthamoeba castellanii*. *Parasitology*. 2021;143(12):1656–64.
82. Alves DSM, Moraes AS, Alves LM, Gurgel-Gonçalves R, Lino Júnior RS, Cuba-Cuba CA, et al. Experimental infection of T4 *Acanthamoeba* genotype determines the pathogenic potential. *Parasitol Res*. 2016;115(9):3435–40.
83. Machado ATP, Silva M, Iulek J. Expression, purification, enzymatic characterization and crystallization of glyceraldehyde-3-phosphate dehydrogenase from *Naegleria gruberi*, the first one from phylum Percolozoa. *Protein Expr Purif*. 2016;127:125–30.
84. Fabres LF, Santos SPR, Benitez LB, Rott MB. Isolation and identification of *Acanthamoeba* spp. from thermal swimming pools and spas in Southern Brazil. *Acta Parasitol*. 2016;61(2):221–7.
85. Corrêa TQ, Geralde MC, Carvalho MT, Bagnato VS, Kurachi C, Souza CWO. Photodynamic inactivation of *Acanthamoeba polyphaga* with curcuminoids: an *in vitro* study. *Opt Methods Tumor Treat Detect Mech Tech Photodyn Ther XXV*. 2016;9694:1–7.
86. Fuciková K, Lahr DJG. Uncovering cryptic diversity in two amoebozoan species using complete mitochondrial genome sequences. *J Eukaryot Microbiol*. 2016;63(1):112–22.
87. Alves DSM, Alves LM, Costa TL, Castro AM, Vinaud MC. Anaerobic metabolism in T4 *Acanthamoeba* genotype. *Curr Microbiol*. 2017;74(6):685–90.
88. Souza TK, Soares SS, Benitez LB, Rott MB. Interaction between methicillin-resistant *Staphylococcus aureus* (MRSA) and *Acanthamoeba polyphaga*. *Curr Microbiol*. 2017;74(5):541–9.
89. Ribeiro NS, Santos FM, Garcia AWA, Ferrareze PAG, Fabres LF, Schrank A, et al. Modulation of zinc homeostasis in *Acanthamoeba castellanii* as a possible antifungal strategy against *Cryptococcus gattii*. *Front Microbiol*. 2017;8(August):1–11.
90. Costa AO, Furst C, Rocha LO, Cirelli C, Cardoso CN, Neiva FS, et al. Molecular diagnosis of *Acanthamoeba* keratitis: evaluation in rat model and application in suspected human cases. *Parasitol Res*. 2017;116(4):1339–44.
91. Panatieri LF, Brazil NT, Faber K, Medeiros-Neves B, von Poser GL, Rott MB, et al. Nanoemulsions containing a coumarin-rich extract from *Pterocaulon balansae* (Asteraceae) for the treatment of ocular *Acanthamoeba* keratitis. *AAPS PharmSciTech*. 2017;18(3):721–8.
92. Soares SS, Souza TK, Berte FK, Cantarelli VV, Rott MB. Occurrence of infected free-living amoebae in cooling towers of southern Brazil. *Curr Microbiol*. 2017;74(12):1461–8.
93. Sant'ana VP, Foronda AS, Freitas D de, Carrijo-Carvalho LC, Carvalho FR de S. Sensitivity of enzymatic toxins from corneal isolate of *Acanthamoeba* protozoan to physicochemical parameters. *Curr Microbiol*. 2017;74(11):1316–23.
94. Faber K, Zorzi GK, Brazil NT, Rott MB, Teixeira HF. siRNA-loaded liposomes: inhibition of encystment of *Acanthamoeba* and toxicity on the eye surface. *Chem Biol Drug Des*. 2017;90(3):406–16.
95. Carrijo-Carvalho LC, Sant'ana VP, Foronda AS, de Freitas D, Ramos F, Carvalho DS. Therapeutic agents and biocides for ocular infections by free-living amoebae of *Acanthamoeba* genus. *Surv Ophthalmol*. 2016;62(2):203–18.
96. Sticca MP, Carrijo-carvalho LC, Silva IMB, Vieira LA, Souza LB, Junior RB, et al. *Acanthamoeba* keratitis in patients wearing scleral contact lenses. *Contact Lens Anterior Eye*. 2018;41(3):307–10.
97. Santos DL, Kwitko S, Marinho DR, Araújo BS, Locatelli CI, Rott MB. *Acanthamoeba* keratitis in Porto Alegre (southern Brazil): 28 cases and risk factors. *Parasitol Res*. 2018;117(3):747–50.
98. Possamai CO, Loss AC, Costa AO, Falqueto A, Furst C. *Acanthamoeba* of three morphological groups and distinct genotypes exhibit variable and weakly inter-related physiological properties. *Parasitol Res*. 2018;117(5):1389–400.
99. Maschio JV, Virgínio VG, Ferreira HB, Rott MB. Comparative proteomic analysis of soluble and surface-enriched proteins from *Acanthamoeba castellanii* trophozoites. *Mol Biochem Parasitol*. 2018;225:47–53.
100. Buchele MLC, Wopereis DB, Casara F, Macedo JP, Rott MB, Monteiro FBF, et al. Contact lens-related polymicrobial keratitis: *Acanthamoeba* spp. genotype T4 and *Candida albicans*. *Parasitol Res*. 2018;117(11):3431–6.

101. Gonçalves DS, Ferreira MS, Liedke SC, Gomes KX, Oliveira GA, Leão PEL, et al. Extracellular vesicles and vesicle-free secretome of the protozoa *Acanthamoeba castellanii* under homeostasis and nutritional stress and their damaging potential to host cells. *Virulence*. 2018;9(1):2150–5608.
102. Machado ATP, Silva M, Iulek J. Structural studies of glyceraldehyde-3-phosphate dehydrogenase from *Naegleria gruberi*, the first one from phylum. *Biochim Biophys Acta Proteins Proteom*. 2018;1866:581–8.
103. Alves DSMM, Gonçalves GS, Moraes AS, Alves LM, Carmo Neto JR, Hecht MM, et al. The first *Acanthamoeba* keratitis case in the Midwest region of Brazil: diagnosis, genotyping of the parasite and disease outcome. *Rev Soc Bras Med Trop*. 2018;51(5):716–9.
104. Bellini NK, Santos TM, da Silva MTA, Thiemann OH. The therapeutic strategies against *Naegleria fowleri*. *Exp Parasitol*. 2018 Apr;187:1–11.
105. Fabres LF, Maschio VJ, Santos DL, Kwitko S, Marinho DR, Araújo BS, et al. Virulent T4 *Acanthamoeba* causing keratitis in a patient after swimming while wearing contact lenses in Southern Brazil. *Acta Parasitol*. 2018;63(2):428–32.
106. Zorzi GK, Schuh RS, Maschio VJ, Brazil NT, Rott MB, Teixeira HF. Box Behnken design of siRNA-loaded liposomes for the treatment of a murine model of ocular keratitis caused by *Acanthamoeba*. *Colloids Surfaces B Biointerfaces*. 2019;173:725–32.
107. Gonçalves DS, Ferreira MS, Guimarães AJ. Extracellular vesicles from the protozoa *Acanthamoeba castellanii*: Their role in pathogenesis, environmental adaptation and potential applications. *Bioengineering*. 2019;6(1):12.
108. Serrano-Solis V, Toscano Soares PE, de Farias ST. Genomic signatures among *Acanthamoeba polyphaga* entoorganisms unveil evidence of coevolution. *J Mol Evol*. 2019;87(1):7–15.
109. Medina G, Leyán P, Silva CV, Flores-Martin S, Manosalva C, Fernández H. Intra-amoebic localization of *Arcobacter butzleri* as an endocytobiont of *Acanthamoeba castellanii*. *Arch Microbiol*. 2019;201(10):1447–52.
110. Henker LC, Cruz RAS, Silva FS, Driemeier D, Sonne L, Uzal FA, et al. Meningoencephalitis due to *Naegleria fowleri* in cattle in southern Brazil. *Rev Bras Parasitol Vet*. 2019;28(3):514–7.
111. Corsaro D, Köhler M, Venditti D, Rott MB, Walochnik J. Recovery of an *Acanthamoeba* strain with two group I introns in the nuclear 18S rRNA gene. *Eur J Protistol*. 2019;68:88–98.
112. Gonçalves DS, Ferreira MS, Gomes KX, La Noval CR, Liedke SC, Costa GCV, et al. Unravelling the interactions of the environmental host *Acanthamoeba castellanii* with fungi through the recognition by mannose - binding proteins. *Cell Microbiol*. 2019;21(10):e13066.
113. Faria LV, Carmo PHF, Costa MC, Peres NTA, Chagas IAR, Furst C, et al. *Acanthamoeba castellanii* as an alternative interaction model for the dermatophyte *Trichophyton rubrum*. *Mycoses*. 2020;63(12):1331–40.
114. Carvalho-Kelly LF, Pralon CF, Rocco-Machado N, Nascimento MT, Carvalho-de-Araújo AD, Meyer-Fernandes JR. *Acanthamoeba castellanii* phosphate transporter (AcPHS) is important to maintain inorganic phosphate influx and is related to trophozoite metabolic processes. *J Bioenerg Biomembr*. 2020;52(2):93–102.
115. Weber-Lima MM, Prado-Costa B, Becker-Finco A, Costa AO, Billilad P, Furst C, et al. *Acanthamoeba* spp. monoclonal antibody against a CPA2 transporter : a promising molecular tool for acanthamoebiasis diagnosis and encystment study. *Parasitology*. 2020;147(14):1678–88.
116. Cirelli C, Mesquita EIS, Chagas IAR, Furst C, Possamai CO, Abrahão JS, et al. Extracellular protease profile of *Acanthamoeba* after prolonged axenic culture and after interaction with MDCK cells. *Parasitol Res*. 2020;119(2):659–66.
117. Wopereis DB, Bazzo ML, De Macedo JP, Casara F, Golfeto L, Venancio E, et al. Free-living amoebae and its relationship with air quality in hospital environments: isolation and characterization of *Acanthamoeba* spp. from air-conditioning system. *Parasitology*. 2020;147(7):789–90.
118. Fonseca JDG, Gómez-Hernández C, Barbosa CG, Rezende-Oliveira K. Identification of T3 and T4 genotypes of *Acanthamoeba* sp. in dust samples isolated from air conditioning equipment of public hospital of Ituiutaba-MG. *Curr Microbiol*. 2020;77(5):890–5.
119. Fabres LF, da Costa Gonçalves F, Duarte EOS, Berté FK, da Conceição DKS Iva, Ferreira LA, et al. *In vitro* amoebicidal activity of imidazolium salts against trophozoites. *Acta Parasitol*. 2020;65(2):317–26.
120. Bellini NK, da Fonseca ALM, Reyes-Batlle M, Lorenzo-Morales J, Rocha O, Thiemann OH. Isolation of *Naegleria* spp. from a Brazilian water source. *Pathogens*. 2020;9(2):90.
121. Marujo FI, Hirai FE, Yu MCZ, Hofling-Lima AL, de Freitas D, Sato EH. Distribution of infectious keratitis in a tertiary hospital in Brazil. *Arq Bras Oftalmol*. 2013;76(6):370–3.
122. Moriyama AS, Hofling-Lima AL. Contact lens-associated microbial keratitis. *Arq Bras Oftalmol*. 2008;71(7):32–6.
123. Santos A, Silva LD, Sousa LB, de Freitas D, Oliveira LA. Results with the Boston Type I keratoprosthesis after *Acanthamoeba* keratitis. *Am J Ophthalmol Case Rep*. 2017;6:71–3.
124. Farias R, Pinho L, Santos R. Epidemiological profile of infectious keratitis. *Rev Bras Oftalmol*. 2017;76(3):116–20.
125. Müller RT, Abedi F, Cruzat A, Witkin D, Baniasadi N, Cavalcanti BM, et al. Degeneration and regeneration of subbasal corneal nerves after infectious keratitis: a longitudinal in vivo confocal microscopy study. *Ophthalmology*. 2015;122(11):2200–9.
126. Frade MTS, Ferreira JS, Nascimento MJR, Aquino VVF, Macêdo IL, Carneiro RS, et al. Central nervous system disorders diagnosed in dogs. *Pesq Vet Bras*. 2018;38(5):935–48.
127. Boratto PVM, Oliveira GP, Machado TB, Andrade ACSP, Baudoin J-P, Klose T, et al. Yaravirus: A novel 80-nm virus infecting *Acanthamoeba castellanii*. *Proc Natl Acad Sci*. 2020;117(28):202001637.
128. Perim LV, Custódio NCC, Lima VCV, Igreja JASL, Alves DSMM, Storchilo HR, et al. Occurrence of parasites in salads in restaurants in Aparecida de Goiânia, Goiás, Brazil. *J Trop Pathol Vol*. 2020;49(3):207–14.
129. Leal DAG, Souza DSM, Caumo KS, Fongaro G, Panatieri LF, Durigan M, et al. Genotypic characterization and assessment of infectivity of human waterborne pathogens recovered from oysters and estuarine waters in Brazil. *Water Res*. 2018;137:273–80.
130. Silva LKDS, Rodrigues RAL, Andrade ACDSP, Hikida H, Andreani J, Levasseur A, et al. Isolation and genomic characterization of a new mimivirus of lineage B from a Brazilian river. *Arch Virol*. 2020;165(4):853–63.
131. Albuquerque P, Nicola AM, Magnabosco DAG, da Silveira Derengowski L, Crisóstomo LS, Xavier LCG, et al. A hidden battle in the dirt: soil amoebae interactions with *Paracoccidioides* spp. *PLoS Negl Trop Dis*. 2019;13(10):e0007742.
132. Rodrigues RAL, Abrahão JS, Drumond BP, Kroon EG. Giants among larges : how gigantism impacts giant virus entry into amoebae. *Curr Opin Microbiol*. 2016;31(31):88–93.

133. Gasparetto EL, Cabral RF, Hygino LC, Domingues RC. Diffusion imaging in brain infections. *Neuroimaging Clin NA*. 2011;21(1):89–113.
134. Abrahão JS, Dornas FP, Silva LCF, Almeida GM, Boratto PVM, Colson P, et al. *Acanthamoeba polyphaga* mimivirus and other giant viruses: an open field to outstanding discoveries. *Virol J*. 2014;11:1–12.
135. Hofstatter PG, Ribeiro GM, Porfirio-Sousa AL, Lahr DJG. The sexual ancestor of all eukaryotes: a defense of the “meiosis tool-kit”: a rigorous survey supports the obligate link between meiosis machinery and sexual recombination. *BioEssays*. 2020;42(9):1–10.
136. Rolland C, Andreani J, Louazani AC, Aherfi S, Francis R, Rodrigues R, et al. Discovery and further studies on Giant viruses at the IHU Mediterranée infection that modified the perception of the virosphere. *Viruses*. 2019;11(1), 312.
137. Raoult D, Abrahão J. Editorial overview : the megaviromes. *Curr Opin Microbiol*. 2016;31:31–3.
138. Yin R, Dai T, Avci P, Jorge AES, Melo WCMA, Vecchio D, et al. Light based anti-infectives: ultraviolet C irradiation, photodynamic therapy, blue light, and beyond. *Curr Opin Pharmacol*. 2013;13(5):731–62.
139. Kollipara R, Peranteau AJ, Nawas ZY, Tong Y, Woc-colburn L, Yan AC, et al. Emerging infectious diseases with cutaneous manifestations: fungal, helminthic, protozoan and ectoparasitic infections. *J Am Acad Dermatol*. 2016;75(1):19–30.
140. Chimelli L. Co-infection of HIV and tropical infectious agents that affect the nervous system. *Rev Neurol (Paris)*. 2012;168(3):270–82.
141. Sullivan KE, Bassiri H, Bousfiha AA, Costa-Carvalho BT, Freeman AF, Hagin D, et al. Emerging infections and pertinent infections related to travel for patients with primary immunodeficiencies. *J Clin Immunol*. 2017;37:650–92.
142. Chimelli L. A morphological approach to the diagnosis of protozoal infections of the central nervous system. *Patholog Res Int*. 2011;2011:29085.
143. Oliveira G, La Scola B, Abrahão J. Giant virus vs amoeba: fight for supremacy. *Virol J*. 2019;16(1):1–12.
144. Abrahão J, La Scola B. Editorial: large and giant DNA viruses. *Front Microbiol*. 2019;10(July):1–2.
145. Mougari S, Bekliz M, Abrahao J, Pinto F Di, Levasseur A, La Scola B. Guarani Virophage , a new sputnik-like isolate from a Brazilian lake. *Front Microbiol*. 2019;10(3).
146. Silva LCF, Araújo R, Rodrigues L, Oliveira GP, Dornas FP, La Scola B, et al. Microscopic analysis of the Tupanvirus cycle in *Vermamoeba vermiformis*. 2019;10(April):1–9.
147. Abrahão J, Silva L, Oliveira D, Almeida G. Lack of evidence of mimivirus replication in human PBMCs. *Microbes Infect*. 2018; May;20(5):281–283.
148. Andrade ACDSP, Rodrigues RAL, Oliveira GP, Andrade KR, Bonjardim CA, La Scola B, et al. Filling gaps about mimivirus entry, uncoating and morphogenesis. *J Virol*. 2017;22:e01355–17.
149. Hori JI, Pereira MSF, Roy CR, Nagai H, Zamboni DS, Celular B, et al. Identification and functional characterization of K(+) transporters encoded by *Legionella pneumophila* kup genes. *Cell Microbiol*. 2013;15(12):2006–19.
150. Medina G, Neves P, Flores-Martin S, Manosalva C, Andaur M, Otth C, et al. Transcriptional analysis of flagellar and putative virulence genes of *Arcobacter butzleri* as an endocytobiont of *Acanthamoeba castellanii*. *Arch Microbiol*. 2019;201(8):1075–83.
151. Dornas FP, Silva LCF, Almeida GM De, Campos RK, Boratto PVM, Franco-Luiz APM, et al. *Acanthamoeba polyphaga* mimivirus stability in environmental and clinical substrates: implications for virus detection and isolation. *PLoS One*. 2014;9(2):16–8.
152. Silva A, Nobre HJr, Sampaio L, Nascimento B, Silva C, de Andrade Neto JB, et al. Antifungal and antiprotozoal green amino acid-based rhamnolipids: mode of action , antibiofilm efficiency and selective activity against resistant *Candida* spp . strains and *Acanthamoeba castellanii*. *Colloids Surfaces B Biointerfaces*. 2020;193:111148.
153. Rodrigues RAL, Louazani AC, Picorelli A, Oliveira GP, Lobo FP, Colson P, et al. Analysis of a Marseillevirus transcriptome reveals temporal gene expression profile and host transcriptional shift. *Front Microbiol*. 2020;11(April):1–17.
154. Tavares PL, Ribeiro AC, Berte FK, Hellwig AH S, Pagani DM, Souza CCT, et al. The interaction between *Sporothrix schenckii* sensu stricto and *Sporothrix brasiliensis* with *Acanthamoeba castellanii*. *Mycoses*. 2020;63(3):302–7.
155. Pereira Andrade ACS, Boratto PVM, Rodrigues RAL, Bastos TM, Azevedo BL, Dornas FP, et al. New isolates of Pandoravirus: contribution to the study of replication cycle steps. *J Virol*. 2019;93(5):1–12.
156. Mougari S, Abrahao J, Oliveira GP, Bou Khalil JY, La Scola B. Role of the R349 gene and its repeats in the MIMIVIRE defense system. *Front Microbiol*. 2019;10(May):1–10.
157. Borges IA, Assis FL, Silva LKDS, Abrahão J. Rio Negro virophage: sequencing of the near complete genome and transmission electron. *Brazilian J Microbiol*. 2018;49(Suppl 1):260–1.
158. Boratto PVM, Dornas FP, Silva LCF, Rodrigues RAL, Oliveira GP, Cortines JR, et al. Analyses of the Kroon virus major capsid gene and its transcript highlight a distinct pattern of gene evolution and splicing among Mimiviruses. *J Virol*. 2018;92(2):1–11.
159. Oliveira GP, Lima MT, Arantes TS, Assis FL, Rodrigues RAL, da Fonseca FG, et al. The investigation of promoter sequences of Marseilleviruses highlights a remarkable abundance of the AAATATTT motif in intergenic regions. *J Virol*. 2017;91(21):1–10.
160. Rizzo J, Albuquerque PC, Wolf JM, Nascimento R, Pereira MD, Nosanchuk JD, et al. Analysis of multiple components involved in the interaction between *Cryptococcus neoformans* and *Acanthamoeba castellanii*. *Fungal Biol*. 2017;121(6–7):602–14.
161. Schrad JR, Young EJ, Abrahão JS, Cortines JR, Parent KN. Microscopic characterization of the Brazilian giant samba virus. *Viruses*. 2017;9(30):1–16.
162. Arantes TS, Rodrigues RAL, dos Santos Silva LK, Oliveira GP, de Souza HL, Khalil JYB, et al. The large Marseillevirus explores different entry pathways by forming giant infectious vesicles. *J Virol*. 2016;90(11):5246–55.
163. Decarli MC, Carvalho MT, Corrêa TQ, Bagnato VS, Souza CWO de. Different photoresponses of microorganisms: from bioinhibition to biostimulation. *Curr Microbiol*. 2016;72(4):473–81.
164. Dornas FP, Assis FL, Aherfi S, Arantes T, Abrahão JS, Colson P, et al. A Brazilian Marseillevirus is the founding member of a lineage in family Marseilleviridae. *Viruses*. 2016;8(76):1–16.
165. Rodrigues RAL, Silva LKS, Dornas FP, Oliveira DB, Magalhães TFF, Santos DA, et al. *Mimivirus fibrils* are important for viral attachment to the microbial world by a diverse glycoside interaction repertoire. *J Virol*. 2015;89(23):11812–9.
166. Boratto PVM, Arantes TS, Silva LCF, Assis FL, Kroon EG, La Scola B, et al. Niemeyer virus: a new mimivirus group A isolate harboring a set of duplicated aminoacyl-tRNA synthetase genes. *Front Microbiol*. 2015;6(NOV):1–11.

167. Campos RK, Boratto P V, Assis FL, Aguiar ERGR, Silva LCF, Albarnaz JD, et al. Samba virus: A novel mimivirus from a giant rain forest, the Brazilian Amazon. *Viol J*. 2014;11(1):1–11.
168. Boratto PVM, Dornas FP, Andrade KR, Rodrigues R, Peixoto F, Silva LCF, et al. Amoebas as mimivirus bunkers: increased resistance to UV light, heat and chemical biocides when viruses are carried by amoeba hosts. *Arch Virol*. 2014;159(5):1039–43.
169. Silva LCF, Almeida GMF, Oliveira DB, Dornas FP, Campos RK, La Scola B, et al. A resourceful giant: APMV is able to interfere with the human type I interferon system. *Microbes Infect*. 2014;16(3):187–95.
170. Lopes FC, Tichota DM, Sauter IP, Meira SMM, Segalin J, Rott MB, et al. Active metabolites produced by *Penicillium chrysogenum* IFL1 growing on agro-industrial residues. *Ann Microbiol*. 2013;63:771–8.
171. Derengowski LS, Paes HC, Albuquerque P, Tavares AHFP, Fernandes L, Silva-Pereira I, et al. The transcriptional response of *Cryptococcus neoformans* to ingestion by *Acanthamoeba castellanii* and macrophages provides insights into. *Eukaryot Cell*. 2013;12(5):761–74.
172. Campos RK, Andrade KR, Ferreira PCP, Bonjardim CA, Scola B La, Kroon EG, et al. Virucidal activity of chemical biocides against mimivirus , a putative pneumonia agent. *J Clin Virol*. 2012;55(4):323–8.
173. Souza F, Rodrigues R, Reis E, Lima M, La Scola B, Abrahão J. In-depth analysis of the replication cycle of *Orpheovirus*. *Viol J*. 2019;16(1):1–11.
174. Borges I, Rodrigues RAL, Dornas FP, Almeida G, Aquino I, Bonjardim CA, et al. Trapping the enemy: *Vermamoeba vermiformis* circumvents *Faustovirus mariensis* dissemination by enclosing viral progeny inside cysts . *J Virol*. 2019;93(14):1–19.
175. Augusto G, Souza P, Queiroz VF, Lima MT, Vinicius E, Reis DS, et al. Virus goes viral: an educational kit for virology classes. *Viol J*. 2020;17(1):1–8.
176. Abrahão J, Silva L, Silva LS, Khalil JYB, Rodrigues R, Arantes T, et al. Tailed giant Tupanvirus possesses the most complete translational apparatus of the known virosphere. *Nat Commun* . 2018;9(1).
177. Dornas FP, Khalil JYB, Pagnier I, Raoult D, Abrahão J, La Scola B. Isolation of new Brazilian giant viruses from environmental samples using a panel of protozoa. *Front Microbiol*. 2015;6(October):1–9.
178. Abrahão JS, Boratto P, Dornas FP, Silva LC, Campos RK, Almeida GMF, et al. Growing a giant: evaluation of the virological parameters for mimivirus production. *J Virol Methods*. 2014;207:6–11.