

Comparative morphometric evaluation of hepatic hemosiderosis in wild Magellanic penguins (*Spheniscus magellanicus*) infected with different *Plasmodium* spp. subgenera

Avaliação morfométrica comparativa da hemossiderose hepática em pinguins-de-Magalhães (*Spheniscus magellanicus*) de vida livre infectados por diferentes subgêneros de *Plasmodium* spp.

Ana Carolina Ewbank^{1*} ; Ricardo de Francisco Strefezzi²; Carlos Sacristán¹; Cristiane Kiyomi Miyaji Kolesnikovas³; Aryse Martins⁴; Luis Felipe Silva Pereira Mayorga⁵; Ralph Eric Thijl Vanstreels¹; José Luiz Catão-Dias¹

¹ Laboratório de Patologia Comparada de Animais Selvagens, Departamento de Patologia, Universidade de São Paulo – USP, São Paulo, SP, Brasil

² Laboratório de Oncologia Comparada e Translacional, Faculdade de Zootecnia e Engenharia de Alimentos, Universidade de São Paulo – USP, Pirassununga, SP, Brasil

³ Associação R3 Animal, Florianópolis, SC, Brasil

⁴ Centro de Recuperação de Animais Marinhos – CRAM, Museu Oceanográfico Professor Eliézer de Carvalho Rios, Universidade Federal do Rio Grande – FURG, Rio Grande, RS, Brasil

⁵ Instituto de Pesquisa e Reabilitação de Animais Marinhos – IPRAM, Cariacica, ES, Brasil

Received: 5 August, 2018

Accepted: 12 November, 2018

Abstract

Avian malaria is one of the most important diseases of captive penguins. We employed morphometric techniques to evaluate hepatic hemosiderosis in rehabilitating wild Magellanic penguins (*Spheniscus magellanicus*) that were negative (n = 9) or naturally infected by different subgenera of *Plasmodium* spp. (n = 24), according with: *Plasmodium* subgenera (*Haemamoeba*, *Huffia*, Other lineages, and Unidentified lineages), severity of *Plasmodium* histopathological lesions, and concurrent diseases, age class (juvenile or adult plumage), sex (male, female or not determined), body score (emaciated, thin, good, excellent, not available), molt, presence or absence of oil contamination upon admission, iron supplementation, and rehabilitation center. The percentage of the area occupied by hemosiderin was called 'Index of Hepatic Hemosiderosis (IHH)'. *Plasmodium*-positive females presented significantly higher IHH values ($17.53 \pm 12.95\%$) than males ($7.20 \pm 4.25\%$; $p = 0.041$). We observed higher levels of congestion ($p = 0.0182$) and pneumonia ($p = 0.0250$) severity between Unidentified lineages *vs.* *Huffia*. We believe that the hepatic hemosiderosis observed in this study was multifactorial, the result of pathological processes caused by malaria, molting, hemoglobin and myoglobin catabolism during migration, anemia, concomitant diseases, and iron supplementation, all possibly potentiated by decreased liver mass. Further studies are needed to clarify the mechanisms of these hypotheses.

Keywords: Avian malaria, iron, hemosiderin, liver, morphometry, penguin.

Resumo

Malária aviária é uma das mais relevantes doenças em pinguins cativos. Foram aplicadas técnicas morfométricas para avaliar a hemossiderose hepática em pinguins-de-Magalhães (*Spheniscus magellanicus*) de vida livre em reabilitação negativos (n = 9) e naturalmente infectados por diferentes subgêneros de *Plasmodium* spp. (n = 24), quanto a: subgênero de *Plasmodium* (*Haemamoeba*, *Huffia*, Outras Linhagens, e Linhagens não identificadas), severidade das lesões histopatológicas causadas por *Plasmodium* e doenças concomitantes, faixa etária (plumagem juvenil ou adulta), sexo (macho, fêmea, indeterminado), condição corporal (emaciado, magro, bom, excelente, indisponível), muda, presença/ausência de óleo a admissão,

*Corresponding author: Ana Carolina Ewbank. Laboratório de Patologia Comparada de Animais Selvagens, Departamento de Patologia, Universidade de São Paulo – USP, Av. Orlando Marques de Paiva, 87, Butantã, CEP 05508-270, São Paulo, SP, Brasil. e-mail: carolewbank@yahoo.com.br



suplementação de ferro, e centro de reabilitação. A porcentagem da área ocupada por hemossiderina foi denominada “Índice de Hemossiderose Hepática (IHH)”. Fêmeas *Plasmodium*-positivas apresentaram IHH significativamente mais elevado que machos, respectivamente, $17,53 \pm 12,95\%$ e $7,20 \pm 4,25\%$ ($p = 0,041$). Níveis mais elevados de congestão ($p = 0,0182$) e pneumonia ($p = 0,0250$) foram observados entre Linhagens não identificadas *vs.* *Huffia*. Possivelmente, a hemossiderose hepática observada nesse estudo seja multifatorial, resultado de processos patológicos causados por malária, muda, catabolismo de hemoglobina e mioglobina durante a migração, anemia, doenças concomitantes e suplementação de ferro, potencialmente intensificados por massa hepática reduzida. Estudos complementares são necessários para esclarecer os mecanismos de tais hipóteses.

Palavras-chave: Malária aviária, ferro, hemosiderina, fígado, morfometria, pinguim.

Introduction

The Magellanic penguin (*Spheniscus magellanicus*) breeds in colonies located on the coasts of Chile, Argentina, and the Falkland (Malvinas) Islands. During their winter migration, individuals from the Atlantic Ocean colonies migrate along the continental shelf of northern Argentina, Uruguay, and southern Brazil (STOKES et al., 2014), where they face many challenges (PETRY & FONSECA, 2002; GARCÍA-BORBOROGLU et al., 2006; BRANDÃO et al., 2011; BALDASSIN et al., 2012; REZENDE et al., 2013; KEHRIG et al., 2015). Birds stranded alive are rescued and referred to local rehabilitation centers along the Brazilian coast (GARCÍA-BORBOROGLU et al., 2010; RODRIGUES et al., 2010; RUOPPOLO et al., 2012).

While under rehabilitation, these birds may be exposed to avian malaria, one of the most significant threats to the conservation of penguins, able to seriously compromise their survival in captivity (JONES & SHELLAM, 1999; VANSTREELS et al., 2016). Avian malaria is transmitted by Culicidae mosquitoes infected with *Plasmodium* spp. protozoans (VALKIŪNAS, 2005), and is relatively asymptomatic in most avian species, but potentially pathogenic for those species that have not co-evolved with the parasite, such as penguins (Spheniscidae) (VALKIŪNAS, 2005; VANSTREELS et al., 2016). Penguins are highly susceptible to avian malaria. Rapid and severe outbreaks with high mortality (VANSTREELS et al., 2016) have been reported worldwide in populations of several penguin species in the wild (FANTHAM & PORTER, 1944; LAIRD, 1950; LEVIN et al., 2009), in captivity (GRINER & SHERIDAN, 1967; BAK et al., 1984; FIX et al., 1988), and undergoing rehabilitation (PARSONS & UNDERHILL, 2005). In Brazil, reports of avian malaria in Magellanic penguins include birds in zoos (BUENO et al., 2010) and rehabilitation centers (SILVEIRA et al., 2013; VANSTREELS et al., 2014, 2015).

Iron homeostasis in birds, as in other vertebrates, is regulated mainly through intestinal absorption and excess iron is stored as intracellular aggregations of ferritin or hemosiderin, particularly in hepatic and splenic cells (NORAMBUENA & BOZINOVIC, 2009). Hemosiderosis is the accumulation of iron in hepatic and splenic sinusoidal macrophages and hepatocytes without significant architectural, cellular, or functional alterations (LOWENSTEIN & MUNSON, 1999). Iron absorption is influenced by age, iron and health status, gastrointestinal conditions, the amount and chemical form of the iron ingested, and the amount and proportions of various other organic and inorganic components of the diet (CRISSEY et al., 2000; SHEPPARD & DIERENFELD, 2002; PEREIRA et al., 2010). Although there is no conclusive evidence

that hepatic hemosiderosis has any clinical significance in birds, it has been associated with concurrent infections, neoplasia, parasitism, anemia, hepatopathies, intoxications, starvation, oil exposure, trauma, lead poisoning, and the physiological cycle (e.g., egg production and laying, migration, and molting) (CORK et al., 1995; GOTTDENKER et al., 2008; PEREIRA et al., 2010; KLASING et al., 2012).

Hemosiderosis, especially hepatic, has been previously reported in *Plasmodium*-infected penguins (FIX et al., 1988; GRILO et al., 2016), believed to be caused by the build up of hemosiderin in visceral macrophages following intravascular hemolysis and phagocytosis of parasitized cells (VANSTREELS & PARSONS, 2014). However, these studies did not evaluate if such condition varied among *Plasmodium* spp. subgenera or considered other possible causes of hemosiderosis aside from *Plasmodium* spp. infection. Considering the importance of avian malaria to penguin conservation and the limited amount of information regarding iron metabolism and the pathogenesis of avian malaria in birds, our objective was to quantify the hepatic hemosiderosis in *Plasmodium* spp.-negative and -naturally infected Magellanic penguins that died while under care in rehabilitation centers in Brazil, in regards to the presence of *Plasmodium* and *Plasmodium* species/lineages, and within the context of these birds' biology and individual history.

Material and Methods

The Magellanic penguins evaluated in this study represent all *Plasmodium*-positive ($n = 24$) and some of the *Plasmodium*-negative ($n = 9$) birds, that died while under rehabilitation along 2500 km of the Brazilian coastline between January 2009 and February 2013 during studies on the prevalence of *Plasmodium* in Magellanic penguins (SILVEIRA et al., 2013; VANSTREELS et al., 2014, 2015). Physical and postmortem examinations, including necropsy and hematology findings, are described elsewhere (VANSTREELS et al., 2015). The positive group comprised 24 *Plasmodium*-positive animals: 21 individuals naturally infected by *Plasmodium* spp., previously diagnosed by morphologic (blood smears, histopathology) and molecular techniques (nested PCR, genetic sequencing) (VANSTREELS et al., 2015); and 3 additional natural infection cases identified using the same methods (cases 3, 14, and 15). Positive birds were divided into 4 study groups: (1) *Haemamoeba* (*P. cathemerium* and *P. tejerai*), (2) *Huffia* (*P. elongatum*), (3) Other lineages (*P. nucleophilum*, *P. unalis*, and *Plasmodium* sp. lineages E, G, and H), and (4) Unidentified lineages (*Plasmodium* sp. that

could not be morphologically or molecularly characterized). The positive birds evaluated in this study either (a) died suddenly, (b) presented antemortem malaria-associated clinical signs, dying after unsuccessful treatment, or (c) were euthanized for reasons unrelated to malaria (e.g., aspergillosis). Nine *Plasmodium*-negative Magellanic penguins were selected based on individual history and a negative diagnosis for *Plasmodium* spp. on both nested PCR and blood smears. Selected *Plasmodium*-negative birds died without presenting any clinical signs of concurrent infections,

and remained in captivity for similar periods as the positive birds. All birds from both groups (positive and negative) were kept under similar infrastructure and husbandry conditions, and diet and iron supplementation varied among rehabilitation facilities (Table 1). All birds were examined *postmortem*, and liver samples were collected and fixed in 10% formalin for a minimum of 72 hours before processing. Paraffin embedded liver fragments were sectioned at 5 µm and stained with haematoxylin-eosin and Perls stains, as described by Luna (1992) (Figure 1). The Index of Hepatic




Figure 1. Hepatic hemosiderosis in a Magellanic penguin (*Spheniscus magellanicus*). A. Areas of intracellular hemosiderin deposition in Kupffer cells (black arrowheads) and hepatocytes (red arrowhead), diffuse cordal derangement (red asterisk) and inflammatory infiltrate (black asterisk). H&E stain. 40 µm. B. Intracellular hemosiderin deposits within Kupffer cells (black arrowheads) and hepatocytes (red arrow). Perls stain. 80 µm.

Table 1. *Plasmodium*-positive and control Magellanic penguins (*Spheniscus magellanicus*) according with rehabilitation center, iron supplementation, administration criteria and diet.

Rehabilitation Center	Iron supplementation (Dose/Route/Frequency)	Administration Criteria	Diet (whole fish-based)
CRAM ^a	Ferrodex ^d → 0.2mg/kg/IM – single dose (until 2013)	Hematocrit < 35%	banded croaker (<i>Paralichthys brasiliensis</i>)
	Ferrodex → 0.2mg/kg/IM – single dose (from 2013 to the present)	Hematocrit < 30% until the parameter reaches the acceptable interval for the species ^f	stripped weakfish (<i>Cynoscion guatucupa</i>) acoupa weakfish (<i>Cynoscion acoupa</i>)
	Ferrodex → 0.2mg/kg/IM – single dose in association with Hemolitan pet ^c → 0.1mg/kg PO – q24h	Hematocrit < 35%until the parameter reaches the acceptable interval for the species ^f	Argentine croaker (<i>Umbrina canosai</i>)
	(from 2013 to the present)		whitemouth croaker (<i>Micropogonias furnieri</i>)
R3 Animal ^b	Ferrodex → 1ml/IM– single dose	Upon admission	Brazilian sardine (<i>Sardinella brasiliensis</i>)
IPRAM ^c	Hemolitan pet → 1ml/PO – q24h	Penguins treated as huddles. Medication administered upon admission and maintained for days to weeks, until the birds exhibited normal colored oral mucosae	Brazilian sardine (<i>Sardinella brasiliensis</i>)

^aCRAM = Centro de Recuperação de Animais Marinhos (Rio Grande do Sul state, Brazil). ^bR3 = Associação R3 Animal (Santa Catarina state, Brazil). ^cIPRAM = Instituto de Pesquisa e Reabilitação de Animais Marinhos (Espírito Santo state, Brazil). ^dFerrodex® (Tortuga, Companhia Zootécnica Agrária, São Amaro, São Paulo, Brazil). ^eHemolitan pet (Indústria e Comércio de Produtos Veterinários Ltda., Louveira, São Paulo, Brazil). ^fReference interval for the species (SILVA-FILHO & RUOPPOLO, 2014).

Hemosiderosis (IHH) was evaluated according with the presence of infection, *Plasmodium* species/lineage, malaria severity (based on the classification of the histopathological findings observed in

heart, lung, liver, and spleen samples: mild, mild to moderate, moderate, moderate to severe, and severe - Table 2), concurrent diseases and/or significant lesions, individual history (age class

Table 2. *Plasmodium*-positive and control Magellanic penguins according with the Index of Hepatic Hemosiderosis (IHH), study group, *Plasmodium* lineage, *Plasmodium* species and malaria severity.

Case	Study group	<i>Plasmodium</i> lineage	IHH (%)	Malaria severity
1	<i>Haemamoeba</i>	<i>P. cathemerium</i>	0.4	Moderate pulmonary congestion Moderate granulocytic pneumonia Moderate-severe monocytic hepatitis
2	<i>Haemamoeba</i>	<i>P. cathemerium</i>	14.8	Moderate-severe pulmonary congestion Moderate to severe granulocytic pneumonia Mixed moderate-severe splenitis Mild-moderate monocytic hepatitis
3	<i>Haemamoeba</i>	<i>P. cathemerium</i>	1.1	Mild-moderate acute diffuse granulocytic pneumonia Parasitic pneumonia compatible with <i>Schistosoma</i> sp. Diffuse necrotic splenitis Splenic lymphocytolysis, histiocytolysis and lymphoid hypoplasia Parasitic granulomatous hepatitis Diffuse renal degeneration Mild cerebellar congestion Mild necrotic focal myocarditis Mild monocytic diffuse myocarditis
4	<i>Haemamoeba</i>	<i>P. cathemerium</i>	18.3	Moderate pulmonary congestion Moderate granulocytic pneumonia Moderate-severe granulocytic splenitis Moderate mixed hepatitis
5	<i>Haemamoeba</i>	<i>P. cathemerium</i>	4.2	Moderate-severe pulmonary congestion Mild granulocytic pneumonia Moderate-severe mixed splenitis Moderate-severe monocytic hepatitis
6	<i>Haemamoeba</i>	<i>P. cathemerium</i>	5.8	Moderate pulmonary congestion Moderate granulocytic pneumonia Mild-moderate granulocytic splenitis Mild-moderate mixed hepatitis
7	<i>Haemamoeba</i>	<i>P. tejerai</i>	10.8	Moderate-severe pulmonary congestion Moderate granulocytic pneumonia Mild-moderate granulocytic splenitis Moderate monocytic hepatitis Hepatic ductal hyperplasia
8	<i>Haemamoeba</i>	<i>P. tejerai</i>	1.5	Moderate-severe pulmonary congestion Moderate mixed hepatitis Hepatic ductal hyperplasia
9	<i>Haemamoeba</i>	<i>P. tejerai</i>	38.2	Moderate pulmonary congestion Moderate granulocytic pneumonia Mild-moderate granulocytic splenitis Mild-moderate granulocytic hepatitis Hepatic ductal hyperplasia
10	<i>Haemamoeba</i>	<i>P. tejerai</i>	31.9	Moderate-severe pulmonary congestion Severe granulocytic pneumonia Mild-moderate mixed splenitis Moderate monocytic hepatitis Hepatic ductal hyperplasia Mild monocytic myocarditis

Table 2. Continued...

Case	Study group	<i>Plasmodium</i> lineage	IHH (%)	Malaria severity
11	<i>Haemamoeba</i>	<i>P. tejeraei</i>	9.4	Mild-moderate pulmonary congestion
				Mild-moderate granulocytic pneumonia
				Mild-moderate mixed splenitis
				Mild granulocytic myocarditis
12	<i>Huffia</i>	<i>P. elongatum</i>	22.8	Moderate pulmonary congestion
				Mild granulocytic pneumonia
				Moderate monocytic hepatitis
13	<i>Huffia</i>	<i>P. elongatum</i>	1.4	Mild-moderate congestion
				Acute granulocytic diffuse myocarditis
				Moderate cerebellar congestion
				Moderate diffuse acute granulocytic pneumonia
				Moderate coalescent multifocal histiocytosis, moderate lymphoid hypoplasia, lymphocytolysis, extramedullary erythropoiesis, mild-moderate mixed diffuse necrotic splenitis
				Mixed multifocal interstitial nephritis
				Moderate-severe multifocal necrotic hepatitis
				Extramedullary hepatic erythropoiesis
14	<i>Huffia</i>	<i>P. elongatum</i>	6.9	Hepatic ductal hyperplasia
				Mild cardiac congestion
				Mild acute diffuse myocarditis
				Mild to moderate cerebellar congestion
				Moderate mixed diffuse pneumonia
				Moderate pulmonary congestion
				Spleen tissue was not available for examination
				Diffuse renal degeneration
15	Other lineages	<i>P. nucleophilum</i>	27.6	Moderate-severe mixed coalescent multifocal necrotic periportal hepatitis and mild extramedullary erythropoiesis
				Mild pulmonary congestion
				Mild-moderate monocytic pneumonia
				Mild-moderate monocytic splenitis
16	Other lineages	<i>P. unalis</i>	15	Moderate-severe mixed hepatitis
				Moderate-severe pulmonary congestion
				Moderate-severe monocytic pneumonia
17	Other lineages	<i>Plasmodium</i> sp. <i>E</i>	6.2	Mild monocytic hepatitis
				Moderate pulmonary congestion
				Moderate granulocytic pneumonia
				Moderate granulocytic splenitis
18	Other lineages	<i>Plasmodium</i> sp. <i>G</i>	4.8	Mild monocytic hepatitis
				Hepatic ductal hyperplasia
				Moderate-severe pulmonary congestion
				Moderate granulocytic pneumonia
19	Other lineages	<i>Plasmodium</i> sp. <i>G</i>	45.4	Moderate granulocytic splenitis
				Moderate-severe monocytic hepatitis
				Moderate-severe pulmonary congestion
				Severe granulocytic pneumonia
20	Other lineages	<i>Plasmodium</i> sp. <i>H</i>	14.6	Mild granulocytic splenitis
				Mild-moderate granulocytic hepatitis
				Severe myocardial necrosis
				Moderate pulmonary congestion
				Moderate mixed pneumonia
				Moderate monocytic hepatitis

Table 2. Continued...

Case	Study group	<i>Plasmodium</i> lineage	IHH (%)	Malaria severity
21	Unidentified lineages	<i>Plasmodium</i> sp.	10.7	Moderate-severe pulmonary congestion
				Moderate-severe monocytic pneumonia
				Moderate granulocytic splenitis
				Moderate-severe monocytic hepatitis
				Mild-moderate monocytic myocarditis
22	Unidentified lineages	<i>Plasmodium</i> sp.	12.8	Moderate-severe pulmonary congestion
				Moderate-severe granulocytic pneumonia
				Moderate-severe granulocytic splenitis
				Moderate granulocytic hepatitis
				Mild granulocytic hepatitis
23	Unidentified lineages	<i>Plasmodium</i> sp.	8.3	Severe pulmonary congestion
				Severe monocytic pneumonia
				Mild-moderate mixed splenitis
				Moderate monocytic hepatitis
				Mild granulocytic myocarditis
24	Unidentified lineages	<i>Plasmodium</i> sp.	13.8	Moderate-severe pulmonary congestion
				Moderate granulocytic pneumonia
				Mild-moderate monocytic hepatitis
				Hepatic ductal hyperplasia
25	<i>Plasmodium</i> - negative	Negative	0.9	-
26	<i>Plasmodium</i> - negative	Negative	18.1	-
27	<i>Plasmodium</i> - negative	Negative	11.6	-
28	<i>Plasmodium</i> - negative	Negative	2.9	-
29	<i>Plasmodium</i> - negative	Negative	0.9	-
30	<i>Plasmodium</i> - negative	Negative	5.6	-
31	<i>Plasmodium</i> - negative	Negative	13.4	-
32	<i>Plasmodium</i> - negative	Negative	6.5	-
33	<i>Plasmodium</i> - negative	Negative	12.5	-

[juvenile or adult plumage] (SILVA-FILHO & RUOPPOLO, 2014), sex [male, female or unknown], body mass on admission (emaciated, thin, good, excellent, not available [N/A]), body score, hematocrit, presence or absence of oil upon admission [oiling], molt and iron supplementation [Fe] (Table 3). To establish the IHH, slides were evaluated by light microscopy and captured images were analyzed by a computerized image analyses software (Image Pro® Plus, version 5.1.2.59, Media Cybernetics) to determine the percentage of liver tissue occupied by hemosiderin pigment. The intersection of the two longest axis of each bird's hepatic section was determined as the center of the sample, and used as the reference field to capture images at 20X magnification. Another 8 images were captured as TIFF files, at 50 µm of this point, and 45° intervals, under the same lighting conditions and magnification. The malarial pigment, called hemozoin, stains brown with Perls stain, while hemosiderin stains bright turquoise blue, promoting an easy visual distinction between the two pigments. Hemosiderin deposits were semi-automatically delineated on the Perls stained slides, starting at the lowest blue intensity to the most intense, to avoid the inclusion of artifacts and the background. When overlapping areas of formolic pigment and hemosiderin

deposition were present, the areas of formolic pigment deposition were individually marked and deducted from such areas. In order to standardize the technique, a maximum digital zoom of 50% was stipulated. The average of the 9 microscopic fields was considered the IHH.

One-way ANOVA/Kruskal-Wallis test followed by Dunn's multiple comparisons test and Mann-Whitney tests were used to verify differences between the evaluated parameters and groups. The Spearman test was used to evaluate the correlation between quantitative variables. The significance level was defined as 5%. Data were analyzed with GraphPad Prism software (Version 4.02 for Windows, GraphPad Software, Inc.).

All samples used in this study were collected in full compliance with specific federal permits issued by the Brazilian Ministry of Environment and approved by the Biodiversity Information and Authorization System (SISBIO 20825-8). All procedures were performed according to the Ethical Committee in Animal Research of the School of Veterinary Medicine and Animal Sciences, University of São Paulo (Process numbers 1757/2009 and 9411100414).

Table 3. *Plasmodium*-positive and control Magellanic penguins according with the Index of Hepatic Hemosiderosis (IHH), rehabilitation center, age class, sex, body mass on admission (g), body score, hematocrit (Ht), presence or absence of oil upon admission (oil), moult, iron supplementation (Fe) and concurrent disease and/or significant lesion.

Case	IHH (%)	Rehab center ^a	Age class ^b	Sex ^c	Body mass (g)	Body score	Ht ^d	Oil	Moult	Fe	Concurrent disease and/or significant lesion
1	0.4	IPRAM	J	F	1915	Thin	N/A	No	No	No	Aspergillosis
2	14.8	IPRAM	J	F	2290	Thin	N/A	No	No	No	Spleen hemorrhage and rupture
3	1.1	CRAM	J	M	N/A	N/A	N/A	No	No	N/A	-
4	18.3	R3	J	F	2560	Good	44%	No	Yes	No	Helminthiasis
5	4.2	R3	J	M	N/A	Good	40%	No	Yes	No	N/A
6	5.8	R3	J	M	3240	Good	N/A	No	Yes	No	Avipoxvirus
7	10.8	R3	J	M	2400	N/A	32%	No	No	Yes	N/A
8	1.5	R3	J	ND	2600	N/A	N/A	No	No	Yes	Disseminated intravascular clotting
9	38.2	R3	J	ND	3450	N/A	N/A	No	Yes	Yes	N/A
10	31.9	R3	A	F	3300	Good	N/A	No	No	No	Helminthiasis (small intestine)
11	9.4	R3	J	M	2100	Good	N/A	No	No	N/A	N/A
12	22.8	IPRAM	J	F	2260	Good	N/A	No	No	No	Biliary stasis, Unidentified cysts (myocardium)
13	1.4	CRAM	A	F	N/A	Good	N/A	No	No	N/A	-
14	6.9	CRAM	A	M	N/A	Good	N/A	No	No	N/A	-
15	27.6	CRAM	J	F	2802	Good	50%	Yes	No	N/A	Amyloidosis, helminthiasis (gastrointestinal, lungs, liver)
16	15	CRAM	J	F	2810	Good	40%	Yes	No	N/A	Aspergillosis, Amyloidosis
17	6.2	R3	A	F	3220	Good	N/A	No	No	No	-
18	4.8	R3	J	M	3060	Good	27%	No	Yes	No	Amyloidosis, helminths (air sacs)
19	45.4	R3	J	F	2700	Bad	19%	No	Yes	No	Aspergillosis, pancarditis
20	14.6	R3	A	M	3260	Good	N/A	No	Yes	No	Helminthiasis
21	10.7	CRAM	J	ND	3890	Good	36%	No	No	Yes	-
22	12.8	CRAM	J	F	2910	N/A	33%	No	No	N/A	Spleen hemorrhage and rupture
23	8.3	CRAM	J	ND	2645	Thin	53%	Yes	No	Yes	-
24	13.8	CRAM	A	F	2750	Thin	46%	Yes	No	Yes	-
25	0.9	IPRAM	J	F	1750	Cachectic	N/A	Yes	No	Yes	-
26	18.1	IPRAM	J	F	1865	Thin	N/A	Yes	No	Yes	-
27	11.6	IPRAM	J	F	2015	Thin	N/A	Yes	No	Yes	-
28	2.9	IPRAM	J	F	2250	Thin	N/A	Yes	No	Yes	-
29	0.9	R3	J	ND	N/A	Good	N/A	Yes	No	Yes	-
30	5.6	R3	A	ND	N/A	Good	N/A	Yes	No	Yes	-
31	13.4	CRAM	J	M	2248	N/A	N/A	Yes	No	N/A	-
32	6.5	CRAM	J	M	2268	N/A	N/A	Yes	No	N/A	-
33	12.5	CRAM	J	M	3110	N/A	N/A	Yes	No	N/A	-

^aIPRAM = Instituto de Pesquisa e Reabilitação de Animais Marinhos (Espírito Santo state, Brazil); R3 = Associação R3 Animal (Santa Catarina state, Brazil); CRAM = Centro de Recuperação de Animais Marinhos (Rio Grande do Sul state, Brazil); ^bJ = Juvenile; A = adult; ^cF = female; M = male; ND = not determined;

^dN/A = Information not available.

Results and Discussion

The pulmonary congestion, intense inflammatory response (especially in liver and spleen), and hemosiderosis found in our birds are in agreement with previous microscopic descriptions of *Plasmodium* lesions in penguins (FIX et al., 1988; VANSTREELS & PARSONS, 2014; GRILO et al., 2016). However, there were no significant differences neither in the presence of moderate to severe hepatic hemosiderosis or in IHH values between

Plasmodium-negative vs. *Plasmodium*-positive individuals or in regards to the *Plasmodium* species/lineage. The epidemiology and pathogenicity of the *Plasmodium* lineages previously diagnosed in our birds (VANSTREELS et al., 2015) raised the question on whether the *Plasmodium* subgenus *Haemamoeba* was more pathogenic to these penguins than other *Plasmodium* subgenera. Unfortunately, we were not able to find a clear answer to that question; these lineages are not completely understood and have been recorded only recently. However, we found higher levels of

congestion ($p = 0.0182$) and pneumonia severity ($p = 0.0250$) between Unidentified lineages *vs.* *Huffia* (Figure 2).

Plasmodium-positive females presented significantly higher IHH values ($17.53 \pm 12.95\%$) than males ($7.20 \pm 4.25\%$; $p = 0.041$) (Figure 3), in contrast with findings in toucans and aracarís (Ramphastidae family) (CUBAS, 2008) and in red-spectacled amazons (*Amazona pretrei*) (Psittacidae family) (PEREIRA et al., 2010), who failed to observe correlation between the rate of hepatic hemosiderosis and sex. Hemosiderosis has been previously associated with egg production (GOTTDENKER et al., 2008), which could not have played a role in our findings based on the age and underdeveloped gonads observed in necropsy. It is not clear why females presented higher IHH values than males; however, one may not eliminate the possibility of sampling influence, either due to the limited number of evaluated birds or higher number of females than males, which reflects previous studies documenting a female-biased sex ratio in Magellanic penguins stranded along the Brazilian coast (VANSTREELS et al., 2013). Comparison between the Pneumonia score (A) and Congestion Score (B) among *Plasmodium* species/lineages, and IHH in female *vs.* male *Plasmodium*-positive Magellanic penguins (*Spheniscus magellanicus*) (C) are shown in Figure 3.

Concurrent diseases were not statistically relevant. According to the institutions' protocols, many of the birds analyzed in this study were considered anemic, and received iron supplementation, which we believe may have also played a role in the hepatic hemosiderosis observed in this study. The physiological values of circulating iron concentration in Magellanic penguins have been previously evaluated (GHEBREMESEKEL et al., 1989); however, to the authors' knowledge, there are no available data regarding iron hepatic levels for this species. Furthermore, the posology of iron supplementation is usually *ad-hoc*, and might not address the physiological requirements of these birds. In addition, iron represents a critical nutrient for pathogens and tight regulation of iron ("iron-withholding") is a paramount defense strategy of

the host's innate immune system (KOSKI & SCOTT, 2003; JOHNSON, 2008; BEERNAERT et al., 2010; CASSAT & SKAAR, 2013). Thus, individuals suffering from iron overload have an enhanced risk of infection (BEERNAERT et al., 2010). Some of our specimens presented gastrointestinal nematodes, air sac parasites, and aspergillosis. It has been suggested that chronic infection or repeated acute infections could lead to greater accumulation of iron in the liver of birds (KLASING et al., 2012). Unfortunately, the low number of concurrent diseases observed in this study prevented statistical evaluation. Aspergillosis and parasitosis could have not only contributed to hemosiderosis, but also thrived upon iron supplementation, further debilitating the affected birds and contributing to their death. Due to all the factors discussed above, we recommend caution in the employment of iron supplementation to Magellanic penguins undergoing rehabilitation.

The IHH values in relation to presence of infection, *Plasmodium* species/lineage and pathology are summarized in Table 3.

Individual history parameters were not statistically relevant to the IHH values either. Magellanic penguins molt soon after breeding season, immediately prior to their winter migration (SILVA-FILHO & RUOPPOLO, 2014). Anseriformes and Passeriformes of the Sturnidae family showed an association between the hepatic level of hemosiderosis in this same period, with increased amounts of positive iron staining primarily located within lysosomal organelles of parenchymal hepatic cells (WARD et al., 1988; CORK et al., 1995). Such findings are possibly related to an increased hematopoietic activity and consequent erythrocyte production required to enhance oxygen uptake during periods of increased thermogenesis or changes in the levels of thyroid hormone involved in the molting process (WARD et al., 1988; CORK et al., 1995; OSBORN, 1979; CORK, 2000). Although not statistically relevant, some birds (4, 5, 6, 9, 18, 19, and 20) were molting during the study, so it is not possible to completely discard the hypothesis that residual iron from molting might have contributed to the hepatic hemosiderosis observed in our birds.

Fat can be severely depleted during migration, forcing birds to use body protein from the heart, pectoral, and leg muscles and the digestive organs (intestine, gizzard, and liver) as a second fuel option (LINDSTRÖM & PIERSMA, 1993; BATTLEY et al., 2000; BAUCHINGER & BIEBACH, 2001; SCHWILCH et al., 2002; MCWILLIAMS et al., 2004). Protein metabolism stems to a large extent from the skeletal muscles, which in penguins are especially rich in myoglobin, releasing the iron present in heme group of myoglobin, and possibly contributing to the development of hemosiderosis (SCHWARTZ et al., 1973; BAUCHINGER & BIEBACH, 2001; PONGANIS et al., 2010). The majority of migrating Magellanic penguins stranded in Brazil are in poor body condition, with markedly atrophied skeletal muscles and no subcutaneous fat deposits (CARDOSO et al., 2011). We assessed the true nutritional condition of these birds: body mass (ranging from 3.5-5 kg in wild specimens) and body condition score (SILVA-FILHO & RUOPPOLO, 2014). The only 2 emaciated birds (1 and 19), however, presented IHH values within the range found in this study. Nevertheless, increased iron deposition has been reported in malnourished black-necked swans (*Cygnus melanocoryphus*) (NORAMBUENA & BOZINOVIC, 2009), in the liver of rats




Figure 2. Moderate to severe pulmonary congestion and moderate to severe diffuse granulocytic pneumonia in a Magellanic penguin (*Spheniscus magellanicus*). H&E stain. 40 μ m.




Figure 3. Comparison between the Pneumonia score (A) and Congestion Score (B) among *Plasmodium* species/lineages, and the index of hepatic hemosiderosis (IHH) in female *vs.* male *Plasmodium*-positive Magellanic penguins (*Spheniscus magellanicus*) (C).

subjected to food restriction (ROZENTAL et al., 1967), and in some severe cases of prolonged malnutrition in obese humans subjected to a variety of fasting periods (MITCHELL & JOHNS, 2008). It is important to consider that malnutrition may have a role in hemosiderosis in migratory bird species.

Intoxication (e.g., petroleum products) and hemoparasitosis are known causes of hemolytic anemia (FRY & LOWENSTINE, 1985). Systemic absorption of petroleum and its derivatives following acute or chronic oil exposure causes hemolytic anemia in birds (LEIGHTON, 1986; KHAN & RYAN, 1991; DINHOPL et al., 2011). Oil pollution is an important conservation threat to Magellanic penguins on wintering grounds (GARCÍA-BORBOROGLU et al., 2006). In this study, however, IHH was not significantly different between individuals with and without history of acute oil contamination upon arrival at the rehabilitation centers, suggesting that (a) oil exposure may have been too superficial or rapidly mitigated and therefore systemic absorption was minimal, and/or (b) other factors may have played a greater role in producing/modulating hemosiderosis in a manner that surpassed or contradicted any detectable effects from oiling. Nevertheless, these birds' history during migration is unknown, so we were not able to completely exclude the potential influence of chronic oil exposure in the hepatic hemosiderosis seen in this study.

Although previous studies (CRANFIELD et al., 1994; ONG et al., 2006) suggested that the low parasitaemia levels seen in *Plasmodium*-infected penguins are not sufficient to cause clinical anemia, *Plasmodium*-infected erythrocytes may suffer hemolysis or induce erythrocyte sequestration and extravascular hemolysis (VALKIÜNAS, 2005), releasing iron, which is subsequently stored in macrophages, as ferritin and hemosiderin (RATLEDGE, 2007). Eventually, cases 2 and 22 presented spleen hemorrhage and rupture, whereas case 8 presented disseminated intravascular coagulation - acute vascular complications likely associated with avian malaria that if not the primary cause, contributed to the development of secondary hemosiderosis. Another possible cause of anemia was the infestation by gastrointestinal nematodes (cases 4, 10, 15, 18 and 20).

Liver mass changes caused by the spontaneous long-term fasting characteristic of the annual cycle of penguins during reproduction and molt (GROSCOLAS & ROBIN, 2001) and/or due to concurrent diseases could have interfered with the morphometric evaluation of hepatic hemosiderin. It is unclear if changes in digestive organs of birds during migration are related to cell size (hypertrophy/atrophy) and/or cell number (hyperplasia/hypoplasia) (WAGENMAKERS et al., 1984). Further morphometric studies are needed in order to test these possibilities.

Both the amount of dietary iron and length of exposure to dietary iron are factors in hepatic iron accumulation in birds (CRISSEY et al., 2000; HELMICK et al., 2011). Unfortunately, data regarding their diet while in the Brazilian continental shelf during the winter migration is limited and there is no such information regarding the species offered to the studied birds while under care. To our knowledge, the only suggested nutritional iron concentration for adult penguin diets in a dry matter basis (60 - 80mg/kg) is based on requirements of domestic poultry, cats, and inferences from wild foods composition (WALLACE & WALSH, 2005; AZA Penguin Taxon Advisory Group, 2014). Further studies are needed to determine the nutritional iron requirements of Magellanic penguins.

Finally, our findings show that other factors aside from avian malaria likely played a role in the presence of hepatic hemosiderosis in *Plasmodium*-infected and *Plasmodium*-negative Magellanic penguins under rehabilitation in Brazil, and the importance of considering the species life cycle and clinical history when evaluating hepatic hemosiderosis. We propose that the increased hepatic hemosiderosis noted in this study was multifactorial, the result of pathological processes caused by malaria, molting, hemoglobin and myoglobin catabolism promoted by the intense energetic demands of Magellanic penguins' winter migration, anemia, concomitant diseases, and iron supplementation while under care, all possibly potentiated by an altered liver mass.

Acknowledgements

We thank the staffs of the Centro de Recuperação de Animais Marinhos da Universidade Federal do Rio Grande (CRAM-FURG), Associação R3 Animal and Instituto de Pesquisa e Reabilitação de Animais Marinhos (IPRAM), for their collaboration and support, and the State Environmental and Water Resources Institute (IEMA) and Brazilian Institute of the Environment and Renewable Natural Resources (IBAMA) for their assistance with the penguins rescued in Espírito Santo. We also thank Jorge Oyakawa, Sândara Sguario and Luciano Bugalho at the School of Veterinary Medicine and Animal Sciences, University of São Paulo for their technical support. This study was supported by the Coordination for the Improvement of Higher Level Personnel (CAPES) and the São Paulo Research Foundation (FAPESP) under Grant [2009/53956-9, 2010/51801-5].

References

- AZA Penguin Taxon Advisory Group. *Penguin (Spheniscidae) care manual* [online]. Silver Spring: Association of Zoos and Aquariums; 2014 [cited 2018 Mar 30]. Available from: https://www.speakcdn.com/assets/2332/penguin_care_manual_aza_final_2014.pdf
- Bak UB, Park JC, Lim YJ. An outbreak of malaria in penguins at the Farm-land Zoo. *Korean J Parasitol* 1984; 22(2): 267-272. <http://dx.doi.org/10.3347/kjp.1984.22.2.267>. PMID:12891023.
- Baldassin P, Taniguchi S, Gallo H, Silva RJ, Montone RC. Persistent organic pollutants in juvenile Magellan penguins (*Spheniscus magellanicus*) found on the northern shore of the state of São Paulo and southern shore of the state of Rio de Janeiro, Brazil. *Mar Pollut Bull* 2012; 64(11): 2502-2506. <http://dx.doi.org/10.1016/j.marpolbul.2012.07.035>. PMID:22910330.
- Battley PF, Piersma T, Dietz MW, Tang S, Dekinga A, Hulsman K. Empirical evidence for differential organ reductions during trans-oceanic bird flight. *Proc Biol Sci* 2000; 267(1439): 191-195. <http://dx.doi.org/10.1098/rspb.2000.0986>. PMID:10687826.
- Bauchinger U, Biebach H. Differential catabolism of muscle protein in garden warblers (*Sylvia borin*): flight and leg muscle act as a protein source during long-distance migration. *J Comp Physiol B* 2001; 171(4): 293-301. <http://dx.doi.org/10.1007/s003600100176>. PMID:11409626.
- Beernaert LA, Pasmans F, Van Waeyenbergh L, Haesebrouck F, Martel A. *Aspergillus* infections in birds: a review. *Avian Pathol* 2010; 39(5): 325-331. <http://dx.doi.org/10.1080/03079457.2010.506210>. PMID:20954008.
- Brandão ML, Braga KM, Luque JL. Marine debris ingestion by Magellanic penguins, *Spheniscus magellanicus* (Aves: Sphenisciformes), from the Brazilian coastal zone. *Mar Pollut Bull* 2011; 62(10): 2246-2249. <http://dx.doi.org/10.1016/j.marpolbul.2011.07.016>. PMID:21864861.
- Bueno MG, Lopez RPG, Menezes RMT, Costa-Nascimento MJ, Lima GFMC, Araújo RAS, et al. Identification of *Plasmodium* relictum causing mortality in penguins (*Spheniscus magellanicus*) from São Paulo Zoo, Brazil. *Vet Parasitol* 2010; 173(1-2): 123-127. <http://dx.doi.org/10.1016/j.vetpar.2010.06.026>. PMID:20638795.
- Cardoso LG, Bugoni L, Mancini PL, Haimovici M. Gillnet fisheries as a major mortality factor of Magellanic penguins in wintering areas. *Mar Pollut Bull* 2011; 62(4): 840-844. <http://dx.doi.org/10.1016/j.marpolbul.2011.01.033>. PMID:21376348.
- Cassat JE, Skaar EP. Iron in infection and immunity. *Cell Host Microbe* 2013; 13(5): 509-519. <http://dx.doi.org/10.1016/j.chom.2013.04.010>. PMID:23684303.
- Cork SC, Alley MR, Stockdale PHG. A quantitative assessment of haemosiderosis in wild and captive birds using image analysis. *Avian Pathol* 1995; 24(2): 239-254. <http://dx.doi.org/10.1080/03079459508419066>. PMID:18645783.
- Cork SC. Iron storage diseases in birds. *Avian Pathol* 2000; 29(1): 7-12. <http://dx.doi.org/10.1080/03079450094216>. PMID:19184782.
- Cranfield MR, Graczyk TK, Beall FB, Ialeggio DM, Shaw ML, Skjoldager ML. Subclinical avian malaria infections in African black-footed penguins (*Spheniscus demersus*) and induction of parasite recrudescence. *J Wildl Dis* 1994; 30(3): 372-376. <http://dx.doi.org/10.7589/0090-3558-30.3.372>. PMID:7933280.
- Crissey SD, Ward AM, Block SE, Maslanka MT. Hepatic iron accumulation over time in European starlings (*Sturnus vulgaris*) fed two levels of iron. *J Zoo Wildl Med* 2000; 31(4): 491-496. [http://dx.doi.org/10.1638/1042-7260\(2000\)031\[0491:HIAOTI\]2.0.CO;2](http://dx.doi.org/10.1638/1042-7260(2000)031[0491:HIAOTI]2.0.CO;2). PMID:11428395.
- Cubas ZS. *Siderose hepática em tucanos e araraçaris* [dissertação]. Paraná: Universidade Federal do Paraná; 2008.
- Dinhopl N, Mostegl MM, Richter B, Nedorost N, Maderner A, Fragner K, et al. Application of in-situ hybridization for the detection and identification of avian malaria parasites in paraffin wax-embedded tissues from captive penguins. *Avian Pathol* 2011; 40(3): 315-320. <http://dx.doi.org/10.1080/03079457.2011.569533>. PMID:21711191.
- Fantham HB, Porter A. On a *Plasmodium* (*Plasmodium* relictum var. spheniscidae, n. var.), observed in four species of penguins. *J Zool* 1944; 114(3): 279-292.

- Fix AS, Waterhouse C, Greiner EC, Stoskopf MF. *Plasmodium relictum* as a cause of avian malaria in wild-caught Magellanic penguins (*Spheniscus magellanicus*). *J Wildl Dis* 1988; 24(4): 610-619. <http://dx.doi.org/10.7589/0090-3558-24.4.610>. PMID:3193555.
- Fry DM, Lowenstine LJ. Pathology of common murre and Cassin's auklets exposed to oil. *Arch Environ Contam Toxicol* 1985; 14(6): 725-737. <http://dx.doi.org/10.1007/BF01055780>.
- García-Borboroglu P, Boersma PD, Ruoppolo V, Pinho-da-Silva-Filho R, Corrado-Adornes A, Conte-Sena D, et al. Magellanic penguin mortality in 2008 along the SW Atlantic coast. *Mar Pollut Bull* 2010; 60(10): 1652-1657. <http://dx.doi.org/10.1016/j.marpolbul.2010.07.006>. PMID:20674946.
- García-Borboroglu P, Boersma PD, Ruoppolo V, Reyes L, Rebstock GA, Griot K, et al. Chronic oil pollution harms Magellanic penguins in the Southwest Atlantic. *Mar Pollut Bull* 2006; 52(2): 193-198. <http://dx.doi.org/10.1016/j.marpolbul.2005.11.004>. PMID:16364372.
- Ghebremeskel K, Williams G, Keymer IF, Horsley D, Gardner DA. Plasma chemistry of rockhopper (*Eudyptes crestatus*), Magellanic (*Spheniscus magellanicus*) and gentoo (*Pygoscelis papua*) wild penguins in relation to moult. *Comp Biochem Physiol A Physiol* 1989; 92(1): 43-47.
- Gottdenker NL, Walsh T, Jiménez-Uzcátegui G, Betancourt F, Cruz M, Soos C, et al. Causes of mortality of wild birds submitted to the Charles Darwin Research Station, Santa Cruz, Galapagos, Ecuador from 2002-2004. *J Wildl Dis* 2008; 44(4): 1024-1031. <http://dx.doi.org/10.7589/0090-3558-44.4.1024>. PMID:18957662.
- Grilo ML, Vanstreels RET, Wallace R, García-Párraga D, Braga ÉM, Chitty J, et al. Malaria in penguins - current perceptions. *Avian Pathol* 2016; 45(4): 393-407. <http://dx.doi.org/10.1080/03079457.2016.1149145>. PMID:27009571.
- Griner LA, Sheridan BW. Malaria (*Plasmodium relictum*) in penguins at the San Diego Zoo. *Am J Vet Clin Pathol* 1967; 1: 7-17.
- Groscolas R, Robin JP. Long-term fasting and re-feeding in penguins. *Comp Biochem Physiol A Mol Integr Physiol* 2001; 128(3): 645-653. [http://dx.doi.org/10.1016/S1095-6433\(00\)00341-X](http://dx.doi.org/10.1016/S1095-6433(00)00341-X). PMID:11246051.
- Helmick KE, Kendrick EL, Dierenfeld ES. Diet manipulation as treatment for elevated serum iron parameters in captive Raggiana bird of paradise (*Paradisaea raggiana*). *J Zoo Wildl Med* 2011; 42(3): 460-467. <http://dx.doi.org/10.1638/2010-0131.1>. PMID:22950319.
- Johnson L. Iron and siderophores in fungal-host interactions. *Mycol Res* 2008; 112(2): 170-183. <http://dx.doi.org/10.1016/j.mycres.2007.11.012>. PMID:18280720.
- Jones HI, Shellam GR. Blood parasites in penguins, and their potential impact on conservation. *Mar Ornithol* 1999; 27: 181-184.
- Kehrig HA, Hauser-Davis RA, Seixas TG, Fillmann G. Trace-elements, methylmercury and metallothionein levels in Magellanic penguin (*Spheniscus magellanicus*) found stranded on the Southern Brazilian coast. *Mar Pollut Bull* 2015; 96(1-2): 450-455. <http://dx.doi.org/10.1016/j.marpolbul.2015.05.006>. PMID:25960272.
- Khan RA, Ryan P. Long term effects of crude oil on common murre (*Uria aalge*) following rehabilitation. *Bull Environ Contam Toxicol* 1991; 46(2): 216-222. <http://dx.doi.org/10.1007/BF01691940>. PMID:2018868.
- Klasing KC, Dierenfeld ES, Koutsos EA. Avian iron storage disease: variations on a common theme? *J Zoo Wildl Med* 2012; 43(3 Suppl): 27-34. <http://dx.doi.org/10.1638/2011-0157.1>. PMID:23156703.
- Koski KG, Scott ME. Gastrointestinal nematodes, trace elements, and immunity. *J Trace Elem Exp Med* 2003; 16(4): 237-251. <http://dx.doi.org/10.1002/jtra.10043>.
- Laird M. Some blood parasites of New Zealand birds. *Zool Pub Victoria Uni College*. 1950; 5: 1-20.
- Leighton FA. Clinical, gross, and histological findings in herring gulls and Atlantic puffins that ingested Prudhoe Bay crude oil. *Vet Pathol* 1986; 23(3): 254-263. <http://dx.doi.org/10.1177/030098588602300305>. PMID:3727312.
- Levin II, Outlaw DC, Vargas FH, Parker PG. *Plasmodium* blood parasite found in endangered Galapagos penguins (*Spheniscus mendiculus*). *Biol Conserv* 2009; 142(12): 3191-3195. <http://dx.doi.org/10.1016/j.biocon.2009.06.017>.
- Lindström Å, Piersma T. Mass changes in migrating birds: the evidence for fat and protein storage re-examined. *Ibis* 1993; 135(1): 70-78. <http://dx.doi.org/10.1111/j.1474-919X.1993.tb02811.x>.
- Lowenstein LJ, Munson L. Iron overload in the animal kingdom. In: Fowler ME, Miller RE. *Zoo and wild animal medicine — current therapy* 4. Philadelphia: WB Saunders; 1999. p. 260-269.
- Luna LG. *Histopathological methods and color atlas of special stains and tissue artifacts*. Gaithersburg: American Histolabs; 1992.
- McWilliams SR, Guglielmo C, Pierce B, Klaassen M. Flying, fasting, and feeding in birds during migration: a nutritional and physiological ecology perspective. *J Avian Biol* 2004; 35(5): 377-393. <http://dx.doi.org/10.1111/j.0908-8857.2004.03378.x>.
- Mitchell EB, Johns J. Avian hematology and related disorders. *Vet Clin North Am Exot Anim Pract* 2008; 11(3): 501-522, vi. <http://dx.doi.org/10.1016/j.cvex.2008.03.004>. PMID:18675731.
- Norambuena MC, Bozinovic F. Effect of malnutrition on iron homeostasis in black-necked swans (*Cygnus melanocoryphus*). *J Zoo Wildl Med* 2009; 40(4): 624-631. <http://dx.doi.org/10.1638/2007-0160.1>. PMID:20063807.
- Ong ST, Ho JZS, Ho B, Ding JL. Iron-withholding strategy in innate immunity. *Immunobiology* 2006; 211(4): 295-314. <http://dx.doi.org/10.1016/j.imbio.2006.02.004>. PMID:16697921.
- Osborn D. Seasonal changes in the fat, protein and metal content of the liver of the starling *Sturnus vulgaris*. *Environ Pollut* 1979; 19(2): 145-155. [http://dx.doi.org/10.1016/0013-9327\(79\)90145-9](http://dx.doi.org/10.1016/0013-9327(79)90145-9).
- Parsons NJ, Underhill LG. Oiled and injured African penguins *Spheniscus demersus* and other seabirds admitted for rehabilitation in the Western Cape, South Africa, 2001 and 2002. *Afr J Mar Sci* 2005; 27(1): 289-296. <http://dx.doi.org/10.2989/18142320509504087>.
- Pereira LQ, Strefezzi RDF, Catão-Dias JL, Trindade MM, Figuera RA, Kommers GD, et al. Hepatic hemosiderosis in red-spectacled amazons (*Amazona pretrei*) and correlation with nutritional aspects. *Avian Dis* 2010; 54(4): 1323-1326. <http://dx.doi.org/10.1637/9418-060710-Case.1>. PMID:21313859.
- Petry MV, Fonseca VSS. Effects of human activities in the marine environment on seabirds along the coast of Rio Grande do Sul, Brazil. *Ornitol Neotrop* 2002; 13: 137-142.
- Ponganis PJ, Welch TJ, Welch LS, Stockard TK. Myoglobin production in emperor penguins. *J Exp Biol* 2010; 213(11): 1901-1906. <http://dx.doi.org/10.1242/jeb.042093>. PMID:20472777.

- Ratledge C. Iron metabolism and infection. *Food Nutr Bull* 2007;28(4 Suppl): 515-523. <http://dx.doi.org/10.1177/15648265070284S405>. PMID:18297890.
- Rezende GC, Baldassin P, Gallo H, Silva RJ. Ecological aspects of helminth fauna of Magellanic penguins, *Spheniscus magellanicus* (Aves: Spheniscidae), from the northern coast of the State of São Paulo, Brazil. *Braz J Biol* 2013; 73(1): 61-66. <http://dx.doi.org/10.1590/S1519-69842013000100008>. PMID:23644789.
- Rodrigues SC, Adornes AC, Santos Filho EC, Silva-Filho RP, Colares EP. Surviving probability indicators of landing juvenile Magellanic penguins arriving along the southern Brazilian coast. *Braz Arch Biol Technol* 2010; 53(2): 419-424. <http://dx.doi.org/10.1590/S1516-89132010000200021>.
- Rozental P, Biava C, Spencer H, Zimmerman HJ. Liver morphology and function tests in obesity and during total starvation. *Am J Dig Dis* 1967; 12(2): 198-208. <http://dx.doi.org/10.1007/BF02233443>. PMID:6016689.
- Ruoppolo V, Vanstreels RET, Woehler EJ, Heredia SA, Adornes AC, Silva-Filho RP, et al. Survival and movements of Magellanic penguins rehabilitated from oil fouling along the coast of South America, 2000–2010. *Mar Pollut Bull* 2012; 64(7): 1309-1317. <http://dx.doi.org/10.1016/j.marpolbul.2012.05.009>. PMID:22683106.
- Schwartz E, Tornabeni JA, Boxill GC. The effects of food restriction on hematology, clinical chemistry and pathology in the albino rat. *Toxicol Appl Pharmacol* 1973; 25(4): 515-524. [http://dx.doi.org/10.1016/0041-008X\(73\)90020-3](http://dx.doi.org/10.1016/0041-008X(73)90020-3). PMID:4741773.
- Schwilch R, Grattarola A, Spina F, Jenni L. Protein loss during long-distance migratory flight in passerine birds: adaptation and constraint. *J Exp Biol* 2002; 205(5): 687-695. PMID:11907058.
- Sheppard C, Dierenfeld E. Iron storage disease in birds: speculation on etiology and implications for captive husbandry. *J Avian Med Surg* 2002; 16(3): 192-197. [http://dx.doi.org/10.1647/1082-6742\(2002\)016\[0192:ISDIBS\]2.0.CO;2](http://dx.doi.org/10.1647/1082-6742(2002)016[0192:ISDIBS]2.0.CO;2).
- Silva-Filho RP, Ruoppolo V. Sphenisciformes (Pinguim). In: Cubas ZS, Silva JCR, Catão-Dias JL. *Tratado de Animais Selvagens*. 2nd ed. São Paulo: Roca; 2014. p. 384-416.
- Silveira P, Belo NO, Lacorte GA, Kolesnikovas CK, Vanstreels RE, Steindel M, et al. Parasitological and new molecular-phylogenetic characterization of the malaria parasite *Plasmodium tejerai* in South American penguins. *Parasitol Int* 2013; 62(2): 165-171. <http://dx.doi.org/10.1016/j.parint.2012.12.004>. PMID:23269202.
- Stokes DL, Boersma PD, Casenave JL, García-Borboroglu P. Conservation of migratory Magellanic penguins requires marine zoning. *Biol Conserv* 2014; 170: 151-161. <http://dx.doi.org/10.1016/j.biocon.2013.12.024>.
- Valkiūnas G. *Avian malaria parasites and other haemosporidia*. Boca Raton: CRC Press; 2005.
- Vanstreels RET, Adornes AC, Canabarro PL, Ruoppolo V, Amaku M, Silva-Filho RP, et al. Female-biased mortality of Magellanic Penguins (*Spheniscus magellanicus*) on the wintering grounds. *Emu* 2013; 113(2): 128-134. <http://dx.doi.org/10.1071/MU12060>.
- Vanstreels RET, Braga EM, Catão-Dias JL. Blood parasites of penguins: a critical review. *Parasitology* 2016; 143(8): 931-956. <http://dx.doi.org/10.1017/S0031182016000251>. PMID:27253438.
- Vanstreels RET, Kolesnikovas CK, Sandri S, Silveira P, Belo NO, Ferreira FC Jr, et al. Outbreak of avian malaria associated to multiple species of *Plasmodium* in Magellanic penguins undergoing rehabilitation in Southern Brazil. *PLoS One* 2014; 9(12): e116554. <http://dx.doi.org/10.1371/journal.pone.0116554>. PMID:24736326.
- Vanstreels RET, Parsons NJ. Malária aviária e outros Hemosporídeos aviários. In: Cubas ZS, Silva JCR, Catão-Dias JL. *Tratado de animais selvagens*. 2nd ed. São Paulo: Roca; 2014. p. 1427-1443.
- Vanstreels RET, Silva-Filho RP, Kolesnikovas CKM, Bhering RCC, Ruoppolo V, Epiphanyo S, et al. Epidemiology and pathology of avian malaria in penguins undergoing rehabilitation in Brazil. *Vet Res* 2015; 46(1): 30. <http://dx.doi.org/10.1186/s13567-015-0160-9>. PMID:25888987.
- Wagenmakers AJ, Schepens JT, Veerkamp JH. Effect of starvation and exercise on actual and total activity of the branched-chain 2-oxo acid dehydrogenase complex in rat tissues. *Biochem J* 1984; 223(3): 815-821. <http://dx.doi.org/10.1042/bj2230815>. PMID:6508743.
- Wallace R, Walsh M. Health. In: Penguin Taxon Advisory Group. *Penguin Husbandry Manual*. 3rd ed. Silver Spring: American Zoo and Aquarium Association; 2005. p. 86-111.
- Ward RJ, Iancu TC, Henderson GM, Kirkwood JR, Peters TJ. Hepatic iron overload in birds: analytical and morphological studies. *Avian Pathol* 1988; 17(2): 451-464. <http://dx.doi.org/10.1080/03079458808436463>. PMID:18766702.