

Nº 75

The generalized Gauss map of minimal
surfaces in H^3 and H^4

Célia C.Gões and Plinio A.Q.
Simões

novembro 1984.

THE GENERALIZED GAUSS MAP OF MINIMAL SURFACES IN H^3 AND H^4

Célia C. Gões and Plínio A.Q. Simões

1. Introduction

The object of this paper is to establish conditions for a C^∞ map of a Riemann surface M into Q_{n-2} , the hyperquadric $z^1 z^1 + \dots + z^n z^n = 0$ of $\mathbb{C}P^{n-1}$, to be the generalized Gauss map of a minimal conformal immersion of M into H^3 and H^4 , the hyperbolic space of dimensions three and four respectively. Using the upper half-hyperplane as model for the hyperbolic space and exploiting the conformality between the metrics induced on M , by the euclidean metric and the hyperbolic metric through the immersion, we can adapt the theory developed by Hoffman and Osserman [H-0,2] to obtain the conditions.

2. Basic facts

Let $\langle \cdot, \cdot \rangle$ be the usual euclidean metric on \mathbb{R}^n and let H^n and \mathbb{R}_+^n , $n=3,4$, the set $\{(x,t) \mid x \in \mathbb{R}^{n-1}, t > 0\}$ endowed with the metrics $(\cdot, \cdot)_{(x,t)} = \frac{1}{t^2} \langle \cdot, \cdot \rangle$ and $\langle \cdot, \cdot \rangle$ respectively. Given $(x,t) \in H^n$, let $\hat{x} = (x^1, \dots, x^{n-1}, 0) \in \mathbb{R}^{n-1}$. Thus $a, b \in H^n$ implies that $L_{a,b}(x,t) = \frac{b^n}{a^n} [(x,t) - \hat{a}] + \hat{b}$ is an isometry of H^n such that $L_{a,b}(a) = b$, $(L_{a,b})_*(v) = \frac{b^n}{a^n} v$ for all v in the

tangent space $T_a(H^n)$. Let M be a Riemann surface and $\tilde{\Theta}(p) = (x(p), t(p))$ be a conformal immersion of M into H^n . If $q = (0, \dots, 0, 1) \in H^n$, $(L_{\tilde{\Theta}(p), q})_*$ sends $T_{\tilde{\Theta}(p)}(H^n)$ isometrically onto $T_q(H^n)$, which is \mathbb{R}^n endowed with its usual inner product. Let $G_2(\mathbb{R}^n)$ be the grassmannian of the oriented 2-vector subspaces of \mathbb{R}^n . The map $\tilde{G}: M \rightarrow G_2(\mathbb{R}^n)$ defined by $\tilde{G}(p) = (L_{\tilde{\Theta}(p), q})_* \tilde{\Theta}_{*p}(T_p(M))$ is the generalized Gauss map of $\tilde{\Theta}$. It is well known that $G_2(\mathbb{R}^n)$ can be identified with the hyperquadric $Q_{n-2} = \{[z] \in \mathbb{C}P^{n-1} / z = (z^1, \dots, z^n) \text{ and } \sum_{k=1}^n z^k z^k = 0\}$ of the $(n-1)$ -dimensional complex projective space. Such identification will be assumed throughout this paper.

Now let $z = u + iv$ be local isothermal parameters for M and let Θ be a conformal immersion of M into \mathbb{R}_+^n given by $\Theta(p) = \tilde{\Theta}(p)$ ($\forall p \in M$). Then

$$(2.1) \quad \tilde{G}(z) = \left[\frac{\partial \Theta}{\partial u} - i \frac{\partial \Theta}{\partial v} \right] = \left[\frac{\partial \Theta}{\partial z} \right],$$

where $\frac{\partial \Theta}{\partial z} = 1/2 \left(\frac{\partial \Theta^1}{\partial u} - i \frac{\partial \Theta^1}{\partial v}, \dots, \frac{\partial \Theta^n}{\partial u} - i \frac{\partial \Theta^n}{\partial v} \right) \in \mathbb{C}^n$.

If $\phi(z) = (\phi^1(z), \dots, \phi^n(z)) \in \mathbb{C}^n$ is a homogeneous local expression of $\tilde{G}(z)$, there is $\psi: M \rightarrow \mathbb{C} - \{0\}$ such that

$$(2.2) \quad \frac{\partial \Theta}{\partial z} = \psi \phi.$$

Let ds^2 be the riemannian metric induced on M by Θ , $\frac{\partial}{\partial z} = 1/2 \left(\frac{\partial}{\partial u} + i \frac{\partial}{\partial v} \right)$, Δ the Laplace-Beltrami operator of M with respect to ds^2 , $\lambda^2 = \left| \frac{\partial \Theta}{\partial u} \right|^2 = \left| \frac{\partial \Theta}{\partial v} \right|^2$, H the mean curvature vector of Θ , it is well known that

$$(2.3) \quad \Delta\theta = \frac{4}{\lambda^2} \frac{\partial}{\partial z} \frac{\partial}{\partial \bar{z}} \theta = 2H .$$

Indicating by $\langle\langle, \rangle\rangle$ the usual hermitian inner product on \mathbb{C}^n , let

$$(2.4) \quad v = \phi_{\bar{z}} - \eta \phi ,$$

$$\text{where } \phi_{\bar{z}} = \frac{\partial \phi}{\partial \bar{z}} \quad \text{and} \quad \eta = \frac{\langle\langle \phi_{\bar{z}}, \phi \rangle\rangle}{|\phi|^2}$$

Then by (2.2), (2.3) and (2.4) we have

$$(2.5) \quad v = |\phi|^2 \bar{\psi}_H ,$$

$$(2.6) \quad (\log \psi)_{\bar{z}} = -\eta .$$

3. The case n=3

Let $v = (v^1, v^2, v^3)$ be the normal unitary vector field to $\theta(M)$ in \mathbb{R}^3 , $\hat{\mathbb{C}}$ the extended complex plane and $f: M \rightarrow \hat{\mathbb{C}}$ the composition of the classical Gauss map of θ with the stereographic projection with respect to the north pole of the euclidean unitary sphere. Then for $z \in U \subset \mathbb{C}$, we have

$$(3.1) \quad v(z) = \frac{1}{1 + |f(z)|^2} (2 \operatorname{Re} f(z), 2 \operatorname{Im} f(z), |f(z)|^2 - 1).$$

Identifying $\hat{\mathbb{C}}$ biholomorphically with Q_1 , through the correspondence $\xi(\omega) = [1 - \omega^2, i(1 + \omega^2), 2\omega]$, $\xi(\infty) = [-1, i, 0]$,

$\xi^{-1}([z^1, z^2, z^3]) = -\frac{z^1 + iz^2}{z^3}$, where $\omega \in \hat{\mathbb{C}}$ and z^1, z^2, z^3 are homogeneous coordinates on $Q_1 \subset \mathbb{C}P^2$, we obtain.

$$(3.2) \quad [\phi] = [1-f^2, i(1+f^2), 2f].$$

Then from (2.4) we obtain

$$(3.3) \quad \eta = \frac{2\bar{f}f_z}{1+|f|^2},$$

$$(3.4) \quad v = -2f_z v.$$

If h is defined by $H = hv$, (2.5) and (3.4) imply

$$(3.5) \quad h = \frac{-f_z}{\bar{\psi}(1+|f|^2)^2}$$

Therefore

$$(3.6) \quad f_z = 0 \iff h = 0$$

Restricting to those $z \in U$, where h is non-zero, we obtain

$$(3.7) \quad (\log h)_z = \frac{f_z z \bar{z}}{f_z} - 2 \frac{\bar{f} f_z}{1+|f|^2} - \bar{\eta} - (\log \bar{\psi})_z$$

Then (2.6) implies

$$(3.8) \quad (\log h)_z = \frac{f_z z \bar{z}}{f_z} - 2 \frac{\bar{f} f_z}{1+|f|^2}$$

Let \tilde{H} be the mean curvature vector of $\tilde{\Theta} = (x^1, x^2, t)$. Then we have $\tilde{H} = t^2 H + t v^3 v$ [G-S]. Hence the minimality of $\tilde{\Theta}$ implies

$$(3.9) \quad h = -\frac{v^3}{t}$$

Then from (3.1) we obtain

$$(3.10) \quad h = - \frac{|f|^2 - 1}{t(|f|^2 + 1)}$$

Therefore

$$(3.11) \quad f_{\bar{z}} = 0 \iff h = 0 \iff |f|^2 = 1 \iff v^3 = 0$$

From the expressions of h given by (3.10) and (3.5) we obtain

$$(3.12) \quad \bar{\psi} = \frac{tf_{\bar{z}}}{|f|^4 - 1}$$

Since $\bar{\psi}$ is well defined on U , we can extend the ratio $\frac{f_{\bar{z}}}{|f|^4 - 1}$ to U . Then it is straightforward to verify that

$$(3.13) \quad \alpha = \frac{\bar{f}_z}{|f|^4 - 1} dz$$

is a differential 1-form globally defined on M .

Also (3.12) and (2.2) imply

$$(3.14) \quad \theta_z = \frac{t\bar{f}_z}{|f|^4 - 1} (1 - f^2, i(1 + f^2), 2f).$$

From (3.10) and (3.13) we obtain

$$(3.15) \quad (|f|^2 - 1)f_{z\bar{z}} - \frac{2\bar{f}|f|^2}{|f|^2 + 1} f_z f_{\bar{z}} = 0$$

Now it is straightforward to see that the differential 2-form

$$(3.16) \quad \omega = \left\{ (|f|^2 - 1) f_{z\bar{z}} - \frac{2\bar{f}|f|^2}{|f|^2+1} f_z f_{\bar{z}} \right\} |dz|^2$$

is globally defined on M and so its vanishing is a necessary condition for \tilde{G} to be the generalized Gauss map of $\tilde{\Theta}$.

Remark: Through the identification of Q_1 with \hat{C} , the existence of ω and α is independent from $\tilde{\Theta}$. ω is defined on all M and α at all points of M where $|f|^2 \neq 1$.

Then assuming the existence of $\tilde{\Theta}$ we have that α is globally defined and so that $\beta_1 = (1-f^2)\alpha$, $\beta_2 = i(1+f^2)\alpha$ and $\beta_3 = 2f\alpha$ are differential 1-forms globally defined on M .

Let us now state the results of this section.

Theorem 1. - Let M be a connected Riemann surface and let $\tilde{G}: M \rightarrow Q_1$ be smooth. Then the following conditions are necessary for \tilde{G} to be the generalized Gauss map of a minimal conformal immersion of M into H^3 .

- 1) $\omega \equiv 0$,
- 2) α is globally defined,

We then have

$$(3.17) \quad \begin{cases} t = \exp\left(2 \int_M \operatorname{Re} \beta_3\right) \\ x^2 = 2 \int_M t \operatorname{Re} \beta_2 \\ x^1 = 2 \int_M t \operatorname{Re} \beta_1 \end{cases}$$

Proof: - The conditions 1) and 2) were already verified and the formulas in (3.17) are consequence of (3.14).

Theorem 2. - Let M be a connected Riemann surface. If $\tilde{\theta}_1$ and $\tilde{\theta}_2$ are two minimal conformal immersions of M into H^3 having the same generalized Gauss map, then there is an isometry $T: H^3 \rightarrow H^3$ such that $T \circ \tilde{\theta}_1 = \tilde{\theta}_2$.

Proof: - From Hoffman and Osserman [H-0,2] there is $c \in \mathbb{R}$, $c \neq 0$ and a vector d such that $\theta_2 = c\theta_1 + d$. If $\theta_1 = (x^1, x^2, t)$, $\theta_2 = (y^1, y^2, s)$ and $d = (d^1, d^2, d^3)$ we have $s = ct + d^3$. Then $s_z = ct_z$, But from (3.14) we obtain

$$\frac{t_z}{t} = \frac{2f}{|f|^{k-1}} \bar{f}_z = \frac{s_z}{s}. \quad \text{Then } \frac{s_z}{s} = \frac{ct_z}{ct+d^3} = \frac{t_z}{t} \quad \text{and so } d^3 = 0.$$

Now it is straightforward to see that $T(x^1, x^2, t) = c(x^1, x^2, t) + (d^1, d^2, 0)$ is an isometry of H^3 .

Theorem 3. - Let M be a simply connected, non compact and non parabolic Riemann surface and let $\tilde{G}: M \rightarrow Q_1$ be smooth.

If $f: M \rightarrow \hat{\mathbb{C}}$ is the expression of \tilde{G} , obtained through the natural

identification of Q_1 with \hat{C} and $|f|^2 \neq 1$ on M , the differential forms α and ω are globally defined and the condition $\omega \equiv 0$ implies the existence of a minimal conformal immersion $\tilde{\theta}$ of M into H^3 that has \tilde{G} as generalized Gauss map.

Remarks: 1) There is no minimal conformal immersion of Riemann surfaces that are either compact or have parabolic type in H^n . [G-S].

2) The condition $|f|^2 \neq 1$ implies the following

- (a) the image of the classical Gauss map of θ is contained in one of the hemisphere of the unitary euclidean sphere;
- (b) the mean curvature vector of θ never vanishes;
- (c) $f_{\bar{z}}$ never vanishes and so f is nowhere conformal.

Proof: - $|f|^2 \neq 1$ implies the global existence of α and $\omega \equiv 0$ is the integrability condition of the system (3.14). This gives the local existence of $\tilde{\theta}$. The global existence follows from theorem 2 and the simply-connectedness of M , through a standard monodromy argument.

Theorem 4. - Let M be a Riemann surface and $\tilde{\theta}: M \rightarrow H^3$ be a minimal conformal immersion. If $f: M \rightarrow \hat{C}$ is the generalized Gauss map (through the identification of Q_1 with \hat{C}) then the quadratic differential form $\gamma = \frac{f_z \bar{f}_{\bar{z}}}{|f|^4 - 1} dz^2$ is globally defined and holomorphic on M .

Proof: - (3.12) implies that γ is globally defined. The

condition $\omega \equiv 0$ implies that $\left(\frac{f \bar{f}}{|f|^4 - 1}\right) \bar{z} = 0$.

Corollary - If $\tilde{\Theta}$ is non totally geodesic its umbilic points are isolated.

Proof: - $p \in M$ is umbilic for $\tilde{\Theta}$ if, and only if, it is umbilic for Θ , that is if, and only if, $f_z(p) = 0$ [K]. Then p is umbilic for $\tilde{\Theta}$ if p is a zero of γ ; but the zeroes of γ are isolated points.

4. The case $n = 4$

In this case, the hyperquadric Q_2 is biholomorphically identified with $\hat{\mathbb{C}} \times \hat{\mathbb{C}}$ through the correspondence given by $\xi(\omega^1, \omega^2) = [1 + \omega^1 \omega^2, i(1 - \omega^1 \omega^2), \omega^1 - \omega^2, -i(\omega^1 + \omega^2)]$,

$\xi(\infty, \omega^2) = [\omega^2, i\omega^2, 1, -i]$, $\xi(\omega^1, \infty) = [\omega^1, -i\omega^1, -1, -i]$,

$\xi(\infty, \infty) = [1, -i, 0, 0]$, $\xi^{-1}([z^1, z^2, z^3, z^4]) = \left(\frac{z^3 - iz^4}{z^1 - iz^2}, \frac{-z^3 + iz^4}{z^1 - iz^2}\right)$

where $(\omega^1, \omega^2) \in \hat{\mathbb{C}} \times \hat{\mathbb{C}}$ and z^1, z^2, z^3, z^4 are homogeneous coordinates

on $Q_2 \subset \mathbb{C}P^3$. Taking on $\hat{\mathbb{C}} \times \hat{\mathbb{C}}$ the metric $ds^2 = \frac{2|d\omega^1|^2}{(1+|\omega^1|^2)^2} + \frac{2|d\omega^2|^2}{(1+|\omega^2|^2)^2}$

and on Q_2 the metric induced by the Fubini-Study's metric on

$\mathbb{C}P^3$, ξ becomes an isometry. Then the generalized Gauss map

decomposes naturally into a pair of functions $(f_1, f_2): M \rightarrow \hat{\mathbb{C}} \times \hat{\mathbb{C}}$

Moreover, if $z = u + iv \in U \subset \mathbb{C}$ are isothermal parameters on M , we

have $\tilde{G}(z) = [\phi(z)]$ where

$$(4.1) \quad \phi(z) = (1+f_1 f_2, i(1-f_1 f_2), f_1 - f_2, -i(f_1 + f_2))$$

If we set

$$(4.2) \quad A(z) = (f_2 - \bar{f}_1, -i(f_2 + \bar{f}_1), 1 + \bar{f}_1 f_2, -i(1 - \bar{f}_1 f_2))$$

than

$$(4.3) \quad \langle\langle \phi(z), A(z) \rangle\rangle = 0 ;$$

$$(4.4) \quad |\phi|^2 = |A|^2 = 2(1+|f_1|^2)(1+|f_2|^2)$$

Moreover

$$(4.5) \quad e_1 = \frac{\sqrt{2} \operatorname{Re} \phi}{|\phi|}, \quad e_2 = \frac{\sqrt{2} \operatorname{Im} \phi}{|\phi|}, \quad e_3 = \frac{\sqrt{2} \operatorname{Re} \bar{A}}{|A|}, \quad e_4 = \frac{\sqrt{2} \operatorname{Im} \bar{A}}{|A|},$$

is an orthonormal frame of \mathbb{R}^4 , adapted to the immersion θ . Then, from (2.4), (4.1) and (4.5), we obtain

$$(4.6) \quad \eta = \bar{f}_1 F_1 + \bar{f}_2 F_2,$$

$$(4.7) \quad \nu = (F_1 - F_2) \frac{|A|}{\sqrt{2}} e_3 - i(F_1 + F_2) \frac{|A|}{\sqrt{2}} e_4,$$

where $F_1 = \frac{(f_1) \bar{z}}{1+|f_1|^2}$ and $F_2 = \frac{(f_2) \bar{z}}{1+|f_2|^2}$

From (2.5), (4.7) and $H = h_1 e_3 + h_2 e_4$ we obtain

$$(4.8) \quad h_1 = \frac{F_1 - F_2}{2\bar{\psi}\sqrt{(1+|f_1|^2)(1+|f_2|^2)}}, \quad h_2 = \frac{-i(F_1 + F_2)}{2\bar{\psi}\sqrt{(1+|f_1|^2)(1+|f_2|^2)}}$$

Then

$$(4.9) \quad H = 0 \iff (f_1)_{\bar{z}} = (f_2)_{\bar{z}} = 0$$

But $\tilde{H} = t^2 H + t(U_4)^N$, where $U_4 = (0, 0, 0, 1)$ and $()^N$ stands for the orthogonal projection over the normal fibre bundle of $\tilde{\Theta}$. Then assuming that $\tilde{\Theta}$ is minimal we have

$$(4.10) \quad H = -\frac{(U_4)^N}{t}, \quad [G-S].$$

This together with (4.5) imply

$$(4.11) \quad h_1 = \frac{\operatorname{Im}(1 - f_1 \bar{f}_2)}{t\sqrt{(1+|f_1|^2)(1+|f_2|^2)}}, \quad h_2 = \frac{-\operatorname{Re}(1 - f_1 \bar{f}_2)}{t\sqrt{(1+|f_1|^2)(1+|f_2|^2)}}$$

Therefore

$$(4.12) \quad (f_1)_{\bar{z}} = (f_2)_{\bar{z}} = 0 \iff H = 0 \iff 1 - f_1 \bar{f}_2 = 0$$

From the expressions of h_1 and h_2 given by (4.11) and (4.8) we obtain

$$(4.13) \quad \bar{\psi} = \frac{itF_1}{1 - f_1 \bar{f}_2} = \frac{it(f_1)_{\bar{z}}}{(1+|f_1|^2)(1 - f_1 \bar{f}_2)};$$

$$(4.14) \quad F_1(1 - \bar{f}_1 f_2) = F_2(1 - f_1 \bar{f}_2) .$$

Since ψ is defined at all points of U , the ratios $\frac{(f_1)_z}{1 - f_1 \bar{f}_2}$ and $\frac{(f_2)_z}{1 - \bar{f}_1 f_2}$ may be extended to all U .

Then, the differential 1-forms

$$(4.15) \quad \alpha_1 = \frac{(\bar{f}_1)_z}{(1 - \bar{f}_1 f_2)(1 + |f_1|^2)} dz, \quad \alpha_2 = \frac{(\bar{f}_2)_z}{(1 - f_1 \bar{f}_2)(1 + |f_2|^2)} dz,$$

are globally defined on M .

From (2.2), (4.13) and (4.14) we obtain

$$(4.16) \quad \theta_z = \frac{-it}{1 - \bar{f}_1 f_2} \frac{(\bar{f}_1)_z}{1 - |f_1|^2} (1 + f_1 f_2, i(1 - f_1 f_2), f_1 - f_2, -i(f_1 + f_2)),$$

$$(4.17) \quad \begin{cases} (f_1 \bar{f}_2 - 1)(f_1)_{z\bar{z}} - \frac{2|f_1|^2 \bar{f}_2}{1 + |f_1|^2} (f_1)_z (f_1)_{\bar{z}} + \frac{\bar{f}_1 - \bar{f}_2}{1 + |f_1|^2} (f_1)_z (f_1)_{\bar{z}} = 0 \\ (\bar{f}_1 f_2 - 1)(f_2)_{z\bar{z}} - \frac{2|f_2|^2 \bar{f}_1}{1 + |f_2|^2} (f_2)_z (f_2)_{\bar{z}} + \frac{\bar{f}_2 - \bar{f}_1}{1 + |f_2|^2} (f_2)_z (f_2)_{\bar{z}} = 0 \end{cases}$$

If we set

$$(4.18) \quad \begin{cases} \omega_1 = [(f_1 \bar{f}_2 - 1)(f_1)_{z\bar{z}} - \frac{2|f_1|^2 \bar{f}_2}{1 + |f_1|^2} (f_1)_z (f_1)_{\bar{z}} + \frac{\bar{f}_1 - \bar{f}_2}{1 + |f_1|^2} (f_1)_z (f_1)_{\bar{z}}] |dz|^2 \\ \omega_2 = [(\bar{f}_1 f_2 - 1)(f_2)_{z\bar{z}} - \frac{2|f_2|^2 \bar{f}_1}{1 + |f_2|^2} (f_2)_z (f_2)_{\bar{z}} + \frac{\bar{f}_2 - \bar{f}_1}{1 + |f_2|^2} (f_2)_z (f_2)_{\bar{z}}] |dz|^2 \end{cases}$$

we have two differential 2-forms globally defined on M.

The existence of the minimal conformal immersion $\tilde{\Theta}$ having \tilde{G} as its generalized Gauss map implies that $\alpha_1, \alpha_2, \omega_1$ and ω_2 are globally defined and $\alpha_1 \equiv \alpha_2$ and $\omega_1 \equiv \omega_2 \equiv 0$. Then the differential 1-forms $\beta_1 = -i(1+f_1f_2)\alpha_1$, $\beta_2 = (1-f_1f_2)\alpha_1$, $\beta_3 = -i(f_1-f_2)\alpha_1$ and $\beta_4 = -(f_1+f_2)\alpha_1$, are globally defined on M.

Let us now state the results of this section.

Theorem 5. - Let M be a connected Riemann surface and let

$\tilde{G}: M \rightarrow Q_2$ be smooth. Then the following conditions are necessary for the existence of a minimal conformal immersion

$\tilde{\Theta} = (x^1, x^2, x^3, t)$ of M into H^4 :

- 1) α_1 and α_2 are globally defined and $\alpha_1 \equiv \alpha_2$,
- 2) $\omega_1 \equiv \omega_2 \equiv 0$.

Moreover, if these conditions are satisfied, we have

$$(4.19) \quad \left\{ \begin{array}{l} t = \exp\left[2 \int_M \operatorname{Re} \beta_4\right] \\ x^3 = 2 \int_M t \operatorname{Re} \beta_3 \\ x^2 = 2 \int_M t \operatorname{Re} \beta_2 \\ x^1 = 2 \int_M t \operatorname{Re} \beta_1 \end{array} \right.$$

Proof: - It remains only to prove the formulas in (4.19), but they follow from (4.16).

Theorem 6. - Let M be a Riemann surface and let $\tilde{\theta}_1, \tilde{\theta}_2$ be minimal conformal immersions of M into H^4 . If $\tilde{\theta}_1$ and $\tilde{\theta}_2$ have the same generalized Gauss map, then there is an isometry $T: H^4 \rightarrow H^4$ such that $T \circ \tilde{\theta}_1 = \tilde{\theta}_2$.

Proof: - It is similar to that one of theorem 2.

Theorem 7. - Let M be simply connected, non compact and non parabolic Riemann surface and let $\tilde{G}: M \rightarrow Q_2$ be smooth.

If $f_1, f_2: M \rightarrow \hat{C}$ are the components of \tilde{G} obtained through the natural identification of Q_2 with $\hat{C} \times \hat{C}$, and $1 - f_1 \bar{f}_2 \neq 0$, the differential forms $\alpha_1, \alpha_2, \omega_1$ and ω_2 are globally defined and the conditions $\alpha_1 \equiv \alpha_2$ and $\omega_1 \equiv \omega_2 \equiv 0$ imply the existence of a minimal conformal immersion $\tilde{\theta}$ of M into H^4 that has \tilde{G} as its generalized Gauss map.

Remark: - The hypothesis imply that the mean curvature vector of $\tilde{\theta}$ never vanishes and that $(f_1)_{\bar{z}}$ and $(f_2)_{\bar{z}}$ never vanish

Proof: $1 - f_1 \bar{f}_2 \neq 0$ implies the global existence of α_1 and α_2 . Besides, the conditions $\alpha_1 \equiv \alpha_2$, $\omega_1 \equiv \omega_2 \equiv 0$ are the integrability conditions of (4.16). Therefore $\tilde{\theta}$ exists locally. The global existence of $\tilde{\theta}$ follows from theorem 6 by a straightforward monodromy argument.

Theorem 8. - Let M be a connected Riemann surface, $\tilde{\theta}$ be a minimal conformal immersion of M into H^4 and $f_1, f_2: M \rightarrow \hat{C}$ the components of its generalized Gauss map, obtained through the natural identification

of Q_2 with $\hat{\mathbb{C}} \times \hat{\mathbb{C}}$. Then the quadratic differential form

$$\gamma = \frac{(f_1)_z (f_2)_z (\bar{f}_1)_z (\bar{f}_2)_z}{(1 - \bar{f}_1 f_2)(1 - f_1 \bar{f}_2)(1 + |f_1|^2)(1 + |f_2|^2)} dz^2$$

is globally defined and holomorphic on M .

Proof: - The global existence follows from (4.13) and the conditions $\alpha_1 \equiv \alpha_2$ and $\omega_1 \equiv \omega_2 \equiv 0$ imply that γ is holomorphic.

Corollary - If $\tilde{\theta}$ is non totally geodesic the umbilic points and the umbilic directions of $\tilde{\theta}$ are isolated.

5. Final remarks

1. If in (3.1) we consider the stereographic projection relative to the south pole of the unitary euclidean sphere, the identification of Q_1 with $\hat{\mathbb{C}}$ is made through the correspondence

$$\xi(\omega) = [\omega^2 - 1, i(\omega^2 + 1), 2\omega], \quad \xi(\infty) = [1, i, 0],$$

$$\xi^{-1}([z^1, z^2, z^3]) = \frac{z^1 - iz^2}{z^3}, \quad \text{where } \omega \in \hat{\mathbb{C}} \text{ and } z^1, z^2, z^3 \text{ are}$$

homogeneous coordinates on $Q_1 \subset \mathbb{C}P^2$. Then instead of (3.14) we have

$$\theta_z = \frac{t\bar{f}_z}{|f|^4 - 1} (f^2 - 1, i(f^2 + 1), 2f)$$

and the conclusions are similar.

2. If we set $f_1 \equiv f_2 \equiv if$, on the case $n=4$, we recapture the case $n=3$ on the totally geodesic submanifold

$$x^3 = 0 \text{ of } H^4.$$

References

- [G-S] C. Gões and P. Simões. Some remarks on minimal immersions in the hyperbolic space. (preprint).
- [H-0,1] D.A. Hoffman and R. Osserman. The Geometry of the generalized Gauss map. AMS-Memoirs 236(1980).
- [H-0,2] _____ The Gauss map of surfaces in \mathbb{R}^n . J. of. Diff. Geometry 18 (1983) 733-794.
- [K] K. Kenmotsu. Weierstrass formula for surfaces of prescribed mean curvature. Math. Ann. 245 (1979) 89-99

Instituto de Matemática e Estatística
 Universidade de São Paulo
 Cidade Universitária "Armando Salles de Oliveira"
 Caixa Postal 20570 - CEP 05508
 São Paulo - Brasil

TRABALHOS

DO

DEPARTAMENTO DE MATEMATICA

TITULOS PUBLICADOS

- 8001 - PLETCH, A. Local freeness of profinite groups. São Paulo, IME-USP, 1980. 10p.
- 8002 - PLETCH, A. Strong completeness in profinite groups. São Paulo, IME-USP, 1980. 8p.
- 8003 - CARNIELLI, W.A. & ALCANTARA, L.P. de. Transfinite induction on ordinal configurations. São Paulo, IME-USP, 1980. 22p.
- 8004 - JONES RODRIGUES, A.R. Integral representations of cyclic p-groups. São Paulo, IME-USP, 1980. 13p.
- 8005 - CORRADA, M. & ALCANTARA, L.P. de. Notes on many-sorted systems. São Paulo, IME-USP, 1980. /25/p.
- 8006 - POLCINO MILIES, F.C. & SEHGAL, S.K. FC-elements in a group ring. São Paulo, IME-USP, 1980. /10/p.
- 8007 - CHEN, C.C. On the Ricci condition and minimal surfaces with constantly curved Gauss map. São Paulo, IME-USP, 1980. 10p.
- 8008 - CHEN, C.C. Total curvature and topological structure of complete minimal surfaces. São Paulo, IME-USP, 1980. 21p.
- 8009 - CHEN, C.C. On the image of the generalized Gauss map of a complete minimal surface in R^4 . São Paulo, IME-USP, 1980. 8p.
- 8110 - JONES RODRIGUES, A.R. Units of ZCp^n . São Paulo, IME-USP, 1981. 7p.
- 8111 - KOTAS, J. & COSTA, N.C.A. da. Problems of modal and discussive logics. São Paulo, IME-USP, 1981. 35p.

- .8112 - BRITO, F.B. & GONÇALVES, D.L. Algebras não associativas, sistemas diferenciais polinomiais homogêneos e classes características. São Paulo, IME-USP, 1981. 7p.
- .8113 - POLCINO MILIES, F.C. Group rings whose torsion units form a subgroup II. São Paulo, IME-USP, 1981. 1v. (não paginado)
- .8114 - CHEN, C.C. An elementary proof of Calabi's theorems on holomorphic curves. São Paulo, IME-USP, 1981. 5p.
- .8115 - COSTA, N.C.A. da & ALVES, E.H. Relations between paraconsistent logic and many-valued logic. São Paulo, IME-USP, 1981. 8p.
- .8116 - CASTILLA, M.S.A.C. On Przymusiński's theorem. São Paulo, IME-USP, 1981. 6p.
- .8117 - CHEN, C.C. & GOES, C.C. Degenerate minimal surfaces in R^4 . São Paulo, IME-USP, 1981. 21p.
- .8118 - CASTILLA, M.S.A.C. Imagens inversas de algumas aplicações fechadas. São Paulo, IME-USP, 1981. 11p.
- .8119 - ARAGONA VALEJO, A.J. & EXEL FILHO, R. An infinite dimensional version of Hartogs' extension theorem. São Paulo, IME-USP, 1981. 9p.
- .8120 - GONÇALVES, J.Z. Groups rings with solvable unit groups. São Paulo, IME-USP, 1981. 15p.
- .8121 - CARNIELLI, W.A. & ALCANTARA, L.P. de. Paraconsistent algebras. São Paulo, IME-USP, 1981. 16p.
- .8122 - GONÇALVES, D.L. Nilpotent actions. São Paulo, IME-USP, 1981. 10p.
- .8123 - COELHO, S.P. Group rings with units of bounded exponent over the center. São Paulo, IME-USP, 1981. 25p.

- 8124 - PARMENTER, M.M. & POLCINO MILIES, F.C. A note on isomorphic group rings. São Paulo, IME-USP, 1981. 4p.
- 8125 - MERKLEN GOLDSCHMIDT, H.A. Hereditary algebras with maximum spectra are of finite type. São Paulo, IME-USP, 1981. 10p.
- 8126 - POLCINO MILIES, F.C. Units of group rings: a short survey São Paulo, IME-USP, 1981. 32p.
- 8127 - CHEN, C.C. & GACKSTATTER, F. Elliptic and hyperelliptic functions and complete minimal surfaces with handles. São Paulo, IME-USP, 1981. 14p.
- 8128 - POLCINO MILIES, F.C. A glance at the early history of group rings. São Paulo, IME-USP, 1981. 22p.
- 8129 - FERRER SANTOS, W.R. Reductive actions of algebraic groups on affine varieties. São Paulo, IME-USP, 1981. 52p.
- 8130 - COSTA, N.C.A. da. The philosophical import of paraconsistent logic. São Paulo, IME-USP, 1981. 26p.
- 8131 - GONÇALVES, D.L. Generalized classes of groups, spaces c -nilpotent and "the Hurewicz theorem". São Paulo, IME-USP, 1981. 30p.
- 8132 - COSTA, N.C.A. da & MORTENSEN, Chris. Notes on the theory of variable binding term operators. São Paulo, IME-USP, 1981. 18p.
- 8133 - MERKLEN GOLDSCHMIDT, H.A. Homogenous λ -hereditary algebras with maximum spectra. São Paulo, IME-USP, 1981. 32p.
- 8134 - PERESI, L.A. A note on semiprime generalized alternative algebras. São Paulo, IME-USP, 1981. 10p.
- 8135 - MIRAGLIA NETO, F. On the preservation of elementary equivalence and embedding by filtered powers and structures of stable continuous functions. São Paulo, IME-USP, 1981. 9p.

- 8136 - FIGUEIREDO, G.V.R. Catastrophe theory: some global theory a full proof. São Paulo, IME-USP, 1981. 91p.
- 8237 - COSTA, R.C.F. On the derivations of gametic algebras. São Paulo, IME-USP, 1982. 17p.
- 8238 - FIGUEIREDO, G.V.R. de. A shorter proof of the Thom-Zeeman global theorem for catastrophes of cod ≤ 5 . São Paulo, IME-USP, 1982. 7p.
- 8239 - VELOSO, J.M.M. Lie equations and Lie algebras: the intransitive case. São Paulo, IME-USP, 1982. 97p.
- 8240 - GOES, C.C. Some results about minimal immersions having flat normal bundle. São Paulo, IME-USP, 1982. 37p.
- 8241 - FERRER SANTOS, W.R. Cohomology of comodules II. São Paulo, IME-USP, 1982. 15p.
- 8242 - SOUZA, V.H.G. Classification of closed sets and diffeos of one-dimensional manifolds. São Paulo, IME-USP, 1982. 15p.
- 8243 - GOES, C.C. The stability of minimal cones of codimension greater than one in R^n . São Paulo, IME-USP, 1982. 27p.
- 8244 - PERESI, L.A. On automorphisms of gametic algebras. São Paulo, IME-USP, 1982. 27p.
- 8245 - POLCINO MILIES, F.C. & SEHGAL, S.K. Torsion units in integral group rings of metacyclic groups. São Paulo, IME-USP, 1982. 18p.
- 8246 - GONÇALVES, J.Z. Free subgroups of units in group rings. São Paulo, IME-USP, 1982. 8p.
- 8247 - VELOSO, J.M.M. New classes of intransitive simple Lie pseudo-groups. São Paulo, IME-USP, 1982. 8p.

- 8248 - CHEN, C.C. The generalized curvature ellipses and minimal surfaces. São Paulo, IME-USP, 1982. 10p.
- 8249 - COSTA, R.C.F. On the derivation algebra of zygotic algebras for polyploidy with multiple alleles. São Paulo, IME-USP, 1982. 24p.
- 8350 - GONÇALVES, J.Z. Free subgroups in the group of units of group rings over algebraic integers. São Paulo, IME-USP, 1983. 8p.
- 8351 - MANDEL, A. & GONÇALVES, J.Z. Free k-triples in linear groups. São Paulo, IME-USP, 1983. 7p.
- 8352 - BRITO, F.G.B. A remark on closed minimal hypersurfaces of S^4 with second fundamental form of constant length. São Paulo, IME-USP, 1983. 12p.
- 8353 - KIIHL, J.C.S. U-structures and sphere bundles. São Paulo, IME-USP, 1983. 8p.
- 8354 - COSTA, R.C.F. On genetic algebras with prescribed derivations. São Paulo, IME-USP, 1983. 23p.
- 8355 - SALVITTI, R. Integrabilidade das distribuições dadas por subálgebras de Lie de codimensão finita no $gh(n,C)$. São Paulo, IME-USP, 1983. 4p.
- 8356 - MANDEL, A. & GONÇALVES, J.Z. Construction of open sets of free k-Tuples of matrices. São Paulo, IME-USP, 1983. 18p.
- 8357 - BRITO, F.G.B. A remark on minimal foliations of codimension two. São Paulo, IME-USP, 1983. 24p.
- 8358 - GONÇALVES, J.Z. Free groups in subnormal subgroups and the residual nilpotence of the group of units of group rings. São Paulo, IME-USP, 1983. 9p.

- 8359 - BELOQUI, J.A. Modulus of stability for vector fields on 3-manifolds. São Paulo, IME-USP, 1983. 40p.
- 8360 - GONÇALVES, J.Z. Some groups not subnormal in the group of units of its integral group ring. São Paulo, IME-USP, 1983. 8p.
- 8361 - GOES, C.C. & SIMÕES, P.A.Q. Imersões mínimas nos espaços hiperbólicos. São Paulo, IME-USP, 1983. 15p.
- 8462 - GIAM BRUNO, A.; MISSO, P. & POLCINO MILIES, F.C. Derivations with invertible values in rings with involution. São Paulo, IME-USP, 1984. 12p.
- 8463 - FERRER SANTOS, W.R. A note on affine quotients. São Paulo, IME-USP, 1984. 6p.
- 8464 - GONÇALVES, J.Z. Free-subgroups and the residual nilpotence of the group of units of modular and p-adic group rings. São Paulo, IME-USP, 1984. 12p.
- 8465 - GONÇALVES, D.L. Fixed points of S^1 -fibrations. São Paulo, IME-USP, 1984. 18p.
- 8466 - RODRIGUES, A.A.M. Contact and equivalence of submanifolds of homogenous spaces. São Paulo, IME-USP, 1984. 15p.
- 8467 - LOURENÇO, M.L. A projective limit representation of (DFC) - spaces with the approximation property. São Paulo, IME-USP, 1984. 20p.
- 8468 - FORNARI, S. Total absolute curvature of surfaces with boundary. São Paulo, IME-USP, 1984. 25p.
- 8469 - BRITO, F.G.B. & WALCZAK, P.G. Totally geodesic foliations with integral normal bundles. São Paulo, IME-USP, 1984. 6p.
- 8470 - LANGEVIN, R. & POSSANI, C. Quase-folheações e integrais de curvatura no plano. São Paulo, IME-USP, 1984. 26p.

- 8471 - OLIVEIRA, M.E.G.G. de Non-orientable minimal surfaces in \mathbb{R}^N . São Paulo, IME-USP, 1984. 41p.
- 8472 - PERESI, L.A. On baric algebras with prescribed automorphisms. São Paulo, IME-USP, 1984. 42p.
- 8473 - MIRAGLIA NETO, F. & ROCHA FILHO, G.C. The measurability of Riemann integrable functions with values in Banach spaces and applications. São Paulo, IME-USP, 1984. 27p.
- 8474 - MERKLEN GOLDSCHMIDT, H.A. Artin algebras wich are equivalent to a hereditary algebra modulo preprojectives. São Paulo, IME-USP, 1984. 38p.