

RESEARCH

Open Access



Malaria mortality in Brazil: an age-period cohort study

Lucas Casagrande Passoni Lopes^{1*}

Abstract

Background Malaria is a parasitic disease caused by protozoa of the genus *Plasmodium*, transmitted through the bites of *Anopheles* mosquitoes. Despite significant progress in malaria control, the disease remains a persistent public health challenge, particularly in specific Brazilian regions where environmental and socioeconomic factors contribute to its transmission. Understanding mortality trends across different age groups, periods, births cohorts, and regions is essential for developing targeted intervention strategies and optimizing resource allocation. This study aimed to analyse malaria mortality trends in Brazil, focusing on regional differences using an age-period cohort (APC) model.

Methods This ecological study analysed malaria mortality data in Brazil from 1980 to 2024, sourced from DATASUS. Population estimates by sex were retrieved from the Instituto Brasileiro de Geografia e Estatística. Mortality data, including age, year of death, Brazilian macrorregions and administrative Brazilian Amazon legal, as well as the cause of death (ICD-9: 084; ICD-10: B50–B53), were analysed using an APC model. A Poisson distribution was assumed for mortality counts, and analyses were conducted using Holford's method and its adaptations.

Results In the North region, malaria mortality showed an age-related increase, with the highest rates observed in individuals over 80 years old. A significant decline in mortality was observed over the study periods, particularly from 1980 to 1985 to 2020–2024. In the Midwest, the period effect showed fluctuations with an overall decline in recent decades. In the Northeast and Legal Amazon regions, age, period, and cohort effects highlighted clear trends of decreasing mortality over time, particularly for younger cohorts.

Conclusion Malaria mortality is influenced by age, period, cohort, and regions. The regional disparities emphasize the need for localized strategies, considering demographic shifts and epidemiological patterns. By integrating these findings into public health planning, policymakers can enhance malaria surveillance, improve healthcare access in vulnerable regions, and refine control measures to further reduce mortality. The study underscores the necessity of continuous investment in malaria prevention, particularly for older adults in endemic areas, to sustain progress and mitigate resurgence risks.

Keywords Malaria, Age-period cohort, Brazil

Background

Malaria is a parasitic disease caused by protozoa of the genus *Plasmodium*, with four main species affecting humans: *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium malariae*, and *Plasmodium ovale* [1]. Among them, *P. falciparum* and *P. vivax* are the most prevalent worldwide [1]. The disease is transmitted

*Correspondence:

Lucas Casagrande Passoni Lopes
lucaspasconi@usp.br

¹ Faculdade de Medicina de Bauru, Universidade de São Paulo, Bauru, SP 17012-901, Brazil



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

primarily through the bites of infected *Anopheles* mosquitoes [1].

Clinically, malaria presents with a range of symptoms with variable severity [2]. The disease typically begins with fever, chills, headache, and muscle pain [2]. In more severe cases, especially those caused by *P. falciparum*, complications such as anaemia, acute respiratory distress, organ failure, and cerebral malaria can occur, which can be life-threatening if untreated [2]. Recurrent episodes are common, particularly with *P. vivax* infections, which can remain dormant in the liver and cause relapses months or years after the initial infection [2].

Globally, malaria remains a major public health issue, with an estimated 247 million cases and 619,000 deaths reported in 2021 across 84 endemic countries [3]. The disease disproportionately affects low-income regions, primarily in sub-Saharan Africa and parts of Southeast Asia, where healthcare infrastructure is often inadequate [3]. The economic costs of malaria are immense, straining healthcare systems and limiting productivity, particularly in regions heavily dependent on agricultural labour [3]. Malaria's toll extends beyond immediate morbidity and mortality, hampering economic development and perpetuating cycles of poverty [3].

In Brazil, malaria is largely confined to the Amazon Basin, which accounts for 99% of the country's cases [4]. *Plasmodium vivax* is responsible for the majority of infections (approximately 83.7%), with *P. falciparum* making up around 16.3% [4]. The epidemiology of malaria in Brazil is shaped by migration, settlement patterns, and environmental factors unique to the Amazon region [4]. The economic burden of the disease is significant, not only in terms of healthcare costs but also in the loss of productivity, which exacerbates poverty in affected communities [4]. Control efforts in Brazil are complicated by the presence of asymptomatic infections, which challenge malaria elimination strategies [4].

A study using age-period cohort analysis on malaria mortality in Brazil may help to understanding the temporal trends and the influence of demographic and societal changes on mortality risk. Such an approach can help clarify the roles of aging populations, shifts in exposure due to migration, and the effectiveness of control interventions over time. By identifying vulnerable cohorts and periods of increased risk, this study can inform targeted interventions and enhance public health strategies aimed at reducing malaria mortality in endemic regions. Hence, we developed a study with the aim to investigate malaria mortality in Brazil using an age-period cohort model.

Methods

Study characterization

This is an ecologic study that evaluated malaria mortality by an age-period cohort model.

Data source

All data were provided by the Sistema de Informação de Mortalidade (SIM), that is, a Brazilian national system that controls all data regarding death causes in the country. This system feeds the DATASUS, a database that encompasses data from SIM and others Brazilian data health surveillance programs and present them open access to public allowing their access and analysis. The data were categorized into two groups: demographic and clinical. The demographic data refer to the total population of Brazilian macrorregions evaluated along the observed period. Such data were obtained from the census and demographic projections of the Instituto Brasileiro de Geografia e Estatística available on the aforementioned platform.

The clinical data of the patients refer to the age at which they deceased, the year of their death and the cause of death. In this case, we chose patients whose cause of death was identified by the code 084 according to the standardization of the International Classification of Diseases in its ninth edition (ICD9) and codes B50, B51, B52, and B53 according to ICD10 were evaluated. It is worth to mention that this code include all species of protozoa that cause malaria. An inclusion deaths for which the species could not be specified as there is not available data on the subject.

A 45 year period of time, 1980 and 2024, were evaluated since it was the period in DATASUS with all the evaluated data available.

The Brazilian macrorregions of North, Northeast and Central-West of Brazil, as well as the administrative Brazilian Legal Amazon region, were evaluated. It was decided to evaluate these regions because they concentrate practically all the country's cases related to malaria, since they are linked to the Amazon rainforest region. The focus on Legal Amazon derives as the Brazilian government consider it as an endemic region to malaria, presenting a particular environmental, and population dynamics that contributes to malaria occurrence. Furthermore, the South and Southeast regions of Brazil did not present complete data to enable their analysis (Fig. 1).

Data analysis

The data were arranged in Excel® spreadsheets according to Lexis diagram.

Birth cohorts were calculated by subtracting the patient's death year by their age at death, according



Fig. 1 Brazilian macroregions. (Legal Amazon is highlighted by a red edges)

to the classical method. The periods were categorized as: 1980–1984; 1985–1989; 1990–1994; 1995–1999; 2000–2004; 2005–2009; 2010–2014; 2015–2019; and 2020–2024. Age was categorized in 5 years groups as: 0–4, 5–9, 10–14, 15–19, 20–24, 25–29, 30–34, 35–39, 40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, and more than 80 years. This divisions follows the classic model for an APC analysis. In case, it divides time and periods into groups of 5 years starting from the initial time/period, regardless external factors that may influence them, which must be addressed after the analysis [5–7].

It can be assumed that the number of deaths, resulting from a counting process, follows a Poisson distribution. Hence, age-period cohort allows to assign a more satisfactory distribution for the response variable, and consider different forms for the relationship between

the number of deaths and the explanatory variables (age, period, and cohort), which is often non-linear.

The effects were analysed using the age-period-cohort model proposed by Holford and adapted by Clayton and Schifflers and Carstensen [5–7]. The mortality rate (λ_{ijk}) for age (i), period (j) and cohort (k) was modelled according to the formula below. The μ represents the global average mortality rate; α represents the average age effect; β represents the average period effect and γ represents the average cohort effect.

Likelihood ratio tests were used to compare different submodels objecting to assess the effects of age, period, and cohort on mortality rates. Such submodels were adjusted in a conveniently organized sequence in order to provide the tests for the mentioned effects as a comparison between them. Based on these comparisons, made by using the Akaike Information Criterion, the best

model was obtained. The adequacy of the final model fit was verified via deviance statistics. The analysis was performed using the R 4.4.2 software[®] by using the function Epy.

Results

North

The analysis of mortality by age shows a distinct upward trend, with rates per 100,000 individuals increasing steadily from individuals who deacease between 0 and 4 years old (1.657, 95% CI 1.476–1.861) to those whose deacease with more than 80 years old (3.653, 95% CI 3.076–4.338). In terms of mortality risk across periods, a clear decline is observed when compared the period between 1980 and 1985 (Relative Risk–RR: 8.451; 95% confidence Interval–95%CI 7.973–8.958) to period between 2015 and 2019 (RR: 0.143; 95%CI 0.130–0.158).

Finally, cohort analysis reveals a U-shaped pattern, with higher risks in 1950's cohorts (RR: 1.102; 95%CI 1.038–1.169), while younger cohorts demonstrate a decreasing trend, as observed for 2015's cohort (RR: 0.660; 95%CI 0.553–0.788)—Tables 1, 2, 3, and Fig. 2.

Midwest

The age-specific mortality rates show a growing trend between individuals who deacease between 0 and 4 years old (RR: 0.116; 95%CI 0.086–0.157) to those who deacease with more than 80 years (RR: 0.355; 95%CI 0.219–0.575). The period presents fluctuations in its effects. There is a escalating tend between 1980 and 1984 (RR: 5.917; 95%CI 4.995–7.009) till 1985 and 1989 (RR: 15.786; 95%CI 13.746–18.130). Since then, a marked decline was observed till 2015 to 2019 period (RR: 0.029; 95%CI 0.023–0.037). Most of the cohorts evaluated did not

Table 1 - Mortality rate by 100.000 individuals by age

Age (years)	North	Midwest	Northeast	Legal amazon
0–4	1.657 (1.476–1.861)	0.116 (0.086–0.157)	0.003 (0.002–0.005)	1.137 (1.023–1.263)
5–9	0.617 (0.554–0.687)	0.114 (0.088–0.146)	0.006 (0.005–0.009)	0.577 (0.525–0.635)
10–14	0.471 (0.428–0.518)	0.112 (0.089–0.142)	0.012 (0.009–0.015)	0.394 (0.359–0.432)
15–19	0.607 (0.564–0.655)	0.113 (0.091–0.141)	0.022 (0.018–0.027)	0.437 (0.406–0.470)
20–24	0.867 (0.813–0.925)	0.118 (0.098–0.142)	0.039 (0.032–0.046)	0.630 (0.595–0.668)
25–29	0.987 (0.932–1.044)	0.129 (0.111–0.150)	0.053 (0.045–0.061)	0.630 (0.595–0.668)
30–34	0.943 (0.891–0.997)	0.149 (0.128–0.173)	0.050 (0.043–0.059)	0.865 (0.821–0.912)
35–39	0.875 (0.817–0.937)	0.182 (0.155–0.215)	0.043 (0.036–0.052)	0.773 (0.723–0.826)
40–44	0.886 (0.819–0.959)	0.226 (0.185–0.275)	0.038 (0.031–0.048)	0.753 (0.695–0.815)
45–49	0.980 (0.903–1.064)	0.268 (0.211–0.340)	0.038 (0.029–0.049)	0.813 (0.748–0.883)
50–54	1.150 (1.059–1.248)	0.292 (0.228–0.374)	0.040 (0.031–0.052)	0.939 (0.865–1.020)
55–59	1.388 (1.272–1.515)	0.304 (0.238–0.388)	0.045 (0.039–0.057)	1.123 (1.033–1.222)
60–64	1.685 (1.528–1.857)	0.314 (0.241–0.408)	0.050 (0.039–0.064)	1.351 (1.233–1.480)
65–69	2.044 (1.826–2.288)	0.323 (0.239–0.438)	0.056 (0.042–0.074)	1.625 (1.466–1.801)
70–74	2.480 (2.176–2.827)	0.334 (0.234–0.476)	0.063 (0.045–0.086)	1.955 (1.737–2.200)
75–79	3.010 (2.589–3.500)	0.344 (0.227–0.522)	0.070 (0.048–0.102)	2.351 (2.053–2.692)
More than 80	3.653 (3.076–4.338)	0.355 (0.219–0.575)	0.007 (0.051–0.120)	2.828 (2.424–3.300)

Table 2 - Relative risk between periods

Period	North	Midwest	Northeast	Legal amazon
1980–1984	8.451 (7.973–8.958)	5.917 (4.995–7.009)	3.968 (3.327–4.731)	7.351 (6.973–7.749)
1985–1989	9.746 (9.211–10.312)	15.786 (13.746–18.130)	10.405 (9.018–12.006)	10.122 (9.634–10.635)
1990–1994	4.427 (4.260–4.601)	9.860 (8.594–11.313)	4.230 (3.850–4.647)	4.830 (4.661–5.004)
1995–1999	1.957 (1.903–2.012)	3.257 (2.998–3.538)	1.738 (1.621–1.863)	2.076 (2.023–2.131)
2000–2004	–	–	–	–
2005–2009	0.524 (0.507–0.541)	0.307 (0.283–0.334)	0.608 (0.561–0.659)	0.491 (0.477–0.506)
2010–2014	0.274 (0.257–0.292)	0.094 (0.080–0.111)	0.370 (0.315–0.434)	0.241 (0.227–0.256)
2015–2019	0.143 (0.130–0.158)	0.029 (0.023–0.037)	0.225 (0.177–0.286)	0.118 (0.108–0.129)
2020–2024	0.125 (0.115–1.35)	0.037 (0.019–0.045)	0.139 (0.105–0.183)	2.107 (1.820–2.439)

RR (CI955); RR Relative risk, CI95% Confidence interval 95%

Table 3 - Relative risk between cohorts

Cohort	North	Midwest	Northeast	Legal amazon
1900	0.469 (0.390–0.564)	0.612 (0.356–1.053)	0.222 (0.130–0.378)	0.390 (0.327–0.464)
1905	0.515 (0.438–0.607)	0.640 (0.399–1.027)	0.283 (0.178–0.449)	0.442 (0.380–0.515)
1910	0.566 (0.490–0.653)	0.669 (0.446–1.005)	0.360 (0.242–0.536)	0.502 (0.440–0.573)
1915	0.621 (0.549–0.703)	0.700 (0.495–0.989)	0.459 (0.328–0.643)	0.570 (0.509–0.638)
1920	0.682 (0.614–0.757)	0.732 (0.546–0.980)	0.586 (0.440–0.780)	0.647 (0.587–0.713)
1925	0.749 (0.686–0.818)	0.765 (0.595–0.983)	0.747 (0.581–0.959)	0.734 (0.675–0.799)
1930	0.822 (0.763–0.886)	0.800 (0.636–1.006)	0.952 (0.750–1.207)	0.833 (0.772–0.899)
1935	0.903 (0.845–0.965)	0.837 (0.665–1.054)	1.204 (0.940–1.542)	0.943 (0.875–1.017)
1940	0.988 (0.926–1.055)	0.883 (0.698–1.116)	1.455 (1.130–1.873)	1.048 (0.971–1.132)
1945	1.062 (0.995–1.134)	0.944 (0.765–1.166)	1.603 (1.289–1.993)	1.125 (1.045–1.211)
1950	1.102 (1.038–1.169)	1.021 (0.884–1.178)	1.550 (1.342–1.791)	1.145 (1.079–1.215)
1955	1.082 (1.042–1.123)	1.061 (0.994–1.133)	1.313 (1.200–1.436)	1.093 (1.061–1.127)
1960	–	–	–	–
1965	0.907 (0.879–0.936)	0.832 (0.763–0.909)	0.772 (0.696–0.858)	0.900 (0.881–0.920)
1970	0.843 (0.805–0.884)	0.688 (0.575–0.822)	0.675 (0.579–0.787)	0.818 (0.780–0.858)
1975	0.804 (0.750–0.862)	0.635 (0.510–0.791)	0.655 (0.558–0.770)	0.762 (0.706–0.824)
1980	0.778 (0.709–0.854)	0.646 (0.516–0.808)	0.670 (0.565–0.793)	0.735 (0.674–0.802)
1985	0.758 (0.687–0.837)	0.689 (0.548–0.865)	0.690 (0.566–0.842)	0.723 (0.663–0.788)
1990	0.741 (0.671–0.817)	0.740 (0.575–0.953)	0.712 (0.557–0.909)	0.714 (0.653–0.780)
1995	0.724 (0.652–0.803)	0.796 (0.594–1.067)	0.734 (0.543–0.990)	0.705 (0.639–0.776)
2000	0.707 (0.629–0.795)	0.856 (0.606–1.209)	0.756 (0.528–1.084)	0.695 (0.623–0.776)
2005	0.691 (0.604–0.790)	0.920 (0.614–1.379)	0.780 (0.511–1.190)	0.686 (0.606–0.778)
2010	0.675 (0.578–0.788)	0.990 (0.620–1.579)	0.804 (0.494–1.308)	0.677 (0.587–0.781)
2015	0.660 (0.553–0.788)	1.064 (0.624–1.814)	0.828 (0.477–1.440)	0.669 (0.569–0.786)
2020	0.629 (0.573–0.721)	1.029 (0.658–1.349)	0.909 (0.535–1.549)	0.701 (0.549–0.729)

RR (CI95%); RR Relative Risk, CI95% Confidence interval 95%




Fig. 2 - Summary of key findings for each assessed region. (A refers to North region; B refers do Midwest region; C refers to Northeast region; D refers to legal Amazon region)

demonstrate statistical significance of their effect on mortality from malaria—Tables 1, 2, 3, and Fig. 2.

Northeast

Age's effect under mortality demonstrated a increasing pattern, among individuals who deacease between 0 and 4 years old (0.003; 95%CI 0.002–0.005) and those who deacease with more than 80 years (0.007; 95%CI 0.051–0.120). Period's effect showed a growing tendency between 1980 and 1984 (RR: 3.968; 95%CI 3.327–4.731) till 1985 and 1989 (RR: 10.405; 95%CI: 9.018–12.006), followed by a sharp decline over time, reaching a low peak between 2015 and 2019 (RR: 0.225; 95%CI 0.177–0.286). Cohort trends reveal a gradual rise in relative risk, with cohorts born in 1945 reaching a peak (RR: 1.603, 95% CI 1.289–1.993) before gradually declining for later birth cohorts—Tables 1 to 3, and Fig. 2.

Legal amazon

A peak in mortality rate between deceased individuals with 0 to 4 years (1.137; 95%CI 1.023–1.263) with a decreasing pattern till individuals who deceased between 10 and 14 years (RR: 0.394; 95%CI 0.359–0.432) and a growing pattern since then till the rates for deceased individuals with more than 80 years (2.828; 95%CI 2.424–3.300). Period effects shoes a growing tendency between 1980 and 1984 (RR: 7.351; 95%CI: 6.973–7.749) till 1985 and 1989 (RR = 10.122; 95%CI 9.634–10.635) and significantly decreased over time, with the lowest risk observed period between 2015 and 2019 (RR = 0.118; 95% CI 0.108–0.129). Cohort trends highlight an increase in mortality risk for those born between 1900 and 1945, with the highest relative risk observed for the 1950 cohort (RR = 1.145; 95% CI 1.079–1.215). After 1960, the risk declined steadily, reaching minor values for 2015 cohort (RR: 0.669; 95%CI 0.569–0.786)—Tables 1, 2, 3, and Fig. 2.

Supplementary material

In order to provide better information and make the readers'assessment more complete, the link below contains 4 supplementary figures that translate with greater specificity the regional assessment of the effects of age, period, and cohort on malaria mortality: <https://docs.google.com/document/d/1mXyTNvgh-SLFU1N23zk5o6HnQ1eu1GQ9jPX7MaDwEKI/edit?tab=t.0>.

Discussion

A Brazilian study, along with an analysis from the Global Burden of Disease, highlighted that children under four face a disproportionately higher risk of malaria-related mortality compared to other age groups [8, 9]. This vulnerability likely stems from the immaturity of their

immune system, which limits their ability to mount an effective response against the parasite [8, 9]. Additionally, factors such as higher parasite loads, delayed diagnosis, and limited access to prompt treatment further contribute to the increased mortality in this age group [8, 9].

In parallel, as proposed by another studies, were observed a notable burden of malaria mortality in children [10, 11]. However, significant death rates were observed among adults, particularly those over 50 years old [10, 11]. This trend is likely driven by the presence of comorbidities such as cardiovascular disease, diabetes, and impaired renal function, which are more prevalent with aging and can exacerbate malaria's effects [10, 11]. Moreover, the natural decline of immune function with age, known as immunosenescence, diminishes the body's ability to effectively clear infections [10, 11]. This deterioration not only weakens the immune response to malaria but also heightens the risk of severe complications, including multi-organ failure, which is more frequently observed in older populations [10, 11].

As noted in the performed analysis and supported by others Brazilian, Indian and African studies, malaria mortality has been steadily declining over the past decades [8, 12, 13]. In Brazil, a large-scale vector control campaigns—such as insecticide spraying and swamp drainage—were implemented to reduce mosquito breeding sites since 1980 s [8, 14, 15]. Furthermore, the establishment of the National Malaria Control Programme (NMCP) in the 1990 s marked a turning point, introducing widespread early diagnosis, improved treatment protocols, and expanded healthcare access in remote areas [8, 14]. The introduction of artemisinin-based combination therapy (ACT) in 2006, alongside the expansion of rapid diagnostic tests (RDTs), further accelerated progress by ensuring faster and more effective treatment [8, 15]. Additionally, large-scale distribution of insecticide-treated mosquito nets (ITNs) and indoor residual spraying (IRS) significantly reduced transmission [14, 15]. In the 2010 s, efforts shifted toward addressing the environmental and socioeconomic drivers of malaria, particularly in the Amazon Basin, where deforestation and land-use changes fueled outbreaks [14, 15].

Globally, a similar decline in malaria mortality has been observed, driven by strengthened international cooperation and advancements in disease management [12, 13, 16, 17]. The Roll Back Malaria (RBM) initiative, launched in 1998 by the WHO, UNICEF, UNDP, and the World Bank, provided a coordinated framework for malaria control, prioritizing access to essential tools, such as insecticide-treated nets and antimalarial medications [12, 16]. In high-burden regions, investments in healthcare infrastructure, vaccination

research, and community-based intervention programs have contributed to reducing mortality rates [13, 16]. The widespread implementation of ACT, particularly in sub-Saharan Africa, has dramatically improved survival rates for *Plasmodium falciparum* infections [16, 17]. Additionally, novel approaches, such as genetically modified mosquitoes and malaria vaccines like RTS,S/AS01, represent promising strategies for long-term control [16, 17].

Temporal factors, including shifts in economic activities, have a profound influence on malaria transmission and mortality across birth cohorts [18–20]. Over the past decades, Brazil and many other regions worldwide have undergone significant transformations, such as urbanization, industrialization, and improved healthcare systems [18–20]. These changes have led to a general decline in malaria incidence and mortality, particularly in urban centers where vector control measures, sanitation, and access to medical care have improved [18–20].

However, these advancements have not been equitably distributed. Many rural and remote communities, especially in the Amazon and sub-Saharan Africa regions, continue to experience high malaria burdens due to persistent socioeconomic disparities, limited healthcare infrastructure, and environmental conditions favorable to vector proliferation [18, 19]. While industrialization and economic development have reduced malaria risk in some areas, they have also led to deforestation and migration patterns that create new vulnerabilities to transmission [20]. Globally, similar trends have been observed. Countries that have successfully industrialized and urbanized often exhibit lower malaria mortality rates, while regions with slow economic growth and persistent inequalities continue to struggle with high transmission levels [18, 19].

Interestingly, studies from 2000 s predominantly suggested that older individuals were more susceptible to malaria-related mortality [10, 11]. However, 2020 s research has shifted focus toward younger cohorts, indicating that those born in later decades may now face a heightened risk of severe disease and death [8, 9]. Several hypotheses could explain this trend, including waning population-level immunity, shifts in vector-host interactions, and broader changes in epidemiological patterns [10, 11].

For instance, in recent eras, a shift in climatic patterns has been increasingly documented, with reports of malaria cases and fatalities emerging in subtropical and temperate regions [20, 21]. This expanding geographic distribution of malaria underscores the complex interplay between environmental changes and vector dynamics, raising concerns about the disease's potential

resurgence in areas previously considered at low risk [20–22].

Additionally, the impact of co-infections has gained attention, as diseases such as COVID-19 and H1 N1 influenza have been shown to compromise immune responses, potentially exacerbating malaria severity in previously lower-risk groups [23, 24]. These findings call for further investigation into the evolving landscape of malaria mortality and the multifaceted factors driving its demographic shifts [23, 24].

Comparing across regions, the North and Legal Amazon exhibit the highest malaria mortality rates, whereas the Midwest and Northeast show lower but still significant trends [25, 26]. This is possibly due to the fact that the Northeast and Midwest regions have higher urbanization rates, which correlate with greater healthcare coverage, ensuring timely treatments [25]. At the same time, the Legal Amazon, which largely overlaps with the North, has healthcare services concentrated in major urban centers, such as state capitals [26]. This geographic distribution hinders access to preventive measures and delays treatment for patients requiring intensive care, ultimately contributing to worse prognoses [26].

The period effect highlights nationwide improvements in malaria control over the decades, though the decline has been more pronounced in urbanized and non-Amazonian regions [26, 28]. This likely reflects the impact of public policies and regional health measures implemented by the Brazilian government in recent decades, as previously discussed [27]. Moreover, the more dispersed population distribution in the North and Legal Amazon, along with the unique characteristics of riverine and Indigenous communities in these areas, difficulties the adaptation and implementation of similar policies [27].

The cohort effects suggest that older generations faced higher malaria-related mortality risks, likely due to historically limited healthcare access and less effective control measures [25, 28]. However, other factors may have contributed to this pattern [25]. Shifts in occupational and migratory patterns may also be relevant, as past generations were more frequently engaged in high-exposure activities, such as agriculture and extractives, in endemic regions [28]. Indeed, Brazil observed an extensive populational migration towards Midwest and North along 1960 s to 1980 s, which may increased the local population, which supplements and house conditions may be not adequate to avoid malaria infection [28]. In parallel, the Northwest observed a individual's migration towards Brazilian Southeast region, which reduced its population, and possible, malaria transmission and mortality [27, 28].

This study is limited by the ecological design, which may not capture individual-level risk factors. Incomplete data from the South and Southeast regions of Brazil also restricted a nationwide analysis. Additionally, the analysis relied on public health databases, which may have reporting inaccuracies.

Nonetheless, this study has some valuable strengths that must be recognized. Indeed, after a literature review, this study observed the largest time range, a 45-year approach, regarding malaria mortality. Furthermore, over the years an improvement in notification system have been documented, which improved data quality and reliability. By focusing on regions with the highest malaria burden, it offers relevant findings for public health strategies, as well as evaluating specific effects of age, period, and cohort, it contribute to understanding the dynamics of malaria mortality in Brazil.

Conclusions

The study provides a comprehensive analysis of malaria mortality in Brazil from 1980 to 2024, evaluating the impact of age, period, and birth cohort across different regions. The findings reveal a consistent pattern in age-specific mortality, with higher rates observed in the older population across all regions. Period effects consistently showed a significant reduction in malaria mortality over time across all regions. Cohort analyses highlighted region-specific trends, with earlier birth cohorts (especially those born around the 1940 s and 1950 s) experiencing higher risks, particularly in the Legal Amazon and Northeast. However, a steady decline in risk was observed for more recent cohorts, reflecting the impact of improved malaria control measures on younger generations.

Overall, the results indicate significant progress in malaria control in Brazil, marked by substantial reductions in mortality over time. Despite these advancements, the persistent age-related risk suggests a need for targeted strategies to protect vulnerable age groups, particularly the elderly, to achieve further reductions in malaria-related deaths.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12936-025-05373-y>.

Supplementary material 1

Acknowledgements

To the Malaria Journal editors and reviewers by the support along all manuscript's refinement

Declaration of generative AI in scientific writing

No AI was used to develop this manuscript.

Author contribution

LCPL was the unique author from this research. He was in charge of the conceptualization, data collection, data analysis, discussion of the results obtained and writing of this manuscript.

Funding

Not applicable.

Availability of data and materials

No datasets were generated or analysed during the current study.

Declarations

Competing Interests

The authors declare no competing interests.

Received: 27 November 2024 Accepted: 15 April 2025

Published online: 25 April 2025

References

1. Poespoprodjo JR, Douglas NM, Ansong D, Kho S, Anstey NM. Malaria. *Lancet*. 2023;402:2328–45.
2. Nordling L. Malaria's modelling problem. *Nature*. 2023;618:S34–5.
3. Verma K, Razack SA, Schneider KA, Bharti PK. Microsatellites markers fostering the understanding of malaria parasite biology, epidemiology and population genetics. *Diagn Microbiol Infect Dis*. 2025;111: 116667.
4. Garcia K, Soremekun S, Abrahão AA, Marchesini PB, Drakeley C, Ramalho WM, et al. Is Brazil reaching malaria elimination? A time series analysis of malaria cases from 2011 to 2023. *PLoS Global Public Health*. 2024;4: e0002845.
5. Holford TR. The estimation of age, period and cohort effects for vital rates. *Biometrics*. 1983;39:311–24.
6. Clayton D, Schifflers E. (1987). Models for Temporal Variation in Cancer Rates. II: Age-period-cohort Models. *Stat Med*. 1987;6:469–81.
7. Carstensen B. Age–period–cohort models for the Lexis diagram. *Stat Med*. 2007;26:3018–45.
8. Farias MF, Figueiredo ERL, Silva RNSD, Galhardo DDR, da Silva CL, Moreira EMF, et al. Malaria mortality in Brazil: age-period-cohort effects, sociodemographic factors, and sustainable development indicators. *Trop Med Infect Dis*. 2025;10:41.
9. Cao Y, Wu H, Zhang Y, Wu X, Li J, Chen H, et al. Time trends in malaria incidence from 1992 to 2021 in high-risk regions: an age-period-cohort analysis based on the global burden of disease study 2021. *Int J Infect Dis*. 2025;153: 107770.
10. Mühlberger N, Jelinek T, Behrens RH, Gjørup I, Coulaud JP, Clerinx J, et al. Age as a risk factor for severe manifestations and fatal outcome of falciparum malaria in European patients: observations from TropNetEurop and SIMPID surveillance data. *Clin Infect Dis*. 2003;36:990–5.
11. Dondorp AM, Lee SJ, Faiz MA, Mishra S, Price R, Tjitra E, et al. The relationship between age and the manifestations of and mortality associated with severe malaria. *Clin Infect Dis*. 2008;47:151–7.
12. Chalageri VH, Marinaik SB, Nath SN, Singhal R, Rawat S, Ravikumar K, et al. Malaria control - lessons learned from trends of malaria indices over three decades in Karnataka. *India Malar J*. 2023;22:353.
13. Leal Filho W, May J, May M, Nagy GJ. Climate change and malaria: Some recent trends of malaria incidence rates and average annual temperature in selected sub-Saharan African countries from 2000 to 2018. *Malar J*. 2023;22:248.
14. Melo JO, Padilha MAO, Barbosa RTA, et al. Evaluation of the malaria elimination policy in Brazil: a systematic review and epidemiological analysis study. *Trop Biomed*. 2020;37:513–35.
15. Garcia KKS, Abrahão AA, Oliveira AFM, Henriques KMD, de Pina-Costa A, Siqueira AM, et al. Malaria time series in the extra-Amazon region of Brazil: epidemiological scenario and a two-year prediction model. *Malar J*. 2022;21:157.

16. Mangani C, Mzilahowa T, Cohee L, Kayange M, Ntenda P, Sixpence A, et al. Malawi ICMR Malaria Research: interactions and results influencing health policies and practices. *Am J Trop Med Hyg.* 2022;107(4_Suppl):49–54.
17. Perera R, Wickremasinghe R, Newby G, Caldera A, Fernando D, Mendis K. Malaria control, elimination, and prevention as components of health security: a review. *Am J Trop Med Hyg.* 2022;107:747–53.
18. Awasthi KR, Jancey J, Clements ACA, Sah RK, Koirala MP, Chalise B, et al. Traditional beliefs, practices, and migration: a risk to malaria transmission in rural Nepal. *Int J Environ Res Public Health.* 2022;19:16872.
19. Ngonghala CN, Enright H, Prosper O, Zhao R. Modeling the synergistic interplay between malaria dynamics and economic growth. *Math Biosci.* 2024;372: 109189.
20. Canelas T, Castillo-Salgado C, Baquero OS, Ribeiro H. Environmental and socioeconomic analysis of malaria transmission in the Brazilian Amazon, 2010–2015. *Rev Saude Publica.* 2019;53:49.
21. Chen TT, Kim Y, Ljungqvist FC, Jarsjö J, Hesson JC, Linderholm HW. Climate and malaria: Modeling non-linear dynamics in the nordic countries during the 18th and 19th centuries. *Scand J Public Health.* 2025;53:162–71.
22. Rahmani AA, Susanna D, Febrian T. The relationship between climate change and malaria in South-East Asia: a systematic review of the evidence. *F1000Res.* 2023;11:1555.
23. Mohamed AH, Eltyeb E, Said B, Eltayeb R, Algaissi A, Hober D, et al. COVID-19 and malaria co-infection: a systematic review of clinical outcomes in endemic areas. *PeerJ.* 2024;12: e17160.
24. Küpper-Tetzel CP, Idris R, Kessel J, Schüttfort G, Hoehl S, Kohmer N, et al. Coinfections and antimicrobial treatment in a cohort of falciparum malaria in a non-endemic country: a 10-year experience. *Infection.* 2024;52:461–9.
25. Souza PF, Xavier DR, Suarez Mutis MC, da Mota JC, Peiter PC, de Matos VP, et al. Spatial spread of malaria and economic frontier expansion in the Brazilian Amazon. *PLoS ONE.* 2019;14: e0217615.
26. Laporta GZ, Grillet ME, Rodovalho SR, Massad E, Sallum MAM. Reaching the malaria elimination goal in Brazil: a spatial analysis and time-series study. *Infect Dis Poverty.* 2022;11:39.
27. Ferreira MU, Gamboa D, Torres K, Rodríguez-Ferrucci H, Soto-Calle VE, Pardo K, et al. 2022. Evidence-based malaria control and elimination in the Amazon: input from the International Center of Excellence in Malaria Research Network in Peru and Brazil. *Am J Trop Med Hyg.* 107(4_Suppl):160–7.
28. Gonzalez Daza W, Muylaert RL, Sobral-Souza T, Lemes LV. Malaria risk drivers in the Brazilian Amazon: land use-land cover interactions and biological diversity. *Int J Environ Res Public Health.* 2023;20:6497.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.