

Taylor & Francis  
Taylor & Francis Group

CRANIO®

The Journal of Craniomandibular & Sleep Practice

ISSN: 0886-9634 (Print) 2151-0903 (Online) Journal homepage: [www.tandfonline.com/journals/ycra20](http://www.tandfonline.com/journals/ycra20)

## Associations between temporomandibular disorders and tinnitus – a systematic review

Giancarlo De La Torre Canales, Nikolaos Christidis, Anastasios Grigoriadis, Tilde Strandberg, Veronica Montan, Dyanne Medina Flores, Essam Ahmed Al-Moraissi & Maria Christidis

To cite this article: Giancarlo De La Torre Canales, Nikolaos Christidis, Anastasios Grigoriadis, Tilde Strandberg, Veronica Montan, Dyanne Medina Flores, Essam Ahmed Al-Moraissi & Maria Christidis (2025) Associations between temporomandibular disorders and tinnitus – a systematic review, CRANIO®, 43:6, 969-985, DOI: [10.1080/08869634.2024.2404270](https://doi.org/10.1080/08869634.2024.2404270)

To link to this article: <https://doi.org/10.1080/08869634.2024.2404270>



© 2024 The Author(s). Published with license by Taylor & Francis Group, LLC.



[View supplementary material](#)



Published online: 17 Sep 2024.



[Submit your article to this journal](#)



Article views: 5024



[View related articles](#)



[View Crossmark data](#)



Citing articles: 7 [View citing articles](#)

## Associations between temporomandibular disorders and tinnitus – a systematic review

Giancarlo De La Torre Canales Ph.D.<sup>a,b</sup>, Nikolaos Christidis Ph.D.<sup>a</sup>, Anastasios Grigoriadis Ph.D.<sup>a</sup>, Tilde Strandberg DDS<sup>a</sup>, Veronica Montan DDS<sup>a</sup>, Dyanne Medina Flores MSc<sup>a,c</sup>, Essam Ahmed Al-Moraissi Ph.D.<sup>d</sup> and Maria Christidis Ph.D.<sup>e,f</sup>

<sup>a</sup>Division of Oral Rehabilitation, Department of Dental Medicine, Karolinska Institutet, Huddinge, Sweden; <sup>b</sup>Egas Moniz Center for Interdisciplinary Research (CiiEM); Egas Moniz School of Health & Science, Caparica, Almada, Portugal; <sup>c</sup>Department of Prosthodontics, Bauru School of Dentistry, University of Sao Paulo, Sao Paulo, Brazil; <sup>d</sup>Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Thamar University, Thamar, Yemen; <sup>e</sup>The Institute of Health Sciences, The Swedish Red Cross University, Huddinge, Sweden; <sup>f</sup>Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Huddinge, Sweden

### ABSTRACT

**Objectives:** Temporomandibular disorders (TMD) and tinnitus are highly prevalent conditions, that affects about 10–30% of the adult population and seem to co-exist. The primary objective of this systematic review was to investigate any associations between TMD and tinnitus. The secondary objective was to investigate if the associations differ between painful and non-painful TMDs.

**Methods:** An electronic literature search in five databases was performed, from the inception of the databases until 26th of October 2022. This was to identify clinical trials with prevalence numbers of patients with TMD, with and without tinnitus and vice versa. From 1240 studies, a total number of 32 studies were included in the meta-analysis. A risk of bias analysis was made using the Methodological Evaluation of Observational Research (MORE).

**Results:** Seventeen studies showed low risk of bias, while fifteen studies showed some risk of bias. Among patients with TMD, 57.5% also displayed tinnitus. In contrast, among patients with tinnitus, 92.9% also suffered from TMD. There was a strong association between patients with TMD that also had tinnitus, and patients with tinnitus that also had TMD ( $p$ 's < 0.001). The odds ratio for TMD-patients also having tinnitus was 1.556 ( $p$  < .05), while it for tinnitus-patients also having TMD was 2.859 ( $p$  < .05). Six studies examined the psychological status, and there was a higher degree of psychosocial distress among patients with TMD and TMD/tinnitus.

**Conclusions:** There is a strong significant association between TMD and tinnitus, but further research is needed to unravel the nature of this association and its clinical implication.

### KEYWORDS

Temporomandibular disorders; tinnitus; systematic review; human; association


## Introduction

Temporomandibular disorders (TMD) encompass various conditions affecting the masticatory muscles, temporomandibular joints (TMJ), and related structures [1]. It is the leading non-odontogenic source of orofacial pain [2]. Currently, approximately 5–31% of individuals experience TMD, with an incidence in children reaching 11%. In addition, TMD is more prevalent in women, especially those of childbearing age [3,4].

While the exact cause of TMD remains uncertain, it has been proposed that the psychological profile, state of pain amplification, as well as general health and global symptoms play an important role in the etiology of painful TMDs [5,6]. Thus, TMDs seem to

be biopsychosocial in nature, i.e. caused by several risk factors such as psychosocial, autonomic and genetic factors [7–11]. The biopsychosocial model includes the interactions between biological, psychological, and social perspectives on pain [12]. When it comes to the biological part of TMD genetic factors, physical health, pain modulation, hormonal changes, and sex are included in the model. Further, the included psychological factors on TMD can involve individual beliefs, coping abilities, anxiety, fear, depression, sleep disturbances, and mood changes. Finally, the social factors of the model can include relationships, communication and intimacy, culture, socioeconomic status, as well as school or work environments [9–11,13,14].

**CONTACT** Giancarlo De La Torre Canales  [giancarlo.de.la.torre.canales@ki.se](mailto:giancarlo.de.la.torre.canales@ki.se)  Department of Dental Medicine, Karolinska Institutet, Alfred Nobels Allé 8, Huddinge SE-141-52, Sweden

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/08869634.2024.2404270>

© 2024 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

Clinically, the major signs and symptoms related to TMD are pain in the temporomandibular joint (TMJ) and masticatory muscles, impaired jaw movements, TMJ sounds and otologic symptoms such as hearing loss, vertigo, dizziness, ear pain, and tinnitus [6]. In addition, studies have demonstrated a high frequency of otologic symptoms in TMD patients with tinnitus representing 52.1% among the otologic symptoms [6].

Tinnitus is defined as hearing noises without an external source causing the noise [15]. The exact mechanisms of tinnitus have not yet been determined, although it seems to be generally accepted that tinnitus often exists as a consequence of destruction of outer auditory structures, resulting in a disruption of the intrinsic neural activity in the central auditory pathway [16]. The pooled prevalence estimates of any tinnitus in young and older adults have been reported to be of 13.7% and 23.6%, respectively, which impairs their quality of life [17,18].

Notwithstanding, the clinical potential connections between these conditions remain under-researched, despite existing evidence suggesting their existence as well as co-existence [19,20]. In this direction, studies have identified bi-directional links between tinnitus and TMD, with varying findings related to the severity of TMD and tinnitus [20–22]. A meta-analysis revealed that individuals with TMD faced a 4.45 times higher likelihood of experiencing tinnitus compared to those without TMD, yet only three studies were included in this quantitative assessment [23]. Further, there are some previous systematic reviews, however, there is a need to broaden the search-terms and make them more inclusive since the previous ones only included few studies and did not do risk-assessments [23–25]. Therefore, a review with broader but still specific search-terms, including all diagnostic systems of TMD and tinnitus is necessary to be conducted in order to bring to light what possible cause-relationships exist. When this knowledge increases, it could provide clinicians and researchers points of departure for future research projects but also for choosing individualized treatment approaches. Based on this, the primary objective of this systematic review was to investigate if there are any associations between tinnitus and TMD, and the secondary objective to investigate if these associations differ between painful and non-painful TMDs.

## Materials and methods

### Protocol and research question

This systematic review was registered beforehand in Prospero (the International Prospective Register of

Systematic Reviews), registration number (CRD42022356896). The research question was formed using the PECO framework [26], being an acronym for: *p* = Patients, *E* = Exposure, *C* = Control, *O* = Outcome. The population comprised patients (*P*) suffering from TMD, while the investigated condition/exposure (*E*) was tinnitus. The outcome (*O*) was which associations there are between patients suffering from TMD in general or divided in painful as well as non-painful TMDs (*P*) and patients suffering from tinnitus. The control (*C*) was patients without TMD or tinnitus. The present systematic review followed the Preferred Reporting Items for The PRISMA Extension Statement for Reporting of Systematic Reviews Incorporating Network Meta-Analyses of Health Care Interventions (the PRISMA-P checklist) [27] (Supplemental file 1).

The inclusion criteria were **a)** clinical trials, cross-sectional, case-control studies, and observational studies; **b)** adult individuals (over the age of 18) having TMD; **c)** adult individuals (over the age of 18) having tinnitus; **d)** adult individuals (over the age of 18) having TMD and tinnitus.

The following exclusion criteria were applied: **a)** Studies that cannot be found in other languages other than English or Scandinavian languages (Danish, Norwegian and Swedish); **b)** publications irrelevant to the research question; **c)** editorials, letters, legal cases, interviews, case-series, case reports and reviews.

### Search strategy and selection criteria

The search strategy was developed and performed in MEDLINE (Ovid) in collaboration with the librarians LL and ELS at the University Library at (blinded). The search strategies were developed by LL in collaboration with GDC and NC, then peer-reviewed by another librarian (ELS) before LL performed the final searches. Together with the authors NC, EA, and MC each search concept was identified using the Medical Subject Headings (MeSH-terms) and free text terms. Finally, in order to translate the search into the other databases the Polyglot Search Translator was used [28]. The final electronic search was performed in the databases MEDLINE, EMBASE, CINAHL, the Cochrane Central Registry of Controlled Trials (CENTRAL), and Web of Science from the inception of each database to the 26<sup>th</sup> of October 2022. Also, a de-duplication was performed using the method described by Bramer et al. (2016) [29]. Lastly, one final step was added to compare digital object identifiers (DOI) as well as a search in the reference-lists of the included studies and in several systematic reviews

found in the search was performed. The complete search strategy (for all databases) is available in Supplemental file 2.

### Selection of studies

In order to avoid any risk of biasness in the process of screening the studies, the web-based tool Rayyan was used [30]. This was done independently and in a blinded mode by two of the authors (TS and VM). In cases when there was disagreement regarding eligibility this was resolved by discussion with the author (NC) who served as a judge making the final decision. When all disagreements were resolved the authors (TS and VM) attempted to retrieve the full texts of the included and potentially eligible studies. The studies that were retrieved were then reviewed in full text by the same authors (TS and VM) to determine whether they aligned with the inclusion criteria or not. As before, any disagreement was resolved through a discussion with the author (NC). Finally, additional articles were identified by means of citation search. The full texts of these identified studies were then retrieved and reviewed in the same manner as presented before.

### Analysis of risk of bias

Risk of bias and quality assessment of the included articles were performed with the Methodological Evaluation of Observational Research protocol (MORE) [31] that distinguishes between external and internal validity. While external validity refers to the degree that the outcomes of the study can be applied to the general population, internal validity refers to the degree that the outcomes of the study are accurate for the studied subjects. *External validity* is assessed by six parameters, which are: 1) sampling of subjects; 2) assessment of sampling bias; 3) estimate of sampling bias; 4) exclusion rate from the analysis; 5) address of sampling bias in the analysis; and 6) subject flow. *Internal validity* is assessed by five parameters, and these are 1) source to measure outcomes; 2) definition of outcomes; 3) measurements of outcomes; 4) outcomes in subpopulations; and 5) reporting of outcomes. Thus, the assessment was done on the level of parameters where minor or major flaws could be identified. Two authors (TS and VM) evaluated the risk of bias for each study blinded and independently. In cases where conflict aroused it was resolved by discussion with the author (MC), who served as a judge. The individual quality assessment of each included study is presented in Table 1. Even though “Healthcare-based sampling frame” is considered a major flaw according to the

MORE quality assessment tool, these studies were decided beforehand to be included in this systematic review since it is not possible to completely avoid this in the present area.

### Extraction of data

After the risk of bias and quality assessment, data extraction followed. A data extraction form was designed, developed by the authors (EA, AG, and GDC), and pilot-tested independently on three randomly selected studies by two of the authors (TS and VM) to ensure consistency in extraction. The extracted data included information on the characteristics of the included studies and study participants, such as the authors, study type, diagnosis/criteria that had been used, number of participants, mean age of participants, male-female ratio, prevalence of tinnitus, prevalence of TMD, presence of tinnitus among patients with TMD, and presence of TMD among patients with tinnitus. Any disagreement in the data extracting process was resolved by the author (EA), who served as a judge.

### Statistical analysis

All of the tests were performed in SPSS (v.25 IBM, New York, USA), by the author GTC. Descriptive analyses were used to report frequencies. The Chi-square test was used to compare the prevalence ratios of both the presence of tinnitus among patients with TMD, as well as the presence of TMD among patients with tinnitus. The odds ratio was also determined in the two groups to determine the probability of patients with tinnitus to have TMD and the probability of patients with TMD to have tinnitus. The odds ratio was used as a measure in a random effects model with a 95% confidence interval. The Chi-square test of association was performed to assess the association between TMD and tinnitus. The strength of the associations was also determined using Phi. A significance level of  $p < .05$  was applied when performing all of the tests. The results of the individual studies that differed between painful and non-painful TMD were narratively described.

## Results

### Literature search outcome

The entire literature search from all databases resulted in a total of 1240 articles, but after the removal of duplicates 618 publications remained. The title and abstracts of those articles were then screened which resulted in an exclusion of 533 articles, leaving 85 full

Table 1. Table illustrating the extracted study characteristics of the seventeen included studies.


| Author, year                  | Study type     | Number of participants | Male: Female ratio | Mean age of participants | Diagnosical system for TMD | Type of TMD   | Prevalence of TMD (%)   | Prevalence of tinnitus (%) | Prevalence of tinnitus in patients with TMD (%)      | Presence of TMD in patients with tinnitus (%) |
|-------------------------------|----------------|------------------------|--------------------|--------------------------|----------------------------|---|---|----------------------------|--|---|
| Akhter et al., 2013 [45]      | CSS            | 1930                   | 1:1                | 19                       | Questionnaire              | I: Clicking only<br>II: Pain in TMJ only<br>III: Difficulty in mouth opening only<br>IV: Clicking and pain in TMJ<br>V: Clicking and difficulty in mouth opening<br>VI: Pain in TMJ and difficulty in mouth opening<br>VII: Clicking, pain in TMJ and difficulty in mouth opening | I = 58.7%<br>II = 3.9%<br>III = 3.3%<br>IV = 5.3%<br>V = 8.8%<br>VI = 2.0%<br>VII = 17.8% | Total 7%                   | 6%<br>47.6%<br>5.6%<br>55.2%<br>4.2%<br>9.1%<br>4.1% | Total 39%                                     |
| Algieri et al., 2017 [50]     | CSS            | 200                    | 1:4                | 44                       | Clyde H Wilke              | Not specified   | 100%  | 60.0%                      | 60.0%  | 100%  |
| Bush, 1987 [32]               | CSS            | 105                    | 1:6                | 39                       | Questionnaire              | Not specified   | 100%  | 33.0%                      | 33.0%  | 100%  |
| Bernhardt et al., 2011 [42]   | CSS            | 3300                   | 1:10               | NA                       | Questionnaire              | TMJ pain  | 5.8%  | 9.9%                       | NA   | 13.1%   |
| Burgers et al., 2014 [46]     | Prospective CT | 951                    | 1:1                | 54                       | RDC/TMD                    | Not specified   | 6.8%  | 7.2%                       | 36.6%  | 44.1%   |
| Cebj, 2020 [47]               | CSS            | 288                    | 1:3                | 33                       | DC/TMD                     | (1) disk displacement with reduction<br>(2) disk displacement without reduction   | Total 100%  | 8.7%<br>2.8%               | 75.8%<br>24.2%                                       | 100%  |
| Ciancaglini et al., 1994 [39] | CSS            | 797                    | 1:2                | 37                       | Not specified              | TMJ   | 100%  | 6.4%                       | 6.4%   | 100%  |
| Edvall et al., 2019 [36]      | CSS            | 2482                   | 1:1                | NA                       | Questionnaire              | Not specified   | 19.6%   | 100%                       | 100%   | 19.6%   |
| Fernandes et al., 2013 [44]   | CSS            | 224                    | 1:5                | 38                       | RDC/TMD                    | (1) Painful TMD<br>(2) Nonpainful TMD   | (1) 72.3%<br>(2) 27.7%  | Total 57.6%                | 70.4%<br>24.2%                                       | 88.4%<br>11.6%                                |
| Fernandes et al., 2014 [60]   | CSS            | 261                    | 0:1                | 37                       | RDC/TMD                    | (1) Painful TMD<br>(2) Nonpainful TMD   | (1) 82.8%<br>(2) 17.2%  | Total 62.5%                | 70.3%<br>24.4%                                       | Total 93.3%                                   |
| Henderson et al., 1992 [38]   | CSS            | 21                     | 1:20               | 28                       | Not specified              | Myofascial pain and internal derangement  | 100%  | 85.7%                      | 85.7%  | 100%  |
| Hilgenberg et al., 2012 [43]  | CSS            | 200                    | 1:2                | 39                       | RDC/TMD                    | Not specified   | 85.0%PG, 55%CG  | 100%PG, 0%CG               | 100%PG, 0%CG   | 85%PG, 55%CG                                  |
| Kim et al., 2017 [35]         | CSS            | 11745                  | 1:1                | NA                       | Questionnaire              | Not specified   | NA  | 21.8%                      | NA   | 9.9%  |
| Kong et al., 2021 [37]        | CSS            | 5786                   | 2:1                | NA                       | Questionnaire              | Not specified   | 21.9%   | 24.9%                      | 35.0%  | 31.0%   |
| Kusdra et al., 2018 [51]      | CSS            | 485                    | 1:3                | NA                       | RDC/TMD                    | Not specified   | 100%  | 46.0%                      | 46.0%  | 100%  |
| Kuttilla et al., 2005 [41]    | CSS            | 1720                   | 3:4                | 45                       | Questionnaire              | TMJ pain  | 8.0%  | 15.0%                      | 53.0%  | 20.5%   |

(Continued)

Table 1. (Continued).

| Author, year                 | Study type       | Number of participants | Male: Female ratio | Mean age of participants | Diagnosical system for TMD | Type of TMD   | Prevalence of TMD (%)   | Prevalence of tinnitus (%) | Prevalence of tinnitus in patients with TMD (%)                | Presence of TMD in patients with tinnitus (%) |
|------------------------------|------------------|------------------------|--------------------|--------------------------|----------------------------|---|---|----------------------------|--|---|
| Lam et al., 2001 [40]        | Retrospective CT | 776                    | 3:2                | 39                       | RDC/TMD                    | (1) Myalgia<br>(2) TMJ pain<br>(3) Combination of both  | Total 43.0%   | Total 24.7%                | NA   | Total 64.1%                                   |
| Lee et al., 2016 [33]        | Retrospective CS | 37925                  | 19:1               | NA                       | Not specified              | Not specified   | 100%PG, 0%CG  | 4.8%PG, 1.7%CG             | 4.8%PG, 0%CG   | 100%PG, 0%CG                                  |
| Maciel et al., 2018 [52]     | CSS              | 251                    | 1:3                | NA                       | Questionnaire              | Not specified   | 70.5%   | 15.9%                      | 18%  | 80%   |
| Manfredini et al., 2015 [20] | CSS              | 238                    | 10:3               | 49                       | Not specified              | (1) Myalgia<br>(2) Nonpainful internal derangement<br>(3) Nonpainful TMJ arthrosis  | Total 100%  | Total 30.4%                | 32.8%<br>29.3%<br>33.7%  | Total 100%                                    |
| Michiels et al., 2019 [54]   | CSS              | 6115                   | 1:10               | NA                       | Questionnaire              | Not specified   | 7.0%  | 100%                       | 100%   | 7%  |
| Oliveira et al., 2023 [48]   | Retrospective CT | 57                     | 9:1                | NA                       | Not specified              | TMJ pain  | 100%  | 72%                        | 72%  | 100%  |
| Parker & Chole, 1995 [56]    | CSS              | 1032                   | 1:2                | NA                       | Questionnaire              | (1) TMJ click and TMJ pain<br>(2) TMJ click or TMJ pain   | Total 32.8%   | Total 19.6%                | 54.8%<br>23.7%   | Total 58.9%                                   |
| Pavani et al., 2019 [53]     | CSS              | 230                    | 1:1                | 53                       | Not specified              | Not specified   | 48.5%PG, 33.0%CG  | 100%PG, 0%CG               | NA   | 48.0%PG, 0%CG                                 |
| Peleg et al., 2022 [4]       | CSS              | 51                     | 9:8                | 52                       | DC/TMD                     | TMD   | 27.5%   | 100%                       | 100%   | 27.5%   |
| Riga et al., 2010 [59]       | Prospective CT   | 40                     | 1:9                | 43                       | Questionnaire              | Arthralgia  | 100%  | 70%                        | 70%  | 100%  |
| Ren & Isberg, 1995 [57]      | CSS              | 135                    | 1:2                | NA                       | Questionnaire              | disk displacement with reduction and disk displacement without reduction  | Total 100%  | Total 39.3%                | Total 39.3%  | Total 100%                                    |
| Rubinstein et al., 1990 [55] | CSS              | 102                    | 3:2                | 56                       | Helkimo's index            | TMJ click and TMJ pain  | 82.4%   | 100%                       | 100%   | 82.4%   |
| Saldanha et al., 2012 [43]   | CSS              | 200                    | 1:3                | 37                       | RDC/TMD                    | la: myofascial pain<br>lb: myofascial pain with limited opening<br>lla: disc displacement with reduction<br>llb: disc displacement without reduction with limited opening<br>llc: disc displacement without reduction without limited opening<br>llla: arthralgia<br>lllb: osteoarthritis<br>lllc: osteoarthritis | la = 27.0%<br>lb = 27.5%<br>lla = 36.5%<br>llb = 0.5%<br>llc = 5.0%<br>llla = 38.5%<br>lllc = 1.5%<br>lllc = 1.0% | Total 50%                  | 32.0%<br>39.0%<br>44.3%<br>0%<br>6.2%<br>53.5%<br>3.0%<br>1%   | Total 85%                                     |
| Song et al., 2018 [34]       | CSS              | 17575                  | 9:1                | 43                       | Questionnaire              | Not specified   | 11.7%   | 22.0%                      | NA   | 17%   |
| Tuz et al., 2003 [58]        | Prospective CT   | 250                    | 1:4                | 30                       | Unspecified                | (1) Myofascial pain<br>(2) Internal derangement<br>(3) Painful TMD and internal derangement<br>(4) No TMD   | (1) 11.0%PG, 0%CG<br>(2) 77%PG, 0%CG<br>(3) 12%PG, 0%CG<br>(4) 0%PG, 100%CG                                       | Total 41.6%                | 59.1%PG, 0%CG<br>44.2%PG, 0%CG<br>41.6%PG, 0%CG<br>0%PG, 26%CG | 100%PG, 0%CG                                  |
| Yap et al., 2022 [21]        | CSS              | 200                    | 1:3                | 19                       | DC/TMD                     | Not specified   | 59.5%   | 35.0%                      | 46.0%  | NA  |

Abbreviations: TMD = temporomandibular disorders; CSS = cross sectional study; CS = cohort study; CT = clinical trial; NA = not available; PG = Patient group; CG = Control group; TMD = temporomandibular disorders; TMJ = temporomandibular joint.



\*Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/registers). \*\*If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools.

**Figure 1.** The figure illustrates the PRISMA flow-chart of the database search strategy. \*Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/registers). \*\*If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools. From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi:10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>

texts that were sought for retrieval, of which 72 were successfully retrieved. Additionally, 6 articles were found by citation searching. Out of these 78 full-texts, 35 of them did not meet the inclusion criteria and were excluded in this stage. After the risk of bias/quality assessment an additional 11 full-texts were excluded, which resulted in a total of 32 full-texts included in this systematic review [4,20,21,32–60]. The PRISMA flow-diagram including the process of evaluating full-texts for inclusion is shown in Figure 1.

### Risk of bias and study characteristics

Sixteen of the thirty-two included articles displayed a “Healthcare-based sampling frame” which is considered a major flaw according to the MORE quality assessment. However, as described in the materials and methods section these studies were included in this systematic review. A total of 17 studies showed a low risk of bias (displayed in green) [32–48] while 15 studies showed some risk of bias (displayed in orange) [4,20,21,49–60], as

shown in Table 1. The characteristics of the 32 included studies were compiled into a table containing the type of TMD (painful or non-painful) or non-TMD, mean age of participants, male-female ratio, prevalence of tinnitus, prevalence of TMD, presence of tinnitus among patients with TMD, and presence of TMD among patients with tinnitus, are reported in Table 2.

### Relations and associations between tinnitus and TMD

Thirteen of the included studies had tinnitus as the “main group” and assessed the number of participants with TMDs among the tinnitus population. Regarding the TMD studies, 19 of them assessed the number of participants with tinnitus among the TMD populations.

When it comes to the patient group suffering from tinnitus the vast majority of the patients have a diagnosis of TMD (Table 3). However, when it comes to the patient-group suffering from TMD approximately half

**Table 2.** Table summarizing quality assessment and risk of bias.

| Author, year                  | External validity   | Internal validity  | Comments  | Risk of bias      |
|-------------------------------|---|--|---|-------------------|
| Akhter et al., 2013 [45]      | <i>Minor flaw:</i><br>Sampling of the subjects by the investigators   | <i>Minor flaw:</i><br>Source of measure incidence/<br>prevalence of chronic diseases<br><i>Poor reporting:</i><br>Validation of outcomes<br>measurements not reported        | –   | Low risk of bias  |
| Algieri et al., 2017 [50]     | <i>Minor flaw:</i><br>Sampling of the subjects by the investigators<br>Nongeneral population based sampling method<br>No assessment of sampling bias<br><i>Poor reporting:</i><br>Exclusion rate from the analysis not reported<br>Estimate bias not reported<br>Exclusion rate from analysis not reported<br><i>Major flaw:</i><br>Healthcare-based sampling frame<br><i>Poor reporting:</i><br>Not reported estimate bias.<br>The exclusion rate from the analysis not reported | –  | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Bush, 1987 [32]               | <i>Major flaw:</i><br>Healthcare-based sampling frame<br><i>Poor reporting:</i><br>Not reported estimate bias.<br>The exclusion rate from the analysis not reported   | –  | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Bernhardt et al., 2011 [42]   | <i>Poor reporting:</i><br>Assessment of sampling bias   | –  | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Buergers et al., 2014 [46]    | <i>Major flaw:</i><br>Healthcare-based sampling frame   | –  | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Cebi, 2020 [47]               | <i>Minor flaw:</i><br>Sampling of the subjects by the investigators<br><i>Major flaw:</i><br>Healthcare-based sampling frame<br><i>Minor flaw:</i><br>Nongeneral population based sampling method   | <i>Minor flaw:</i><br>Source of measure incidence/<br>prevalence of chronic diseases<br><i>Minor flaw:</i><br>Source of measure incidence/<br>prevalence of chronic diseases | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Ciancaglini et al., 1994 [39] | <i>Major flaw:</i><br>Healthcare-based sampling frame<br><i>Minor flaw:</i><br>Nongeneral population based sampling method<br>Healthcare-based sampling frame   | <i>Minor flaw:</i><br>Source of measure incidence/<br>prevalence of chronic diseases   | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Edvall et al., 2019 [36]      | <i>Minor flaw:</i><br>Sampling of the subjects by the investigators   | –  | –   | Low risk of bias  |
| Fernandes et al., 2013 [44]   | <i>Minor flaw:</i><br>Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i><br>Healthcare-based sampling frame   | <i>Minor flaw:</i><br>Source of measure incidence/<br>prevalence of chronic diseases   | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Fernandes et al., 2014        | <i>Minor flaw:</i><br>Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Poor reporting:</i><br>Estimate bias not reported<br>Exclusion rate from the analysis not reported<br><i>Major flaw:</i><br>Healthcare-based sampling frame<br><i>Minor flaw:</i><br>Random sampling restricted to geographic area. <i>Major flaw:</i><br>Healthcare-based sampling frame <i>Minor flaw:</i> sampling bias not addressed in analysis     | <i>Minor flaw:</i><br>Source of measure incidence/<br>prevalence of chronic diseases   | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |
| Henderson et al., 1992 [38]   | <i>Minor flaw:</i><br>Random sampling restricted to geographic area. <i>Major flaw:</i><br>Healthcare-based sampling frame <i>Minor flaw:</i> sampling bias not addressed in analysis   | –  | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |

(Continued)

Table 2. (Continued).

| Author, year                 | External validity  | Internal validity  | Comments  | Risk of bias      |
|------------------------------|--|--|---|-------------------|
| Hilgenberg et al., 2012 [43] | <i>Minor flaw:</i> Random sampling restricted to geographic area. <i>Major flaw:</i> Healthcare-based sampling frame <i>Poor reporting:</i> Assessment of sampling bias not reported<br>Response rate in total sample not reported<br>Response rate in subgroups not reported<br>Exclusion rate from the analysis not reported<br>Sampling bias not reported in the analysis<br><i>Minor flaw:</i> Sampling of the subjects by the investigators   | –  | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |
| Kim et al., 2017 [35]        | <i>Minor flaw:</i> Sampling of the subjects by the investigators   | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases  | –   | Low risk of bias  |
| Kong et al., 2021 [37]       | <i>Poor reporting:</i> Exclusion rate from the analysis not reported   | –  | –   | Low risk of bias  |
| Kusdra et al., 2018 [51]     | <i>Poor reporting:</i> Sampling of the subjects by the investigators not reported<br>Exclusion rate from the analysis not reported<br>Address Bias not reported<br><i>Minor flaw:</i> Nongeneral population based sampling method<br><i>Major flaw:</i> Medical records based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area <i>Poor reporting:</i> exclusion rate from the analysis not reported<br>Exclusion rates in subgroups not reported<br><i>Minor flaw:</i> Random sampling restricted to geographic area<br><i>Major flaw:</i> Medical records based sampling frame (they used medical records for dataextraction) | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases<br><i>Poor reporting:</i> Reliability of the estimates not reported  | –   | Some risk of bias |
| Kuttila et al., 2005 [41]    | Random sampling restricted to geographic area <i>Poor reporting:</i> exclusion rate from the analysis not reported   | –  | –   | Low risk of bias  |
| Lam et al., 2001 [40]        | Exclusion rates in subgroups not reported<br><i>Minor flaw:</i> Random sampling restricted to geographic area<br><i>Major flaw:</i> Medical records based sampling frame (they used medical records for dataextraction)  | <i>Minor flaw:</i> Data obtained from medical records  | –   | Low risk of bias  |
| Lee et al., 2016 [33]        | <i>Poor reporting:</i> Aim of study not reported<br><i>Minor flaw:</i> Random sampling restricted to geographic area <i>Major flaw:</i> insurance claims based sampling frame  | <i>Minor flaw:</i> Source of measure incidence/prevalence obtained from administrative database.<br>Frequency can be relevant but not assessed in the study  | –   | Low risk of bias  |
| Maciel et al., 2018 [52]     | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported   | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases: proxy reported<br><i>Poor reporting:</i> Validation of outcomes not reported<br>measurements<br>Reliability of the estimates not reported | –   | Some risk of bias |
| Manfredini et al., 2015 [20] | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Poor reporting:</i> Assessment of sampling bias not reported<br>Estimate bias not reported<br>Address Bias not reported<br><i>Major flaw:</i> Healthcare-based sampling frame  | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases  | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |

(Continued)

Table 2. (Continued).

| Author, year                 | External validity  | Internal validity   | Comments  | Risk of bias      |
|------------------------------|--|---|---|-------------------|
| Michiels et al., 2019 [54]   | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases | –   | Some risk of bias |
| Oliveira et al., 2023 [48]   | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |
| Parker & Chole, 1995 [56]    | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | –   | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |
| Pavaci et al., 2019 [53]     | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | –   | –   | Some risk of bias |
| Peleg et al., 2022 [4]       | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | Source of measure incidence/prevalence of chronic diseases: minor flaw        | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |
| Riga et al., 2010 [59]       | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | –   | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |
| Ren & Isberg, 1995 [57]      | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | –   | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |
| Rubinstein et al., 1990 [55] | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Some risk of bias |
| Saldanha et al., 2012 [43]   | <i>Minor flaw:</i> Sampling of the subjects by the investigators<br>Nongeneral population-based sampling method<br><i>Major flaw:</i> Healthcare-based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area. self selection- based sampling method. <i>Major flaw:</i> Health care-based sampling frame<br><i>Minor flaw:</i> Sampling of the subjects by the investigators<br><i>Poor reporting:</i> Estimate bias not reported<br>Exclusion rate from the analysis not reported  | <i>Minor flaw:</i> Source of measure incidence/prevalence of chronic diseases | Healthcare-based sampling frame is accepted since the research area makes it impossible to exclude such studies | Low risk of bias  |

(Continued)

**Table 2.** (Continued).

| <i>Author, year</i>    | <i>External validity</i>   | <i>Internal validity</i>   | <i>Comments</i> | <i>Risk of bias</i> |
|------------------------|--|--|-----------------|---------------------|
| Song et al., 2018 [34] | <i>Minor flaw:</i><br>Sampling of the subjects by the investigators  | <i>Minor flaw:</i><br>Source of measure incidence/prevalence of chronic diseases | –               | Low risk of bias    |
| Tuz et al., 2003 [58]  | <i>Major flaw:</i> Medical records based sampling frame<br><i>Minor flaw:</i> Random sampling restricted to geographic area<br><i>Poor reporting:</i> Response rate in total sample not reported<br>Response rate in other subgroups not reported<br>Exclusion rate from analysis not reported<br>Exclusion rate from analysis in subgroups not reported<br>Sampling bias not addressed in analysis<br>Number screened not reported<br><i>Minor flaw:</i><br>Sampling of the subjects by the investigators<br>Nongeneral population based sampling method<br>Assessment of sampling bias | <i>Minor flaw:</i><br>Data obtained from medical records                         | –               | Some risk of bias   |
| Yap et al., 2022 [21]  |  |  | –               | Some risk of bias   |

**Table 3.** The prevalence ratios of the presence of tinnitus among patients with temporomandibular disorders as well as the presence of temporomandibular disorders in patients with tinnitus are presented in this table both in number of patients as well as in percentage.

|          |     |               | Tinnitus   |        | Total        |
|----------|-----|---------------|------------|--------|--------------|
|          |     |               | No         | Yes    |              |
| TMD      | No  | cases         | 17160      | 14908  | 32068        |
|          |     | % in TMD      | 53.5%      | 46.5%  | 100.0%       |
|          | Yes | cases         | 2156       | 2915*  | 5071         |
|          |     | % in TMD      | 42.5%      | 57.5%* | 100.0%       |
| Total    |     | cases         | 19316      | 17823  | 37139        |
|          |     | % in TMD      | 52.0%      | 48.0%  | 100.0%       |
|          |     |               | <b>TMD</b> |        | <b>Total</b> |
| TINNITUS | No  | cases         | 2172       | 9994   | 12166        |
|          |     | % in TINNITUS | 17.9%      | 82.1%  | 100.0%       |
|          | Yes | cases         | 134        | 1763*  | 1897         |
|          |     | % in TINNITUS | 7.1%       | 92.9%* | 100.0%       |
| Total    |     | cases         | 2306       | 11757  | 14063        |
|          |     | % in TINNITUS | 16.4%      | 83.6%  | 100.0%       |

\*=  $p < .05$  (Chi-square test).

TMD = temporomandibular disorders.

of the patients also have a diagnosis of tinnitus (Table 1). Further, there is a strong association of having tinnitus if the patient also has a diagnosis of TMD (Chi-square value:  $[X^2(1) = 212.075; p < .001]$ ) as well as the opposite (Chi-square value:  $[x^2(1) = 139.354; p < .001]$ ). This association was strong for both groups (Phi value for TMD in tinnitus is 0.076;  $p < .001$ , and Phi value for tinnitus in TMD is 0.100;  $p < .001$ ). The odds ratio for TMD is 1.556 ( $p < .05$ ) and for tinnitus 2.859 ( $p < .05$ ).

### Narrative description of painful TMD vs non-painful TMD

In this article search 7 articles that clearly differed between painful and non-painful TMD were revealed. In one study by Henderson et al. (1992) there was no significant difference in the prevalence of tinnitus among patients with painful TMD and non-painful TMD [38]. However, in the other six studies there were two-eight times more patients with painful TMD also having tinnitus (range between 65% and 93% of the patients with tinnitus) than those with non-painful TMDs (range between 7% and 35% of the patients with tinnitus) [20,43–45,58,60].

### Psychosocial findings of Tinnitus and TMD

Six of the studies included in the analysis examined the psychological status of individuals with tinnitus and TMD, with a primary focus on depression, assessed through validated tools. Among these, four studies revealed that individuals with both tinnitus (mild or severe) and TMD experienced higher levels of depression compared to those with tinnitus alone, as per their severity [4,36,37,44]. In a study by Hilgerberg et al. (2012), it was found that the presence of tinnitus was

associated with significantly elevated levels of depression when compared to a control group (no tinnitus) [49]. Furthermore, a study identified a positive relationship between the severity of tinnitus, chronic pain, and depression, highlighting a significant correlation between chronic pain intensity and tinnitus [43].

### Discussion

This study aimed to investigate potential associations between the presence of tinnitus and TMD or vice versa, but also the strength of this association. Secondly, it aimed to distinguish between the associations of tinnitus with painful and/or non-painful TMD. Our results revealed a strong bi-directional association between TMD and tinnitus, with the highest prevalence of tinnitus reported in patients diagnosed with painful TMD.

A clinically interesting finding that was revealed was that 92.9% of the patients with tinnitus displayed a TMD diagnosis, which is a significantly higher prevalence when compared to the general population where literature suggests that only about 5–31% has a TMD diagnosis [3,4]. Furthermore, while tinnitus is represented by 12–30% in the general population [61], our meta-analysis showed that 57.5% of the patients with TMD also suffered from tinnitus. This indicates that the presence of TMD is 3 to 15 times higher among tinnitus patients than in a non-TMD population, while the increase of the prevalence of TMD – 2 to 5 times higher – is not that distinct in a healthy population when compared to patients suffering from tinnitus. This finding is in line with a recent cross-sectional study reporting that 67% of tinnitus population presented TMD [62]. In contrast to previous systematic reviews, our study reported a higher bi-directional prevalence of TMD and tinnitus. This could be explained

by different factors. One is because of the higher number of studies included in the analysis that could be a contributing factor only by time. However, when analyzing the previous ones, this was not the fact. Another reason was that the present study used broader and more inclusive search terms than the previous [23–25]. One of the previous reviews did not do any odds-ratio, risk-ratio and chi-square analysis, but only reported the findings [24]. Further, the studies assessing odds-and-risk ratios just included studies using clinically diagnosed TMD, resulting in only five [23] and eight studies [25] for their analysis, respectively. Taken together, the results from this study further contribute to knowledge about the potential cause-relationship, where tinnitus seems rather to be a consequence of TMD instead of TMD being a consequence of tinnitus.

Further, the results from the meta-analysis also indicate that the strong bi-directional association between both conditions points towards the fact that patients presenting tinnitus are more likely to have TMD (odds ratio = 2.85) than TMD patients having tinnitus (odds ratio = 1.55). Even though an exact cause for this association is not yet fully understood, some hypotheses can be raised. TMD and tinnitus may be linked due to the close proximity of the middle ear and TMJ [1,19,43]. This close anatomical and functional relationship between these two structures is mediated by the anterior malleolar ligament (AML), the sphenomandibular ligament (SML), and the discomalleolar ligament (DML). In addition, DML, distinct from AML, is a key player in the mobility of the malleus, a bone in the middle ear, with its structure potentially affecting outcomes in TMJ surgery and conditions leading to anterior displacement of the articular disc. These ligaments can be stretched during TMJ use affecting middle ear structures like the eardrum and malleus [63]. The relationship between the TMJ and the middle ear depends on which ligament is present, the structure of the ligament (whether collagenous or elastic), and the width of the petrotympanic fissure. Also, the overactivity of masticatory muscles could influence the association between tinnitus and TMD, since these muscles which are involved in TMD share the trigeminal nerve with muscles involved in the function of the eustachian tube, which indeed includes the lateral and medial pterygoids muscles [64]. Disruption of the pressure regulation of the auditory tube may be an important factor contributing to the increased risk of developing tinnitus risk [64]. Furthermore, this association could also be explained by the somatosensory theory, in which chronic pain or impaired psychosocial

status like depression or anxiety induce changes in the central nervous system, producing a sensorial hypersensitivity and an altered perception of the auditive stimuli [43,65,66]. A recent review also imposes that this link can be contributed by an increased gut permeability, a disturbed gut microbiome and disturbed metabolism of tryptophan – a well-known biomarker for chronic musculoskeletal pain. This is a substrate for serotonin and melatonin, that both play a key role in microglia activation (e.g. gastrointestinal tract) and regulation of chronic orofacial pain that leads to an increased sensibility to pain and depression, which in turn could affect both TMD and tinnitus [67]. In addition, another study found an association between depression and TMDs as well as tinnitus [4]. Finally, TMJ herniation – even though it is rare – could also be the cause of tinnitus [68].

The present systematic review also investigated if the associations differ between painful and non-painful TMDs among patients with tinnitus. However, due to an insufficient number of studies distinguishing between painful and non-painful TMDs, a meta-analysis was not feasible. Notwithstanding, when the prevalence was compared there was a higher prevalence of tinnitus among patients with painful TMDs than in patients with non-painful TMDs. In addition, one of the included studies reported that the severity of tinnitus was correlated with the severity of chronic pain, which could be explained by the influence that the central nervous system exerts on these two symptoms [43].

In this direction, since the complexity of chronic pains such as TMDs and their multifactorial biopsychosocial etiology are well known [69,70], and that tinnitus could be triggered by psychosocial distress [71], the assessment of the psychosocial status in patients suffering of TMD and/or tinnitus is of main importance. However, it is often neglected. In the present study, only 13 of the included studies assessed the psychosocial status, mainly focusing on depression symptoms and severity; however, only six studies used validated tools for this assessment. These studies found that the single presence of tinnitus is associated with higher levels of depression and that there is a relationship between the presence of chronic pain such as TMDs, tinnitus and depression (43,49). In addition, it has been reported that tinnitus could be associated with periods of stress and anxiety, as a result of enhanced activity in the limbic and autonomic nervous systems [71]. These variables are also usually impaired in TMD patients [72,73], fact that could strengthen the relationship between psychosocial impairment, TMD and tinnitus.

### Study strengths and limitations

One limitation is the diversity in the system used for TMD diagnostics in the included articles. Only a few used a standardized and validated clinical protocol; most used questionnaires or did not even specify how TMD was diagnosed. Based on this, there is an uncertainty about the TMD group, what TMD condition is elaborated, i.e. it could be a painful condition, or just an internal derangement of the temporomandibular joint. However, both types of TMD could be associated with tinnitus, but with different cause-relationships. Therefore, even though there is a strong reported association in the present study, the outcome might have been different if only validated systems would have been used and the different TMDs analyzed separately. As for TMD, the same accounts for tinnitus. Even though patient-history/anamnesis remains the easiest and most cost-effective method for assessing tinnitus, there are instrumental examinations available to assess the presence and severity of tinnitus [74]. Another limitation could be that only adults were included in this review. However, this was chosen since children are growing and therefore growth itself could be a possible confounding factor. Further, the present validated diagnostic criteria for TMD (DC/TMD) [75], as well as the previous one (research diagnostic criteria for TMD; RDC/TMD) [76] are not validated for children which therefore would be a confounding factor itself. Even though, the Axis I of the DC/TMD was adapted for children and recently published in a Delphi study, there is still no validation of this tool [77]. RDC/TMD is validated for adolescents, but not children. A third limitation was that this study could not investigate if there are any gender differences when it comes to frequency of both TMD and tinnitus. This could be relevant since the vast majority of TMD patients are women, and sex itself seems to be one of the important etiological factors [3,4]. Based on this, a deeper understanding on possible sex differences would also contribute to the knowledge of the TMD-tinnitus cause-relationship. However, when it comes to sex differences, present knowledge is inconsistent. For instance, one of the included studies reported that there are no significant differences between men and women according to either TMD or tinnitus [4]. On the other hand, another study reported that the prevalence of tinnitus is more frequent among men than women [61]. Therefore, in future studies a possible gender difference needs to be investigated, since it would have a great clinical implication once the vast majority of the patients with TMD are women [78]. This would also contribute to a deeper understanding of the connection and causality of both conditions.

One of the strengths of this study is the inclusion of a relatively large number of full texts ( $n = 32$ ). Another strength is that most included studies showed a low risk of bias and high quality, which indicates that the outcome of the present study is valid and reliable.

### Clinical implications and generalizability

The results of this study indicate that it is important to identify patients with tinnitus and TMD by health care professionals, since both conditions seem to be comorbid. This could lead to the fact that these patients will receive individualized treatments that affect both conditions. However, the available literature is inconclusive about the effects of treating one condition on the improvement of the other condition, which should be a matter of assessment for future researches [79,80]. This would lead to less patient suffering, less costs for the society and increased quality of life in these two common patient-groups (10–20%) [78,81]. Thus, based on our results, the authors suggest that health care professionals in clinical practice that have patients that suffer from either tinnitus or TMD should also consider assessing the respective condition by for instance the validated 3Q-TMD [82] in patients with tinnitus or the questionnaires FiveQ, THI, THQ etc. in patients with TMD complaining of ear symptoms [74] or the PHQ-4 in both patients, since anxiety and depression could increase both conditions [83]. This screening would then help clinicians to individualize and choose the treatment-approach based on both conditions.

Additionally, as tinnitus could be a red flag of other potentially life-threatening diseases, assessing tinnitus in clinical practice is of main importance, since comorbidities other than TMD like, multiple sclerosis, vestibular schwannoma, meningitis, syphilis, Lyme disease and medications like salicylates, nonsteroidal anti-inflammatory drugs, chemotherapy agents between others are associated with it [71]. Therefore, the orofacial pain specialist should be aware of these other associations, in order to make differential diagnosis and consequently to refer the patient to a proper health care provider.

### Future applications

Based on this review, one could recommend that future studies investigate not just the prevalence but also the biopsychosocial aspects among these patients in order to find common etiological factors. Further, there is a need for studies based on the validated diagnostic system, dividing the patients in the specified sub-

groups of DC/TMD in order to present associations based on cause-relationships and not only the TMD group as an entity. The same accounts for tinnitus. Anamnesis remains the easiest and most cost-effective method for assessing tinnitus. However, instrumental examinations are also available to assess its severity, which then also could be used to correlate tinnitus severity with the presence of TMD. Then, studies could also specify whether tinnitus was bilateral or unilateral and if it is present on the same side of each specific TMD diagnosis. When common etiological factors are presented, then randomized, controlled clinical treatment studies are warranted with for instance standard treatments of TMD and tinnitus compared to combination of treatments looking at both pain reduction, physical functioning, and reduction in tinnitus. In addition, since conservative treatments for TMD include cognitive behavioral therapies, it would be interesting to investigate the role of these therapies in tinnitus related to TMD. It would also be intriguing to distinguish if there could be gender-specific causes or treatment approaches.

## Conclusion

It can be concluded that there is a strong significant association between tinnitus and TMD and that the prevalence of TMD among patients with tinnitus is significantly higher than the prevalence of tinnitus among patients with TMD. Further research is needed to unravel the nature of this association and its clinical implication.

## Acknowledgments

We would like to thank the librarians Lovisa Liljegren and Emma-Lotta Säätelä of the University Library at Karolinska Institutet.


## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

The author(s) reported there is no funding associated with the work featured in this article.

## ORCID

Giancarlo De La Torre Canales Ph.D.  <http://orcid.org/0000-0002-0921-342X>

## Data availability statement

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

## References

- [1] Scrivani SJ, Keith DA, Kaban LB. Temporomandibular disorders. *N Engl J Med.* 2008;359(25):2693–2705. doi: [10.1056/NEJMra0802472](https://doi.org/10.1056/NEJMra0802472)
- [2] Kapos FP, Exposto FG, Oyarzo JF, et al. Temporomandibular disorders: a review of current concepts in aetiology, diagnosis and management. *Oral Surg.* 2020;13(4):321–334. doi: [10.1111/ors.12473](https://doi.org/10.1111/ors.12473)
- [3] Valesan LF, Da-Cas CD, Réus JC, et al. Prevalence of temporomandibular joint disorders: a systematic review and meta-analysis. *Clin Oral Investig.* 2021;25(2):441–453. doi: [10.1007/s00784-020-03710-w](https://doi.org/10.1007/s00784-020-03710-w)
- [4] Peleg O, Haddad L, Kleinman S, et al. Temporomandibular disorders and bruxism in patients attending a tinnitus clinic. *Appl Sci.* 2022;12(10):1. doi: [10.3390/app12104970](https://doi.org/10.3390/app12104970)
- [5] Slade GD, Fillingim RB, Sanders AE, et al. Summary of findings from the OPPERA prospective cohort study of incidence of first-onset temporomandibular disorder: implications and future directions. *J Pain.* 2013 Dec;14(12 Suppl):T116–24. doi: [10.1016/j.jpain.2013.09.010](https://doi.org/10.1016/j.jpain.2013.09.010)
- [6] Bair E, Gaynor S, Slade GD, et al. Identification of clusters of individuals relevant to temporomandibular disorders and other chronic pain conditions: the OPPERA study. *Pain.* 2016 Jun;157(6):1266–1278. doi: [10.1097/j.pain.0000000000000518](https://doi.org/10.1097/j.pain.0000000000000518)
- [7] Fillingim RB, Slade GD, Diatchenko L, et al. Summary of findings from the OPPERA baseline case-control study: implications and future directions. *J Pain.* 2011 Nov;12(11 Suppl):T102–7. doi: [10.1016/j.jpain.2011.08.009](https://doi.org/10.1016/j.jpain.2011.08.009)
- [8] Dworkin SF. Perspectives on the interaction of biological, psychological and social factors in TMD. *J Am Dent Assoc.* 1994 Jul;125(7):856–863. doi: [10.14219/jada.archive.1994.0212](https://doi.org/10.14219/jada.archive.1994.0212)
- [9] Manfredini D, Winocur E, Ahlberg J, et al. Psychosocial impairment in temporomandibular disorders patients. RDC/TMD axis II findings from a multicentre study. *J Dent.* 2010 Oct;38(10):765–772. doi: [10.1016/j.jdent.2010.06.007](https://doi.org/10.1016/j.jdent.2010.06.007)
- [10] Manfredini D, di Poggio AB, Romagnoli M, et al. Mood spectrum in patients with different painful temporomandibular disorders. *Cranio.* 2004 Jul;22(3):234–240. doi: [10.1179/crn.2004.028](https://doi.org/10.1179/crn.2004.028)
- [11] Kandasamy S, Greene CS. The evolution of temporomandibular disorders: a shift from experience to evidence. *J Oral Pathol Med.* 2020 Jul;49(6):461–469. doi: [10.1111/jop.13080](https://doi.org/10.1111/jop.13080)
- [12] Suvinen TI, Reade PC, Kempainen P, et al. Review of aetiological concepts of temporomandibular pain disorders: towards a biopsychosocial model for integration of physical disorder factors with psychological and psychosocial illness impact factors. *Eur J Pain.* 2005 Dec;9(6):613–613. doi: [10.1016/j.ejpain.2005.01.012](https://doi.org/10.1016/j.ejpain.2005.01.012)

- [13] Durham J, Steele JG, Wassell RW, et al. Living with uncertainty: temporomandibular disorders. *J Dent Res*. 2010 Aug;89(8):827–830. doi: [10.1177/0022034510368648](https://doi.org/10.1177/0022034510368648)
- [14] McGrath PJ, Stevens BJ, Walker SM, et al. Psychological theories and biopsychosocial models in paediatric pain. In: *Oxford textbook of paediatric pain*. 1st ed. Oxford: Oxford University Press; 2014. p. 85–87.
- [15] Baguley D, McFerran D, Hall D. Tinnitus. *Lancet*. 2013;382(9904):1600–1607. doi: [10.1016/S0140-6736\(13\)60142-7](https://doi.org/10.1016/S0140-6736(13)60142-7)
- [16] Eggermont JJ, Roberts LE. The neuroscience of tinnitus. *Trends Neurosci*. 2004;27(11):676–682. doi: [10.1016/j.tins.2004.08.010](https://doi.org/10.1016/j.tins.2004.08.010)
- [17] Jarach CM, Lugo A, Scala M, et al. Global prevalence and incidence of tinnitus: a systematic review and meta-analysis. *JAMA Neurol*. [2022 Sep 1];79(9):888–900. doi: [10.1001/jamaneurol.2022.2189](https://doi.org/10.1001/jamaneurol.2022.2189)
- [18] Amini R, Haghani H, Masoumi M. Quality of life in the Iranian blind war survivors in 2007: a cross-sectional study. *BMC Int Health Hum Rights*. [2010 Aug 21];10(1):21. doi: [10.1186/1472-698X-10-21](https://doi.org/10.1186/1472-698X-10-21)
- [19] Esmaili AA, Renton J. A review of tinnitus. *Aust J Gen Pract*. 2018;47(4):205–208. doi: [10.31128/AJGP-12-17-4420](https://doi.org/10.31128/AJGP-12-17-4420)
- [20] Manfredini D, Olivo M, Ferronato G, et al. Prevalence of tinnitus in patients with different temporomandibular disorders symptoms. *Int Tinnitus J*. 2015;19(2):47–51. doi: [10.5935/0946-5448.20150008](https://doi.org/10.5935/0946-5448.20150008)
- [21] Yap AU, Peng BMH, Lim SY, et al. Association of temporomandibular disorders severity with otologic and concomitant pain symptoms in Asian youths. *Cranio*. 2022;1–7. doi: [10.1080/08869634.2022.2088455](https://doi.org/10.1080/08869634.2022.2088455)
- [22] De Toledo I P, Stefani FM, Porporatti AL, et al. Prevalence of otologic signs and symptoms in adult patients with temporomandibular disorders: a systematic review and meta-analysis. *Clin Oral Investig*. 2017 Mar;21(2):597–605. doi: [10.1007/s00784-016-1926-9](https://doi.org/10.1007/s00784-016-1926-9)
- [23] Mottaghi A, Menéndez-Díaz I, Cobo JL, et al. Is there a higher prevalence of tinnitus in patients with temporomandibular disorders? A systematic review and meta-analysis. *J Oral Rehabil*. 2019;46(1):76–86. doi: [10.1111/joor.12706](https://doi.org/10.1111/joor.12706)
- [24] Skog C, Fjellner J, Ekberg E, et al. Tinnitus as a comorbidity to temporomandibular disorders-A systematic review. *J Oral Rehabil*. 2019 Jan;46(1):87–99. doi: [10.1111/joor.12710](https://doi.org/10.1111/joor.12710)
- [25] Omidvar S, Jafari Z. Association between tinnitus and temporomandibular disorders: a systematic review and meta-analysis. *Ann Otol Rhinol Laryngol*. 2019 Jul;128(7):662–675. doi: [10.1177/0003489419842577](https://doi.org/10.1177/0003489419842577)
- [26] Riva JJ, Malik KMP, Burnie SJ, et al. What is your research question? An introduction to the PICOT format for clinicians. *J Can Chiropr Assoc*. 2012;56(3):167–171.
- [27] Hutton B, Salanti G, Caldwell DM, et al. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Ann Intern Med*. [2015 Jun 2];162(11):777–784. doi: [10.7326/M14-2385](https://doi.org/10.7326/M14-2385)
- [28] Clark J, Sanders S, Carter M, et al. Improving the translation of search strategies using the polyglot search Translator: a randomized controlled trial. *J Med Lib Assoc*. 2020;108(2):12. doi: [10.5195/jmla.2020.834](https://doi.org/10.5195/jmla.2020.834)
- [29] Bramer W, Giustini D, de Jonge G, et al. De-duplication of database search results for systematic reviews in EndNote. *J Med Lib Assoc*. 2016;104(3):1. doi: [10.5195/jmla.2016.24](https://doi.org/10.5195/jmla.2016.24)
- [30] Ouzzani M, Hammady H, Fedorowicz Z, et al. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. [2016 Dec 5];5(1):210. doi: [10.1186/s13643-016-0384-4](https://doi.org/10.1186/s13643-016-0384-4)
- [31] Shamliyan TA, Kane RL, Ansari MT, et al. Development quality criteria to evaluate nontherapeutic studies of incidence, prevalence, or risk factors of chronic diseases: pilot study of new checklists. *J Clin Epidemiol*. 2011;64(6):637–657. doi: [10.1016/j.jclinepi.2010.08.006](https://doi.org/10.1016/j.jclinepi.2010.08.006)
- [32] Bush F. Tinnitus and otalgia in temporomandibular disorders. *J Prosthet Dent*. 1987;58(4):4. doi: [10.1016/0022-3913\(87\)90282-4](https://doi.org/10.1016/0022-3913(87)90282-4)
- [33] Lee C, Lin M-C, Lin H-T, et al. Increased risk of tinnitus in patients with temporomandibular disorders: a retrospective population-based cohort study. *Eur Arch Otorhinolaryngol*. 2016;273(1):203–208. doi: [10.1007/s00405-015-3491-2](https://doi.org/10.1007/s00405-015-3491-2)
- [34] Song H-S, Shin J-S, Lee J, et al. Association between temporomandibular disorders, chronic diseases, and ophthalmologic and otolaryngologic disorders in Korean adults: a cross-sectional study. *PLoS One*. 2018;13(1):e0191336. doi: [10.1371/journal.pone.0191336](https://doi.org/10.1371/journal.pone.0191336)
- [35] Kim Y, Park Y, Han K, et al. Prevalence of tinnitus according to temporomandibular joint disorders and dental pain: the Korean National population-based study. *J Oral Rehabil*. 2018;45(3):6. doi: [10.1111/joor.12604](https://doi.org/10.1111/joor.12604)
- [36] Edvall N, Gunan E, Genitsaridi E, et al. Impact of temporomandibular joint complaints on tinnitus-related distress. *Front Neurosci*. 2019;13:1. doi: [10.3389/fnins.2019.00879](https://doi.org/10.3389/fnins.2019.00879)
- [37] Kong G-S, Lee S-H, Park K, et al. Association of temporomandibular disorders and tinnitus with health-related quality of life: a cross-sectional study using the fifth Korea national health and nutrition examination survey. *J Oral Rehabil*. 2022;49(3):12. doi: [10.1111/joor.13287](https://doi.org/10.1111/joor.13287)
- [38] Henderson D, Cooper J, Bryan G, et al. Otolgic complaints in temporomandibular joint syndrome. *Arch Otolaryngol Head Neck Surg*. 1992;118(11):5. doi: [10.1001/archotol.1992.01880110076015](https://doi.org/10.1001/archotol.1992.01880110076015)
- [39] Ciancaglini R, Loreti P, Radaelli G. Ear, nose, and throat symptoms in patients with TMD: the association of symptoms according to severity of arthropathy. *J Orofac Pain*. 1994;8(3):293–297.
- [40] Lam D, Lawrence H, Tenenbaum H. Aural symptoms in temporomandibular disorder patients attending a craniofacial pain unit. *J Orofac Pain*. 2001;15(2):146–157.
- [41] Kuttilla S, Kuttilla M, Bell Y, et al. Recurrent tinnitus and associated ear symptoms in adults. *Int J Audiol*. 2005;44(3):7. doi: [10.1080/14992020500057608](https://doi.org/10.1080/14992020500057608)
- [42] Bernhardt O, Mundt T, Welk A, et al. Signs and symptoms of temporomandibular disorders and the incidence of tinnitus. *J Oral Rehabil*. 2011;38(12):11. doi: [10.1111/j.1365-2842.2011.02224.x](https://doi.org/10.1111/j.1365-2842.2011.02224.x)

- [43] Saldanha ADD, Hilgenberg PB, Pinto LMS, et al. Are temporomandibular disorders and tinnitus associated? *Cranio*. 2012;30(3):166–171. doi: [10.1179/crn.2012.026](https://doi.org/10.1179/crn.2012.026)
- [44] Fernandes G, Goncalves D, Siqueira J, et al. Painful temporomandibular disorders, self reported tinnitus and depression are highly associated. *Arquivos de Neuro psiquiatria*. 2013;71(12):5. doi: [10.1590/0004-282X20130191](https://doi.org/10.1590/0004-282X20130191)
- [45] Akhter R, Morita M, Ekuni D, et al. Self-reported aural symptoms, headache and temporomandibular disorders in Japanese young adults. *BMC Musculoskeletal Disord*. 2013;14(58). doi: [10.1186/1471-2474-14-58](https://doi.org/10.1186/1471-2474-14-58)
- [46] Buegers R, Kleinjung T, Behr M, et al. Is there a link between tinnitus and temporomandibular disorders? *J Prosthet Dent*. 2014;111(3):6. doi: [10.1016/j.prosdent.2013.10.001](https://doi.org/10.1016/j.prosdent.2013.10.001)
- [47] Cebi A. Presence of tinnitus and tinnitus-related hearing loss in temporomandibular disorders. *Cranio*. 2023;41(2):5. doi: [10.1080/08869634.2020.1829290](https://doi.org/10.1080/08869634.2020.1829290)
- [48] Oliveira M, Barros C, Caixeirinho P, et al. Is there an association between tinnitus and temporomandibular disorders? *Portuguese J Otorhinolaryngology- Head Neck Surg*. 2023;61(3):5.
- [49] Hilgenberg P, Saldanha A, Cunha C, et al. Temporomandibular disorders, otologic symptoms and depression levels in tinnitus patients. *J Oral Rehabil*. 2012;39(4):6. doi: [10.1111/j.1365-2842.2011.02266.x](https://doi.org/10.1111/j.1365-2842.2011.02266.x)
- [50] Algieri GMA, Leonardi A, Arangio P, et al. Tinnitus in temporomandibular joint disorders: Is it a specific somatosensory tinnitus subtype? *Int Tinnitus J*. 2017;20(2):5. doi: [10.5935/0946-5448.20160016](https://doi.org/10.5935/0946-5448.20160016)
- [51] Kusdra P, Stechman-Neto J, Leao B, et al. Relationship between otological symptoms and TMD. *Int J Tinnitus*. 2018;22(1):5. doi: [10.5935/0946-5448.20180005](https://doi.org/10.5935/0946-5448.20180005)
- [52] Maciel L, Landim F, Vasconcelos B. Otological findings and other symptoms related to temporomandibular disorders in young people. *Br J Oral Maxillofacial Surg*. 2018;56(8):5. doi: [10.1016/j.bjoms.2018.08.005](https://doi.org/10.1016/j.bjoms.2018.08.005)
- [53] Pavaci S, Tortorella F, Fioretti A, et al. Analysis of the audiological characteristics and comorbidity in patients with chronic tinnitus. *Audiol Res*. 2019;9(2):1. doi: [10.4081/audiores.2019.231](https://doi.org/10.4081/audiores.2019.231)
- [54] Michiels S, Harrison S, Vesala M, et al. The presence of physical symptoms in patients with tinnitus: international web-based survey. *Interact J Med Res*. 2019;8(3):e14519. doi: [10.2196/14519](https://doi.org/10.2196/14519)
- [55] Rubinstein B, Axelsson A, Carlsson G. Prevalence of signs and symptoms of craniomandibular disorders in tinnitus patients. *J Craniomandib Disord*. 1990;4(3):186–192.
- [56] Parker W, Chole R. Tinnitus, vertigo, and temporomandibular disorders. *Am J Orthod Dentofac Orthopedics*. 1995;107(2):6. doi: [10.1016/S0889-5406\(95\)70130-3](https://doi.org/10.1016/S0889-5406(95)70130-3)
- [57] Ren Y, Isberg A. Tinnitus in patients with temporomandibular joint internal derangement. *Cranio*. 1995;13(2):6. doi: [10.1080/08869634.1995.11678048](https://doi.org/10.1080/08869634.1995.11678048)
- [58] Tuz H, Onder E, Kisnisci R. Prevalence of otologic complaints in patients with temporomandibular disorder. *Am J Orthod Dentofacial Orthop*. 2003;123(6):4. doi: [10.1016/S0889-5406\(03\)00153-7](https://doi.org/10.1016/S0889-5406(03)00153-7)
- [59] Riga M, Xenellis J, Peraki E, et al. Aural symptoms in patients with temporomandibular joint disorders: multiple frequency tympanometry provides objective evidence of changes in middle ear impedance. *Otology And Neurotology*. 2010;31(9):6. doi: [10.1097/MAO.0b013e3181edb703](https://doi.org/10.1097/MAO.0b013e3181edb703)
- [60] Fernandes G, Siqueira J, Goncalves D, et al. Association between painful temporomandibular disorders, sleep bruxism and tinnitus. *Braz Oral Res*. 2014;28(1):1–7. doi: [10.1590/1807-3107BOR-2014.vol28.0003](https://doi.org/10.1590/1807-3107BOR-2014.vol28.0003)
- [61] McCormack A, Edmondson-Jones M, Somerset S, et al. A systematic review of the reporting of tinnitus prevalence and severity. *Hear Res*. 2016;337:70–79. doi: [10.1016/j.heares.2016.05.009](https://doi.org/10.1016/j.heares.2016.05.009)
- [62] Cimino R, Bucci R, Ambrosio A, et al. Temporomandibular disorders, neck disability, and oral parafunctions in tinnitus patients: a cross-sectional epidemiological study from southern Italy. *Cranio*. 2022 Nov;40(6):485–493. doi: [10.1080/08869634.2020.1781499](https://doi.org/10.1080/08869634.2020.1781499)
- [63] Sencimen M, Varol A, Baykal B, et al. Histological characteristics of ligaments between middle ear and temporomandibular joint. *Eur J Dent*. 2009 Oct;3(4):280–284. doi: [10.1055/s-0039-1697445](https://doi.org/10.1055/s-0039-1697445)
- [64] Myrhaug H. The incidence of ear symptoms in cases of malocclusion and temporo-mandibular joint disturbances. *Br J Oral Surg*. 1964;2(1):28–32. doi: [10.1016/S0007-117X\(64\)80004-4](https://doi.org/10.1016/S0007-117X(64)80004-4)
- [65] Moller AR. Similarities between severe tinnitus and chronic pain. *J Am Acad Audiol*. 2000;11(3):115–124. doi: [10.1055/s-0042-1748038](https://doi.org/10.1055/s-0042-1748038)
- [66] Ramirez L, Ballesteros L, Sandoval G. Topical review: temporomandibular disorders in an integral otic symptom model. *Int J Audiol*. 2008;47(4):215–227. doi: [10.1080/14992020701843137](https://doi.org/10.1080/14992020701843137)
- [67] Lassmann Ł, Pollis M, Żółtowska A, et al. Gut bless your pain—roles of the gut microbiota, sleep, and melatonin in chronic orofacial pain and depression. *Biomedicines*. 2022;10(7):1528. doi: [10.3390/biomedicines10071528](https://doi.org/10.3390/biomedicines10071528)
- [68] Ko SY, Baeg MS, Kwon HR, et al. Temporomandibular joint herniation into the external auditory canal presenting as clicking tinnitus: a case report. *Cranio*. 2024 Jan;42(1):90–93. doi: [10.1080/08869634.2023.2221990](https://doi.org/10.1080/08869634.2023.2221990)
- [69] Greene CS. Diagnosis and treatment of temporomandibular disorders: emergence of a new care guidelines statement. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2010 Aug;110(2):137–139. doi: [10.1016/j.tripleo.2010.04.032](https://doi.org/10.1016/j.tripleo.2010.04.032)
- [70] Christidis N, Al-Moraissi EA, Al-Ak'hali MS, et al. Psychological treatments for temporomandibular disorder pain-A systematic review. *J Oral Rehabil*. 2024 Jul;51(7):1320–1336. doi: [10.1111/joor.13693](https://doi.org/10.1111/joor.13693)
- [71] Han BI, Lee HW, Kim TY, et al. Tinnitus: characteristics, causes, mechanisms, and treatments. *J Clin Neurol*. 2009 Mar;5(1):11–19. doi: [10.3988/jcn.2009.5.1.11](https://doi.org/10.3988/jcn.2009.5.1.11)
- [72] Kmeid E, Nacouzi M, Hallit S, et al. Prevalence of temporomandibular joint disorder in the Lebanese population, and its association with depression, anxiety, and stress. *Head Face Med*. [2020 Sep 4];16(1):19. doi: [10.1186/s13005-020-00234-2](https://doi.org/10.1186/s13005-020-00234-2)
- [73] Yap AU, Sultana R, Natu VP. Stress and emotional distress: their associations with somatic and

- temporomandibular disorder-related symptoms. *Psychol Health Med.* 2022 Apr;27(4):876–887. doi: [10.1080/13548506.2021.1908571](https://doi.org/10.1080/13548506.2021.1908571)
- [74] Connell J, Harrison E, Bassiouni A, et al. FiveQ: a new easy-to-use validated clinical instrument for tinnitus severity. *Clin Otolaryngol.* 2022 Nov;47(6):672–679. doi: [10.1111/coa.13973](https://doi.org/10.1111/coa.13973)
- [75] Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic criteria for temporomandibular disorders (DC/TMD) for clinical and research applications: recommendations of the international RDC/TMD consortium Network\* and orofacial pain special interest Group† [article]. *J Oral Facial Pain Headache.* 2014;28(1):6–27. doi: [10.11607/jop.1151](https://doi.org/10.11607/jop.1151)
- [76] Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord.* 1992 Fall;6(4):301–355.
- [77] Rongo R, Ekberg E, Nilsson IM, et al. Diagnostic criteria for temporomandibular disorders (DC/TMD) for children and adolescents: an international delphi study-part 1-development of axis I. *J Oral Rehabil.* 2021 Jul;48(7):836–845. doi: [10.1111/joor.13175](https://doi.org/10.1111/joor.13175)
- [78] Lövgren A, Häggman-Henrikson B, Visscher CM, et al. Temporomandibular pain and jaw dysfunction at different ages covering the lifespan—A population based study. *Eur J Pain.* 2016 Apr;20(4):532–540. doi: [10.1002/ejp.755](https://doi.org/10.1002/ejp.755)
- [79] Stechman-Neto J, Porporatti AL, Porto de Toledo I, et al. Effect of temporomandibular disorder therapy on otologic signs and symptoms: a systematic review. *J Oral Rehabil.* 2016 Jun;43(6):468–479. doi: [10.1111/joor.12380](https://doi.org/10.1111/joor.12380)
- [80] Morell GFC. Insufficient evidence to support or reject effect of conservative TMD therapies on otologic signs and symptoms. *Evid Based Dent.* [2018 Mar 23];19(1):26–27. doi: [10.1038/sj.ebd.6401293](https://doi.org/10.1038/sj.ebd.6401293)
- [81] Paulino MR, Moreira VG, Lemos GA, et al. Prevalence of signs and symptoms of temporomandibular disorders in college preparatory students: associations with emotional factors, parafunctional habits, and impact on quality of life. *Cien Saude Colet.* 2018 Jan;23(1):173–186. doi: [10.1590/1413-81232018231.18952015](https://doi.org/10.1590/1413-81232018231.18952015)
- [82] Lövgren A, Visscher CM, Häggman-Henrikson B, et al. Validity of three screening questions (3Q/TMD) in relation to the DC/TMD. *J Oral Rehabil.* 2016 Oct;43(10):729–736. doi: [10.1111/joor.12428](https://doi.org/10.1111/joor.12428)
- [83] Kroenke K, Spitzer RL, Williams JB, et al. An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics.* 2009 Nov-Dec;50(6):613–621. doi: [10.1176/appi.psy.50.6.613](https://doi.org/10.1176/appi.psy.50.6.613)