



Available online at www.sciencedirect.com

ScienceDirect

Nuclear Physics A 932 (2014) 432–436

**NUCLEAR
PHYSICS**

A

www.elsevier.com/locate/nuclphysa

Jet quenching effects on the anisotropic flow at RHIC

R.P.G. Andrade ^{a,*}, J. Noronha ^b, Gabriel S. Denicol ^c

^a *Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, Santo André, SP, 09210-170, Brazil*

^b *Instituto de Física, Universidade de São Paulo, C.P. 66318, 05315-970 São Paulo, SP, Brazil*

^c *Department of Physics, McGill University, 3600 University Street, Montreal, Quebec, H3A 2T8, Canada*

Received 7 April 2014; accepted 10 July 2014

Available online 30 July 2014

Abstract

In this article we investigate how the energy and momentum deposited by partonic dijets in the quark–gluon plasma may perturb the geometry-induced hydrodynamic expansion of the bulk nuclear matter created in heavy ion collisions at the RHIC. The coupling between the jets and the medium is done through a source term in the energy–momentum conservation equations for ideal hydrodynamics. We concentrate our attention at mid-rapidity and solve the equations event-by-event imposing boost-invariance. For $p_T \gtrsim 1$ GeV the anisotropic flow is found to be considerably enhanced, if the dijets deposit on average more than 12 GeV in the medium (or equivalently 6 GeV for each jet of the pair), which corresponds, in our model, to an average suppression greater than 65% of the initial jet transverse energy.

© 2014 Elsevier B.V. All rights reserved.

Keywords: Relativistic hydrodynamics; Jet quenching; Fourier coefficients of the flow

1. Introduction

The jet suppression observed in relativistic heavy ion collisions performed at the Relativistic Heavy Ion Collider (RHIC) and the Large Hadron Collider (LHC) [1–7] is a strong evidence that a hot and dense nuclear matter, usually called quark–gluon plasma (QGP), is created in these experiments. The observation of such a phenomenon ushered in an ample scenario for the study of how the jet spectra are modified by the interaction with the medium (see, for instance, [8] and references therein). However, the counterpart, i.e., the study of how the medium is modified

* Corresponding author.

by the interaction with the jets, has been poorly explored. The aim of this article is to improve our knowledge of the effects of the dijets on the hydrodynamic evolution of the QGP. Using a 2 + 1 hydrodynamic model on an event-by-event basis, we try to understand such effects through of the anisotropic flow parameters $\{v_n, \Psi_n\}$, namely the n th Fourier coefficient of the azimuthal distribution of hadrons and the respective phase. Naturally, depending on the amount of energy-momentum deposited in the medium by the dijets, the jet-induced anisotropic flow may represent a non-negligible fraction of the total anisotropic flow. All the results presented in this article correspond to Au+Au collisions at 200 A GeV in the (0–5)% centrality window. We use hyperbolic coordinates, i.e., $\tau = \sqrt{t^2 - z^2}$, $\eta = 0.5 \ln[(t+z)/(t-z)]$ and $\vec{r} = (x, y)$. The initial time at which we begin the hydrodynamic evolution $\tau_0 = 1$ fm. In addition, $\hbar = k_B = c = 1$.

2. Hydrodynamic model

Based on the assumption that the energy-momentum lost by the jets quickly thermalizes in the QGP [9], the coupling between the dijets and the medium, in our model, is done through a source term in the energy-momentum conservation equations. Thus, in the ideal fluid approximation, one finds that

$$D_\mu T^{\mu\nu} = J^\nu, \quad (1)$$

where $T^{\mu\nu} = \omega u^\mu u^\nu - pg^{\mu\nu}$ is the ideal fluid energy-momentum tensor, ω is the enthalpy, p the pressure, u^μ the fluid 4-velocity, D_μ the covariant derivative, and $g^{\mu\nu}$ is the metric tensor. In our model, this equation is solved assuming boost-invariance. The 4-current density J^ν (the source) is parameterized, in the laboratory frame, as (see, for instance, Refs. [9–15])

$$J^\nu(\tau, \vec{r}) = \sum_{n=1}^{n_p} \frac{s(\vec{r}_n^{\text{jet}}(\tau))}{s_0} \frac{dE}{dl} \Big|_0 F(\vec{r} - \vec{r}_n^{\text{jet}}(\tau), \tau; \sigma) (1, \vec{v}_n^{\text{jet}}, 0), \quad (2)$$

where n_p is the number of partonic jets, \vec{r}_n^{jet} and \vec{v}_n^{jet} (with $|\vec{v}_n^{\text{jet}}| = 1$) are the position and velocity of the n th parton that moves in a straight line on the mid-rapidity transverse plane, $s(\vec{r}_n^{\text{jet}})$ is the entropy density computed at the position of the n th parton, and the function F [9,15] corresponds to a Gaussian shaped source of width σ (we set $\sigma = 0.6$ fm). We assume only one dijet per event, consequently, $n_p = 2$, $\vec{r}_1^{\text{jet}}(\tau_0) = \vec{r}_2^{\text{jet}}(\tau_0)$ and $\vec{v}_1^{\text{jet}} = -\vec{v}_2^{\text{jet}}$. The parameters $dE/dl|_0$ and s_0 are the reference energy loss rate ($l = \tau - \tau_0$ is transverse distance traveled by the partons) and the reference entropy density. The latter corresponds to the maximum of the average entropy density distribution in the (0–5)% centrality window (in our model, $s_0 = 70$ fm⁻³). The former is a free parameter, which is varied from 5 to 20 GeV/fm. Naturally, the bigger this parameter is, the more energy-momentum the jets deposit in the medium.

We use the equation of state EOS S95n-v1 [16], which combines results from lattice QCD at high temperatures and the hadron resonance gas equation at low temperatures. To compute the particle spectrum, we use the Cooper–Frye prescription [17]. In this method, the particles escape from the fluid after crossing a hyper-surface of constant temperature, usually called freeze-out temperature, T_{fo} (we set $T_{\text{fo}} = 0.14$ GeV). Finally, all the results presented in this paper correspond to positively charged pions directly emitted from the freeze-out hyper-surface.

In few words, the procedure to compute an observable event-by-event, including the jet parametrization, is the following: (i) the initial conditions for hydrodynamics are computed using an implementation of the Monte Carlo Glauber model [18,19]; (ii) the initial position of the dijet (one dijet per event) is chosen on the mid-rapidity transverse plane (see the details in Ref. [15]);




Fig. 1. (Color online.) Left: average energy deposited in the medium, around mid-rapidity, by the dijet, $\langle E_d^{\text{jet}} \rangle$, in the (0–5)% centrality window, as a function of the reference energy loss rate $dE/dl|_0$ (using the jet ensemble). Right: distribution of the ratio $\delta E = E_d^{\text{jet}} / E_T^{\text{jet}}$ for four values of the parameter $dE/dl|_0$. The respective average value $\langle \delta E \rangle$ is shown on the plot.

(iii) the dijet azimuthal angle is chosen isotropically; (iv) the initial jet transverse energy E_T^{jet} (the same for both jets in the pair) is chosen according to the jet yield per event in p+p collisions scaled by the number of binary collisions in Au+Au collisions [15,20]; (v) the hydrodynamic evolution is computed through the SPH method [21,22] and (vi) the final spectra (for direct positively charged pions) is computed using the Cooper–Frye prescription [17]. At the end of the simulation, the average value of a given observable is calculated over an ensemble of events. We define “mixed ensemble” as the ensemble composed by 750 events without dijets and 250 with dijets (totaling 1000 events). This proportion is fixed by the jet yield per event [15,20]. On the other side, the “jet ensemble” corresponds only to events with dijets (250 events).

3. Results

In Fig. 1 (left) we show the average energy deposited in the medium by the dijet, $\langle E_d^{\text{jet}} \rangle$, in the (0–5)% centrality window, as a function of the reference energy loss rate $dE/dl|_0$. To compute the curve labeled “smooth” (squares), the fluctuating initial energy density distribution was replaced, in each event, by a smooth one while keeping unchanged the initial position of the dijet. As one can see, the fluctuations slightly enhance the suppression of jets in the medium. In the same figure (right), we show the distribution of the ratio $\delta E = E_d^{\text{jet}} / E_T^{\text{jet}}$, i.e., the relative amount of energy (with respect to the initial jet transverse energy E_T^{jet}) that is lost to the medium, for four values of the parameter $dE/dl|_0$. The respective average value $\langle \delta E \rangle$ is shown on the plot. Observe that $\langle \delta E \rangle$ gets close to unity when $dE/dl|_0$ is increased. In fact, depending on the magnitude of the coupling between the jets and the QGP, a considerable fraction of the jets may be completely absorbed by the medium. These distributions survey, in our model, an estimative of the suppression of the jets in the medium and can be used to calibrate the free parameter $dE/dl|_0$. As we are going to see in the next plot, for $dE/dl|_0 \gtrsim 15$ GeV/fm, which corresponds to a suppression on average greater than 65%, the jet quenching effect may create relevant additional anisotropic flow.

In Fig. 2, we show the v_n coefficients ($n = 1, 2, 3$), as a function of the transverse momentum, for four values of the parameter $dE/dl|_0$. The left panels correspond to the Event Plane method




Fig. 2. (Color online.) Transverse momentum dependence of the v_n coefficients ($n = 1, 2, 3$) for four values of the parameter $dE/dl|_0$. The left panels correspond to the event plane method. On the right panels we show the same observables computed using $\Psi_n = \Psi_n(p_T)$, i.e., the phase is computed for each p_T bin [22]. The panels labeled “mixed” correspond to an ensemble of 1000 events that includes 750 events without and 250 events with dijets. The panels labeled “jet” correspond to an ensemble of 250 events that includes only events with dijets.

(EP) where the phase Ψ_n is computed using all the hadrons of the event [23]. The right panels correspond to method used in Ref. [22] where $\Psi_n = \Psi_n(p_T)$, i.e., the phase is computed for each p_T bin. Naturally, the latter procedure maximizes the anisotropy. The negative sign observed in the coefficient $v_1(p_T)$, computed using the event plane method, is a consequence of momentum conservation: if the low p_T particles move in one direction, the higher p_T particles must move in the opposite direction to conserve momentum. Note that, in the majority of the cases, for $dE/dl|_0 > 5$ GeV/fm, the effects of the jets are important in the region of intermediate p_T ($1 \lesssim p_T \lesssim 3$ GeV). In the region of low p_T ($p_T < 1$ GeV) the effects are negligible. On the other side, using our lower limit for the coupling between the QGP and the jets, $dE/dl|_0 = 5$ GeV/fm, the results are nearly identical to the results without jets. Finally, the anisotropy is enhanced, as expected, when only events with dijets are included (the jet ensemble).

4. Conclusions

In our simplified model we found that the effects of the dijets on the hydrodynamic evolution of the QGP are less important in the region of low p_T ($p_T < 1$ GeV). Even for the highest value of energy loss used in this paper, $dE/dl|_0 = 20$ GeV/fm, which corresponds to an average suppression of 83% of the initial jet transverse energy, we found that the dijets affect mainly the region of intermediate p_T ($1 \lesssim p_T \lesssim 3$ GeV). For a more detailed analysis see [15].

Acknowledgements

R.P.G. Andrade and J. Noronha thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 2012/14057-1, 2011/21173-5) and Conselho Nacional de Desenvolvimento Científico

e Tecnológico (CNPq 306285/2011-8) for financial support. G.S. Denicol acknowledges the support of a Banting fellowship provided by the Natural Sciences and Engineering Research Council of Canada.

References

- [1] J. Adams, et al., Transverse momentum and collision energy dependence of high p_T hadron suppression in Au+Au collisions at ultrarelativistic energies, *Phys. Rev. Lett.* 91 (2003) 172302, <http://dx.doi.org/10.1103/PhysRevLett.91.172302>.
- [2] K. Adcox, et al., Suppression of hadrons with large transverse momentum in central Au+Au collisions at $\sqrt{s_{NN}} = 130$ GeV, *Phys. Rev. Lett.* 88 (2002) 022301, <http://dx.doi.org/10.1103/PhysRevLett.88.022301>.
- [3] J. Adams, et al., Evidence from d+Au measurements for final state suppression of high p_T hadrons in Au+Au collisions at RHIC, *Phys. Rev. Lett.* 91 (2003) 072304, <http://dx.doi.org/10.1103/PhysRevLett.91.072304>.
- [4] S. Adler, et al., Absence of suppression in particle production at large transverse momentum in $\sqrt{s_{NN}} = 200$ GeV d+Au collisions, *Phys. Rev. Lett.* 91 (2003) 072303, <http://dx.doi.org/10.1103/PhysRevLett.91.072303>.
- [5] S. Chatrchyan, et al., Study of high- p_T charged particle suppression in PbPb compared to pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Eur. Phys. J. C* 72 (2012) 1945, <http://dx.doi.org/10.1140/epjc/s10052-012-1945-x>.
- [6] B. Abelev, et al., Centrality dependence of charged particle production at large transverse momentum in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Lett. B* 720 (2013) 52–62, <http://dx.doi.org/10.1016/j.physletb.2013.01.051>.
- [7] K. Aamodt, et al., Suppression of charged particle production at large transverse momentum in central Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Lett. B* 696 (2011) 30–39, <http://dx.doi.org/10.1016/j.physletb.2010.12.020>.
- [8] K.M. Burke, A. Buzzatti, N. Chang, C. Gale, M. Gyulassy, et al., Extracting jet transport coefficient from jet quenching at RHIC and LHC, [arXiv:1312.5003](http://arxiv.org/abs/1312.5003).
- [9] A. Chaudhuri, U. Heinz, Effect of jet quenching on the hydrodynamical evolution of QGP, *Phys. Rev. Lett.* 97 (2006) 062301, <http://dx.doi.org/10.1103/PhysRevLett.97.062301>.
- [10] B. Betz, J. Noronha, G. Torrieri, M. Gyulassy, D.H. Rischke, Universal flow-driven conical emission in ultrarelativistic heavy-ion collisions, *Phys. Rev. Lett.* 105 (2010) 222301, <http://dx.doi.org/10.1103/PhysRevLett.105.222301>.
- [11] B. Betz, J. Noronha, G. Torrieri, M. Gyulassy, D.H. Rischke, Conical correlations, Bragg peaks, and transverse flow deflections in jet tomography, *Nucl. Phys. A* 830 (2009) 777C–780C, <http://dx.doi.org/10.1016/j.nuclphysa.2009.10.070>.
- [12] G. Torrieri, B. Betz, J. Noronha, M. Gyulassy, Mach cones in heavy ion collisions, *Acta Phys. Pol. B* 39 (2008) 3281–3308.
- [13] B. Betz, J. Noronha, G. Torrieri, M. Gyulassy, I. Mishustin, D.H. Rischke, Universality of the diffusion wake from stopped and punch-through jets in heavy-ion collisions, *Phys. Rev. C* 79 (2009) 034902, <http://dx.doi.org/10.1103/PhysRevC.79.034902>.
- [14] B. Betz, M. Gyulassy, J. Noronha, G. Torrieri, Anomalous conical di-jet correlations in pQCD vs AdS/CFT, *Phys. Lett. B* 675 (2009) 340–346, <http://dx.doi.org/10.1016/j.physletb.2009.04.026>.
- [15] R.P.G. Andrade, J. Noronha, G.S. Denicol, Jet quenching effects on the direct, elliptic, and triangular flow at RHIC, [arXiv:1403.1789](http://arxiv.org/abs/1403.1789).
- [16] P. Huovinen, P. Petreczky, QCD equation of state and hadron resonance gas, *Nucl. Phys. A* 837 (2010) 26–53, <http://dx.doi.org/10.1016/j.nuclphysa.2010.02.015>.
- [17] F. Cooper, G. Frye, Comment on the single particle distribution in the hydrodynamic and statistical thermodynamic models of multiparticle production, *Phys. Rev. D* 10 (1974) 186, <http://dx.doi.org/10.1103/PhysRevD.10.186>.
- [18] H.-J. Drescher, Y. Nara, Effects of fluctuations on the initial eccentricity from the Color Glass Condensate in heavy ion collisions, *Phys. Rev. C* 75 (2007) 034905, <http://dx.doi.org/10.1103/PhysRevC.75.034905>.
- [19] H.-J. Drescher, Y. Nara, Eccentricity fluctuations from the color glass condensate at RHIC and LHC, *Phys. Rev. C* 76 (2007) 041903, <http://dx.doi.org/10.1103/PhysRevC.76.041903>.
- [20] S. Salur, First direct measurement of jets in $\sqrt{s_{NN}} = 200$ GeV heavy ion collisions by STAR, *Eur. Phys. J. C* 61 (2009) 761–767, <http://dx.doi.org/10.1140/epjc/s10052-009-0880-y>.
- [21] C. Aguiar, T. Kodama, T. Osada, Y. Hama, Smoothed particle hydrodynamics for relativistic heavy ion collisions, *J. Phys. G* 27 (2001) 75–94, <http://dx.doi.org/10.1088/0954-3899/27/1/306>.
- [22] R. Andrade, J. Noronha, A study of the di-hadron angular correlation function in event by event ideal hydrodynamics, *Phys. Rev. C* 88 (2013) 034909, <http://dx.doi.org/10.1103/PhysRevC.88.034909>.
- [23] A.M. Poskanzer, S. Voloshin, Methods for analyzing anisotropic flow in relativistic nuclear collisions, *Phys. Rev. C* 58 (1998) 1671–1678, <http://dx.doi.org/10.1103/PhysRevC.58.1671>.