

## STABILITY FOR THE VERTICAL ROTATION INTERVAL OF TWIST MAPPINGS

SALVADOR ADDAS-ZANATA

Instituto de Matemática e Estatística  
Universidade de São Paulo  
Rua do Matão 1010, Cidade Universitária  
05508-090 São Paulo, SP, Brazil

(Communicated by Zhihong Jeff Xia)

ABSTRACT. In this paper we consider twist mappings of the torus,  $\bar{T} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ , and their vertical rotation intervals  $\rho_V(T) = [\rho_V^-, \rho_V^+]$ , which are closed intervals such that for any  $\omega \in ]\rho_V^-, \rho_V^+[$  there exists a compact  $\bar{T}$ -invariant set  $\bar{Q}_\omega$  with  $\rho_V(\bar{x}) = \omega$  for any  $\bar{x} \in \bar{Q}_\omega$ , where  $\rho_V(\bar{x})$  is the vertical rotation number of  $\bar{x}$ . In case  $\omega$  is a rational number,  $\bar{Q}_\omega$  is a periodic orbit (this study began in [1] and [2]). Here we analyze how  $\rho_V^-$  and  $\rho_V^+$  behave as we perturb  $\bar{T}$  when they assume rational values. In particular we prove that for analytic area-preserving mappings these functions are locally constant at rational values.

**1. Introduction and main result.** The dynamics of a homeomorphism of the circle has a well-known and very important invariant, the so called rotation number. Roughly speaking, it measures the average speed the orbit of a point in the circle rotates around it and its rationality or not has strong implications on the dynamics of the homeomorphism (see [15] for a nice didactic exposition). In particular, given a mapping with a rational rotation number which is not conjugate to a rigid rotation; if the rotation number changes when one composes the mapping with a small positive rotation, it does not change with a small negative one and vice-versa.

Similar results for circle endomorphisms have been proved by Boyland in [9]. As it is well-known, endomorphisms of the circle do not have a single rotation number, they have a closed interval of rotation numbers (the rotation interval), with many interesting properties (see [14] and [21]). The results proved by Boyland in [9] concern the behavior of the extreme points of the rotation interval.

The aim of the present paper is to study a similar problem in the context of twist mappings of the torus. First we remember that twist mappings of the torus have an invariant called the vertical rotation interval (see [1], [2], [4] and [5]). To be more precise, given a twist mapping  $T : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}$  which induces a diffeomorphism  $\bar{T} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$  homotopic to the Dehn twist  $(\phi, I) \rightarrow (\phi + I \bmod 1, I \bmod 1)$ , there exists a closed interval  $\rho_V(T) = [\rho_V^-, \rho_V^+]$  with the following property:

---

2000 *Mathematics Subject Classification.* 37E40, 37E45, 37C25, 37G15.

*Key words and phrases.* twist mappings, vertical rotation set, topological methods, analyticity, bifurcations.

partially supported by CNPq, grant: 301485/03-8.

- for every  $\bar{x} \in \mathbb{T}^2$  such that its vertical rotation number

$$\rho_V(\bar{x}) = \lim_{n \rightarrow \infty} \frac{p_2 \circ T^n(x) - p_2(x)}{n}$$

exists ( $x$  is any point in the cylinder that projects on  $\bar{x}$  and  $p_2 : S^1 \times \mathbb{R} \rightarrow \mathbb{R}$  is the projection in the vertical coordinate), we have  $\rho_V(\bar{x}) \in \rho_V(T)$

Moreover, for  $\omega \in \text{int}(\rho_V(T))$ , there are 2 different situations.

1)  $\omega = \frac{p}{q}$  is a rational number. In this case there is a  $q$ -periodic point  $\bar{x}$  for  $\bar{T}$  (in fact there are at least two such points) that lifts to a point  $x \in S^1 \times \mathbb{R}$  such that  $T^q(x) = x + (0, p)$ .

2)  $\omega$  is an irrational number. Then there is a compact,  $\bar{T}$ -invariant set  $\bar{Q} \subset \mathbb{T}^2$  that lifts to a set  $Q \subset S^1 \times \mathbb{R}$  such that for any  $x \in Q$  we have:

$$\lim_{n \rightarrow \infty} \frac{p_2 \circ T^n(x) - p_2(x)}{n} = \omega$$

In [4] we have shown that  $\rho_V^+$  and  $\rho_V^-$  are continuous functions of  $T$  in the  $C^1$  topology (in fact a stronger result was proved there). Note that this is in complete similarity with results for circle homeomorphisms and endomorphisms.

As we already said, the main result of this paper is inspired by the one we mentioned for circle homeomorphisms with a rational rotation number. If  $T : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}$  is an analytic area-preserving twist mapping which induces a diffeomorphism  $\bar{T} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$  homotopic to the Dehn twist and  $\rho_V^-(T) < \rho_V^+(T) \in \mathbb{Q}$ , then  $\rho_V^+(T)$  is locally constant in the  $C^1$  topology, that is, it can not be increased and decreased by appropriate  $C^1$  small perturbations. More precisely, at most one of these possibilities is allowed. A similar result holds for  $\rho_V^-(T)$  when it assumes a rational value and the vertical rotation interval is not degenerated to a point.

In [5] we solved the complementary problem. There we proved that for a  $C^1$  twist mapping  $T$ , if  $\rho_V^-(T) < \rho_V^+(T) \notin \mathbb{Q}$ , then  $\rho_V^+(T + (0, \alpha)) > \rho_V^+(T)$  for any  $\alpha > 0$ . Note that this result is also analogous to what happens for a circle homeomorphism with an irrational rotation number. For a similar result in the context of torus homeomorphisms homotopic to the identity see [7].

In [6] a partial version of the main result of this paper was proved. The class of mappings studied there was, in a certain sense, a little restrictive, as for instance, it did not contain the standard mapping (in fact, it did not contain any area-preserving mapping):

$$S_M : \begin{cases} \phi' = \phi + I' \text{ mod } 1 \\ I' = I + \frac{k}{2\pi} \sin(2\pi\phi) \text{ mod } 1 \end{cases} \tag{1}$$

Clearly, as  $S_M$  is analytic and area-preserving, the result proved here can be applied to it.

Our main goal with all this analysis of how the vertical rotation interval changes as we perturb  $T$  is a better understanding of the bifurcations that happen in a family like  $S_M$ .

As the research in this subject has shown, this is a very difficult task. The main problems with our analysis are the following:

1. The particular family may not be "versal" for the bifurcation we are considering. For instance, even when  $\rho_V^+(T)$  can be increased by appropriate perturbations, it may be the case that it remains constant or even decreases if the perturbations are restricted to a particular family.

2. There is not a nice topological characterization (at least not to our knowledge) of all the possible dynamics in the neighborhood of an index one periodic point of an area-preserving mapping. This problem is related to the following one: If  $\rho_V^+(T)$  can be increased by appropriate perturbations, then new index one periodic orbits appear. Is the appearance of these orbits associated to invariant sets of positive Lebesgue measure? In other words, is the appearance of these orbits associated to non-ergodicity with respect to Lebesgue measure for the perturbed torus mapping?
3. When  $\rho_V^+(T)$  is locally constant at a rational value, are there non-trivial invariant sets with this vertical rotation number? (by non-trivial we mean with positive Lebesgue measure)

This paper is organized as follows. In the first section we present some basic definitions and the precise statements of our results, in the second section we present a brief summary of the results used in the paper and in the third section we prove our main theorem.

**Notation and definitions:**

0) Let  $(\phi, I)$  denote the coordinates for the cylinder  $S^1 \times \mathbb{R} = (\mathbb{R}/\mathbb{Z}) \times \mathbb{R}$ , where  $\phi$  is defined modulo 1. Let  $(\tilde{\phi}, \tilde{I})$  denote the coordinates for the universal cover of the cylinder,  $\mathbb{R}^2$ . For all mappings  $T : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}$  we define  $(\phi', I') = T(\phi, I)$  and  $(\tilde{\phi}', \tilde{I}') = \tilde{T}(\tilde{\phi}, \tilde{I})$ , where  $\tilde{T} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a lift of  $T$ .

1)  $D_r^1(\mathbb{R}^2) = \{\tilde{T} : \mathbb{R}^2 \rightarrow \mathbb{R}^2 / \tilde{T} \text{ is a } C^1 \text{ diffeomorphism of the plane, } \tilde{I}'(\tilde{\phi}, \tilde{I}) \xrightarrow{\tilde{I} \rightarrow \pm\infty} \pm\infty, \partial_{\tilde{I}} \tilde{\phi}' > 0 \text{ (twist to the right), } \tilde{\phi}'(\tilde{\phi}, \tilde{I}) \xrightarrow{\tilde{I} \rightarrow \pm\infty} \pm\infty \text{ and } \tilde{T} \text{ is the lift of a } C^1 \text{ diffeomorphism } T : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}\}$ .

2)  $\text{Diff}_r^1(S^1 \times \mathbb{R}) = \{T : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R} / T \text{ is induced by an element of } D_r^1(\mathbb{R}^2)\}$ .

3) Let  $p_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $p_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the standard projections, respectively in the  $\tilde{\phi}$  and  $\tilde{I}$  coordinates ( $p_1(\tilde{\phi}, \tilde{I}) = \tilde{\phi}$  and  $p_2(\tilde{\phi}, \tilde{I}) = \tilde{I}$ ). We also use  $p_1$  and  $p_2$  for the standard projections of the cylinder.

4) Let  $D_{\text{Dehn}} = \{T \in \text{Diff}_r^1(S^1 \times \mathbb{R}) : T \text{ induces a mapping } \bar{T} : \mathbb{T}^2 \rightarrow \mathbb{T}^2 \text{ homotopic to the Dehn twist } (\phi, I) \rightarrow (\phi + I \text{ mod } 1, I \text{ mod } 1)\}$ , where  $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$  is the 2-torus (coordinates in the torus are denoted by  $(\bar{\phi}, \bar{I})$ )

5) Let  $pro : S^1 \times \mathbb{R} \rightarrow \mathbb{T}^2$  and  $\pi : \mathbb{R}^2 \rightarrow S^1 \times \mathbb{R}$  be given by:

$$\begin{aligned} pro(\phi, I) &= (\phi, I \text{ mod } 1) \\ \pi(\tilde{\phi}, \tilde{I}) &= (\tilde{\phi} \text{ mod } 1, \tilde{I}) \end{aligned}$$

6) As we already said, given a point  $\bar{x} \in \mathbb{T}^2$ , we define its vertical rotation number as (when the limit exists):

$$\rho_V(\bar{x}) = \lim_{n \rightarrow \infty} \frac{p_2 \circ T^n(x) - p_2(x)}{n}, \text{ for any } x \in pro^{-1}(\bar{x}) \tag{2}$$

Now we are ready to state our main result:

**Theorem 1.** *Let  $T \in D_{\text{Dehn}}$  be an analytic area-preserving mapping such that  $\rho_V(T) = [\rho_V^-, \frac{p}{q}]$ , with  $\frac{p}{q} \in \mathbb{Q}$ ,  $(p, q) = 1$  and  $\rho_V^- < \frac{p}{q}$ . Then, if  $V \subset D_{\text{Dehn}}$  is any sufficiently small neighborhood of  $T$  in the  $C^1$  topology, one of the following possibilities holds:*

- 1) for all  $G \in V$ ,  $\rho_V^+(G) \leq p/q$
- 2) for all  $G \in V$ ,  $\rho_V^+(G) = p/q$
- 3) for all  $G \in V$ ,  $\rho_V^+(G) \geq p/q$

The next result takes care of the remaining case. It describes what happens when  $\rho_V^- = \rho_V^+ = \frac{p}{q}$ .

**Theorem 2.** *Let  $T \in D_{\text{Dehn}}$  be an area-preserving mapping such that  $\rho_V(T) = \{\frac{p}{q}\}$ , with  $\frac{p}{q} \in \mathbb{Q}$ ,  $(p, q) = 1$ . Then, for any  $\alpha > 0$ ,  $\rho_V^+(T + (0, \alpha)) > \frac{p}{q}$  and  $\rho_V^-(T - (0, \alpha)) < \frac{p}{q}$ .*

**2. Basic tools.** In this part of the paper we state some results we use and present references.

**2.1. Some results for twist mappings.** First we recall some topological results for twist mappings essentially due to Le Calvez (see [16] and [17]). Let  $T \in \text{Diff}_r^1(S^1 \times \mathbb{R})$  and  $\tilde{T} \in D_r^1(\mathbb{R}^2)$  be a lift of  $T$  to the plane. For every pair  $(s, q)$ ,  $s \in \mathbb{Z}$  and  $q \in \mathbb{N}^*$  we define the following sets:

$$\begin{aligned}
 K_{\text{lift}}(s, q) &= \left\{ (\tilde{\phi}, \tilde{I}) \in \mathbb{R}^2: p_1 \circ \tilde{T}^q(\tilde{\phi}, \tilde{I}) = \tilde{\phi} + s \right\} \\
 &\quad \text{and} \\
 K(s, q) &= \pi \circ K_{\text{lift}}(s, q)
 \end{aligned}
 \tag{3}$$

Then we have the following:

**Lemma 1.** *For every  $s \in \mathbb{Z}$  and  $q \in \mathbb{N}^*$ ,  $K(s, q) \supset C(s, q)$ , a connected compact set that separates the cylinder.*

For all  $s \in \mathbb{Z}$  and  $q \in \mathbb{N}^*$  we can define the following functions on  $S^1$ :

$$\begin{aligned}
 \mu^-(\phi) &= \min\{p_2(z): z \in K(s, q) \text{ and } p_1(z) = \phi\} \\
 \mu^+(\phi) &= \max\{p_2(z): z \in K(s, q) \text{ and } p_1(z) = \phi\}
 \end{aligned}$$

And we can define similar functions for  $T^q(K(s, q))$ :

$$\begin{aligned}
 \nu^-(\phi) &= \min\{p_2(z): z \in T^q \circ K(s, q) \text{ and } p_1(z) = \phi\} \\
 \nu^+(\phi) &= \max\{p_2(z): z \in T^q \circ K(s, q) \text{ and } p_1(z) = \phi\}
 \end{aligned}$$

**Lemma 2.** *Defining  $\text{Graph}\{\mu^\pm\} = \{(\phi, \mu^\pm(\phi)) : \phi \in S^1\}$  we have:*

$$\text{Graph}\{\mu^-\} \cup \text{Graph}\{\mu^+\} \subset C(s, q)$$

*So for all  $\phi \in S^1$  we have  $(\phi, \mu^\pm(\phi)) \in C(s, q)$ .*

The next lemma is a fundamental result in all this theory:

**Lemma 3.**  *$T^q(\phi, \mu^-(\phi)) = (\phi, \nu^+(\phi))$  and  $T^q(\phi, \mu^+(\phi)) = (\phi, \nu^-(\phi))$ .*

For proofs of the previous results see Le Calvez [16] and [17].

**2.2. Results previously obtained by the author.** The first theorem we present asserts the existence of the vertical rotation interval and we present some of its properties:

**Theorem 3.** *To each mapping  $T \in D_{\text{Dehn}}$ , we can associate a closed interval  $\rho_V(T) = [\rho_V^-, \rho_V^+]$ , possibly degenerated to a single point, such that for every  $\bar{x} \in \mathbb{T}^2$  for which  $\rho_V(\bar{x})$  exists, we have  $\rho_V(\bar{x}) \in \rho_V(T)$  and for all  $\omega \in ]\rho_V^-, \rho_V^+[$  there is a compact  $\bar{T}$ -invariant set  $\bar{Q}_\omega \subset \mathbb{T}^2$  with  $\rho_V(\bar{x}) = \omega$ , for all  $\bar{x} \in \bar{Q}_\omega$ . If  $\omega$  is a rational number  $\frac{p}{q}$ , then  $\bar{Q}_\omega$  is a  $q$ -periodic orbit. In fact, in this case there are at least two such orbits. If they are finite, then at least one has positive index and another one has negative index.*

For a proof, see theorem 6 of the appendix of [3] and theorem 5 of [2].

**Theorem 4.** *The functions  $\rho_V^+, \rho_V^- : D_{\text{Dehn}} \rightarrow \mathbb{R}$  are continuous.*

**Lemma 4.** *Given  $T \in D_{\text{Dehn}}$ , let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be given by:  $f(\alpha) = \rho_V^+(T + (0, \alpha))$ . Then  $f$  is a non-decreasing function of  $\alpha$ .*

For proofs of the above results, see [4]. To conclude we present a lemma from [5] similar to the main result of [18], which says that we do not need to think of general perturbations in this setting. Vertical translations are enough for most applications. In the following,  $T_\alpha(\phi, I) \stackrel{\text{def.}}{=} T(\phi, I) + (0, \alpha)$ .

**Lemma 5.** *Let  $T \in D_{\text{Dehn}}$  be such that  $\rho_V(T) = [\omega^-, \omega^+]$ . Suppose that by an arbitrarily  $C^1$ -small perturbation applied to  $T$ , we can change  $\omega^+$ , that is, there exists  $T^*$  arbitrarily  $C^1$ -close to  $T$ , such that  $\rho_V^+(T^*) \neq \omega^+$ . Then for any given  $\epsilon > 0$ , at least one of the following inequalities must hold:*

- 1)  $\rho_V^+(T_\epsilon) \neq \omega^+$ , or
- 2)  $\rho_V^+(T_{-\epsilon}) \neq \omega^+$ .

*Moreover, given  $T \in D_{\text{Dehn}}$  with  $\rho_V(T) = [\omega^-, \omega^+]$ , there exists a neighborhood  $U \subset D_{\text{Dehn}}$  such that for any  $T^* \in U$ ,  $\rho_V^+(T^*) = \omega^+$ , if and only if,  $\exists \epsilon > 0$  such that  $\rho_V^+(T_\alpha) = \omega^+$ , for all  $\alpha \in [-\epsilon, \epsilon]$ .*

**2.3. On the dynamics near fixed points of analytic diffeomorphisms of the plane.** The dynamics near singularities of analytic vector fields of the plane is very well understood (see for instance [11]). It can be proved that, if the singularity is not a focus or a center, then the dynamics can be obtained from a finite number of sectors, glued in an adequate way. Topologically, these sectors can be classified in 4 types: elliptic, hyperbolic, expanding and attracting. Dumortier et al. studied this problem for planar diffeomorphisms near fixed points, see [12].

The situation we want to understand in this section is the following: Is there a topological picture of the dynamics near an index zero isolated fixed point of an analytic area-preserving diffeomorphism of the plane?

It turns out that the area-preservation together with the zero index hypotheses imply that the eigenvalues of the derivative of the diffeomorphism at the fixed point are both equal to 1. The area-preservation implies that in a sufficiently small neighborhood of the fixed point, the diffeomorphism is the time one mapping of a formal vector field (defined by a formal series), see [20]. So we are able to apply results from [12], which say that at least in a topological sense, the dynamics near the fixed point can be obtained as in the vector field setting, gluing a finite number of sectors. As we are supposing that area is preserved by the diffeomorphism, there can not be elliptic, expanding and attracting sectors. As the topological index of the fixed point is zero, it can not be a center and so there must be exactly 2 hyperbolic sectors and the dynamics is topologically as in figure 1.

**3. Proof of the main theorem.** In this section we are going to prove the main theorem. Our proof is by contradiction. We suppose that there is a mapping  $T$  satisfying the theorem hypothesis and such that the following holds:

$$\text{For every } \alpha \neq 0, \rho_V^+(T_\alpha) \neq p/q, \text{ where } T_\alpha(\phi, I) = T(\phi, I) + (0, \alpha) \tag{4}$$

Using lemma 5, we get that theorem 1 is true if and only if the above assertion is impossible. Thus we are left to show that (4) is impossible. As  $\rho_V^- < p/q$  and




FIGURE 1. Diagram showing the dynamics near a fixed point of zero index

$\rho_V^+(T_\alpha) > p/q$  for all  $\alpha > 0$  (see lemma 4), we get from theorems 3 and 4 that  $\bar{T}_0$  has  $q$ -periodic points with  $\rho_V = p/q$ .

Lemma 4 again implies that for any  $\alpha < 0$ ,  $\rho_V^+(T_\alpha) < p/q \Rightarrow$  the  $q$ -periodic points of  $\bar{T}_0$  with  $\rho_V = p/q$  are degenerate. Moreover, in the following we are going to prove that they are finite. As  $T$  is an analytic mapping, the same argument used in the proof of theorem 3 of [3] implies that if the  $q$ -periodic points of  $\bar{T}$  with  $\rho_V = p/q$  are infinite, then there exists a simple closed curve  $\gamma \subset S^1 \times \mathbb{R}$  such that for all  $x \in \gamma$ ,  $T^q(x) = x + (0, p)$ . Clearly  $\gamma$  belongs to  $K(s, q)$ , for a certain  $s \in \mathbb{Z}$ . And this curve can not be homotopically non-trivial, because  $\rho_V^-(T) < p/q$ . To see this, note that if  $\gamma$  is homotopically non-trivial, and  $T^q(\gamma) = \gamma + (0, p)$ , as  $T$  induces a mapping on the torus homotopic to the Dehn twist, then the following is true:

$$T^q(\gamma + (0, m)) = \gamma + (0, m) + (0, p), \text{ for all } m \in \mathbb{Z}$$

As  $\rho_V^-(T) = \delta < p/q$ , there exists a point  $z \in S^1 \times \mathbb{R}$  such that  $\rho_V(z) < p/q$ . Now choose  $m \in \mathbb{Z}$  such that  $z$  is above  $\gamma + (0, m) \Rightarrow T^{n \cdot q}(z)$  is above  $\gamma + (0, m) + (0, n \cdot p)$  for all  $n > 0$ , which contradicts the fact that  $\rho_V(z) < p/q$ . So  $\gamma$  is homotopically trivial and thus crosses some vertical twice, at say,  $x$  and  $x'$  ( $p_1(x) = p_1(x') = \phi$ ). Suppose that  $p_2(x) < p_2(x')$ . The definition of  $\mu^-$  and lemma 3, which says that  $T^q(\phi, \mu^-(\phi)) = (\phi, \nu^+(\phi))$ , imply that  $\mu^-(\phi) < p_2(x)$ . Thus we get that  $\nu^+(\phi) - \mu^-(\phi) > p + (p_2(x') - p_2(x))$ , something that contradicts the fact that  $\rho_V^+(T_\alpha) < p/q$  for all  $\alpha < 0$ , because  $C(s, q)$  and  $T^q(C(s, q))$  are upper semi-continuous functions of  $T$  in the Hausdorff topology (see [18]). So the  $q$ -periodic points of  $\bar{T}$  with  $\rho_V = p/q$  are finite.

Let us denote them

$$\{\bar{p}_0^1, \bar{p}_1^1, \dots, \bar{p}_{q-1}^1\}, \{\bar{p}_0^2, \bar{p}_1^2, \dots, \bar{p}_{q-1}^2\}, \dots, \{\bar{p}_0^N, \bar{p}_1^N, \dots, \bar{p}_{q-1}^N\}, \tag{5}$$

where  $\bar{p}_j^i = (\bar{\phi}_j^i, \bar{I}_j^i)$  for  $i \in \{1, 2, \dots, N\}$  and  $j \in \{0, 1, \dots, q - 1\}$ .

Define  $F_\alpha(\phi, I) \stackrel{\text{def.}}{=} (T_\alpha)^q(\phi, I) - (0, p)$ . Clearly  $F_\alpha$  induces a torus diffeomorphism  $\bar{F}_\alpha$  with  $\rho_V^-(F_\alpha) < 0 < \rho_V^+(F_\alpha)$ , for all sufficiently small  $\alpha > 0$  and all the points in (5) are fixed points for  $\bar{F}_0$  with zero vertical rotation number.

Now for each point of (5) let us choose open neighborhoods  $\bar{W}_k$ ,  $k = 1, 2, \dots, N \cdot q$ , such that:

1.  $\text{closure}(\bar{F}_0(\bar{W}_k) \cup \bar{W}_k \cup \bar{F}_0^{-1}(\bar{W}_k)) \cap \text{closure}(\bar{W}_l) = \emptyset$ , for all  $k \neq l$ ,  $k, l \in \{1, 2, \dots, N \cdot q\}$ .
2. As each  $\bar{p}_j^i$  is fixed for  $\bar{F}_0$  and has zero topological index, the dynamics inside each  $\bar{W}_k$  is as in figure 1. So  $\bar{W}_k \supset \bar{E}n_{k,0} \cup \bar{E}x_{k,0}$ , where  $\bar{E}n_{k,0}$  and  $\bar{E}x_{k,0}$  are two open connected sets, free under  $F_0$  (disjoint from their images), such

that every time the orbit of a point enters in  $\overline{W}_k$ , it must enter through  $\overline{E}n_{k,0}$  and leave through  $\overline{E}x_{k,0}$ . Moreover, if we choose  $\overline{W}_k$  properly, we can suppose that  $\text{closure}(\overline{E}n_{k,0}) \cap \text{closure}(\overline{E}x_{k,0}) = \emptyset$  and that there exists a point  $\overline{z}_k \in B_r(\overline{z}_k) \subset \overline{E}n_{k,0}$ , where  $r > 0$  is the same for all  $k \in \{1, 2, \dots, N.q\}$ , such  $\overline{F}_0(\overline{z}_k) \in B_s(\overline{F}_0(\overline{z}_k)) \subset \overline{E}x_{k,0}$ , also for some fixed  $s > 0$ , see figure 2. The reason why such a choice is possible is the following. Just to have a mental picture of what is happening, suppose that the invariant lines in figure 1 are horizontal. Start with a small vertical segment  $seg_k$  through the fixed point and consider the following iterates of it:  $\overline{F}_0^{-2}(seg_k)$ ,  $\overline{F}_0^{-1}(seg_k)$ ,  $seg_k$  and  $\overline{F}_0(seg_k)$ . Clearly, all these arcs end at the same horizontal lines. Now, take a simple arc,  $Arc_{left,k}$ , whose end points are in these horizontal lines, very close to the end points of  $\overline{F}_0^{-2}(seg_k)$ , but to the right of them. Moreover, choose the arc in a way that its first iterate is on the left side of  $seg_k$ . Now, choose another simple arc,  $Arc_{right,k}$ , in a way that it is on the right side of  $\overline{F}_0(seg_k)$ , but with the same end points.  $\overline{W}_k$  is obtained by connecting the appropriate end points of  $Arc_{right,k}$  and  $Arc_{left,k}$  by horizontal segments. It is not difficult to see that this construction has the required properties.




FIGURE 2. Diagram showing the sets  $\overline{W}_k \supset \overline{E}n_{k,0} \cup \overline{E}x_{k,0} \cup \overline{B}_k$

If we choose sufficiently small neighborhoods  $\overline{W}_k$ , then there exists  $\alpha_1 > 0$ , such that for  $0 \leq \alpha \leq \alpha_1$ ,  $\overline{F}_\alpha$  has periodic orbits with negative vertical rotation number, disjoint from  $\cup_{k=1}^{N.q} \overline{W}_k$ . This follows from  $\rho_V^-(F_0) < 0 = \rho_V^+(F_0)$  and the continuity of these functions with respect to  $\alpha$ .

Let  $\overline{B}_k \subset \overline{W}_k$  be a open neighborhood of the  $\overline{F}_0$ -fixed point which is inside  $\overline{W}_k$  that satisfies:

$$\overline{W}_k \supset \text{closure}(\overline{F}_0^{-2}(\overline{B}_k) \cup \overline{F}_0^{-1}(\overline{B}_k) \cup \overline{B}_k \cup \overline{F}_0(\overline{B}_k) \cup \overline{F}_0^2(\overline{B}_k)) \tag{6}$$

The above condition implies that

$$\text{closure}(\overline{B}_k) \cap \text{closure}(\overline{F}_0(\overline{E}n_{k,0}) \cup \overline{E}n_{k,0} \cup \overline{E}x_{k,0}) = \emptyset. \tag{7}$$

Following figure 2, let us denote the boundary of  $\overline{W}_k$  as follows:  $\partial \overline{W}_k = \partial l_k \cup \partial b_k \cup \partial r_k \cup \partial t_k$ . Let  $\overline{V}_{b,k}, \overline{V}_{t,k} \subset \overline{W}_k$  be two small open neighborhoods, respectively

of  $\partial b_k$  and  $\partial t_k$  such that the following conditions hold ( $\eta > 0$  is a sufficiently small number and  $B_\eta(\bar{V}_{b,k} \cup \bar{V}_{t,k})$  is the  $\eta$ -neighborhood of  $\bar{V}_{b,k} \cup \bar{V}_{t,k}$ ):

- 1)  $\{\bar{z}_k, \bar{F}_0(\bar{z}_k)\} \cap \text{closure}(\bar{V}_{b,k} \cup \bar{V}_{t,k}) = \emptyset$
- 2) there exists  $L > 0$  such that for all  $\bar{x} \in B_\eta(\bar{V}_{b,k} \cup \bar{V}_{t,k})$ ,  
for some  $l = l(\bar{x}) < L$ ,  $\bar{F}_0^l(\bar{x}) \notin \bar{W}_k$  and  
for  $0 \leq i \leq l$   $\bar{F}_0^i(\bar{x}) \notin \text{closure}(\bar{B}_k)$  (8)
- 3) there exists  $M > 0$  such that for all  $\bar{x} \in B_\eta(\bar{V}_{b,k} \cup \bar{V}_{t,k})$ ,  
for some  $m = m(\bar{x}) < M$ ,  $\bar{F}_0^{-m}(\bar{x}) \notin \bar{W}_k$  and  
for  $0 \leq i \leq m$ ,  $\bar{F}_0^{-i}(\bar{x}) \notin \text{closure}(\bar{B}_k)$

Now choose  $\alpha_2 > 0$  such that for  $0 \leq \alpha \leq \alpha_2$ ,

1. All fixed points of  $\bar{F}_\alpha$  with zero vertical rotation number are contained in  $\cup_{k=1}^{N,q} \bar{B}_k$  and there is an isotopy supported in  $\cup_{k=1}^{N,q} \bar{B}_k$  between  $\bar{F}_\alpha$  and a homeomorphism without fixed points of zero vertical rotation number. To construct this isotopy, one just needs to note that if  $\alpha_2 > 0$  is small enough, then for  $0 \leq \alpha \leq \alpha_2$ , the image of  $\cup_{k=1}^{N,q} \bar{B}_k$  under  $\bar{F}_\alpha$  (which is still an analytic mapping!) is very similar to the case when  $\alpha = 0$ . So, as in that case the fixed points are removable by local isotopies, the same happens for small  $\alpha$ 's (see for instance theorem 2.4 of [8], theorem 2.1 of [23] and the main theorem of [22])
2. The three conditions in (8) hold when, instead of  $\bar{F}_0$ , we use  $\bar{F}_\alpha$ .
3.  $\bar{F}_\alpha^{\pm 1}(\partial b_k) \cap \bar{W}_k \subset \bar{V}_{b,k}$  and  $\bar{F}_\alpha^{\pm 1}(\partial t_k) \cap \bar{W}_k \subset \bar{V}_{t,k}$
4. Condition (6) holds when instead of  $\bar{F}_0$  we use  $\bar{F}_\alpha$ . As before, this implies that  $\text{closure}(\bar{B}_k) \cap \text{closure}(\bar{F}_\alpha(\bar{E}n_{k,\alpha}) \cup \bar{E}n_{k,\alpha} \cup \bar{E}x_{k,\alpha}) = \emptyset$ , where  $\bar{E}n_{k,\alpha}$  and  $\bar{E}x_{k,\alpha}$  are the analogues of  $\bar{E}n_{k,0}$  and  $\bar{E}x_{k,0}$  for  $\alpha > 0$ , that is, every time the orbit of a point under  $\bar{F}_\alpha$  enters in  $\bar{W}_k$ , it must enter through  $\bar{E}n_{k,\alpha}$  and leave through  $\bar{E}x_{k,\alpha}$
5.  $\bar{z}_k \in \bar{E}n_{k,\alpha}$  and  $\bar{F}_\alpha(\bar{z}_k) \in \bar{E}x_{k,\alpha}$
6. If we denote the connected component of  $\bar{E}n_{k,\alpha} \setminus \text{closure}(\bar{V}_{b,k} \cup \bar{V}_{t,k})$  whose boundary intersects  $\partial \bar{W}_k$  by  $\bar{E}n'_{k,\alpha}$  and, in the same way, the connected component of  $\bar{E}x_{k,\alpha} \setminus \text{closure}(\bar{V}_{b,k} \cup \bar{V}_{t,k})$  whose boundary intersects  $\partial \bar{W}_k$  is denoted by  $\bar{E}x'_{k,\alpha}$ , then  $\bar{E}n_{k,\alpha} \subset \bar{E}n'_{k,\alpha} \cup B_\eta(\bar{V}_{b,k} \cup \bar{V}_{t,k})$  and  $\bar{E}x_{k,\alpha} \subset \bar{E}x'_{k,\alpha} \cup B_\eta(\bar{V}_{b,k} \cup \bar{V}_{t,k})$ . Moreover,  $\text{closure}(\bar{E}n'_{k,\alpha}) \cap \text{closure}(\bar{E}x'_{k,\alpha}) = \emptyset$ , see figure 3
7.  $\text{closure}(\bar{F}_\alpha(\bar{W}_k) \cup \bar{F}_\alpha^{-1}(\bar{W}_k)) \cap \text{closure}(\bar{W}_l) = \emptyset$ , for all  $k \neq l$
8.  $\bar{z}_k \in \bar{E}n'_{k,\alpha}$  and  $\bar{F}_\alpha(\bar{z}_k) \in \bar{E}x'_{k,\alpha}$

A simple continuity argument implies that all the above conditions are satisfied for sufficiently small  $\alpha > 0$ .

Now fix  $0 < \alpha < \min\{\alpha_1, \alpha_2\}$ . Our hypothesis imply that  $\rho_V^-(F_\alpha) < 0 < \rho_V^+(F_\alpha)$ . From the choice of  $\alpha_1 > 0$ , there exists  $\bar{x}_- \in \mathbb{T}^2$  and  $n > 0$  such that  $(\bar{F}_\alpha)^n(\bar{x}_-) = \bar{x}_-$ ,  $\rho_V(\bar{x}_-) < 0$  and the orbit of  $\bar{x}_-$  is disjoint from  $\cup_{k=1}^{N,q} \bar{W}_k$ . Let  $m > 0$  be an integer such that  $0 < 1/m < \rho_V^+(F_\alpha)$  and let  $\bar{x}_+ \in \mathbb{T}^2$  (the existence of  $\bar{x}_+$  follows from theorem 3) be such that  $(\bar{F}_\alpha)^m(\bar{x}_+) = \bar{x}_+$  and  $\rho_V(\bar{x}_+) = 1/m$ . If the orbit of  $\bar{x}_+$  is disjoint from  $\cup_{k=1}^{N,q} \bar{B}_k$ , then we are done, because we can destroy the fixed




FIGURE 3. Diagram showing the sets  $\bar{E}n'_{k,\alpha} \cup \bar{E}x'_{k,\alpha}$

points of  $\bar{F}_\alpha$  by a perturbation supported in  $\cup_{k=1}^{N.q} \bar{B}_k$ , something that contradicts theorem 5.

So, suppose that  $Orbit(\bar{x}_+) = \{\bar{x}_+, \bar{F}_\alpha(\bar{x}_+), (\bar{F}_\alpha)^2(\bar{x}_+), \dots, (\bar{F}_\alpha)^{m-1}(\bar{x}_+)\}$  intersects  $\cup_{k=1}^{N.q} \bar{B}_k$ . From the fact that the three conditions in (8) hold for  $\bar{F}_\alpha$ , without loss of generality we can suppose that  $\bar{x}_+ \in \bar{E}x'_{1,\alpha} \subset \bar{W}_1$ . As  $(\bar{F}_\alpha)^m(\bar{x}_+) = \bar{x}_+$ , there exists  $m' < m$  such that  $(\bar{F}_\alpha)^{m'}(\bar{x}_+) \in \bar{E}n'_{1,\alpha} \subset \bar{W}_1$ . Now let

$$\{x_+, W_1, W_2, \dots, W_{N.q}\} \subset S^1 \times [0, 1]$$

be lifts of  $\bar{x}_+$  and  $\bar{W}_1, \bar{W}_2, \dots, \bar{W}_{N.q}$  to the cylinder. From the choice of  $\bar{x}_+$  we get that  $(F_\alpha)^m(x_+) = x_+ + (0, 1)$ . If we consider all the intermediate pieces of the orbit of  $x_+$ , then the following observation is true:

**Observation 1:** There exists  $r_f = (F_\alpha)^i(x_+)$  and  $r_l = (F_\alpha)^j(x_+)$ , for some  $0 \leq i < j \leq m'$  such that  $r_f \in Ex'_{k_1,\alpha} + (0, s) \subset W_{k_1} + (0, s)$  and  $r_l \in En'_{k_1,\alpha} + (0, s+l) \subset W_{k_1} + (0, s+l)$ , for some  $1 \leq k_1 \leq N.q$  and  $s, l \in \mathbb{Z}$ , with  $l > 0$ . And when for 2 points  $z = (F_\alpha)^d(x_+)$ ,  $y = (F_\alpha)^e(x_+)$ ,  $i \leq d < e < j$ ,  $pro(z)$  and  $pro(y)$  belong to the same  $\bar{B}_t$ , then  $z \in B_t + (0, m)$  and  $y \in B_t + (0, m+a)$ , for  $m, a \in \mathbb{Z}$  with  $a \leq 0$ . Moreover,  $\{\bar{F}_\alpha^i(\bar{x}_+), \dots, (\bar{F}_\alpha)^j(\bar{x}_+)\} \cap \bar{B}_{k_1} = \emptyset$ .

In order to find points  $r_f$  and  $r_l$  as above, we just have to start with  $x_+ \in Ex'_{1,\alpha} \subset W_1$  and  $(F_\alpha)^{m'}(x_+) \in En'_{1,\alpha} + (0, 1) \subset W_1 + (0, 1)$  and check if the conditions in the observation are satisfied for the intermediate points of the orbit. If they are not, then we pick the convenient subset of the orbit from  $x_+$  to  $(F_\alpha)^{m'}(x_+)$  and check again. After a finite number of attempts,  $r_f$  and  $r_l$  are found.

Let us denote the orbit segment from  $\bar{r}_f$  to  $\bar{r}_l$  by

$$\bar{r}_f = \bar{x}_0, \bar{x}_1, \dots, \bar{x}_{j-i} = \bar{r}_l, \text{ where } \bar{x}_n = (\bar{F}_\alpha)^n(\bar{r}_f) \tag{9}$$

and suppose that it intersects  $\cup_{k=1}^{N.q} \bar{B}_k$  in  $\bar{B}_{k_2}, \bar{B}_{k_3}, \dots, \bar{B}_{k_L}$ . From the fact that expression (8) is true also for  $\bar{F}_\alpha$ , in each  $\bar{W}_{km}$ ,  $2 \leq m \leq L$ , there are points  $\bar{x}_{km-first} \in \bar{E}n'_{km,\alpha} \subset \bar{W}_{km}$  and  $\bar{x}_{km-last} \in \bar{E}x'_{km,\alpha} \subset \bar{W}_{km}$ , defined as follows:

- $\bar{x}_{km\_first}$  is the first point of the sequence (9) such that  $(\bar{F}_\alpha)^n(\bar{x}_{km\_first})$  belongs to  $\bar{W}_{km}$  for  $n = 0, 1, \dots, n_{max}$ ,  $(\bar{F}_\alpha)^{n_{crit}}(\bar{x}_{km\_first}) \in \bar{B}_{km}$  for some  $0 < n_{crit} < n_{max}$  and of course  $(\bar{F}_\alpha)^{n_{max}}(\bar{x}_{km\_first}) \in \bar{E}x'_{km,\alpha}$
- $\bar{x}_{km\_last}$  is the last point of the sequence (9) such that  $(\bar{F}_\alpha)^{-l}(\bar{x}_{km\_last})$  belongs to  $\bar{W}_{km}$  for  $l = 0, 1, \dots, l_{max}$ ,  $(\bar{F}_\alpha)^{-l_{crit}}(\bar{x}_{km\_last}) \in \bar{B}_{km}$  for some  $0 < l_{crit} < l_{max}$  and of course  $(\bar{F}_\alpha)^{-l_{max}}(\bar{x}_{km\_last}) \in \bar{E}n'_{km,\alpha}$

To understand the above definitions, one just has to remember that every time the orbit of a point falls into some  $\bar{B}_k$ , then it must have entered  $\bar{W}_k$  through  $\bar{E}n'_{k,\alpha}$  and it will leave  $\bar{W}_k$  through  $\bar{E}x'_{k,\alpha}$ .

The next step is to perform 2 deformations in each  $\bar{W}_{km}$  ( $m \geq 1$ ), one supported in  $\bar{E}n'_{km,\alpha}$  and the other supported in  $\bar{E}x'_{km,\alpha}$ . For  $2 \leq m \leq L$ , let  $\mu_{f\_km} \subset \bar{E}n'_{km,\alpha}$  be a simple arc whose endpoints are  $\bar{x}_{km\_first}$  and  $\bar{z}_{km}$ , such that, apart from  $\bar{x}_{km\_first}$ , it does not contain any other point of sequence (9). Let  $V_{f\_km} \subset \bar{E}n'_{km,\alpha}$  be an open neighborhood of  $\mu_{f\_km}$  which also does not contain any other point of sequence (9). In the same way, let  $\mu_{l\_km} \subset \bar{E}x'_{km,\alpha}$  be a simple arc whose endpoints are  $\bar{x}_{km\_last}$  and  $\bar{F}_\alpha(\bar{z}_{km})$ , such that, apart from  $\bar{x}_{km\_last}$ , it does not contain any other point of sequence (9). Let  $V_{l\_km} \subset \bar{E}x'_{km,\alpha}$  be an open neighborhood of  $\mu_{l\_km}$  which also does not contain any other point of sequence (9). As  $\bar{E}n'_{km,\alpha}$  and  $\bar{E}x'_{km,\alpha}$  are disjoint and free under  $\bar{F}_\alpha$ , the same is true for  $V_{f\_km}$  and  $V_{l\_km}$ . So, let us define the following homeomorphisms  $\bar{h}_{V_{f\_km}} : T^2 \rightarrow T^2$  and  $\bar{h}_{V_{l\_km}} : T^2 \rightarrow T^2$ :

$$\begin{aligned} \bar{c}h_{V_{f\_km}} |_{(V_{f\_km})^c} &= id \\ \bar{h}_{V_{f\_km}}(\bar{x}_{km\_first}) &= \bar{z}_{km} \\ &\text{and} \\ \bar{h}_{V_{l\_km}} |_{(V_{l\_km})^c} &= id \\ \bar{h}_{V_{l\_km}}(\bar{F}_\alpha(\bar{z}_{km})) &= \bar{x}_{km\_last} \end{aligned} \tag{10}$$

For  $m = 1$ , we perform an analogous construction. Let  $\mu_{f\_k1} \subset \bar{E}n'_{k1,\alpha}$  be a simple arc whose endpoints are  $\bar{r}_l$  and  $\bar{z}_{k1}$ , which, apart from  $\bar{r}_l$ , avoids all points of sequence (9). Let  $V_{f\_k1} \subset \bar{E}n'_{k1,\alpha}$  be an open neighborhood of  $\mu_{f\_k1}$  which also does not contain any other point of sequence (9). In the same way, let  $\mu_{l\_k1} \subset \bar{E}x'_{k1,\alpha}$  be a simple arc whose endpoints are  $\bar{r}_f$  and  $\bar{F}_\alpha(\bar{z}_{k1})$ , which, apart from  $\bar{r}_f$ , avoids all points of sequence (9). Let  $V_{l\_k1} \subset \bar{E}x'_{k1,\alpha}$  be an open neighborhood of  $\mu_{l\_k1}$  which also does not contain any other point of sequence (9). Then we define:

$$\begin{aligned} \bar{c}h_{V_{f\_k1}} |_{(V_{f\_k1})^c} &= id \\ \bar{h}_{V_{f\_k1}}(\bar{r}_l) &= \bar{z}_{k1} \\ &\text{and} \\ \bar{h}_{V_{l\_k1}} |_{(V_{l\_k1})^c} &= id \\ \bar{h}_{V_{l\_k1}}(\bar{F}_\alpha(\bar{z}_{k1})) &= \bar{r}_f \end{aligned} \tag{11}$$

Denote by  $\bar{h} : T^2 \rightarrow T^2$  the homeomorphism which agrees with all the  $\bar{h}_{V_{f\_km}}$  and  $\bar{h}_{V_{l\_km}}$ ,  $1 \leq m \leq L$ . From the choice of  $\alpha_2 > 0$ , we know that there is an isotopy supported in  $\cup_{k=1}^{N,q} \bar{B}_k$  between  $\bar{F}_\alpha$  and a homeomorphism without fixed points of

zero vertical rotation number. Let us denote the final mapping of this isotopy by  $\overline{F}_\alpha^*$ . By construction  $\overline{F}_\alpha^*$  coincides with  $\overline{F}_\alpha$  outside  $\cup_{k=1}^{N,q} \overline{B}_k$ , so we get that the mapping  $\overline{G} \stackrel{def}{=} \overline{h} \circ \overline{F}_\alpha^*$  has a periodic point with positive vertical rotation number, that comes from the orbit segment between  $\overline{r}_f$  and  $\overline{r}_l$ , see Observation 1. The deformation  $\overline{h}$  is supported in a disjoint union of free sets for  $\overline{F}_\alpha^*$ , so as  $\overline{F}_\alpha^*$  has no fixed points of zero vertical rotation number, the same is true for  $\overline{G}$ .

As  $\overline{G}$  coincides with  $\overline{F}_\alpha$  outside  $\cup_{k=1}^{N,q} \overline{W}_k$ ,  $\overline{G}$  also has periodic points with negative vertical rotation number. But this contradicts a generalization of some results of [13] to this context. To be more precise, it contradicts the next theorem due to H.E. Doeff (see [10], theorem 5.3).

**Theorem 5.** : *If  $\overline{G} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$  is a homeomorphism homotopic to  $(\phi, I) \rightarrow (\phi + n.I \text{ mod } 1, I \text{ mod } 1)$ , for a certain  $n \in \mathbb{N}^*$  and there are points  $\overline{x}_+, \overline{x}_- \in \mathbb{T}^2$  such that  $\rho_V(\overline{x}_-) < 0 < \rho_V(\overline{x}_+)$ , then  $\overline{G}$  has fixed points of zero vertical rotation number.*

So it is not possible that  $\rho_V^+(T_\alpha) \neq \frac{p}{q}$  for all  $\alpha \neq 0$  and the proof is complete.  $\square$

**4. Proof of theorem 2.** As  $T \in D_{\text{Dehn}}$  is area-preserving, it makes sense to compute the vertical rotation number of the Lebesgue measure:

$$\rho_V(\text{Leb}, T) = \int_{S^1 \times [0,1]} [p_2 \circ T(\phi, I) - I] d\phi dI$$

As  $\rho_V(T) = \{\frac{p}{q}\}$ , the Birkhoff ergodic theorem implies that  $\rho_V(\text{Leb}, T) = \frac{p}{q}$ . But then,  $\rho_V(\text{Leb}, T_\alpha) = \frac{p}{q} + \alpha$ , for any  $\alpha \in \mathbb{R}$ . So, as  $\rho_V^-(T_\alpha) \leq \rho_V(\text{Leb}, T_\alpha) \leq \rho_V^+(T_\alpha)$ , the theorem is proved.  $\square$

**Acknowledgements:** I would like to thank the referee for a very careful reading of the paper, for all his suggestions and for his patience in reading the second version of the proof of the main theorem.

## REFERENCES

- [1] Addas-Zanata S. (2000): *On the dynamics of twist maps of the torus*. Doctoral Thesis. In Portuguese. IMEUSP.
- [2] Addas-Zanata S. (2002): On the existence of a new type of periodic and quasi-periodic orbits for twist maps of the torus. *Nonlinearity* **15** (5), 1399-1416
- [3] Addas-Zanata S. and Grotta-Ragazzo C. (2002): On the stability of some periodic orbits of a new type for twist maps. *Nonlinearity* **15** (5), 1385-1397
- [4] Addas-Zanata S. (2005): A note on a standard family of twist maps. to appear in *Qual. Th. of Dynam. Sys.*
- [5] Addas-Zanata S. (2005): On properties of the vertical rotation interval for twist mappings. *Erg. Th. & Dynam. Sys.* **25**, 641-660
- [6] Addas-Zanata S. (2003): On properties of the vertical rotation interval for twist mappings II. *Qual. Th. of Dynam. Sys* **4**, 125-137
- [7] Addas-Zanata S. (2004): Instability for the rotation set of homeomorphisms of the torus homotopic to the identity. *Erg. Th. & Dynam. Sys* **24**, 319-328
- [8] Bonino M. (2001): A dynamical property for planar homeomorphisms and an application to the problem of canonical position around an isolated fixed point. *Topology* **40**, 1241-1257
- [9] Boyland P. (1986): Bifurcations of circle maps: Arnol'd tongues, bistability and rotation intervals. *Comm. Math. Phys.* **106**, 353-381
- [10] Doeff H.E. (1997): Rotation Measures for homeomorphisms of the torus homotopic to a Dehn twist. *Ergod. Th. & Dynam. Sys.* **17**, 575-591
- [11] Dumortier F. (1977): Singularities of vector fields on the plane. *J. Diff. Eq.* **23**, 53-106

- [12] Dumortier F., Rodrigues P. and Roussarie R. (1981): Germs of Diffeomorphisms in the Plane. *Lecture Notes in Mathematics* **902**
- [13] Franks J. (1989): Realizing rotation vectors for torus homeomorphisms. *Trans. Amer. Math. Soc.* **311**, 107–115.
- [14] Ito R. (1981): Rotation sets are closed. *Math. Proc. Camb. Phil. Soc.* **89**, 107-11
- [15] Katok A. and Hasselblatt B. (1995): *Introduction to Modern Theory of Dynamical Systems*. Cambridge University Press
- [16] Le Calvez P. (1986): Existence d'orbites quasi-périodiques dans les attracteurs de Birkhoff. *Comm. Math. Phys.* **106**, 383-394
- [17] Le Calvez P. (1991): Propriétés Dynamiques des Difféomorphismes de L'Anneau et du Tore. *Astérisque* **204**
- [18] Le Calvez P. (1995): Construction d'orbites périodiques par perturbation d'un difféomorphisme de l'anneau déviant la verticale. *C. R. Acad. Sci. Paris*, t **321**, p. 463-468
- [19] Milnor J. (1968): *Singular Points of Complex Hypersurfaces*. Annals of Mathematics Studies no 61. Princeton University Press
- [20] Moser J. (1968): Lectures on Hamiltonian systems. *Mem. Amer. Math. Soc.* **81**, 60 pp
- [21] Newhouse S., Palis J. and Takens F. (1983): Bifurcations and stability of families of diffeomorphisms. *Inst. Hautes Études Sci. Publ. Math.* **57**, 5-71
- [22] Schmitt B. (1975): Sur les plongements, admettant zéro ou un point fixé, du disque dans le plan. *Topology* **14**, 357–365.
- [23] Slaminka E. (1993): Removing index 0 fixed points for area preserving maps of two-manifolds. *Trans. Amer. Math. Soc.* **340**, 429-444

Received November 2004; revised October 2005.

*E-mail address:* sazanata@ime.usp.br