



**IDENTIFICATION AND BIOSYNTHESIS OF PEPTAIBOLS
TRICHOKONINS VI AND VIII WITH ANTIPLASMODIAL ACTIVITY**

Matheus Gotha^{1*}, Ariane F. Bertonha², David E. Williams³, Karen J. Nicacio⁴, Marcelo R. de Amorim¹, Cauê A. W. Zuccarino¹, Sydney M. Schoellhorn⁵, Anna Caroline C. Aguiar⁶, Lamonielli F. Michaliski¹, Lara D. Sette⁷, Antônio G. Ferreira⁸, Raymond J. Andersen³, Rafael V. C. Guido⁹, Roberto G. S. Berlinck¹

matheusgotha@alumni.usp.br, rgsberlinck@iqsc.usp.br

1-São Carlos Institute of Chemistry, University of São Paulo, 13560-970, São Carlos, Brazil. 2-LNBio, Brazilian Center for Research in Energy and Materials, 13083-970, Campinas, Brazil. 3-Departments of Chemistry and EOAS, UBC, 2036 Main Mall, Vancouver, Canada. 4-UFMT, 78060-900, Cuiabá, MT, Brazil. 5-Department of Chemistry and BioDiscovery Institute, UNT, Denton, TX, 76203, USA. 6-DMIP, Unifesp, 04023-062, São Paulo, SP, Brazil. 7-Departamento de Biologia Geral e Aplicada, UNESP, IB, Rio Claro, Brazil. 8-Departamento de Química, UFSCar, 13565-905, São Carlos, SP, Brazil. 9-São Carlos Institute of Physics, USP, 13566-590, São Carlos, SP, Brazil.

The fungus genus *Trichoderma* is found in a wide range of habitats, with more than 200 described species.¹ Secondary metabolites produced by *Trichoderma* spp. have mainly been used commercially in biocontrol.² Modified peptides known as peptaibols are commonly produced by *Trichoderma* species. Typical features of peptaibols include an acyl fragment connected at the N-terminal, incorporation of α -aminoisobutyric (Aib) residues, amino alcohol of the reduced amino acid at C-terminal, and presence of other non-proteinogenic amino acids. *Trichoderma* sp. L2-2 strain was obtained from a lichen sample collected at Admiralty Bay in Antarctica. Cultures of *Trichoderma* sp. L2-2 yielded trichokonins VI and VIII after several separation steps. Both peptaibols presented 20 amino acid residues and were completely identified by NMR, HRMS, MS/MS and dichroism circular spectra analyses. The full genome sequencing and analysis of *Trichoderma* sp. L2-2 enabled the identification of a peptaibol biosynthetic gene cluster and the proposal for a biosynthetic assembly of trichokonins VI and VIII. Trichokonins VI and VIII also exhibited antiplasmodial activities at the sub-micromolar range against *Plasmodium falciparum*.

1- Dou, K. *et al.* Appl Environ Microbiol, 86 (18), 2020.

2- Pedrero-Méndez, A. *et al.* Microbiol. Res. 2025, 296, 128153.

Keywords: Trichokonin VI and VIII, Peptaibols, ¹H and ¹³C NMR analyzes, Secondary metabolite, Antiplasmodial

