

Nº 35

On the preservation of elementary equivalence
and embedding by filtered powers and structures
of stable continuous functions

Francisco Miraglia Neto

Outubro 1981

ON THE PRESERVATION OF ELEMENTARY
EQUIVALENCE AND EMBEDDING BY BOUNDED FILTERED POWERS
AND STRUCTURES OF STABLE CONTINUOUS
FUNCTIONS (*)

Francisco Miraglia

All notation as well as definitions of the concepts mentioned in the title will be as in [M]. In that paper can also be found bibliographical references sustaining the usefulness of filtered powers and of structures of stable continuous functions. The author acknowledges helpful conversations, on the methods used here, with A.M. Sette. We recall definitions briefly.

If X is a topological space and M a set, the symbol $C(X, M)$ will denote the set of all continuous functions from X to M , where M is given the discrete topology. Whenever it appears this symbol will have this meaning and the structures mentioned in the title, in particular, will have as domain subsets of sets of this form. The symbol X indicates a topological space in all that follows.

If L is a first order language with equality, an L -structure and I a set, let $\Sigma = \langle \{F_i\}_{i \in I}, \{M_i\}_{i \in I} \rangle$ be a pair of families of subsets of X and M respectively, such that each F_i is closed in X and each M_i is a substructure of M . The symbol $C(X, M, \Sigma)$ denotes the substructure of

* Partially supported by CNPq grant 300.214/79.

$C(X, M)$, consisting of all continuous $f: X \rightarrow M$ such that for all $i \in I$ $f(F_i) \subseteq M_i$. $C(X, M; \Sigma)$ is called a *filtered power* of M by X .

Let G be a group operating by homeomorphisms on X and by automorphisms on an L -structure M . That is, we have group homomorphisms $g \in G \rightarrow g_* \in \text{Hom}_L(X)$ and $g \in G \rightarrow g^* \in \text{Aut}_L(M)$. The symbol $SC(X, M; G)$ denotes the substructure of $C(X, M)$ consisting of all continuous $X \xrightarrow{f} M$ such that for all $g \in G$ and all $x \in X$ $f(g_* x) = g^* f(x)$. $SC(X, M; G)$ is called the structure of *stable continuous functions* from X to M under the action of G .

In the rest of this note it will be assumed that X is boolean (compact, with a basis of clopens) and that G and I are finite.

If X is a boolean space, $B(X)$ denotes the boolean algebra of clopens in X . To each filtered power $C(X, M; \Sigma)$ we can associate an expansion of M , $\langle M; M_1, \dots, M_n \rangle \equiv M^\#$ and by Stone duality, the augmented boolean algebra $\langle B(X); I_1, \dots, I_n \rangle \equiv B_\#(X)$ where I_k is the ideal in $B(X)$ determined by the closed set $F_k \subseteq X$ (see [M] or [MR] for more details). We denote by L the expansion of L appropriate to structures of type $\langle M; M_1, \dots, M_n \rangle$. Thus $L^\#$ is obtained from L by adding n new unary relation symbols P_1, \dots, P_n . Similarly, to each structure of stable continuous functions $SC(X, M; G)$ we can associate an expansion of M , $M^* = \langle M, \{g^*\}_{g \in G} \rangle$ and an expansion $B_*(X) = \langle B(X) : \{g_*\}_{g \in G} \rangle$ of $B(X)$. Here we make

no distinction between the homeomorphism g_* of X and the automorphism induced by g_* in $B(X)$. L^* denotes the expansion of L , appropriate for structures of type M^* .

In all that follows X will be an arbitrary but fixed boolean space, F_1, \dots, F_n a fixed (but again arbitrary) family of closed subsets of X and G a finite group. If M_1^* and M_2^* are two expansions of L -structures M_1 and M_2 as described above, then $\Sigma_i = \langle \{F_1, \dots, F_n\}, \{M_{i1}, \dots, M_{in}\} \rangle$, $i = 1, 2$. We wish to present proofs of the following:

Theorem 1. If $M_1^* \underset{L}{\#} M_2^* \quad (M_1^* \underset{L}{\equiv} M_2^*)$ then $C(X, M_1; \Sigma_1)$ is an elementary L -substructure of $C(X, M_2; \Sigma_2)$ (resp. $C(X, M_1; \Sigma_1) \equiv_L C(X, M_2; \Sigma_2)$).

Theorem 2. Same as above, with M_i^* and $SC(X, M_i; G)$ $i = 1, 2$ in the place of M_i^* and $C(X, M_i, \Sigma_i)$ respectively.

For this purpose we will use the characterization, due to Fraïssé ([Fr]) of elementary equivalence through the existence of certain families of partial isomorphisms ("back and forth" techniques). All results we need are in [F], whose notation will be followed closely. In particular if M_1 and M_2 are structures, $P(M_1, M_2)$ will denote the set of partial isomorphisms from M_1 to M_2 .

We have the following

Fundamental Lemma : With notational conventions as above, there is a natural injection $C(X, P(M_1, M_2)) \xrightarrow{\Delta} P(C(X, M_1, M_2)) = T$ such that

a. If $S^\# = P(M_1^\#, M_2^\#)$, define $\Delta^\# : C(X, S^\#) \rightarrow T$ by
 $\Delta^\#(f) = \Delta(f)|_{\text{dom } \Delta(f) \cap C(X, M_1; \Sigma_1)}$. Then $\text{Im } \Delta^\# \subseteq P(C(X, M_1; \Sigma_1), C(X, M_2; \Sigma_2)) \cong T^\#$.

b. If $S^* = P(M_1^*, M_2^*)$, let G operate on it trivially and define $\Delta^* : SC(X, S^*; G) \rightarrow T$ by $\Delta^*(f) = \Delta(f)|_{\text{dom } \Delta(f) \cap SC(X, M_1; G)}$. Then $\text{Im } \Delta^* \subseteq P(SC(X, M_1; G), SC(X, M_2; G)) \cong T^*$.

c. Define, for $f, g \in C(X, P(M_1, M_2))$, $f \leq g$ if for all $x \in X$, $f(x) \subseteq g(x)$ (i.e., $\text{dom } f(x) \subseteq \text{dom } g(x)$ and $g(x)|_{\text{dom } f(x)} = f(x)$). Then, if $f \leq g$, $\Delta(f) \subseteq \Delta(g)$.

d. If $(S_n)_{n \in \omega}$ has the back and forth property in $P(M_1, M_2)$ (as in definition 1.4, p. 252 of [F]), then $(\Delta(C(X, S_n)))_{n \in \omega}$ has the same property in T .

e. If $(S_n)_{n \in \omega}$ has the back and forth property in $P(M_1^*, M_2^*)$, let $S_n^* = S_n|_G$. Then $(\Delta^*(SC(X, S_n^*; G)))_{n \in \omega}$ has the same property in T^* .

Proof. Given $f : X \rightarrow P(M_1, M_2)$ put $\text{dom } \Delta(f) = \{\alpha \in C(X, M_1) : \forall x \in X \alpha(x) \in \text{dom } f(x)\}$. Then define for $\alpha \in \text{dom } \Delta(f)$

$$\Delta(f)(\alpha)(x) = f(x)(\alpha(x)).$$

For notational simplicity if $X \xrightarrow{h} A$ is constant on a clopen $u \subseteq X$ we will denote by $h|_u$ both the restriction of h to u and its value on u .

Fact 1 : $\Delta(f)(\alpha)$ is a continuous: choose, given $\alpha \in \text{dom } \Delta(f)$, a partition u_1, \dots, u_s of X such that both $f|_{u_j}$ and $\alpha|_{u_j}$ are constant. Then for $m \in M_2$, $[\Delta(f)(\alpha)]^{-1}(m) = \bigcup \{u_j : f|_{u_j}(\alpha|_{u_j}) = m\}$, which is clopen in X .

It is straightforward to verify that Δ is injective.

Notice now that if $f(x) \in S$ for all $x \in X$ and $\alpha \in C(X, M_1; \Sigma_1)$, then $(f)(\alpha) \in C(X, M_2; \Sigma_2)$: if $x \in F_k$, then $\alpha(x) \in M_{1k}$ and so $\Delta(f)(\alpha)(x) = f(x)(\alpha(x)) \in M_{2k}$. Similarly if $f \in C(X, S^*, G)$ and $\alpha \in \text{dom } (f) \cap SC(X, M_1; G)$ then $\Delta(f)(\alpha) \in SC(X, M_2; G)$: given $g \in G$ and $x \in X$

$$f(g_*x)(\alpha(g_*x)) = f(x)(\alpha(g_*x)) = g^* f(x)(\alpha(x)).$$

Thus a and b will be verified, as well as the assertion as to the range of Δ in the statement of the Lemma, as soon as we ascertain that $\Delta(f)$ is a partial isomorphism. If R is a n -ary relation symbol in L and $\alpha_1, \dots, \alpha_n$ are in $\text{dom } \Delta(f)$ then $C(X, M_1) \models R(\alpha_1, \dots, \alpha_n) \iff \forall x \in X M_1 R(\alpha_1(x), \dots, \alpha_n(x)) \iff \forall x \in X M_2 \models R(f(x)(\alpha(x)), \dots, f(x)(\alpha(x)))$. Function and constant symbols may be handled analogously. The verification of c. is straightforward. For d., suppose we are given $f \in C(X, S_n^*)$ and $\alpha \in C(X, M_2, \Sigma_2)$. Let u_1, \dots, u_s be a partition of X such that $f|_{u_j}$ and $\alpha|_{u_j}$ are constant. For each $1 \leq j \leq s$, there is $g_j \in S_{n-1}$ such that $g_j \supseteq f|_{u_j}$ and $\alpha|_{u_j} \in \text{Im } g_j$, $g_j(\beta_j) = \alpha|_{u_j}$. Define $g: X \rightarrow S_{n-1}$ such that $g|_{u_j} = g_j$ $1 \leq j \leq s$ and $\beta: X \rightarrow M_1$ by $\beta|_{u_j} = \beta_j$. Then, since $g \supseteq f$, $\Delta(g) \supseteq \Delta(f)$ and we have $\alpha \in \text{Im } \Delta(g)$. The case in which $\alpha \in C(X, M_1; \Sigma_1)$ is similarly disposed of. Thus d. is verified. For e. we need

Fact 2 : If $\alpha \in SC(X, M; G)$ we can find a partition u_1, \dots, u_s of X into invariant clopens (i.e. $g_* u_j = u_j \ \forall g \in G$) such that $\alpha(u_j) \leq |G| = \text{order of } G$.

Proof : On M we have an equivalence relation where $m_1 \sim m_2$ iff $m_2 = g^* m_1$ for some $g \in G$. Let $[m_1], \dots, [m_s]$ be a

partition of $\text{Im } \alpha$ into disjoint equivalence classes, and put

$$u_j = \{x \in X : \alpha(x) \in [m_j]\}, \quad 1 \leq j \leq s.$$

It is easy to verify that u_1, \dots, u_s satisfy the conclusions of Fact 2. Now given $f \in \text{SC}(X, S_n^*; G)$ and $\alpha \in \text{SC}(X, M_2; G)$ we can find a partition v_1, \dots, v_t of $\text{Im } \alpha$ into invariant clopens in such a way that $f|_{v_k}$ is constant and $\alpha(v_k) \subseteq |G|$. To obtain such a partition it is sufficient to consider the common refinement of the partition obtained in Fact 2 and that determined by requiring that f be constant on each of its members. Note that since the action of G on S^* is trivial, the set of points of X where f is constant is clopen invariant. We can find for each $1 \leq k \leq t$, $h_k \in S_{n-1}^*$ such that $\text{Im } h_k \supseteq \alpha(v_k)$. Define $h : X \rightarrow S_{(n-1)|G|}$ by $h|_{v_k} = h_k$. Since each v_k is invariant, $h \in \text{SC}(X, S_{n-1}^*; G)$ and $h \supseteq f$. Define $\beta : X \rightarrow M_1$ by the following rule : If $x \in v_k$, $1 \leq k \leq t$, $\beta(x) = h_k^{-1}(\alpha(x))$. Observe that once it is shown that β is in $\text{SC}(X, M_1; G)$ the proof is finished : it is obvious that $\alpha \in \text{Im } \Delta^*(h)$, $\Delta^*(h)(\beta) = \alpha$. We have for $m \in M_1$ $\beta^{-1}(m) = \bigsqcup_{1 \leq k \leq t} \{x \in X : \beta(x) = h_k^{-1}(m)\}$, clearly clopen. Thus β is continuous. On the other hand given $x \in X$, there is $1 \leq k \leq t$, such that $x \in v_k$. Then $\beta(g_*x) = h_k^{-1}(\alpha(g_*x)) = h_k^{-1}(g^*\alpha(x)) = g^*h_k^{-1}(\alpha(x)) = g^*\beta(x)$. In the above computations we have used the fact that v_k is invariant. This completes the proof of the Lemma. Δ

One can immediately conclude.

Theorem 1 b : If $M_1 \stackrel{\#}{\sim} M_2$, then $C(X, M_1; \Sigma_1) \stackrel{L}{\sim} C(X, M_2; \Sigma_2)$.

and

Theorem 2 b : If $M_1^* \underset{L}{\equiv} M_2^*$, then $SC(X, M_1; G) \underset{L}{\equiv} SC(X, M_2; G)$.

The proofs come from the fundamental Lemma plus Theorem 1.13, p.257 of [F] in the case that L is of finite type. In case L is not of finite type, observe that any sentence in L is a sentence in a finite reduct of L . Thus if it is true in $C(X, M_1; \Sigma_1)$ it will be true in $C(X, M_2; \Sigma_2)$ by the result above. A similar argument may, of course, be applied to obtain Theorem 2b in full generality. We have, therefore, Theorems 1b and 2b for all languages L of any type.

In the case that $M_1^{\#} \underset{L}{<} M_2^{\#}$ (or $M_1^* \underset{L}{<} M_2^*$) we have $C(X, M_1; \Sigma_1) \underset{L}{\equiv} E_1 \subseteq C(X, M_2; \Sigma_2) \underset{L}{\equiv} E_2$, canonically. To show that one is an elementary substructure of the other, we will prove.

Fact 3 : For any reduct of L of finite type, L' , including the unary predicate symbols P_1, \dots, P_n , for any $t \in \omega$ and for any $\{\alpha_1, \dots, \alpha_t\} \subseteq E_1$, $\langle E_1, \alpha_1, \dots, \alpha_t \rangle \underset{L'}{\equiv} \langle E_2, \alpha_1, \dots, \alpha_t \rangle$.

Proof: Let $\{m_1, \dots, m_s\} = \bigsqcup_{j=1}^t \text{Im } \alpha_j$. Since $M_1^{\#} \underset{L}{<} M_2^{\#}$, we have $\langle M_1^{\#}, m_1, \dots, m_s \rangle \underset{L'}{\equiv} \langle M_2^{\#}, m_1, \dots, m_s \rangle$. Let $\langle M_i^{\#}, \vec{m} \rangle$ denote $\langle M_i^{\#}, m_1, \dots, m_s \rangle$, $i=1, 2$. Observe that in Fraïssé's result we can assume that the constants belong to the domain of all partial isomorphisms involved in the back and forth arguments. Observe also that if $f \in C(X, P(\langle M_1^{\#}, \vec{m} \rangle, M_2^{\#}, \vec{m}))$ then $\Delta^{\#}(f)(\alpha_j) = \alpha_j$ $1 \leq j \leq t$. By the fundamental Lemma, we may conclude (again using Theorem 1.13 in [F]) that $\langle E_1, \vec{\alpha} \rangle \underset{L'}{\equiv} \langle E_2, \vec{\alpha} \rangle$ which proves Fact 3. Δ

From Fact 3 comes imediately, that if $M_1^{\#} \triangleleft_L M_2^{\#}$, then $E_1 \triangleleft_L E_2$. And Theorem 1 is proven in its entirety. Arguments entirely analous to those described above will prove Theorem 2.

Theorems 1 and 2 generalise the corresponding statements for bounded boolean powers in [B] (Theorem 6.5(i) and (ii)), and even in the case of boolean powers provide a new proof of the aforementioned result. In closing we would like to point out some related questions :

1) If $\langle B(X), I_1, \dots, I_n \rangle \equiv \langle B(Y), I'_1, \dots, I'_n \rangle$ is it true that $C(X, M; \Sigma) \equiv C(Y, M; \Sigma')$?

2) Same as above for structures of stable continuous functions, that is If $B_*(X) \equiv B_*(Y)$ is it true that $SC(X, M; G) \equiv SC(Y, M; G)$?

3) Same as 1) and 2) above, for elementary embedding.

4) If $\{M_i\}_{i \in I}$ is an inductive system of L-structures and $M = \lim_{\rightarrow} M_i$ is its inductive limit, under what circumstances $M_i \triangleleft_L M_i'$ implies $\lim_{\rightarrow} M_i = M \triangleleft_L \lim_{\rightarrow} M_i'$?

When M is finite, questions 1), 2) and 3) have affirmative answers (see [M]).

REF E R E N C E S

[B] S. Burris, "Boolean Powers", Algebra Universalis 5, 1975, 341-360.

[F] J. Flum, "First Order Logic and its Extensions", Proc. of the Intern. Summer Institute and Logic Colloquium, G. H. Muller, A. Obershelp, K. Potthoff (eds.), Kiel, 1974, Springer Lect. Notes 499, 249-310.

[Fr] R. Fraïssé, "Sur Quelques Classifications des Systèmes de Relations", Publ. Sci. Univ. Alger Ser. A, 1, 35-182.

[Mr] A. Macintyre and J. Rosenstein, " λ -Categoricity for Rings without Nilpotent Elements and for Boolean Structures" Journal of Algebra, 43, 1976, 129-154.

[M] F. Miraglia, "Relations between Structures of Stable Continuous Functions and Filtered Powers", Proceedings of the Third Brazilian Conference on Mathematical Logic, A.I. Arruda, A.M. Sette, N.C.A. da Costa, (eds.) Sociedade Bras. de Lógica, 1980, 225-244

Universidade de São Paulo
Departamento de Matemática
C.P. 20.570 - São Paulo

B R A S I L

TRABALHOS

DO

DEPARTAMENTO DE MATEMATICA

TITULOS PUBLICADOS

8001 - PLETCH, A. Local freeness of profinite groups. São Paulo, IME-USP, 1980. 10p.

8002 - PLETCH, A. Strong completeness in profinite groups. São Paulo, IME-USP, 1980. 8p.

8003 - CARNIELLI, W.A. & ALCANTARA, L.P. de. Transfinite induction on ordinal configurations. São Paulo, IME-USP, 1980. 22p.

8004 - JONES RODRIGUES, A.R. Integral representations of cyclic p-groups. São Paulo, IME-USP, 1980. 13p.

8005 - CORRADA, M. & ALCANTARA, L.P. de. Notes on many-sorted systems. São Paulo, IME-USP, 1980. /25/p.

8006 - POLCINO MILIES, F.C. & SEHGAL, S.K. FC-elements in a group ring. São Paulo, IME-USP, 1980. /10/p.

8007 - CHEN, C.C. On the Ricci condition and minimal surfaces with constantly curved Gauss map. São Paulo, IME-USP, 1980. 10p.

8008 - CHEN, C.C. Total curvature and topological structure of complete minimal surfaces. São Paulo, IME-USP, 1980. 21p.

8009 - CHEN, C.C. On the image of the generalized Gauss map of a complete minimal surface in R^4 . São Paulo, IME-USP, 1980. 8p.

8110 - JONES RODRIGUES, A.R. Units of ZCp^n . São Paulo, IME-USP, 1981. 7p.

8111 - KOTAS, J. & COSTA, N.C.A. da. Problems of modal and discussive logics. São Paulo, IME-USP, 1981. 35p.

8112 - BRITO, F.B. & GONCALVES, D.L. Algебras não associativas, sistemas diferenciais polinomiais homogeneos e classes características. São Paulo, IME-USP, 1981. 7p.

8113 - POLCINO MILIES, F.C. Group rings whose torsion units form a subgroup II. São Paulo, IME-USP, 1981. 1v. (não paginado)

8114 - CHEN, C.C. An elementary proof of Calabi's theorems on holomorphic curves. São Paulo, IME-USP, 1981. 5p.

8115 - COSTA, N.C.A. da & ALVES, E.H. Relations between para-consistent logic and many-valued logic. São Paulo, IME-USP, 1981. 8p.

8116 - CASTILLA, M.S.A.C. On Przymusinski's theorem. São Paulo, IME-USP, 1981. 6p.

8117 - CHEN, C.C. & GOES, C.C. Decenerate minimal surfaces in \mathbb{R}^4 . São Paulo, IME-USP, 1981. 21p.

8118 - CASTILLA, M.S.A.C. Imagers inversas de algumas aplicações fechadas. São Paulo, IME-USP, 1981. 11p.

8119 - ARAGONA VALEJO, A.J. & EXPL. FILHO, P. An infinite dimensional version of Hartogs' extension theorem. São Paulo, IME-USP, 1981. 9p.

8120 - GONÇALVES, J.Z. Groups rings with solvable unit groups. São Paulo, IME-USP, 1981. 15p.

8121 - CARNIELLI, W.A. & ALCANTARA, L.P. de. Paraconsistent algebras. São Paulo, IME-USP, 1981. 16p.

8122 - GONÇALVES, D.L. Nilpotent actions. São Paulo, IME-USP, 1981. 10p.

8123 - COELHO, S.P. Group rings with units of bounded exponent over the center. São Paulo, IME-USP, 1981. 25p.

8124 - PARMENTER, M.M. & POLCINO MILIES, F.C. A note on isomorphic group rings. São Paulo, IME-USP, 1981. 4p.

8125 - MERKLEN, H.A. Hereditary algebras with maximum spectra are of finite type. São Paulo, IME-USP, 1981. 10p.

8126 - POLCINO MILIES, F.C. Units of group rings: a short survey. São Paulo, IME-USP, 1981. 32p.

8127 - CHEN, C.C. & GACKSTATTER, F. Elliptic and hyperelliptic functions and complete minimal surfaces with handles. São Paulo, IME-USP, 1981. 14p.

8128 - POLCINO MILIES, F.C. A glance at the early history of group rings. São Paulo, IME-USP, 1981. 22p.

8129 - FERRER SANTOS, W.R. Reductive actions of algebraic groups on affine varieties. São Paulo, IME-USP, 1981. 52p.

8130 - COSTA, N.C.A. da. The philosophical import of paraconsistent logic. São Paulo, IME-USP, 1981. 26p.

8131 - GONÇALVES, D.L. Generalized classes of groups, spaces c-nilpotent and "the Hurewicz theorem". São Paulo, IME-USP, 1981. 30p.

8132 - COSTA, N.C.A. da & MORTENSEN, Chris. Notes on the theory of variable binding term operators. São Paulo, IME-USP, 1981. 18p.

8133 - MERKLEN, H.A. Homogeneous ℓ -hereditary algebras with maximum spectra. São Paulo, IME-USP, 1981. 32p.

8134 - PERESI, L.A. A note on semiprime generalized alternative algebras. São Paulo, IME-USP, 1981. 10p.