

RT-MAE-8917

CRITICAL POINTS OF TWO DIMENSIONAL
BOOTSTRAP PERCOLATION LIKE
CELLULAR AUTOMATA

by

Roberto H. Schonmann

Palavras Chaves: Bootstrap percolation; Cellular automata; Critical points
(Key words)

Classificação AMS: 60k35
(AMS Classification)

**Critical points of two dimensional Bootstrap Percolation
like Cellular Automata**

by

Roberto H. Schonmann

**Instituto de Matemática e Estatística da
Universidade de São Paulo.
Caixa Postal 20570
01498 São Paulo, SP, Brazil.
RHSCHONM@BRUSP.BITNET.**

Abstract

We consider cellular automata in two dimensions that generalize the Bootstrap Percolation dynamics. We focus on the threshold, p_c , of the initial density for convergence to total occupancy to occur, and classify these models according to p_c being 0, 1 or strictly between these extreme values. Explicit upper and lower bounds are provided in the third case.

In this paper we consider further the cellular automata studied in [Sch] in the two dimensional case. The models are defined as follows. Each site x of \mathbb{Z}^2 is at time $t = 0, 1, 2, \dots$ in state $\eta_t(x) \in \{0, 1\}$. At $t = 0$ the sites are independently in state 1 (occupied) with probability p or in state 0 (vacant) with probability $1 - p$. The evolution in time is deterministic, with the following features. Let $N = (0, 1)$, $W = (-1, 0)$, $S = (0, -1)$ and $E = (1, 0)$ be the nearest neighbors of the origin of \mathbb{Z}^2 . Set also $\mathcal{N} = \{N, W, S, E\}$. To define the dynamics, take a set \mathcal{D} of subsets of \mathcal{N} with the property of being increasing, i.e., if $A \subset B$ and $A \in \mathcal{D}$, then $B \in \mathcal{D}$. Let the translation by x of a set $A \subset \mathbb{Z}^2$ be written as

$$A + x = \{y \in \mathbb{Z}^2 : y = z + x \text{ for some } z \in A\}.$$

The evolution is given by the rules:

- (i) If $\eta_t(x) = 1$ then $\eta_{t+1}(x) = 1$.
- (ii) If $\eta_t(x) = 0$ and $\eta_t(y) = 1$ for all $y \in A + x$ for some $A \in \mathcal{D}$, then $\eta_{t+1}(x) = 1$.
- (iii) Otherwise $\eta_{t+1}(x) = 0$.

We denote by $P_p(\cdot)$ the law of the process starting from density p .

Bootstrap Percolation are the models for which a set is or not in \mathcal{D} depending only on its cardinality, that is, the cases of the sort

$$\mathcal{D} = \{A \subset \mathcal{N} : |A| \geq l\},$$

for some fixed $l = 1, 2, 3, 4$.

Adler and Aharony introduced in [AA] models that they call Diffusion Percolation, which are closely related to the models that we are considering.

For each one of these models defined above the density at time t ,

$$\rho_t(p) = P_p(\eta_t(0) = 1),$$

is non-decreasing in t and hence converges to the asymptotic density

$$\rho_\infty(p) = \lim_{t \rightarrow \infty} \rho_t(p).$$

When $\rho_\infty(p) = 1$ one says that the process converges to total occupancy. Due to monotonicity in p one naturally defines the critical point

$$p_c = \inf\{p \geq 0 : \rho_\infty(p) = 1\}.$$

In this paper we will classify all these two dimensional models according to p_c being 0, 1 or strictly between these two extreme values. Some explicit upper and lower bounds will also be provided in the third case.

First we recall results from [Sch]. The four oriented models are those with $\mathcal{D} = \{A \subset \mathcal{N} : S \subset A\}$, where S is respectively $\{E, N\}$, $\{N, W\}$, $\{W, S\}$, or $\{S, E\}$. We (arbitrarily) call the first one the basic oriented model. In Section 4 of [Sch] it was proven that for the oriented models

$$p_c = 1 - p^*,$$

where p^* is the critical point for oriented site percolation on \mathbb{Z}^2 . We say that a model I dominates a model II if (with a self explanatory notation) $\mathcal{D}_{II} \subset \mathcal{D}_I$, i.e., if it is easier for a vacant site to become occupied in I as compared to II. In Section 5 of [Sch] a simple argument was given, showing that if a model does not dominate any oriented model, then this model has $p_c = 1$. In other words, if no one of the sets $\{E, N\}$, $\{N, W\}$, $\{W, S\}$ and $\{S, E\}$ is in \mathcal{D} , then $p_c = 1$. (In our case it is enough to observe that squares of side 2 of vacant sites at time 0 will remain vacant forever.)

Now we consider models that dominate one of the oriented models, and with no loss of generality (due to symmetry) we can suppose that the dominated model is the basic oriented model, i.e.,

$$\{E, N\} \in \mathcal{D}. \quad (0.1)$$

There are two cases:

Case I: (0.1) holds and at least one of the following sets is also in \mathcal{D} : $\{N, W\}$, $\{S, E\}$.

Case II: (0.1) holds but no one of the sets $\{N, W\}$, $\{S, E\}$ is in \mathcal{D} .

Theorem:

- (i) In Case I, $p_c = 0$.
- (ii) In Case II, $1 - (p^*)^{1/4} \leq p_c \leq 1 - p^*$, in particular $p_c \in (0, 1)$.

Proof: i) By symmetry there is no loss in generality in assuming that $\{\{E, N\}, \{S, E\}\} \subset \mathcal{D}$.

Define the subsets of \mathbb{Z}^2 :

$$\begin{aligned} V(i; j) &= \{(x, y) \in \mathbb{Z}^2 : x = i, 0 \leq y \leq j\}, \\ H(i_1, i_2; j) &= \{(x, y) \in \mathbb{Z}^2 : i_1 \leq x \leq i_2, y = j\}. \end{aligned}$$

Consider now the numerical sequence $(m_k)_{k=1,2,\dots}$ given by

$$\begin{aligned} m_1 &= -1, \\ m_{k+1} &= m_k - k. \end{aligned}$$

And for $k = 1, 2, \dots$ define the events

$$\begin{aligned} F_k &= \{\text{For all } i \in \{m_{k+1} + 1, \dots, m_k\} \text{ there is at least one site } x \in V(i; k-1) \\ &\quad \text{such that } \eta_0(x) = 1\}, \\ G_k &= \{\text{For some } x \in H(m_{k+1} + 1, m_k; k) \text{ } \eta_0(x) = 1\}. \end{aligned}$$

If F_1 and G_1 happen then the region $V(m_2 + 1, 1)$ is completely occupied. If also F_2 and G_2 happen, then it is easy to see that $V(m_3 + 1, 2)$ and $H(m_3 + 1, m_3; 0)$ will eventually become completely occupied. Reasoning inductively we see that if all the events $F_k, G_k, k = 1, 2, \dots$ happen, then in particular all sites of the form $(-h, 0), h = 1, 2, \dots$ will become eventually occupied. The probability that all these events happen is

$$\alpha(p) = \prod_{k=1}^{\infty} [P(F_k) \cdot P(G_k)] = \prod_{k=1}^{\infty} (1 - (1 - p)^k)^{k+1}.$$

Hence, for every $p > 0$

$$\alpha(p) > 0. \tag{0.2}$$

We say that the origin is a good site if all the events $F_k, G_k, k = 1, 2, \dots$ happen. More generally we say that the site $x \in \mathbb{Z}^2$ is a good site if all these events happen when η_0 is replaced by the shifted configuration $\theta_{-x}\eta_0$ given by

$$(\theta_{-x}\eta_0)(y) = \eta_0(y + x).$$

By ergodicity and (0.2), there are with probability one good sites on the x-axis to the right of the origin. Therefore, by the remarks above, the origin will almost surely become eventually occupied and

$$\rho_{\infty}(p) = P_p(\eta_t(0) = 1 \text{ for some } t \geq 0) = 1.$$

This completes the proof of (i).

ii) Consider the squares of side 2

$$Q(x, y) = \{(2x, 2y), (2x + 1, 2y), (2x + 1, 2y + 1), (2x, 2y + 1)\}.$$

Think of a new lattice \mathbb{Z}^2 and declare the site (x, y) of this new lattice to be vacant at time t if and only if the four sites in $Q(x, y)$ are vacant at time t in the original lattice.

We say that double oriented percolation of vacant sites occurs in the configuration ζ if there is a doubly infinite chain of sites $\dots, z_{-1}, z_0, z_1, \dots$ such that $z_0 = 0$, $\zeta(z_i) = 0$ and $z_{i+1} \in \{z_i + (1, 0), z_i + (0, 1)\}$, for $i \in \mathbb{Z}$.

Suppose that at $t = 0$ double oriented percolation of vacant sites occurs in the new lattice. We will argue that then the same will occur at any later time and in particular $\eta_t(0) = 0$ for every $t \geq 0$. Indeed, if $z = (x, y)$ belongs to the doubly infinite chain of vacant new sites which is present at $t = 0$, then in the original lattice the four sites of $Q(z)$ are vacant and

- a) $(2x, 2y)$ and $(2x+1, 2y+1)$ have at most one occupied neighbor,
- b) $(2x, 2y + 1)$ has at most the sites to its north and west occupied,
- c) $(2x + 1, 2y)$ has at most the sites to its south and east occupied.

Hence, by the hypothesis, $Q(z)$ will still be completely vacant at time $t = 1$. (Observe that if \mathcal{D} contains a singleton, then it must contain one of the sets $\{N, W\}$, or $\{S, E\}$.) Induction on t completes now the argument which leads to

$$\begin{aligned} & 1 - \rho_\infty(p) \\ & \geq P_p(\text{at } t = 0 \text{ there is double oriented percolation of vacant sites in the new lattice}). \end{aligned} \tag{0.3}$$

But if $(1 - p)^4 > p^*$, then the probability in the r.h.s. of (0.3) is strictly positive, since double oriented percolation is equivalent to oriented percolation in the first quadrant and simultaneous oriented percolation with inverted orientations in the third quadrant. Hence

$$p_c \geq 1 - (p^*)^{1/4}.$$

The other bound,

$$p_c \leq 1 - p^*$$

is a trivial consequence of the fact that the model dominates the basic oriented model and that, as remarked before, the critical point of this latter model coincides with $1 - p^*$. \square

Part (i) of our theorem applies to bootstrap percolation with $l = 2$, for which the result $p_c = 0$ had been proven before by van Enter [vEn] based on an unpublished idea by Straley. Our proof was in fact inspired by these arguments, but in a sense it is more robust, since it applies to certain models in which the orientation is relevant and for which the Straley-van Enter argument does not apply. For instance Duarte [Dua] considered the model for which a vacant site becomes occupied when at least two among its three neighbors to the North, East and South are occupied. He observed that the Straley-van Enter argument does not apply and obtained from simulations on lattices of linear size between 5 and 15360 the numerical prediction $p_c = 0.034 \pm 0.01$. For this model $\{(N, E), (S, E)\} \subset \mathcal{D}$ so that our theorem implies $p_c = 0$.

As a consequence of our bounds on p_c , we conclude in particular that no model in the class that we consider can have p_c in the intervals $(0, 1 - (p^*)^{1/4})$ and $(1 - p^*, 1)$. The oriented models have $p_c = 1 - p^*$ and we conjecture that there are models that dominate an oriented model and have $0 < p_c < 1 - p^*$. A candidate for this seems to be the model defined by

$$\mathcal{D} = \{A \subset \mathcal{N} : \{N, E\} \subset A \text{ or } \{W, E\} \subset A\}.$$

Acknowledgements: *This work was partially done while I was visiting UCLA; I would like to express my gratitude for the warm hospitality of the Mathematics Department and specially of Tom Liggett. I am also indebted to Joel Lebowitz for letting me know of Duarte's work.*

References

- [AA] J.Adler, A.Aharony: Diffusion percolation: 1. Infinite time limit and bootstrap percolation. J. Phys. A: Math. Gen **21**, 1387-1404 (1988).
- [Dua] J.A.M.S. Duarte: Simulation of a cellular automat with an oriented bootstrap rule. Physica A **157**, 1075-1079 (1989).
- [vEn] A.C.D. van Enter: Proof of Straley's argument for bootstrap percolation. J. Stat. Phys. **48**, 943-945 (1987).
- [Sch] R.H. Schonmann: On the behavior of some cellular automata related to bootstrap percolation. Preprint (1989).

RELATÓRIO TÉCNICO
DO
DEPARTAMENTO DE ESTATÍSTICA

TÍTULOS PUBLICADOS EM 1987

- 8701 - ACHCAR, J.A. & BOLFARINE, H.; Constant Hazard Against a Change-Point Alternative: A Bayesian Approach with Censored Data, São Paulo, IME-USP, 1987, 20p.
- 8702 - RODRIGUES, J.; Some Results on Restricted Bayes Least Squares Predictors for Finite Populations, São Paulo, IME-USP, 1987, 16p.
- 8703 - LEITE, J.G., BOLFARINE, H. & RODRIGUES, J.; Exact Expression for the Posterior Mode of a Finite Population Size: Capture-Recapture Sequential Sampling, São Paulo, IME-USP, 1987, 14p.
- 8704 - RODRIGUES, J., BOLFARINE, H. & LEITE, J.G.; A Bayesian Analysis in Closed Animal Populations from Capture Recapture Experiments with Trap Response, São Paulo, IME-USP, 1987, 21p.
- 8705 - PAULINO, C.D.M.; Analysis of Categorical Data with Full and Partial Classification: A Survey of the Conditional Maximum Likelihood and Weighted Least Squares Approaches, São Paulo, IME-USP, 1987, 52p.
- 8706 - CORDEIRO, G.M. & BOLFARINE, H.; Prediction in a Finite Population under a Generalized Linear Model, São Paulo, IME-USP, 1987, 21p.
- 8707 - RODRIGUES, J. & BOLFARINE, H.; Nonlinear Bayesian Least-Squares Theory and the Inverse Linear Regression, São Paulo, IME-USP, 1987, 15p.
- 8708 - RODRIGUES, J. & BOLFARINE, H.; A Note on Bayesian Least-Squares Estimators of Time-Varying Regression Coefficients, São Paulo, IME-USP, 1987, 11p.

- 8709 - ACHCAR, J.A., BOLFARINE, H. & RODRIGUES, J.; Inverse Gaussian Distribution: A Bayesian Approach, São Paulo, IME-USP, 1987, 20p.
- 8710 - CORDEIRO, G.M. & PAULA, G.A.; Improved Likelihood Ratio Statistics for Exponential Family Nonlinear Models, São Paulo, IME-USP, 1987, 26p.
- 8711 - SINGER, J.M.; PERES, C.A. & HARLE, C.E.; On the Hardy -Weinberg Equilibrium in Generalized ABO Systems, São Paulo, IME-USP, 1987, 16p.
- 8712 - BOLFARINE, H. & RODRIGUES, J.; A Review and Some Extensions on Distribution Free Bayesian Approaches for Estimation and Prediction, São Paulo, IME-USP, 1987, 19p.
- 8713 - RODRIGUES, J.; BOLFARINE, H. & LEITE, J.G.; A Simple Nonparametric Bayes Solution to the Estimation of the Size of a Closed Animal Population, São Paulo, IME-USP, 1987, 11p.
- 8714 - BUENO, V.C.; Generalizing Importance of Components for Multistate Monotone Systems, São Paulo, IME-USP, 1987, 12p.
- 8801 - PEREIRA, C.A.B. & WECHSLER, S.; On the Concept of P-value, São Paulo, IME-USP, 1988, 22p.
- 8802 - ZACKS, S., PEREIRA, C.A.B. & LEITE, J.G.; Bayes Sequential Estimation of the Size of a Finite Population, São Paulo, IME-USP, 1988, 23p.
- 8803 - BOLFARINE, H.; Finite Population Prediction Under Dynamic Generalized Linear Models, São Paulo, IME-USP, 1988, 21p.
- 8804 - BOLFARINE, H.; Minimax Prediction in Finite Populations, São Paulo, IME-USP, 1988, 18p.
- 8805 - SINGER, J.M. & ANDRADE, D.F.; On the Choice of Appropriate Error Terms for Testing the General Linear Hypothesis in Profile Analysis, São Paulo, IME-USP, 1988, 23p.
- 8806 - DACHS, J.N.W. & PAULA, G.A.; Testing for Ordered Rate Ratios in Follow-up Studies with Incidence Density Data, São Paulo, IME-USP, 1988, 18p.
- 8807 - CORDEIRO, G.M. & PAULA, G.A.; Estimation, Significance Tests and Diagnostic Methods for the Non-Exponential Family Nonlinear Models, São Paulo, IME-USP, 1988, 29p.

- 8808 - RODRIGUES, J. & ELIAN, S.N.; The Coordinate - Free Estimation in Finite Population Sampling, São Paulo, IME-USP, 1988, 5p.
- 8809 - BUENO, V.C. & CUADRADO, R.Z.B.; On the Importance of Components for Continuous Structures, São Paulo, IME-USP, 1988, 14p.
- 8810 - ACHCAR, J.A., BOLFARINE, H & PERICCHI, L.R.; Some Applications of Bayesian Methods in Analysis of Life Data, São Paulo, IME-USP, 1988, 30p.
- 8811 - RODRIGUES, J.; A Bayesian Analysis of Capture-Recapture Experiments for a Closed Animal Population, São Paulo, IME-USP, 1988, 10p.
- 8812 - FERRARI, P.A.; Ergodicity for Spin Systems, São Paulo, IME-USP, 1988, 25p.
- 8813 - FERRARI, P.A. & MAURO, E.S.R.; A Method to Combine Pseudo-Random Number Generators Using Xor, São Paulo, IME-USP, 1988, 10p.
- 8814 - BOLFARINE, H. & RODRIGUES, J.; Finite Population Prediction Under a Linear Functional Superpopulation Model a Bayesian Perspective, São Paulo, IME-USP, 1988, 22p.
- 8815 - RODRIGUES, J. & BOLFARINE, H.; A Note on Asymptotically Unbiased Designs in Survey Sampling, São Paulo, IME-USP, 1988, 6p.
- 8816 - BUENO, V.C.; Bounds for the Availabilities in a Fixed Time Interval for Continuous Structures Functions, São Paulo, IME-USP, 1988, 22 p.
- 8817 - TOLOI, C.M.C. & MORETTIN, P.A.; Spectral Estimation for Time Series with Amplitude Modulated Observations: A Review, São Paulo, IME-USP, 1988, 16p.
- 8818 - CHAYES, J.T.; CHAYES, L.; GRIMMETT, G.R.; KESTEN, H. & SCHONMANN, R.H.; The Correlation Length for the High Density Phase of Bernoulli Percolation, São Paulo, IME-USP, 1988, 46p.
- 8819 - DURRETT, R.; SCHONMANN, R.H. & TANAKA, N.I.; The Contact Process on a Finite Set, III: The Critical Case, São Paulo, IME-USP, 1988, 31p.

- 8820 - DURRET, R.; SCHONMANN, R.H. & TANAKA, N.I.; Correlation Lengths for Oriented Percolation, São Paulo, IME-USP, 1988, 18p.
- 8821 - BRICMONT, J.; KESTEN, H.; LEBOWITZ, J.L. & SCHONMANN, R.H.; A Note on the Ising Model in High Dimensions, São Paulo, IME-USP, 1988, 21p.
- 8822 - KESTEN, H. & SCHONMANN, R.H.; Behavior in Large Dimensions of the Potts and Heisenberg Models, São Paulo, IME-USP, 1988, 61p.
- 8823 - DURRET, R. & TANAKA, N.I.; Scaling Inequalities for Oriented Percolation, São Paulo, IME-USP, 1988, 21p.
- 3901 - RODRIGUES, J.; Asymptotically Design - Unbiased Predictors to Two-Stage Sampling, São Paulo, IME-USP, 1989, 9p.
- 8902 - TOLOI, C.M.C. & MORETTIN, P.A.; Spectral Analysis for Amplitude Modulated Time Series, São Paulo, IME-USP, 1989, 24p.
- 8903 - PAULA, G.A.; Influence Measures for Generalized Linear Models with Restrictions in Parameters, São Paulo, IME-USP, 1989, 18p.
- 8904 - MARTIN, M.C. & BUSSAB, W.O.; An Investigation of the Properties of Raking Ratio Estimators for Cell Frequencies with Simple Random Sampling, São Paulo, IME-USP, 1989, 11p.
- 8905 - WECHSLER, S.; Yet Another Refutation of Allais' Paradox, São Paulo, IME-USP, 1989, 6p.
- 8906 - BARLOW, R.E. & PEREIRA, C.A.B.; Conditional Independence and Probabilistic Influence Diagrams, São Paulo, IME-USP, 1989, 21p.
- 8907 - BARLOW, R.E.; PEREIRA, C.A.B. & WECHSLER, S.; The Bayesian Approach to Ess, São Paulo, IME-USP, 1989, 20p.
- 8908 - PEREIRA, C.A.B. & BARLOW, R.E.; Medical Diagnosis Using Influence Diagrams, São Paulo, IME-USP, 1989, 13p.
- 8909 - BOLFARINE, H.; A Note on Finite Population Prediction Under Asymmetric Loss Functions, São Paulo, IME-USP, 1989, 8p.

- 8910 - BOLFARINE, H.; Bayesian Modelling in Finite Populations, São Paulo, IME-USP, 1989, 8p.
- 8911 - NEVES, M.M.C & MORETTIN, P.A.; A Generalized Cochrane-Orcutt-Type Estimator for Time Series Regression Models, São Paulo, IME-USP, 1989, 29p.
- 8912 - MORETTIN, P.A., TOLOI, C.M.C., GAIT, N. & MESQUITA, A.R.: Analysis of the Relationships Between Some Natural Phenomena: Atmospheric Precipitation, Mean Sea Level and Sunspots, São Paulo, IME-USP, 1989, 35p.
- 8913 - BOLFARINE, H.; Population Variance Prediction Under Normal Dynamic Superpopulation Models, São Paulo, IME-USP, 1989, 7p.
- 8914 - BOLFARINE, H.; Maximum Likelihood Prediction in Two Stage Sampling, São Paulo, IME-USP, 1989, 4p.
- 3915 - WECHSLER, S.; Exchangeability and Predictivism, São Paulo, IME-USP, 1989, 10p.
- 8916 - BOLFARINE, H.; Equivariant Prediction in Finite Populations, São Paulo, IME-USP, 1989, 13p.