



Genetic variation and cryptic lineage among the sergestid shrimp *Acetes americanus* (Decapoda)

Sabrina Morilhas Simões^{1,2}, Rogério Caetano Costa¹, Fabricio Lopes Carvalho³, Abner Carvalho-Batista⁴, Sarah de Souza Alves Teodoro⁵ and Fernando L. Mantelatto²

¹ Department of Biological Sciences, Faculty of Sciences, University of State of São Paulo (UNESP), Bauru, São Paulo, Brazil

² Department of Biology, Faculty of Philosophy, Science and Letters at Ribeirão Preto (FFCLRP), University of São Paulo (USP), Ribeirão Preto, São Paulo, Brazil

³ Federal University of Southern Bahia (UFSB), Ilhéus, Bahia, Brazil

⁴ Paulista University (UNIP), Bauru, Bauru, São Paulo, Brazil

⁵ Rio Grande Federal University (FURG), Rio Grande, Rio Grande do Sul, Brazil

ABSTRACT

The taxonomic status of the sergestid shrimp, *Acetes americanus*, has been questioned for several decades. No specific study has been performed thus far to resolve the incongruences. This species has a wide geographical range in the western Atlantic and is represented by two formally accepted subspecies: *Acetes americanus carolinae*, distributed in North America, and *Acetes americanus americanus*, present in South America. However, there are regions where the coexistence of both subspecies has been reported, such as Central America. This study aimed to genetically compare specimens of *A. a. americanus* collected in South America with *A. a. carolinae* sampled in North America to check for possible differences and the existence of more than one subspecies of *A. americanus* on the Brazilian coast. Based on the sequences of two informative markers, the cytochrome oxidase I region (COI) and 16S rRNA, phylogenetic reconstruction demonstrated well-defined clades with high support values, reinforcing the idea that *A. a. americanus* is genetically different from *A. a. carolinae*. Our hypothesis was corroborated as the specimens collected in Brazil were divided into two distinct lineages: the first composed of *A. a. americanus sensu stricto* (Brazil 1) and the second by *Acetes americanus* (Brazil 2). The three groups evidenced in the haplotype network were the same as those observed in the phylogenetic tree. The morphometric character (height/length of the thelycum) was effective in distinguishing *A. a. Brazil 1* from *A. a. carolinae*. However, more detailed and conclusive studies comprising other characteristics to propose and describe a possible new entity are necessary. To the best of our knowledge, for the first time, the results of this study provide some insights into the taxonomic status of the sergestid shrimp *A. americanus* in the western Atlantic.

Submitted 1 August 2022
Accepted 27 December 2022
Published 13 February 2023

Corresponding author
Fernando L. Mantelatto,
flmantel@usp.br

Academic editor
Khor Waiho

Additional Information and
Declarations can be found on
page 19

DOI 10.7717/peerj.14751

© Copyright
2023 Morilhas Simões et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Biodiversity, Genetics, Marine Biology, Molecular Biology, Zoology

Keywords Crustacea, Molecular phylogeny, Western Atlantic, 16S rDNA, Sergestidae, Cytochrome Oxidase I (COI), Sergestoidea, Phylogenetics

INTRODUCTION

Genus *Acetes* H. Milne Edwards, 1830 comprises small planktonic shrimps (Wong, 2013), which are essential components of marine systems (Xiao & Greenwood, 1993). For a long time, the genus remained poorly understood among decapods concerning phylogeny. Recently, species have been contextualized in a global phylogeny (Vereshchaka, Lunina & Olesen, 2016a; Vereshchaka, 2017). Fourteen species of *Acetes* are recognized worldwide (De Grave & Fransen, 2011; WoRMS, 2022). Three species occur in the western Atlantic: *Acetes americanus* Ortmann, 1893; *Acetes marinus* (Omori, 1975); and *Acetes paraguayensis* Hansen, 1919.

Historically, four subspecies of *A. americanus* have been identified: *A. a. carolinae* Hansen, 1933 (type locality: Cove Beaufort, South Carolina, USA), *A. a. louisianensis* (Burkenroad, 1934) (type locality: Louisiana coast, from the west of the Mississippi River to Timbalier Island, Gulf of Mexico, USA), *A. a. limonensis* (Burkenroad, 1934) (type locality: Sweetwater River mouth, Panama), and *A. a. americanus* (Ortmann, 1893) (type locality: Tocantins River mouth, Brazil) (Burkenroad, 1934).

However, the subspecies *A. a. louisianensis* and *A. a. limonensis* presented intermediate characteristics of the other two subspecies. Therefore, they were traditionally considered clinal variants, which are not considered valid (Holthuis, 1948). Currently, *A. a. louisianensis* and *A. a. limonensis* are accepted as synonyms for *A. a. americanus* (WoRMS, 2022).

The existing taxonomy considers only *A. a. carolinae* and *A. a. americanus* as valid subspecies (Holthuis, 1948). Although these two subspecies are very similar, careful examination reveals minute morphological differences, as the body and cornea lengths of the southern representatives (*A. a. americanus*) are slightly larger than those of the northern representatives (*A. a. carolinae*) (Omori, 1975). Taxonomic inconsistencies in the subspecies of *A. americanus* have been reported since the 1970s (Omori, 1975). However, this remains unsolved.

Acetes americanus carolinae is distributed from North Carolina, Florida to the Gulf of Mexico, Panama, Suriname, and French Guiana (Omori, 1975); *A. a. americanus* Ortmann, 1893 occurs in Puerto Rico, Panama, Venezuela, Suriname, French Guiana, and Brazil (D'Incao & Martins, 2000; Mantelatto et al., 2022) (Fig. 1).

The uncertainty regarding the difference in the extant subspecies results from: (a) insufficient information about the subspecies' habitats, especially where the two subspecies co-occur, (b) the small magnitude of morphological divergence among species of the genus *Acetes* and (c) the distribution.

Molecular tools were used to elucidate the taxonomic status of several marine shrimp species, whose morphology is insufficient to clarify their species identity (Carvalho, Magalhães & Mantelatto, 2014; Tavares & Gusmão, 2016; Carvalho, Magalhães & Mantelatto, 2020). Combined with morphological data, genetic data allowed the interpretation of variability patterns along the distribution of some taxa (Silva, Mesquita & Paula, 2010; Terossi & Mantelatto, 2012; Rossi & Mantelatto, 2013; Teodoro et al., 2016; Carvalho-Batista et al., 2019; França et al., 2019; Terossi & Mantelatto, 2020; França et al., 2021). The mitochondrial genes cytochrome c oxidase I (COI) and 16S rRNA have




Figure 1 Distribution of studied species of *Acetes* in the Western Atlantic. Geographic distribution of the *Acetes americanus carolinae* (*) (Omori, 1975) and *Acetes americanus americanus* (+) (D’Incao and Martins, 2000). The highlighted circles indicate the sites studied by molecular analysis, *Acetes americanus* USA (pink), *Acetes americanus* Brazil 1 (yellow), *Acetes americanus* Brazil 2 (blue), *Acetes petrunkevitchi* (orange) and *Acetes paraguayensis* (black). Map created by authors using Qgis 3.24.3 (<https://www.qgis.org/en/site/index.html>).

Full-size  DOI: 10.7717/peerj.14751/fig-1

been considered powerful markers for molecular analysis in several studies of decapod shrimp at species and population levels (Gusmão, Lazoski & Solé-Cava, 2000; Maggioni et al., 2001; Lavery et al., 2004; Vergamini, Pileggi & Mantelatto, 2011; Terossi & Mantelatto, 2012; Teodoro et al., 2016; Carvalho-Batista et al., 2018; França et al., 2019).

In this study, we compared the *A. a. americanus* specimens collected from South America with those of *A. a. carolinae* sampled in North America using COI and 16S rRNA markers to test the genetic validity of both subspecies and the possible existence of other entities distributed along the western Atlantic.

MATERIALS & METHODS

Sampling of the biological material

The specimens used in this study were obtained through samplings and loans. All individuals of *A. americanus* from Penha/Santa Catarina, Ubatuba, Cananéia, São Vicente/São Paulo, Macaé/Rio de Janeiro, Baía Formosa/Rio Grande do Norte, and *A. petrunkevitchi* from Ubatuba/São Paulo were sampled using a fishing boat equipped with an otter trawl net with 2 m opening and 3 m in length. The mesh diameter (interknot distance) was five mm in the first half of the net and two mm in the final part. The obtained specimens were collected under field permit approval by Instituto Chico Mendes de Biodiversidade/ICMBio, number 23008-1 and Permanent License to FLM 11777-2) and deposited in the Crustacean Collection of the Department of Biology of the Faculty of Philosophy, Sciences, and Letters of Ribeirão Preto of the University of São Paulo, Brazil (CCDB/FFCLRP/USP) and in the Crustacean Collection of the Laboratory of Biology of Marine and Freshwater Shrimp, UNESP, Bauru, Brazil (CCLC/FC/UNESP) (Table 1 and Table S1).

All other specimens used in this study were loaned from scientific collections: Zoology Museum of the University of São Paulo, São Paulo, Brazil (MZUSP); National Museum of Natural History, Smithsonian Institution, United States (USNM); National Museum of the Federal University of Rio de Janeiro (MNRJ); Oceanographic Museum of the University of Pernambuco, Brazil (MOUFPE); Federal University of Rio Grande (FURG); Crustacean Collection from the Department of Biology of the Faculty of Philosophy, Sciences and Letters of Ribeirão Preto, University of São Paulo (CCDB); Collection of Crustaceans of the Federal University of Sergipe (UFS); Collection of Crustaceans from the Federal University of Espírito Santo (UFES); Collection of Crustaceans from the PUCRS Museum of Science and Technology (MCP); Carcinological Collection of the Institute of Scientific and Technological Research of the State of Amapá, Macapá, AP (IEPA); National Crustacean Collection, UNAM, Mexico (CNCR); Peabody Museum of Natural History Yale University, United States (YPM); University of Louisiana at Lafayette, USA (ULLZ); and Zoological Museum Kiel, Germany (ZMK) (Table 1 and Table S1).

Specimens were morphologically identified following the identification key proposed by *Omori (1975)*, *D’Incao & Martins (2000)*, *Vereshchaka, Lunina & Olesen (2016a)*, and *Vereshchaka, Olesen & Lunina (2016b)*, using the shape of the third thoracic sternite (thelycum) in females and the petasma shape in males.

The names currently used for extant taxa follow the most recent literature (*Vereshchaka, Lunina & Olesen, 2016a*; *Vereshchaka, Olesen & Lunina, 2016b*) and are currently adopted by *WoRMS (2022)*. Genus *Peisos* is dealt with differently. A review of the shrimp genera *Acetes*, *Peisos*, and *Sicyonella* (*Vereshchaka, Lunina & Olesen, 2016a*) proposed to include *Peisos* within the genus *Acetes* using morphological and phylogenetic information. Later, in a global phylogeny, *Vereshchaka (2017)* proposed new families, including Acetidae, to accommodate all species of *Acetes*. *WoRMS (2022)* does not adopt the proposed changes, that is, *Peisos* is an accepted genus, and Acetidae does not exist. We can conjecture a lack of taxonomic reassignment with a proper nomenclatural act in the previous paper. Considering that we were not able to find a clear justification and to avoid taxonomic

Table 1 Size dimension relationship of females *Acetes americanus*. Sample size and ratio between height and length of the third thoracic sternite of the females of *Acetes americanus* Ortmann, 1893.

Subspecies	Number of specimens	Locality (Latitud)	Catalog number	Average \pm Standard Deviation
<i>Acetes americanus</i> USA	28	Off Beaufort Inlet, North Carolina, United States (34 °N)	USNM74550	0.64 \pm 0.08
<i>Acetes americanus</i> USA	2	Cape Lookout, North Carolina, United States (34°N)	USNM258701	0.65 \pm 0.07
<i>Acetes americanus</i> USA	6	Beaufort, North Carolina, United States (34°N)	YPM005386	0.48 \pm 0.08
<i>Acetes americanus</i> USA	5	Off mouth of North Edisto River, South Carolina, United States (32°N)	USNM258707	0.60 \pm 0.11
<i>Acetes americanus</i> USA	9	Louisiana, Gulf of Mexico, United States (29°N)	YPM005387	0.57 \pm 0.08
<i>Acetes americanus</i> USA	17	Laguna de Términos, Frente el Faro de Xicalango, México (19°N)	CNCR2402	0.56 \pm 0.11
<i>Acetes americanus</i> Brazil 1	6	Channel at mouth of Bay, Puerto Rico (18°N)	USNM134695 to USNM134697	0.24 \pm 0.04
<i>Acetes americanus</i> Brazil 1	17	Playa de Guayanes, Puerto Yabucoa, Puerto Rico (18°N)	USNM186645 to USNM186647	0.22 \pm 0.06
<i>Acetes americanus</i> Brazil 1	20	Off Surinam Coast, Suriname (6°N)	USNM103101 to USNM103106	0.29 \pm 0.08
<i>Acetes americanus</i> Brazil 1	26	Maceió,AL, Brazil (09°S)	MZUSP21210	0.38 \pm 0.08
<i>Acetes americanus</i> Brazil 1	35	Macaé, RJ, Brazil (22°S)	CCLC0254	0.36 \pm 0.07
<i>Acetes americanus</i> Brazil 1	28	Ubatuba, SP, Brazil (23°S)	CCLC0253	0.40 \pm 0.34
<i>Acetes americanus</i> Brazil 1	13	Rio Grande, RS, Brazil (32°S)	MZUSP 9079	0.33 \pm 0.07
<i>Acetes americanus</i> Brazil 2	35	Macaé, RJ, Brazil (22°S)	CCLC0261	0.79 \pm 0.19
<i>Acetes americanus</i> Brazil 2	34	Cananéia, SP, Brazil (25°S)	CCLC0262	1.03 \pm 0.29

Notes.

CCLC, Crustacean Collection of the Laboratory of Biology of Marine and Freshwater Shrimp, UNESP, Bauru, Brazil; MZUSP, Zoology Museum of the University of São Paulo, São Paulo, Brazil; USNM, National Museum of Natural History, Smithsonian Institution, United States; CNCR, National Crustacean Collection, UNAM, Mexico; YPM, Peabody Museum of Natural History Yale University, United States.

instability, we maintained the proposed taxonomic status of *Peisos* as part of *Acetes* following [Vereshchaka, Lunina & Olesen \(2016a\)](#).

DNA extraction and amplification

DNA extraction from the muscular abdominal tissue was performed following [Mantelatto, Robles & Felder \(2007\)](#); [Mantelatto et al. \(2009a\)](#); [Mantelatto et al. \(2009b\)](#) and [Pileggi & Mantelatto \(2010\)](#), with specific modifications ([Carvalho-Batista et al., 2014](#)). The regions of interest were amplified by the polymerase chain reactions (PCR) using the primers described in [Table S2](#). Primers specific for this region were designed due to the difficulty in the amplification of COI (see [Mantelatto et al., 2016](#) for details) ([Table S2](#)).

PCR reactions were performed with a total volume of 25 μ L, containing 5 μ L of betaine (5M) (Acros Organics), 4 μ L of dNTPs (5 mM), 3 μ l of MgCl₂ (25 mM), 3 μ l of 10X Taq buffer with KCl (Thermo Scientific), 2 μ l of 1% bovine albumin (Sigma), 1 μ l of each

primer (10 μ M), 1 μ l of resuspended DNA (50 ng/ml), and 0.5 μ l of recombinant Taq DNA Polymerase (Thermo Scientific). The missing volume (25 μ l) was filled with ultrapure water. PCR was performed using an Applied Biosystems[®] Veriti 96 well thermal cycler. PCR steps comprised an initial denaturation period of 3 min at 95 °C, followed by 40 thermal cycles [30 s of denaturation at 95 °C, 45 s for annealing at variable temperature (42–44 °C for 16S; 44–50 °C for COI), 1 min for the extension at 72 °C, and final extension for 10 min at 72 °C. The obtained results were observed by 1.5% agarose gel electrophoresis and photographed with Olympus[®] C-7070 digital camera in a UV M20 UV transilluminator.

The PCR products were purified using the Sureclean purification kit following the manufacturer's protocol. The PCR-purified products were sequenced bidirectionally in automated sequencers (ABI 3100 Genetic Analyzer) at the Department of Technology of the Faculty of Sciences Agricultural and Veterinary Sciences of Jaboticabal, São Paulo State University.

The generated sequences were confirmed and edited (sequence consensus obtained from sense and antisense) in BioEdit 7.0.7.1 software (Hall, 1999) and aligned using CLUSTAL W (Thompson, Higging & Gibson, 1994).

Each genetic sample obtained for the analyses was deposited in the scientific collection of origin (Table S1).

Molecular analyses

A total of 118 sequences used in this study were generated for this project. For the 16S rRNA region, 52 sequences of 518 bp were obtained, of which 29 were *A. a.* Brazil 1, nine were *A. a.* Brazil 2, five were *A. a. carolinae*, five were *A. paraguayensis*, and four were *A. petrunkevitchi*. For the COI region, 66 sequences of 589 bp were generated of which 36 were *A. a.* Brazil 1, 15 were *A. a.* Brazil 2, six were *A. a. carolinae*, six were *A. paraguayensis*, and three were *A. petrunkevitchi*.

However, an additional 20 sequences (COI) of *A. sibogae*, *A. japonicus*, *A. serrulatus*, and *A. indicus*, two sequences (16S and COI) of *A. petrunkevitchi*, and two sequences (16S and COI) of *B. faxoni* (outgroup) were retrieved from the GenBank database and used for phylogenetic and genetic distance analyses to complement this study (Table S1). As there are COI sequences for more *Acetes* species than 16S sequences available on GenBank, phylogenetic analysis with the COI gene generated a larger number of clades.

The mean nucleotide composition and genetic distances were estimated using MEGA 5.0 software (Tamura et al., 2011) and the Neighbor-Joining dendrogram based on the Kimura 2-parameter substitution model (Kimura, 1980).

The appropriate models of nucleotide evolution HKY + G for 16S and TPM1uf + G for COI were selected by Bayesian information criterion (BIC) in jModeltest 2.1.4 (Darriba et al., 2012). Selected models and estimated parameters (Table S3) were implemented in the Bayesian inferences and considered for the choice of the closest models in the maximum likelihood analyses. *Belzebub faxoni* (Borradaile, 1915) (Superfamily Sergestoidea, Family Luciferidae) was included as an outgroup following the most recent global phylogeny (Vereshchaka, Lunina & Olesen, 2016a; Vereshchaka, 2017). Bayesian inference was used to reconstruct the phylogenetic relationships of the species analyzed, with the two genes

as distinct partitions in MrBayes v. 3.2.2 (Ronquist et al., 2012). Bayesian inference was performed with 30 million generations in two independent analyses, with five parallel chains each, one “cold” and four “hot”. The parameters were saved every 1,000 simulations. The analysis was completed on attaining stationarity (mean standard deviation <0.01) after the stipulated number of generations. The first quarter of the parameters and trees were discarded as burn-in (Ronquist, Van Der Mark & Huelsenbeck, 2009). The support values of the branches were obtained using the *a posteriori* probability method.

This analysis was performed for the 16S and COI genes separately and a concatenated matrix of both the trees was generated and edited in the program Figtree v.1.3.1 (Rambaut, 2007).

Population analyses

Only the COI gene was used for population analyses. The use of this gene has been shown to be efficient in decapod population studies (Gusmão, Lazoski & Solé-Cava, 2005; Laurenzano, Mantelatto & Schubart, 2013; Rossi & Mantelatto, 2013; Carvalho-Batista et al., 2014; Teodoro et al., 2015) because it is considered to be more variable (Schubart & Huber, 2006).

The number of haplotypes was calculated in the DnaSP 4.10.9 software (Rozas & Rozas, 1999). Haplotype networks were constructed using the median-joining method, using the network program (Bandelt, Forster & Röhl, 1999), based on the data prepared in the DnaSP. Haplotype and nucleotide diversities, analysis of molecular variance (AMOVA), and pairwise fixation indices (Fst) (Excoffier, Smouse & Quattro, 1992) were calculated using Arlequin 3.11 (Excoffier, Laval & Schneider, 2005).

Morphometric analyses

We used only females (281 individuals) of *A. americanus* for morphometric analyses. Previous observations focused on the difference in size and shape of the female thelycum between the subspecies, motivating us to obtain potential information to complement our study.

Individuals were sexed based on the presence of petasma (first pleopod) in males and the thelycum (third thoracic sternite) in females (Xiao & Greenwood, 1993).

Morphometric measurements were obtained using a stereo microscope Zeiss[®] Stemi 2000C connected to an imaging system Zeiss[®] AxioVision, with an error of up to 0.01 mm. We measured the height (HT) and length (LT) of the thelycum and carapace length (CL).

Following the methodology proposed by Marramà & Kriwet (2017), data were standardized by calculating the ratio between each measure and carapace length (CL), removing the effect of size. The log transformation of data (Log (X+1)) was used to overcome the issue of the non-normal distribution of data by unstretching large-scale values. Moreover, log transformation is useful to considerably reduce variation due to ontogeny (allometric effect) since we assumed that specimens of different developmental stages were subjected to our analyses. A Euclidean distance matrix was constructed using the log-transformed data. Multivariate analysis of variance (PERMANOVA) was applied

to test similarities within subspecies and localities ($P < 0.005$) (Anderson, 2001) using PRIMER software (version 6; Clarke & Gorley, 2006). A Permanova pairwise post-hoc test was performed to further investigate the differences between the subspecies ($P < 0.005$). Principal component analysis (PCA) was performed to characterize the differences between groups using the measured parameters (HT/CL and LT/CL ratios). Similarity percentage tests (SIMPER) were used to evaluate which measure contributed more to the differentiation between subspecies.

RESULTS

Morphological identification of taxonomic entities

All specimens loaned from the United States of America presented characteristics of the subspecies *A. a. carolinae* (hereafter named *A. a. USA*) (Fig. 2C). However, Brazilian samples fell into two categories: (i) samples exhibiting characteristics of the subspecies *A. a. americanus* sensu stricto (hereafter named “*A. a. Brazil 1*”; Fig. 2A) and (ii) samples that exhibited morphological divergence from the nominal subspecies (hereafter named “*A. a. Brazil 2*” (Fig. 2B)). The sampling localities of *A. a. Brazil 1* and *A. a. Brazil 2* are shown in Fig. 1.

Genetic distance

16S rRNA gene: The intraspecific distances varied from 0% (*A. a. USA*, *A. a. Brazil 2*, and *A. petrunkevitchi*) to 0.20% (*A. paraguayensis*) (Table 2). The interspecific distances between congeneric species varied from 0.99 to 11.8%. The distance to the outgroup was 25.9 to 28.5% (Table 3). Regarding the *A. americanus* subspecies, the lowest interspecific distance was between *A. a. Brazil 2* and *A. a. USA* (0.99%). The highest was *A. a. Brazil 1* and *A. a. USA* (2.26%) (Table 3).

COI gene: The intraspecific distances varied from 0.02% (*A. a. Brazil 1*) to 0.97% (*A. paraguayensis*) (Table 2). The interspecific distances between the congeneric species varied from 4.86 to 22.3%. The distance to the outgroup was 21.8 to 25.5% (Table 3). Regarding the *A. americanus* subspecies, the lowest interspecific distance was between *A. a. Brazil 1* and *A. a. USA* (4.86%). The highest was *A. a. Brazil 2* and *A. a. USA* (8.08%) (Table 3).

Phylogenetic analyses

The 16S rRNA phylogenetic analyses showed the following clades (Fig. 3): Clade 1, formed by *A. paraguayensis* sampled in north Brazil (Santarém and Porto de Moz/Pará); Clade 2, formed by *A. petrunkevitchi* sampled in southeast Brazil (São Vicente and Ubatuba/São Paulo); Clade 3, formed by *A. a. USA* sampled in the United States (Lumcon/LA and Horn Island/MS); Clade 4, formed by *A. a. Brazil 2* sampled in southeast Brazil (Cananéia/São Paulo and Macaé/Rio de Janeiro); Clade 5, formed by *A. a. Brazil 1* sampled in the south (Penha/Santa Catarina), southeast (Ubatuba, São Vicente and Cananéia/São Paulo, Macaé/Rio de Janeiro, Anchieta/Espirito Santo) and northeast Brazil (Baía Formosa/Rio Grande do Norte).

Based on the COI sequences, the phylogenetic analysis revealed the following clades (Fig. 4): Clade 1, formed by *A. a. Brazil 1* sampled in the south (Penha/Santa Catarina), southeast




Figure 2 Color photos of studied species of *Acetes*. Lateral view of *Acetes americanus* Ortmann, 1893. (A) Female of the *Acetes americanus* Brazil 1 (CCLC 0260); (B) female of the *Acetes americanus* Brazil 2 (CCLC 0255); (C) male of the *Acetes americanus* USA (CCLC 0268). CCLC: Crustacean Collection of the Laboratory of Biology of Marine and Freshwater Shrimp. Photo credit: Régis Augusto Pescinelli.

[Full-size](#)  DOI: [10.7717/peerj.14751/fig-2](https://doi.org/10.7717/peerj.14751/fig-2)

Table 2 Intraspecific genetic distance from 16S and COI genes. Average intraspecific genetic distance (%) from 16S and Cytochrome Oxidase 1 (COI) gene \pm standard deviation.

Subspecies and species	16S Genetic distance (\pm standard deviation)	COI Genetic distance (\pm standard deviation)
<i>Acetes americanus</i> USA	0.00 (\pm 0.000)	0.18 (\pm 0.001)
<i>Acetes americanus</i> Brazil 1	0.01 (\pm 0.001)	0.02 (\pm 0.000)
<i>Acetes americanus</i> Brazil 2	0.00 (\pm 0.000)	0.26 (\pm 0.001)
<i>Acetes paraguayensis</i>	0.20 (\pm 0.0012)	0.97 (\pm 0.003)
<i>Acetes petrunkevitchi</i>	0.00 (\pm 0.000)	0.26 (\pm 0.001)

Table 3 Interspecific distance from 16S and COI genes between species of *Acetes*. Matrix of average interspecific distance (%) from 16S (below) and Cytochrome Oxidase 1 (COI) gene (above) between species of *Acetes* (numbers on top) \pm standard deviation (values on bottom).

	<i>Acetes americanus</i> Brazil 1	<i>Acetes americanus</i> Brazil 2	<i>Acetes americanus</i> USA	<i>Acetes paraguayensis</i>	<i>Acetes petrunkevitchi</i>	<i>Belzebub faxoni</i>
<i>Acetes americanus</i> Brazil 1		6.44 (\pm 0.011)	4.86 (\pm 0.010)	22.1 (\pm 0.023)	19.0 (\pm 0.019)	21.8 (\pm 0.021)
<i>Acetes americanus</i> Brazil 2	1.97 (\pm 0.006)		8.08 (\pm 0.013)	22.0 (\pm 0.023)	20.2 (\pm 0.020)	22.9 (\pm 0.021)
<i>Acetes americanus</i> USA	2.26 (\pm 0.007)	0.99 (\pm 0.004)		23.3 (\pm 0.023)	21.6 (\pm 0.022)	23.7 (\pm 0.022)
<i>Acetes paraguayensis</i>	7.93 (\pm 0.012)	8.38 (\pm 0.013)	8.04 (\pm 0.013)		21.4 (\pm 0.021)	25.5 (\pm 0.021)
<i>Acetes petrunkevitchi</i>	11.8 (\pm 0.016)	11.1 (\pm 0.015)	11.4 (\pm 0.015)	10.0 (\pm 0.015)		25.4 (\pm 0.019)
<i>Belzebub faxoni</i>	26.6 (\pm 0.026)	25.9 (\pm 0.026)	26.3 (\pm 0.027)	28.5 (\pm 0.028)	27.5 (\pm 0.026)	

(Ubatuba and Cananéia/São Paulo, Macaé/Rio de Janeiro, Anchieta/Espírito Santo) and northeast Brazil (Baía Formosa/Rio Grande do Norte, Maceió/Alagoas); Clade 2, formed by *A. a.* USA sampled in the United States (Lumcon/LA and Horn Island/MS); Clade 3, formed by *A. a.* Brazil 2 sampled in southeast Brazil (Cananéia/SP and Macaé/RJ); Clade 4, formed by *A. sibogae* sampled in Peninsula Malaysia; Clade 5, formed by *A. japonicus* sampled in Peninsula Malaysia; Clade 6, formed by *A. serrulatus* sampled in Peninsula Malaysia; Clade 7, formed by *A. paraguayensis* sampled in north Brazil (Santarém and Porto de Moz/Pará); Clade 8, formed by *A. petrunkevitchi* sampled in southeast Brazil (Ubatuba/SP); Clade 9, formed by *A. indicus* sampled in Peninsula Malaysia.

The phylogenetic tree constructed based on the concatenated data (16S rRNA and COI) generated the same clades observed in the phylogenetic tree constructed with 16S rRNA and COI separately. In addition, *A. a.* USA and *A. a.* Brazil 2 were sister clades (Fig. 5), whereas COI, *A. a.* USA was a sister clade of *A. a.* Brazil 1 (Fig. 4).

The phylogenetic trees constructed (16S rRNA, COI, and concatenated data) resulted in the formation of two distinct clades of *A. americanus* sampled in Brazil with a high support value, in contrast to the clade formed by *A. a. carolinae* sampled in the United




Figure 3 Phylogenetic reconstruction of *Acetes* based on 16mt marker. Phylogenetic tree of Bayesian inference for the *Acetes* species based on the 16S region with Bayesian posterior probabilities indicated (only posterior probabilities > 50% are shown).

Full-size  DOI: 10.7717/peerj.14751/fig-3

States. Furthermore, the *A. americanus* group appears to be a sister taxon to all other *Acetes* (Fig. 4).

Population analyses

The haplotype network exhibited a genetic structure in the three groups, corresponding to those observed in the phylogenetic trees (Fig. 6). Available COI sequences of *A. americanus* ($N = 57$) resulted in 12 haplotypes of which six were unique, that is, represented by a single individual. “Brazil 1” presented the lowest haplotype ($h = 0.11$) and nucleotide ($\pi = 0.00022$) diversities. This group included 34 individuals sharing one haplotype and two individuals with unique haplotypes. “USA” and “Brazil 2” presented similar high haplotype ($h = 0.73$ and $h = 0.76$, respectively) and nucleotide diversities ($\pi = 0.00181$ and $\pi = 0.00277$, respectively).




Figure 4 Phylogenetic reconstruction of *Acetes* based on COI marker. Phylogenetic tree of Bayesian inference for the *Acetes* species based on the COI region with Bayesian posterior probabilities indicated (only posterior probabilities > 50% are shown).

Full-size [DOI: 10.7717/peerj.14751/fig-4](https://doi.org/10.7717/peerj.14751/fig-4)

Analysis of molecular variance (AMOVA) did not detect genetic structuring of any subspecies among the locations studied ($p > 0.05$) (Table 4).

Morphometric analysis

The ratio between the height and length of the thelycum was determined for several western Atlantic locations (Table 1). The ratio values were higher in *A. a.* USA and *A. a.* Brazil 2 than in *A. a.* Brazil 1 (Table 1). The studied *A. americanus* groups (BR1, BR2, and the USA) were morphologically different ($P = 0.0002$) (Table 5). PerMANOVA Pairwise tests indicated that at least one of the taxonomic entities is different from the others ($P = 0.0001$) (Table 6). However, when comparing *A. americanus* groups with each other, Pairwise tests indicated that there was a statistically significant difference only between Brazil 1 and Brazil 2 ($t = 3.1563$; $p = 0.0018$).




Figure 5 Phylogenetic reconstruction of *Acetes* based on concatenated markers. Phylogenetic tree of Bayesian inference for the *Acetes* species based on 16S and COI concatenated data with Bayesian posterior probabilities indicated (only posterior probabilities > 50% are shown).

Full-size DOI: 10.7717/peerj.14751/fig-5

The PCA visualization plot shows that *A. a.* Brazil 1 is more similar to *A. a.* Brazil 2 than to *A. a.* USA (Fig. 7). A simplified Simper test showed that the LT/CL RATIO was responsible for the differences between subspecies, contributing more than 73% for Brazil 1, more than 53% for Brazil 2 and more than 61% for the USA (Table 7; Fig. 7).

DISCUSSION

Our analyses revealed the existence of three lineages of *A. americanus*: *A. a.* Brazil 1 *sensu stricto*, *A. a.* Brazil 2, and *A. a.* USA. Through molecular analysis, we were also able to identify and contextualize another species of *Acetes* that occurs in Brazil, *A. paraguayensis*. Therefore, mitochondrial DNA can be considered an efficient tool for solving taxonomic identification of the genus *Acetes* at species level.

One of the species concepts widely accepted in systematics (Tsoi, Wang & Chu, 2005) is defined as a group of mating individuals or having the potential for it differing from other groups because they are reproductively isolated (Mayr, 1942). However, there are several alternative concepts of species (De Queiroz, 2007). The haplotype network results suggest that there are three possible lineages for *A. americanus*: *A. a.* Brazil 1 *sensu stricto*,




Figure 6 Haplotype network (COI region) showing the groups of *Acetes*. Haplotype network (COI region) according to a median-joining analysis, indicating the distribution of 12 haplotypes of *Acetes americanus* Ortman, 1893. The size of the circle of each haplotype is proportionate to its frequency in the sample. Each small dash represents a mutational step.

Full-size DOI: 10.7717/peerj.14751/fig-6

Table 4 Molecular variance with specimens of *Acetes americanus*. Analysis of molecular variance (AMOVA) performed with specimens of *Acetes americanus*.

		<i>Acetes americanus</i> Brazil 1	<i>Acetes americanus</i> Brazil 2	<i>Acetes americanus</i> USA
Variation source (%)	Among locations	−14.82	8.65	−12.50
	Within locations	114.82	91.35	112.50
FST (P)		−0.1482 (0.967)	0.0865 (0.152)	−0.125 (1.000)

Notes.

*Significant values, $P < 0.05$.

A. a. Brazil 2, and *A. a.* USA. Population-based analyses of mitochondrial DNA indicate that entities are reproductively isolated when gene flow is low (Tsoi, Wang & Chu, 2005). The low level of gene flow is essential evidence for speciation (Futuyma, 1998). Therefore, the non-sharing of haplotypes found in this study indicates that *A. a.* Brazil 1, *A. a.* Brazil 2, and *A. a.* USA are genetically distinct, with a low gene flow between them.

Table 5 Permanova analysis of *Acetes americanus* groups. Results of Permanova analysis considering *Acetes americanus* groups (BR1, BR2 and USA).

Source	Df	SS	MS	Pseudo-F	P(perm)	perms
Subspecies	1	1.3898E-2	1.3898E-2	7.1277	0.0058	9927
Locality	12	13.679	1.1399	584.6	0.0001	9909
Res	266	0.51867	1.9499E-3			

Notes.

D.f., degrees of freedom; SS, sum of squares; MS, mean square; Pseudo-F, statistic; P(perm), probability; Perms, permutations performed.

Table 6 Permanova Pairwise test of *Acetes americanus* groups. Results of the Permanova Pairwise test considering *Acetes americanus* groups (BR1, BR2 and USA).

Source	df	SS	MS	Pseudo-F	P(perm)	Unique perms
Subspecies	2	7.843	3.9215	76.786	0.0001	9949
Res	278	14.198	5,11E+02			
Total	280	22.041				

Notes.

D.f., degrees of freedom; SS, sum of squares; MS, mean square; Pseudo-F, statistic; P(perm), probability; Perms, permutations performed.




Figure 7 Principal component analysis depicting the morphometric groups of *Acetes*. (A) Principal Component Analysis (PCA) of the variation in thelycum morphology of the *Acetes americanus* groups (*A. americanus* USA, *A. americanus* Brazil 1 and *A. americanus* Brazil 2); (B) An enlarged area of the image “A”.

Full-size DOI: 10.7717/peerj.14751/fig-7

The *A. a.* Brazil 1 individuals sampled in several Brazilian regions (northeast: RN, AL, SE; southeast: ES, RJ, SP; south: SC) shared haplotypes between them, indicating the existence of gene flow among these populations. Recent studies on other decapod crustaceans have

Table 7 Results of similarity percentages analysis of body dimensions. Results of SIMPER (Similarity Percentages) analysis.

Species	Average Abundance	Average Similarity	Sim/SD	Contrib%	Cum.%
Group <i>Acetes americanus carolinae</i> (USA)					
Average similarity: 67.12					
LT/CL ratio	0.46	41.37	2.03	61.64	61.64
HT/CL ratio	0.30	25.75	1.89	38.36	100.00
Group <i>Acetes americanus</i> "BRAZIL 1"					
Average similarity: 63.52					
LT/CL ratio	0.24	46.81	1.91	73.69	73.69
HT/CL ratio	0.08	16.71	1.89	26.31	100.00
Group <i>Acetes americanus</i> "BRAZIL 2"					
Average similarity: 89.83					
LT/CL ratio	0.07	47.63	7.97	53.03	53.03
HT/CL ratio	0.06	42.20	7.71	46.97	100.00

Notes.

CL, Carapace Length; HT, height of the third thoracic sternite; LT, length of the third thoracic sternite; SD, Standard Deviation.

also shown genetic homogeneity among populations sampled along the western Atlantic, along the Brazilian coast (Laurenzano, Farias & Schubart, 2012; Terossi & Mantelatto, 2012; Rossi & Mantelatto, 2013; Wieman et al., 2013; Laurenzano, Mantelatto & Schubart, 2013; Carvalho-Batista et al., 2014; Teodoro et al., 2015; Nishikawa, Negri & Mantelatto, 2021). A potential reason for population homogeneity can be ascribed to the high dispersion capacity of planktonic larvae and the absence of barriers to gene flow. Although nothing is known about the larval dispersal of *A. americanus*, the dispersal power is high for species of the genus *Acetes*, as they exhibit long planktonic larval stages (~6 weeks) before becoming juveniles and adults (Rao, 1968). This premise has also been proposed for the other decapods tested (see references above).

The levels of genetic divergence (COI) among congeneric species of the crustaceans may vary up to 17%, a high value compared to other animal groups (Costa et al., 2007). Lepidopteran insects have a genetic divergence among congeneric species of only 6.1% (Hebert, Ratnasingham & Waard, 2003). Bird species show a variation of 7.93% (Hebert et al., 2004a), and fish have a 9.93% divergence (Ward et al., 2005). For sergestid shrimp, the rate of genetic divergence (COI) was also high. The genetic divergence found between *A. indicus*, *A. serrulatus*, *A. japonicus*, and *A. sibogae* ranged from 14.6 to 20.47% (Wong, 2013).

Our results indicated genetic divergence values (COI) from 4.86 to 8.08% between the *A. americanus* subspecies, which are low when compared with the studies mentioned above. However, if we compare these results to those of studies of cryptic or closely related shrimp species, these values are similar. Species morphologically similar displayed genetic divergences of 2.4 to 7% and were considered different (Gusmão, Lazoski & Solé-Cava, 2000; Lavery et al., 2004; Tsoi, Wang & Chu, 2005; Lagrue et al., 2014). Carvalho-Batista et al. (2019) found higher genetic divergence values (up to 13.5%) among the genus *Seabob*

Xiphopenaeus. However, among some closely related *Xiphopenaeus* species, the variation ranged from 2.7–3.3%.

The intraspecific distance (COI) (0.02–0.97%) was lower than the interspecific (4.86–22.3%) for *Acetes americanus* subspecies. This difference, known as the gap, is used by the DNA barcode technique to differentiate species (Hebert et al., 2004b; Ward, 2009; Carvalho-Batista et al., 2019), which reinforces that the three lineages analyzed can potentially be considered as different taxonomic entities, pending future morphological characterization. This study showed variation between *A. americanus* subspecies (0.99–2.26%) similar to the ones previously reported for penaeid shrimps (Costa et al., 2007; Silva, Mesquita & Paula, 2010).

Comparing only the measures of the height/length ratio of the female thelycum (Table 1), our results corroborated Omori (1975), which also found that this measurement differed between the two subspecies (*A. a. americanus* e *A. a. carolinae*). The values found by Omori (1975) for *A. a. carolinae* were 0.56–0.80 (mean 0.68) for North Carolina, 0.50–0.83 (0.66) for Louisiana and Texas, 0.70 and 0.53 for Panama and Suriname specimens, respectively, and values of 0.21–0.31 for specimens of *A. a. americanus* collected in Santos/SP. In our study, *A. a.* USA also presented higher values of height/length of the thelycum when compared to *A. a.* Brazil 1. However, we revealed the presence of *A. a.* Brazil 2 on the Brazilian coast showing high values of the height/length of the thelycum (Table 1). Therefore, if only this character is analyzed, the individual could be misidentified with *A. a.* USA.

PCA analysis showed a separation of the three subspecies groups (Brazil 1, Brazil 2 and USA) when considering thelycum measurements and the carapace length. In the transition areas where the subspecies occur, however, individuals from Suriname and Puerto Rico identified as *A. a.* Brazil 1 were close to *A. a.* USA, and individuals from Mexico identified as *A. a.* USA were close *A. a.* Brazil 1 and *A. a.* Brazil 2. These results point to an interesting geographic pattern, with the separation between individuals collected in the south (Terminos Mexico) and the north of the Gulf of Mexico - GOM (Louisiana). The two portions of the GOM have different water temperatures and have been separated into distinct biogeographic provinces by different authors (Boschi, 2000; Briggs & Bowen, 2012). Additionally, within the GOM, different cyclonic and anti-cyclonic flows occur separating the circulation in each locality (Schmitz Jr et al., 2005) which may be responsible for maintaining the isolation within each one of them. Therefore, this result can be associated with responses to the environmental conditions of the region, as phenotypic variations can be caused by both genetic information and environmental variations (Templeton, 2006). Considering these results, the molecular identification of the subspecies from these transition areas, using the protocols of this study, would be recommended to clarify this issue.

Acetes americanus Brazil 1 is genetically different from *A. a.* USA. In addition, the specimens sampled in Brazil formed two distinct clades: the first was composed of *A. a.* Brazil 1 and the second of *A. a.* Brazil 2. As stated earlier, *A. a.* Brazil 1 exhibits the diagnostic characteristics of *A. americanus americanus*, whereas *A. a.* Brazil 2 exhibits characteristics similar to those of *A. amecicanus carolinae*. Phylogenetic and population analyses pointed

to the divergence of *A. a.* Brazil 2 from *A. a.* USA. Usually, a single characteristic to be fixed after reproductive isolation is sufficient for the diagnosis of a species (Mink & Sites Jr, 1996). However, no morphological characteristics that can discriminate these subspecies have been detected thus far. Several studies have shown that many species with few or no morphological characteristics are distinguished by genetic differences (Reuschel, Cuesta & Schubart, 2010; Puillandre et al., 2011; Carvalho, Magalhães & Mantelatto, 2014; Delić et al., 2017; Mandai et al., 2018). A study focusing on the morphological characteristics used to discriminate *A. americanus* in Brazil should clarify and provide more information about this new possible taxonomic entity. Furthermore, the hypothesis that *A. a.* Brazil 2 is a different entity, previously described (*A. a. louisianensis* or *A. a. limonensis*), cannot be disregarded.

General phylogeny

A previous phylogenetic study of *Acetes* proposed that the *Acetes* clade without *A. petrunkevitchi* (former *Peisos petrunkevitchi*) never gained robust support, thus considering *Peisos* as a junior synonym of *Acetes* (Vereshchaka, Lunina & Olesen, 2016a). Although we carried out an analysis with a robust but limited number of samples, our results confirmed the phylogenetic positioning recovered by both 16S rRNA and COI genes and indicated that *A. petrunkevitchi* is part of the *Acetes* group, but in all analyses forming a single clade of “*Acetes paraguayensis* + *A. petrunkevitchi*”, with a high support value for 16S rRNA and concatenated data. As only mitochondrial genes were used, further studies adding nuclear markers should be carried out to test the topology recovered herein.

However, recent studies about the morphology of the male reproductive system and spermatophore of *A. petrunkevitchi* differed from those of *A. americanus*, *A. marinus* and *A. paraguayensis*, which remains open the discussion about the inclusion of *Peisos* in the *Acetes* group (Salti, 2020).

The global morphological phylogeny proposed for the superfamily Sergestoidea showed significant changes in taxonomy, with the description of three new families, particularly because the family Sergestidae was not considered monophyletic (Vereshchaka, 2017). As a result, *A. americanus* was classified in the new family Acetidae, as proposed by Vereshchaka (2017). The author proposed that a single clade of *A. marinus* and *A. paraguayensis* within Acetidae received high bootstrap support. Our concatenated molecular phylogeny indicated a close relationship between *A. paraguayensis* and *A. petrunkevitchi* in a separate clade, as mentioned above. Additional samples of these species, including *A. marinus*, confirm this hypothesis. This molecular reconstruction sheds light on the unsolved evolutionary relations between the species of the genus *Acetes*, which should be investigated using more comprehensive integrated studies and the addition of nuclear markers.

ACKNOWLEDGEMENTS

We thank the LABCAM and LBSC members for their help during the fieldwork, molecular and morphometric analysis, and their valuable suggestions and contributions. The authors are thankful to the curators of the following crustacean collections for providing valuable helping with samplings, loans and donations samples for this work: Adriane A. Braga

(UFES), Cristiana S. Serejo (MNRJ), Darryl L. Felder (ULLZ), Dirk Brandis (ZMK), Gustavo L. Hirose (UFS), Glauca Pontes (MCP), Inácia M. Vieira (IEPA), Jessor F. de Souza-Filho (MOUFPE), Jose L. Villalobos (CNCR), Luis F. Dumont (FURG), Lourdes Rojas (YPM), Marcos S. Tavares (MZUSP), Rafael Lemaitre (USNM), and Raymond Bauer (ULL). We thank the anonymous reviewers for their suggestions to improve the quality of the manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Financial support for this project was provided by research grants from São Paulo Research Foundation - FAPESP (Temáticos Biota 2010/50188-8 and INTERCRUSTA 2018/13685-5; Post-doctoral fellowship 2014/01632-3 to SSM). Additional support was provided by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES - Código de Financiamento 001 (Ciências do Mar II Proc. 2005/2014 - 23038.004308/201414). Fernando L. Mantelatto, Rogério Caetano Costa, and Fabricio Lopes Carvalho received support from Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq (Research Scholarships PQ 302253/2019-0, 306672/2018-9, and 315997/2021-4, respectively). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

São Paulo Research Foundation - FAPESP (Temáticos Biota): 2010/50188-8, INTERCRUSTA 2018/13685-5; 2014/01632-3 PROTAX 2021/08075-6.

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES - Código de Financiamento 001 (Ciências do Mar II Proc.): 2005/2014 - 23038.004308/201414.

Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq: PQ 302253/2019-0, 306672/2018-9, 315997/2021-4.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Sabrina Morilhas Simões conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Rogério Caetano Costa conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, grant support coordinator, and approved the final draft.
- Fabricio Lopes Carvalho performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Abner Carvalho-Batista performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.

- Sarah de Souza Alves Teodoro performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Fernando L. Mantelatto conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, grant support coordinator, and approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

All specimens used in our analysis were collected under Federal authorization from the Instituto Chico Mendes de Biodiversidade/ICMBio (#11777-2, #23008-1).

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The group of mitochondrial (16S and COI) sequences are available at GenBank: [OP035650–OP035701](#) (16S) and [OP060465–OP060530](#) (COI).

Data Availability

The following information was supplied regarding data availability:

The raw data is available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.14751#supplemental-information>.

REFERENCES

- Anderson M. 2001.** A new method for non-parametric multivariate analysis of variance. *Austral Ecology* **26**:32–46 DOI [10.1111/j.1442-9993.2001.01070.pp.x](https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x).
- Bandelt HJ, Forster P, Röhl A. 1999.** Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* **16**:37–48 DOI [10.1093/oxfordjournals.molbev.a026036](https://doi.org/10.1093/oxfordjournals.molbev.a026036).
- Boschi EE. 2000.** Species of decapod crustaceans and their distribution in the American marine zoogeographic provinces. *Revista de Investigacion y Desarrollo Pesquero* **13**:7–64.
- Briggs JC, Bowen BW. 2012.** A realignment of marine biogeographic provinces with particular reference to fish distributions. *Journal of Biogeography* **39**(1):12–30 DOI [10.1111/j.1365-2699.2011.02613.x](https://doi.org/10.1111/j.1365-2699.2011.02613.x).
- Burkenroad MD. 1934.** Littoral Penaeidae chiefly from the Bingham Oceanographic Collection. *Bulletin of the Bingham Oceanographic Collection* **4**:1–109.
- Carvalho FL, Magalhães C, Mantelatto FL. 2014.** Molecular and morphological differentiation between two Miocene-divergent lineages of Amazonian shrimps, with the description of a new species (Decapoda, Palaemonidae, Palaemon). *ZooKeys* **457**:79–108 DOI [10.3897/zookeys.457.6771](https://doi.org/10.3897/zookeys.457.6771).

- Carvalho FL, Magalhães C, Mantelatto FL. 2020.** A molecular and morphological approach on the taxonomic status of the Brazilian species of *Palaemon* (Decapoda, Palaemonidae). *Zoologica Scripta* **49**:101–116 DOI [10.1111/zsc.12394](https://doi.org/10.1111/zsc.12394).
- Carvalho-Batista A, Negri M, Pileggi LG, Castilho AL, Costa RC, Mantelatto FL. 2014.** Inferring population connectivity across the range of distribution of the stiletto shrimp *Artemesia longinaris* Spence Bate, 1888 (Decapoda, Penaeidae) from DNA barcoding: implications for fishery management. *ZooKeys* **457**:271–288 DOI [10.3897/zookeys.457.6569](https://doi.org/10.3897/zookeys.457.6569).
- Carvalho-Batista A, Negri M, Tamburus AF, Costa RC, Castilho AL, Zara FJ, Mantelatto FL. 2018.** Genetic comparison of the red shrimp *Pleoticus muelleri* (Decapoda:Solenoceridae) using the barcode gene reveals the absence of cryptic speciation along its distribution. *Regional Studies in Marine Science* **24**:392–399 DOI [10.1016/j.rsma.2018.10.003](https://doi.org/10.1016/j.rsma.2018.10.003).
- Carvalho-Batista A, Terossi M, Zara FJ, Mantelatto FL, Costa RC. 2019.** A multigene and morphological analysis expands the diversity of the seabod shrimp *Xiphopenaeus* Smith, 1869 (Decapoda: Penaeidae), with descriptions of two new species. *Scientific Reports* **9**:15281 DOI [10.1038/s41598-019-51484-3](https://doi.org/10.1038/s41598-019-51484-3).
- Clarke KR, Gorley RN. 2006.** Primer. In: *PRIMER-e*. 866. Plymouth.
- Costa FO, de Waard JR, Boutillier J, Ratnasingham S, Dooh RT, Hajibabaei M, Hebert DN. 2007.** Biological identification through DNA barcodes: the case of the Crustacea. *Canadian Journal of Fisheries and Aquatic Sciences* **64**:272–295 DOI [10.1139/f07-008](https://doi.org/10.1139/f07-008).
- Darriba D, Taboada GL, Doallo R, Posada D. 2012.** jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* **9**(8):772 DOI [10.1038/nmeth.2109](https://doi.org/10.1038/nmeth.2109).
- De Grave S, Fransen CHJM. 2011.** Carideorum catalogus: the recent species of the dendrobranchiate, stenopodidean, procarididean and caridean shrimps (Crustacea: Decapoda). *Zoologische Mededeelingen* **85**(9):195–589.
- De Queiroz K. 2007.** Species concepts and species delimitation. *Systematic Biology* **56**(6):879–886 DOI [10.1080/10635150701701083](https://doi.org/10.1080/10635150701701083).
- Delić T, Trontelj P, Rendoš M, Fišer C. 2017.** The importance of naming cryptic species and the conservation of endemic subterranean amphipods. *Scientific Reports* **7**:3391 DOI [10.1038/s41598-017-02938-z](https://doi.org/10.1038/s41598-017-02938-z).
- D’Incao F, Martins STS. 2000.** Brazilian species of the genera *Acetes* H. Milne Edwards, 1830 and *Peisos* Burkenroad, 1945 (Decapoda: Sergestidae). *Journal of Crustacean Biology* **20**(2):78–86 DOI [10.1163/1937240X-90000010](https://doi.org/10.1163/1937240X-90000010).
- Excoffier L, Laval G, Schneider S. 2005.** Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* **1**:47–50.
- Excoffier L, Smouse PE, Quattro JM. 1992.** Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131**(2):479–491 DOI [10.1093/genetics/131.2.479](https://doi.org/10.1093/genetics/131.2.479).

- França NFC, de Moraes AB, Carvalho-Batista A, de Melo MCRB, López-Greco L, Mantelatto FL, de Morais Freire FA. 2019.** *Farfantepenaeus subtilis* (Pérez-Farfante, 1967) and *F. brasiliensis* (Latreille, 1817) (Decapoda, Penaeidae): Ontogenetic comparison using the combined analysis of secondary sexual characters and molecular markers. *Fisheries Research* **216**:89–95 DOI [10.1016/j.fishres.2019.03.024](https://doi.org/10.1016/j.fishres.2019.03.024).
- França NFC, Moraes AB, Carvalho-Batista A, Melo MCRB, Zara FJ, Mantelatto FL, Freire FAM. 2021.** An integrative approach using DNA barcode and scanning electron microscopy for the effective identification of sympatric species of the genus *Farfantepenaeus* Burukovsky, 1997. *Regional Studies in Marine Science* **43**:101670 DOI [10.1016/j.rsma.2021.101670](https://doi.org/10.1016/j.rsma.2021.101670).
- Futuyma DJ. 1998.** *Evolutionary biology*. 3rd ed. Sunderland: Sinauer.
- Gusmão J, Lazoski C, Solé-Cava AM. 2000.** A new species of *Penaeus* (Crustacea: Penaeidae) revealed by allozyme and cytochrome oxidase I analysis. *Marine Biology* **137**:435–446 DOI [10.1007/s002270000365](https://doi.org/10.1007/s002270000365).
- Gusmão J, Lazoski C, Solé-Cava AM. 2005.** Population genetic structure of Brazilian shrimp species (*Farfantepenaeus* sp., *F. brasiliensis*, *F. paulensis* and *Litopenaeus schmitti*: Decapoda: Penaeidae). *Genetics and Molecular Biology* **28**(1):165–171 DOI [10.1590/S1415-47572005000100029](https://doi.org/10.1590/S1415-47572005000100029).
- Hall TA. 1999.** BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**:95–98 Available at <http://www.mbio.ncsu.edu/JWB/papers/1999Hall1.pdf>.
- Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W. 2004b.** Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences of the United States of America* **101**(41):14812–14817 DOI [10.1073/pnas.0406166101](https://doi.org/10.1073/pnas.0406166101).
- Hebert PDN, Ratnasingham S, Waard JR. 2003.** Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society B: Biological Sciences* **270**:S96–S99 DOI [10.1098/rsbl.2003.0025](https://doi.org/10.1098/rsbl.2003.0025).
- Hebert PDN, Stoeckle MY, Zemplak TS, Francis CM. 2004a.** Identification of birds through DNA barcodes. *PLOS Biology* **2**(10):e312 DOI [10.1371/journal.pbio.0020312](https://doi.org/10.1371/journal.pbio.0020312).
- Holthuis LB. 1948.** Notes on some Crustacea Decapoda Natantia from Surinam. *Proceedings Koninklijke Nederlandsche Akademie van Wetenschappen* **51**:1104–1113.
- Kimura M. 1980.** A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* **16**:111–120 DOI [10.1007/BF01731581](https://doi.org/10.1007/BF01731581).
- Lagrange C, Wattier R, Galipaud M, Gauthey Z, Rullmann J-P, Dubreuil C, Rigaud T, Bollache L. 2014.** Confrontation of cryptic diversity and mate discrimination within *Gammarus pulex* and *Gammarus fossarum* species complexes. *Freshwater Biology* **59**:2555–2570 DOI [10.1111/fwb.12453](https://doi.org/10.1111/fwb.12453).
- Laurenzano C, Farias NE, Schubart CD. 2012.** Mitochondrial genetic structure of two populations of *Uca uruguayensis* fails to reveal an impact of the Rio de la Plata on gene flow. *Nauplius* **20**:15–25 DOI [10.1590/S0104-64972012000100003](https://doi.org/10.1590/S0104-64972012000100003).

- Laurenzano C, Mantelatto FL, Schubart CD. 2013.** South American homogeneity versus Caribbean heterogeneity: population genetic structure of the western Atlantic fiddler crab *Uca rapax* (Brachyura, Ocypodidae). *Journal of Experimental Marine Biology and Ecology* **449**:22–27 DOI [10.1016/j.jembe.2013.08.007](https://doi.org/10.1016/j.jembe.2013.08.007).
- Lavery S, Chan TY, Chan YK, Tam YK, Chu HK. 2004.** Phylogenetic relationships and evolutionary history of the shrimp genus *Penaeus* s.l. derived from mitochondrial DNA. *Molecular Phylogenetics and Evolution* **31**:39–49 DOI [10.1016/j.ympev.2003.07.015](https://doi.org/10.1016/j.ympev.2003.07.015).
- Maggioni R, Rogers AD, Maclean N, D’Incao F. 2001.** Molecular phylogeny of Western Atlantic *Farfantepenaeus* and *Litopenaeus* shrimp based on mitochondrial 16S partial sequences. *Molecular Phylogenetics and Evolution* **18**(1):66–73 DOI [10.1006/mpev.2000.0866](https://doi.org/10.1006/mpev.2000.0866).
- Mandai SS, Buranelli RC, Schubart CD, Mantelatto FL. 2018.** Phylogenetic and phylogeographic inferences based on two DNA markers reveal geographic structure of the orange claw hermit crab *Calcinus tibicen* (Anomura: Diogenidae) in the western Atlantic. *Marine Biology Research* **14**(6):565–580 DOI [10.1080/17451000.2018.1497184](https://doi.org/10.1080/17451000.2018.1497184).
- Mantelatto FL, Carvalho FL, Simões SM, Negri M, Souza-Carvalho EA, Terossi M. 2016.** New primers for amplification of cytochrome c oxidase subunit I barcode region designed for species of Decapoda (Crustacea). *Nauplius* **24**:e2016030 DOI [10.1590/2358-2936e2016030](https://doi.org/10.1590/2358-2936e2016030).
- Mantelatto FL, Pardo LM, Pileggi LG, Felder DL. 2009a.** Taxonomic reexamination of the hermit crab species *Pagurus forceps* and *Pagurus comptus* (Decapoda, Paguridae) by molecular analysis. *Zootaxa* **2133**:20–32 DOI [10.11646/zootaxa.2133.1.2](https://doi.org/10.11646/zootaxa.2133.1.2).
- Mantelatto FL, Robles R, Felder DL. 2007.** Molecular phylogeny of the western Atlantic species of the genus *Portunus* (Crustacea: Brachyura, Portunidae). *Zoological Journal of the Linnean Society* **150**(1):211–220 DOI [10.1111/j.1096-3642.2007.00298.x](https://doi.org/10.1111/j.1096-3642.2007.00298.x).
- Mantelatto FL, Robles R, Schubart CD, Felder DL. 2009b.** Molecular phylogeny of the genus *Cronius* Stimpson 1860, with reassignment of *C. tumidulus* and several American species of *Portunus* to the genus *Achelous* de Haan, 1833 (Brachyura: Portunidae). In: Martin JW, Crandall KA, Felder DL, eds. *Crustacean issues: decapod crustacean phylogenetics*. Boca Raton: CRC Press, 567–579.
- Mantelatto FL, Tamburus AF, Carvalho-Batista A, Rossi N, Buranelli RC, Pantaleão JAF, Teles JN, Zara FJ, Carvalho FL, Bochini GL, Terossi M, Robles R, Castilho AL, Costa RC. 2022.** Checklist of decapod crustaceans from the coast of the São Paulo state (Brazil) supported by integrative molecular and morphological data: V. Dendrobranchiata and Pleocyemata [Achelata, Astacidea, Axiidea, Caridea (Alpheoidea and Processoidea excluded), Gebiidea, Stenopodidea. *Zootaxa* **5121**(1):1–074 DOI [10.11646/zootaxa.5121.1.1](https://doi.org/10.11646/zootaxa.5121.1.1).
- Marramà G, Kriwet J. 2017.** Principal component and discriminant analyses as powerful tools to support taxonomic identification and their use for functional and phylogenetic signal detection of isolated fossil shark teeth. *PLOS ONE* **12**(11):e0188806 DOI [10.1371/journal.pone.0188806](https://doi.org/10.1371/journal.pone.0188806).

- Mayr E. 1942.** *Systematics and the origin of species from the viewpoint of a zoologist*. New York: Columbia University Press.
- Mink DG, Sites Jr JW. 1996.** Species limits, phylogenetic relationships, and origins of viviparity in the *Scalaris* complex of the lizard genus *Sceloporus* (Phrynosomatidae). *Herpetologica* 52:551–557.
- Nishikawa KS, Negri M, Mantelatto FL. 2021.** Unexpected absence of population structure and high genetic diversity of the western Atlantic hermit crab *Clibanarius antillensis* Stimpson, 1859 (Decapoda: Diogenidae) based on mitochondrial markers and morphological data. *Diversity* 13(2):56 DOI 10.3390/d13020056.
- Omori M. 1975.** The systematics, biogeography, and fishery of epipelagic shrimps of the genus *Acetes* (Crustacea, Decapoda, Sergestidae). *Bulletin of the Ocean Research Institute, University of Tokyo* 7:1–91.
- Pileggi LG, Mantelatto FL. 2010.** Molecular phylogeny of the freshwater prawn genus *Macrobrachium* (Decapoda, Palaemonidae), with emphasis on the relationships among selected American species. *Invertebrate Systematics* 24(1):194–208 DOI 10.1071/IS09043.
- Puillandre N, Macpherson E, Lambourdière J, Cruaud C, Boisselierdubayle MC, Samadi S. 2011.** Barcoding type specimens helps to identify synonyms and an unnamed new species in *Eumunida* Smith, 1883 (Decapoda: Eumunididae). *Invertebrate Systematics* 25(4):322–333 DOI 10.1071/IS11022.
- Rambaut A. 2007.** FigTree. Available at <http://tree.bio.ed.ac.uk/software/figtree/>.
- Rao PV. 1968.** A new species of shrimp, *Acetes cochinesis* (Crustacea: Decapoda, Sergestidae) from southwest coast of India with an account of its larval development. *Journal of the Marine Biological Association of India* 10(2):298–320.
- Reuschel S, Cuesta JA, Schubart CD. 2010.** Marine biogeographic boundaries and human introduction along the European coast revealed by phytogeography of the prawn *Palaemon elegans*. *Molecular Phylogenetics and Evolution* 55(3):765–775 DOI 10.1016/j.ympev.2010.03.021.
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Huelsenbeck JP. 2012.** MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3):539–542 DOI 10.1093/sysbio/sys029.
- Ronquist F, Van Der Mark P, Huelsenbeck JP. 2009.** Bayesian phylogenetic analysis using MrBays. In: Lemey P, Sameli M, Vandamme A-M, eds. *The phylogenetic handbook*. Cambridge: Cambridge University Press, 210–266.
- Rossi N, Mantelatto FL. 2013.** Molecular analysis of the freshwater prawn *Macrobrachium olfersii* (Decapoda, Palaemonidae) supports the existence of a single species throughout its distribution. *PLOS ONE* 8(1):e54698 DOI 10.1371/journal.pone.0054698.
- Rozas J, Rozas R. 1999.** DnaSP version 3.0: an integrated program for molecular population genetic and molecular evolution analysis. *Bioinformatics* 15:174–175 DOI 10.1093/bioinformatics/15.2.174.

- Salti FC. 2020.** Sistema reprodutor masculino de camarões Dendrobranchiata sob aspectos evolutivos: padrões morfo-histológicos e reconstrução de estado ancestral de caráter. Master Dissertation, Universidade Estadual Paulista Júlio de Mesquita Filho, Instituto de Biociências, Rio Claro, 139.
- Schmitz Jr WJ, Biggs DC, Lugo-Fernandez A, Oey L-Y, Sturges W. 2005.** A synopsis of the circulation in the Gulf of Mexico and on its continental margins. *Washington DC American Geophysical Union Geophysical Monograph Series* **161**:11–29 DOI [10.1029/161GM03](https://doi.org/10.1029/161GM03).
- Schubart CD, Huber MGJ. 2006.** Genetic comparisons of german populations of the stone crayfish. *Austropotamo biustorrentium (Crustacea: Astacidae)*. *Bulletin Francais de la Pêche et de la Pisciculture* **380**:1019–1028.
- Silva IC, Mesquita N, Paula J. 2010.** Genetic and morphological differentiation of the mangrove crab *Perisesarma guttatum* (Brachyura: Sesarmidae) along an East African latitudinal gradient. *Biological Journal of the Linnean Society* **99**:28–46 DOI [10.1111/j.1095-8312.2009.01338.x](https://doi.org/10.1111/j.1095-8312.2009.01338.x).
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011.** MEGA 5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* **28**:2731–2739 DOI [10.1093/molbev/msr121](https://doi.org/10.1093/molbev/msr121).
- Tavares C, Gusmão J. 2016.** Description of a new Penaeidae (Decapoda: Dendrobranchiata) species, *Farfantepenaeus isabelae* sp. nov. *Zootaxa* **4171**(3):505–516 DOI [10.11646/zootaxa.4171.3.6](https://doi.org/10.11646/zootaxa.4171.3.6).
- Templeton AR. 2006.** *Population genetics and microevolutionary theory*. Hoboken: John Wiley & Sons, 705 DOI [10.1002/0470047356](https://doi.org/10.1002/0470047356).
- Teodoro SSA, Terossi M, Costa RC, Mantelatto FL. 2015.** Genetic homogeneity in the commercial pink shrimp *Farfantepenaeus paulensis* revealed by COI barcoding gene. *Estuarine, Coastal and Shelf Science* **166**:124–130 DOI [10.1016/j.ecss.2015.07.009](https://doi.org/10.1016/j.ecss.2015.07.009).
- Teodoro SSA, Terossi M, Mantelatto FL, Costa RC. 2016.** Discordance in the identification of juvenile pink shrimp (*Farfantepenaeus brasiliensis* and *F. paulensis*: family Penaeidae): an integrative approach using morphology, morphometry and barcoding. *Fisheries Research* **183**:244–253 DOI [10.1016/j.fishres.2016.06.009](https://doi.org/10.1016/j.fishres.2016.06.009).
- Terossi M, Mantelatto FL. 2012.** Morphological and genetic variability in *Hippolyte obliquimanus* Dana, 1852 (Decapoda, Caridea, Hippolytidae) from Brazil and the Caribbean Sea. *Crustaceana* **85**:685–712 DOI [10.1163/156854012X643762](https://doi.org/10.1163/156854012X643762).
- Terossi M, Mantelatto FL. 2020.** Revalidation of *Ogyrides occidentalis* (Ortmann, 1893) (Decapoda: Caridea: Ogyrididae) from Brazil based on morphological and mtDNA evidences. *Journal of Crustacean Biology* **40**:627–633 DOI [10.1093/jcabi/ruaa054](https://doi.org/10.1093/jcabi/ruaa054).
- Thompson JD, Higging DG, Gibson TJ. 1994.** CLUSTALW: improving the sensitivity of progressive multiple sequence alignment through sequence weighting specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**:4673e4680 DOI [10.1093/nar/22.22.4673](https://doi.org/10.1093/nar/22.22.4673).

- Tsoi KH, Wang ZY, Chu KH. 2005.** Genetic divergence between two morphologically similar varieties of the kuruma shrimp *Penaeus japonicus*. *Marine Biology* **147**:367–379 DOI [10.1007/s00227-005-1585-x](https://doi.org/10.1007/s00227-005-1585-x).
- Vereshchaka AL. 2017.** The shrimp superfamily Sergestoidea: a global phylogeny with definition of new families and an assessment of the pathways into principal biotopes. *Royal Society Open Science* **4**(9):170221 DOI [10.1098/rsos.170221](https://doi.org/10.1098/rsos.170221).
- Vereshchaka AL, Lunina AA, Olesen J. 2016a.** Phylogeny and classification of the shrimp genera *Acetes*, *Peisos*, and *Sicyonella* (Sergestidae: Crustacea: Decapoda). *Zoological Journal of the Linnean Society* **177**(2):353–377 DOI [10.1111/zoj.12371](https://doi.org/10.1111/zoj.12371).
- Vereshchaka AL, Olesen J, Lunina AA. 2016b.** A phylogeny-based revision of the family Luciferidae (Crustacea: Decapoda). *Zoological Journal of the Linnean Society* **178**:15–32 DOI [10.1111/zoj.12398](https://doi.org/10.1111/zoj.12398).
- Vergamini FG, Pileggi LG, Mantelatto FL. 2011.** Genetic variability of the Amazon River prawn *Macrobrachium amazonicum* (Decapoda, Caridea, Palaemonidae). *Contributions to Zoology* **80**(1):67–83 DOI [10.1163/18759866-08001003](https://doi.org/10.1163/18759866-08001003).
- Ward RD. 2009.** DNA barcode divergence among species and genera of birds and fishes. *Molecular Ecology Resources* **9**:1077–1085 DOI [10.1111/j.1755-0998.2009.02541.x](https://doi.org/10.1111/j.1755-0998.2009.02541.x).
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PD. 2005.** DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences* **360**(1462):1847–1857 DOI [10.1098/rstb.2005.1716](https://doi.org/10.1098/rstb.2005.1716).
- Wieman AC, Berendzen PB, Hampton KR, Jang J, Hopkins MJ, Jurgenson J, McNamara JC, Thurman CL. 2013.** A panmictic fiddler crab from the coast of Brazil? Impact of divergent ocean currents and larval dispersal potential on genetic and morphological variation in *Uca maracoani*. *Marine Biology* **161**:173–185 DOI [10.1007/s00227-013-2327-0](https://doi.org/10.1007/s00227-013-2327-0).
- Wong BW. 2013.** Genetic diversity and morphometric characterization of *Acetes* (Decapoda: Sergestidae) species collected from the West Coast of Peninsular Malaysia. Ph D Thesis, Faculty to the Engineering and Science, 204.
- WoRMS. 2022.** *Acetes* H. Milne Edwards, 1830. Available at <https://www.marinespecies.org/aphia.php?p=taxdetails&id=158343>.
- Xiao Y, Greenwood JG. 1993.** The biology of *Acetes* (Crustacea, Sergestidae). *Oceanography and Marine Biology: An Annual Review* **31**:259–444.

FURTHER READING

- Crandall KA, Fitzpatrick Jr JF. 1996.** Crayfish molecular systematics: using a combination of procedures to estimate phylogeny. *Systematic Biology* **45**:1–26 DOI [10.1093/sysbio/45.1.1](https://doi.org/10.1093/sysbio/45.1.1).
- Schubart CD, Cuesta JA, Felder DL. 2002.** Glyptograpsidae, a new brachyuran family from Central America: larval and adult morphology, and a molecular phylogeny of the Grapsoidea. *Journal of Crustacean Biology* **22**(1):28–44.