

RT-MAT 2003-11

**QUOTIENTS OF INCIDENCE ALGEBRAS
AND THE EULER CHARACTERISTIC**

**Ibrahim Assem, Diane Castonguay,
Eduardo N. Marcos and Sonia Trepode**

Junho 2003

Esta é uma publicação preliminar (“preprint”).

QUOTIENTS OF INCIDENCE ALGEBRAS AND THE EULER CHARACTERISTIC

Ibrahim Assem, Diane Castonguay,
Eduardo N. Marcos and Sonia Trepode

Dedicated to Otto Kerner in his 60th birthday

ABSTRACT. In this note, we show that, if $A \cong kQ_A/I_A$ is a schurian strongly simply connected algebra given by its normed presentation, and Σ is the unique poset whose oriented Hasse diagram coincides with Q_A , then $A \cong k\Sigma$ if and only if I_A has a generating set consisting of exactly $\chi(Q_A)$ elements, where $\chi(Q_A)$ is the Euler characteristic of Q_A . We also prove that a quotient of an incidence algebra $A = k\Sigma/J$ is strongly simply connected if and only if A is simply connected and $k\Sigma$ is strongly simply connected.

1. Introduction.

In this note, we show how the Euler characteristic of a quiver can be used to study the strong simple connectedness of incidence algebras and their quotients.

Our motivation comes from the study of the finite dimensional algebras over an algebraically closed field k . For such an algebra A , there exists a quiver Q_A , and an ideal I of the path algebra kQ_A such that $A \cong kQ_A/I$. For each such pair (Q_A, I) , called a presentation of A , one can define the

2000 Mathematics subject classification: 16G20

Keywords and phrases: incidence algebras, strongly simply connected algebras, Euler characteristic.

fundamental group $\pi_1(Q_A, I)$, see for instance, [12]. The algebra A is called simply connected if Q_A is acyclic and, for each presentation (Q_A, I) , the group $\pi_1(Q_A, I)$ is trivial [5]. If A is the incidence algebra of a (finite) poset Σ , then its quiver Q_A coincides with the (oriented) Hasse diagram of Σ . Moreover, all presentations of the incidence algebra A give rise to isomorphic fundamental groups [6] and A is simply connected if and only if so is the associated simplicial complex [8, 14]. While it is difficult to find criteria for an algebra to be simply connected (see, for instance, [4]), one subclass seems easier to handle: this is the class of strongly simply connected algebras, introduced by Skowroński in [16]. An algebra A is called strongly simply connected if every full convex subcategory of A is simply connected. Characterizations of strongly simply connected algebras have been given in [2]. In particular, an incidence algebra is strongly simply connected if and only if its quiver contains no crowns [9] or, equivalently, if and only if the corresponding poset is dismantlable [15].

It was already observed in [9] that, if A is a schurian strongly simply connected algebra, then there exists a unique poset Σ whose oriented Hasse diagram coincides with Q_A . As a consequence of this result and [2](2.4), there exists a presentation of A , called normed presentation, so that A is the quotient of the incidence algebra $k\Sigma$ by an ideal generated by paths.

In this paper, using properties of the Euler characteristic $\chi(Q_A)$ of the quiver Q_A , we first show that a quotient A of an incidence algebra $k\Sigma$ is strongly simply connected if and only if it is simply connected and $k\Sigma$ is strongly simply connected. We next prove that, if $A = kQ_A/I$ is a schurian strongly simply connected algebra, then the ideal

I has at least $\chi(Q_A)$ generators and it reaches this number if and only if A is an incidence algebra. In a forthcoming paper [1], we give several new characterizations of schurian strongly simply connected algebras.

The paper is organized as follows. Section 2 is devoted to fixing the notation and briefly recalling the necessary concepts and results. In section 3, we show that, if $k\Sigma$ is a strongly simply connected incidence algebra, then $k\Sigma \cong kQ_\Sigma/I_\Sigma$, where I_Σ has a generating set consisting of exactly $\chi(Q_\Sigma)$ elements. We prove our main results in section 4.

2. Preliminaries.

2.1. Notation. In this paper, by algebra, we always mean a basic and connected finite dimensional algebra over an algebraically closed field k . Given a quiver Q , we denote by Q_0 its set of points and by Q_1 its set of arrows. A *relation* in Q from a point x to a point y is a linear combination $\rho = \sum_{i=1}^m \lambda_i w_i$ where, for each i , $\lambda_i \in k$ is non-zero and w_i is a path of length at least two from x to y . A relation ρ as before is called *monomial* if $m = 1$, *binomial* if $m = 2$ and a *commutativity relation* if it equals the difference of two paths. We denote by kQ the path algebra of Q and by $kQ(x, y)$ the k -vector space generated by all paths in Q from x to y . For an algebra A , we denote by Q_A its quiver. For every algebra A , there exists an ideal I in kQ_A , generated by a set of relations, such that $A \cong kQ_A/I$. The pair (Q_A, I) is called a *presentation* of A . An algebra $A = kQ/I$ can equivalently be considered as a k -category of which the object class A_0 is Q_0 , and the set of morphisms $A(x, y)$ from x to y is the quotient of $kQ(x, y)$ by the subspace $I(x, y) = I \cap kQ(x, y)$, see [7]. A full subcategory B of A is called *convex* if any path in A with source and

target in B lies entirely in B . An algebra A is called *triangular* if Q_A is acyclic, and it is called *schurian* if, for all $x, y \in A_0$, we have $\dim_k A(x, y) \leq 1$. In this paper, we deal exclusively with schurian triangular algebras. For a point x in the quiver Q_A of an algebra A , we denote by e_x the corresponding idempotent.

2.2. Simple connectedness. Let Q be a connected acyclic quiver and I be an ideal of kQ generated by relations. A relation $\rho = \sum_{i=1}^m \lambda_i w_i \in I(x, y)$ is called *minimal* if $m \geq 2$ and, for every non-empty proper subset $J \subset \{1, 2, \dots, m\}$, we have $\sum_{j \in J} \lambda_j w_j \notin I(x, y)$. For an

arrow α , we denote by α^{-1} its formal inverse. A *walk* in Q from x to y is a formal composition $\alpha_1^{\varepsilon_1} \alpha_2^{\varepsilon_2} \cdots \alpha_i^{\varepsilon_i}$ (where $\alpha_i \in Q_1$ and $\varepsilon_i \in \{1, -1\}$ for all i) starting at x and ending at y . The *fundamental group* $\pi_1(Q, I)$ is the quotient of the fundamental group $\pi_1(Q)$ of Q by the normal subgroup generated by all elements of the form $[\gamma^{-1} u^{-1} v \gamma]$, where γ is a walk from the base point to x and u, v are paths from x to y such that there exists a minimal relation $\sum \lambda_i w_i \in I(x, y)$ with $u = w_1$ and $v = w_2$, see [12]. It is well known that $\pi_1(Q)$ is the free group in $\chi(Q)$ generators, where $\chi(Q) = 1 - |Q_0| + |Q_1|$ is the *Euler characteristic* of Q , see, for instance, [13].

A triangular algebra A is called *simply connected* if, for any presentation (Q_A, I) of A , the group $\pi_1(Q_A, I)$ is trivial [5]. It is called *strongly simply connected* if every full convex subcategory of A is simply connected [16].

It is shown in [6] that, if an algebra $A \cong kQ_A/I$ is schurian, then the fundamental group $\pi_1(Q_A, I)$ does not depend on

the presentation (Q_A, I) of A . Since, in this paper, we study only schurian triangular algebras, we therefore use the unambiguous notation $\pi_1(A)$ to stand for $\pi_1(Q_A, I)$.

2.3. Strong simple connectedness. We need the following notions and results from [2]. Let Q be a connected acyclic quiver. A *contour* (p, q) in Q from x to y is a pair of parallel paths of positive length from x to y . A contour (p, q) is called *interlaced* if p and q have a common point besides x and y . A contour (p, q) is called *irreducible* if there exists no sequence of paths $p = p_0, p_1, \dots, p_m = q$ in Q from x to y such that, for each i , the contour (p_i, p_{i+1}) is interlaced. A cycle C in Q is called *irreducible* if, either C is an irreducible contour, or C is not a contour, but satisfies the following condition and its dual: for each source x of C , no proper successor of x in Q is also a source of C , and exactly two proper successors of x in Q are sinks of C . This is easily seen to be equivalent to the definition of irreducibility given in [2](1.5). It is proven in [2](2.1) that, if $A = kQ/I$ is any presentation of a schurian strongly simply connected algebra, and if (p, q) is a contour in Q , then the path p lies in I if and only if the path q lies in I . Moreover, we have the following theorem.

THEOREM. [2](2.4) *An algebra A is schurian and strongly simply connected if and only if*

- (a) *all irreducible cycles are irreducible contours, and*
- (b) *there exists a presentation $A \cong kQ_A/I_A$ such that for each irreducible contour (p, q) , we have $p, q \notin I_A$ but $p - q \in I_A$. □*

Such a presentation (in which every irreducible contour, and hence every contour, is commutative) is called a *normed presentation* of A .

3. Quotients of Incidence Algebras.

3.1. Let (Σ, \leq) be a finite poset with n elements. The incidence algebra $k\Sigma$ of Σ is the subalgebra of the algebra $M_n(k)$ of all $n \times n$ matrices over k consisting of the matrices $[a_{ij}]$ satisfying $a_{ij} = 0$ if $j \not\leq i$. The quiver Q_Σ of $k\Sigma$ is the (oriented) Hasse diagram of Σ , and $k\Sigma \cong kQ_\Sigma/I_\Sigma$, where I_Σ is generated by all differences $p - q$, for (p, q) a contour in Q_Σ .

In this paper a quotient of an incidence algebra means an algebra A which admits a presentation $A \simeq kQ_A/I$ such that there exists a poset Σ with $Q_\Sigma = Q_A$ and, furthermore, $I = I_\Sigma + J$, where J is an ideal of kQ_Σ generated by monomials in the radical square of the algebra. By abuse of language, we identify J with an ideal of $k\Sigma$ and write $A = k\Sigma/J$. Such an algebra $A = k\Sigma/J$ is schurian, triangular and satisfies the following condition: for any contour (p, q) in Q_A , we have $p \notin I_\Sigma + J$ if and only if $q \notin I_\Sigma + J$. In this case, we say that (p, q) is a non-zero contour in A . We also observe that such a presentation of $k\Sigma/J$, generated by monomials and differences of parallel paths, is a normed presentation in the sense of (2.3).

In the next subsection, we consider such an algebra $A = k\Sigma/J$ and construct inductively a set of generators of the ideal $I_\Sigma + J$.

3.2. We start by constructing a set of contours, following an idea inspired from [4](5.2). Let x be a source in Q_A , and x^\rightarrow be the set of all arrows of source x . Let \approx be the least equivalence on x^\rightarrow such that $\alpha \approx \beta$ if there exist paths u, v in Q_Σ such that $(\alpha u, \beta v)$ is a contour [3](2.1). Let t be the cardinality of $[\alpha]$. We construct by induction

on $s \leq t$ a family of $t - 1$ irreducible contours of source x .

If $s = 1$, we let $\alpha_1 \in [\alpha]$ be arbitrary. If $t = 1$, we have finished. Otherwise, we consider the set C_1 of all pairs (β, y) , where $\beta \in x^\rightarrow$ and $y \in (Q_\Sigma)_0$ are such that there exists a non-zero non-interlaced contour $(\alpha_1 p, \beta q)$ from x to y . We let $M_1 = \{y \mid (\beta, y) \in C_1\}$. Since $t > 1$, we have $C_1 \neq \emptyset$. We thus choose a pair $(\alpha_2, y_2) \in C_1$ such that y_2 is maximal in the set M_1 and we also choose a non-zero non-interlaced contour $(\alpha_1 p_{1,2}, \alpha_2 q_{1,2})$ from x to y_2 .

We claim that this contour $(\alpha_1 p_{1,2}, \alpha_2 q_{1,2})$ is in fact irreducible. Indeed, if this is not the case, then there exists a sequence $\alpha_1 p_{1,2} = \beta_1 w_1, \beta_2 w_2, \dots, \beta_r w_r = \alpha_2 q_{1,2}$ with $r \geq 3$ such that each pair $(\beta_i w_i, \beta_{i+1} w_{i+1})$ is interlaced. Since $\alpha_1 \neq \alpha_2$, there exists a least i such that $\beta_i \neq \beta_{i+1}$. Also, there exists a point z on both $\beta_i w_i$ and $\beta_{i+1} w_{i+1}$ and a non-interlaced contour (u, v) from x to z such that $\beta_i w_i = u w'_i$, $\beta_{i+1} w_{i+1} = v w'_{i+1}$ for some paths w'_i, w'_{i+1} . It follows from the maximality of y_2 in M_1 that $A(z, y_2) = 0$ or else $u = 0$ in A (and this is equivalent to saying that $A(x, z) = 0$, because A is a quotient of an incidence algebra). In either case, $A(x, y_2) = 0$, a contradiction. This establishes our claim.

Assume inductively that we have a sequence $\{\alpha_1, \dots, \alpha_s\}$ of arrows of $[\alpha]$ as well as $s - 1$ non-zero non-interlaced contours starting at these arrows. If $s = t$, we have finished. Otherwise, we let C_s be the set of all pairs (β, y) , with $\beta \in x^\rightarrow$ and $y \in (Q_\Sigma)_0$ such that $\beta \notin \{\alpha_1, \dots, \alpha_s\}$ and there exists an $i \in \{1, \dots, s\}$ and a non-zero non-interlaced contour $(\alpha_i p, \beta q)$ from x to y . We let $M_s = \{y \mid (\beta, y) \in C_s\}$. Since $t > s$, we have $C_s \neq \emptyset$. We thus choose $(\alpha_{s+1}, y_{s+1}) \in$

C_s such that y_{s+1} is maximal in M_s , and a non-zero non-interlaced contour $(\alpha_{i_{s+1}}p_{i_{s+1},s+1}, \alpha_{s+1}q_{i_{s+1},s+1})$ from x to y_{s+1} , where $1 \leq i_{s+1} \leq s$.

As before, the maximality of the points y_{s+1} in the sets M_s implies that, in fact, each of the contours chosen is irreducible.

We denote by $I_{[\alpha]}$ the subideal of $I_\Sigma + J$ generated by all elements of the form $\alpha_{i_s}p_{i_s,s} - \alpha_s q_{i_s,s}$, where $(\alpha_{i_s}p_{i_s,s}, \alpha_s q_{i_s,s})$ is a contour of the above set. Thus $I_{[\alpha]}$ is generated by exactly $t - 1$ elements and each of these generators is the commutativity relation of an irreducible contour.


LEMMA. *Let x be a source in A , and $(\alpha p, \beta q)$ be a non-zero irreducible contour of source x . Then there exists a set of generators of $I_{[\alpha]}$, constructed as before, containing the element $\alpha p - \beta q$.*

PROOF. In the above construction, let $\alpha_1 = \alpha$. Then $\beta \approx \alpha$. Since $(\alpha p, \beta q)$ is an irreducible contour from x to y (say), it is not interlaced, so that $(\beta, y) \in C_1$ and y is a maximal element in M_1 . We may thus take $(\alpha_2, y_2) = (\beta, y)$ and $(\alpha_1 p_{1,2}, \alpha_2 q_{1,2}) = (\alpha p, \beta q)$. \square

3.3. Let A be a schurian triangular algebra, the interval $[x, y]$ between x and y is the full subcategory of A generated by all points $z \in A_0$ which lie on a non-zero path from x to y , that is, such that $A(x, z)A(z, y) \neq 0$. Clearly, if all paths from x to y in A are non-zero, then $[x, y]$ coincides with the full subcategory of A generated by the convex hull of x and y . This is the case, for instance, whenever A is an incidence algebra.

Generalizing [4](3.1) (see also [1]), we define a notion of a full weak crown in a quotient of an incidence algebra A .

Let C be a full subcategory of A generated by $2n$ points $\{x_1, \dots, x_n, y_1, \dots, y_n\}$, with $n \geq 2$, and of the form:



We say that C is a *full weak crown* in A if:

- (a) $[x_i, y_j] \cap [x_h, y_l] \neq \emptyset$ if and only if $j = i$ and $(h, l) \in \{(i, i), (i-1, i), (i, i+1)\}$ or $j = i+1$ and $(h, l) \in \{(i, i+1), (i, i), (i+1, i+1)\}$ (we agree to set $y_{n+1} = y_1$ and $x_0 = x_n$), and
- (b) the intersection of three distinct $[x_h, y_l]$ is empty.

A full weak crown C as above is called a *full crown* if, for each i , $[x_i, y_i] \cap [x_i, y_{i+1}] = \{x_i\}$ and $[x_i, y_i] \cap [x_{i-1}, y_i] = \{y_i\}$.

As in [4](3.2) (see also [1]), we prove easily that the convex hull of each full weak crown C contains a full crown as a full subcategory.

A point $x \in A_0$ is said to *top* a full weak crown C if x is a direct predecessor of each of the x_i .


Let now $A = k\Sigma/J$ and x be a source in A . Let $B = A / \langle e_x \rangle$. We observe that, if $A = kQ_A/I_A$ is a normed presentation of A , and $B \cong kQ_B/I_B$ is the induced presentation of B (that is, the one such that $I_B = I_A \cap kQ_B$), then the latter is a normed presentation of B . Finally, we recall that the source x is separating if the number of indecomposable summands of $\text{rad}(e_x A)$ equals the number of connected components of B .

LEMMA. Let $A = kQ_A/I_A$ be a quotient of an incidence algebra, given its normed presentation, x be a separating source in A , topping no full weak crown, and $B = A/ \langle e_x \rangle$. If $B \cong kQ_B/I_B$ is the induced presentation, then

$$I_A = I_B + \sum_{[\alpha]} I_{[\alpha]} + \langle m_1, \dots, m_r \rangle$$

where the m_i are paths starting at x , and the sum is taken over all equivalence classes of arrows in x^\rightarrow .

PROOF. We may clearly assume that x^\rightarrow contains only one equivalence class and that $|A_0| \geq 2$. Let $t = |x^\rightarrow|$. If $t = 1$, $I_{[\alpha]} = 0$ and $I_A = I_B + \langle m_1, \dots, m_r \rangle$ for some paths m_i starting at x . Assume $t > 1$, and let (p, q) be a non-zero irreducible contour starting at x . There exist i, j such that $1 \leq i, j \leq t$ and paths u, v such that $p = \alpha_i u$ and $q = \alpha_j v$. By the construction in 3.2, there exists a unique sequence $\{\alpha_i = \alpha_{i_0}, \alpha_{i_1}, \dots, \alpha_{i_s} = \alpha_j\}$ such that each pair $(\alpha_{i_h}, \alpha_{i_{h+1}})$ corresponds to a non-zero irreducible contour of target $y_{i_{h+1}}$ and this contour is one of the generators of $I_{[\alpha]}$. We have the following situation:




If there is no non-zero path between y and one of the y_{i_h} , then either there exists in A a full weak crown topped by x , or else, one of the y_{i_h} is not maximal, a contradiction in either case. Thus, there is a non-zero path between y

and each of the y_{i_h} . Since the y_{i_h} are maximal points in the respective sets M_{i_h} , we deduce that $A(y_{i_h}, y) \neq 0$ for all h . This implies that the relation $p - q$ is induced from relations in $I_{[\alpha]}$ and relations in I_B .

Finally, if (p, q) is a zero contour, then we assume that p and q are among the m_i . This completes the proof. \square

3.4. Example. The lemma above does not hold true if A is a schurian algebra which is not a quotient of an incidence algebra. Let A be given by the quiver



bound by $\gamma\sigma = 0$, $\delta\sigma = 0$, $\alpha\gamma = \beta\delta$ and $\alpha\lambda\mu = \beta\nu\rho$. Clearly, the relation $\alpha\lambda\mu = \beta\nu\rho$ is not induced from relations in I_B and relations in $I_{[\alpha]}$.


3.5. Corollary. Let Σ be a poset, x be a separating source in $k\Sigma$ and $\Sigma' = \Sigma \setminus \{x\}$. Assume x tops no weak crown. Then $I_\Sigma = I_{\Sigma'} + \sum_{[\alpha]} I_{[\alpha]}$. \square

3.6. Proposition. Let $k\Sigma \cong kQ_\Sigma / I_\Sigma$ be a strongly simply connected incidence algebra, and let (p, q) be a given irreducible contour in Q_Σ . Then there exists a set of generators of I_Σ , of cardinality $\chi(Q_\Sigma)$, consisting of the commutativity relations of irreducible contours and such that $p - q$ is one of these generators.

PROOF. By induction on $|\Sigma|$. If $|\Sigma| = 1$, then there is nothing to prove. Assume that $|\Sigma| > 1$ and that the result holds for all posets Ω such that $|\Omega| < |\Sigma|$. Let x be a maximal element of Σ and $\Sigma' = \Sigma \setminus \{x\}$. Without loss of generality, we may assume that Σ' is connected. Since $k\Sigma$ is strongly simply connected, the source x is separating, by [3](2.6). Therefore all arrows of x^\rightarrow are equivalent, by [3](2.2). Moreover, Σ contains no full crown, by [9](3.3). Hence, by [4](3.2), the point x tops no full weak crown. On the other hand, since $k\Sigma'$ is a full convex subcategory of $k\Sigma$, we have that $k\Sigma'$ is strongly simply connected.

Assume first that x is not the source of (p, q) . Then we may, by the induction hypothesis, assume that $p - q$ belongs to a set of generators of $I_{\Sigma'}$, of cardinality $\chi(Q_{\Sigma'})$, and consisting of commutativity relations of irreducible contours lying entirely in Σ' . On the other hand, if x is a source of (p, q) , we may, by the lemma in 3.2, assume that $p - q$ belongs to a set of generators of $I_{[\alpha]}$, of cardinality $|x^\rightarrow| - 1$ (recall that $x^\rightarrow = [\alpha]$). By 3.5, we have $I_\Sigma = I_{\Sigma'} + I_{[\alpha]}$ where $I_{\Sigma'}$ (or $I_{[\alpha]}$) is generated by the commutativity relations of irreducible contours lying entirely in Σ' (or of irreducible contours starting at x , respectively). Since $\chi(Q_{\Sigma'}) + |x^\rightarrow| - 1 = \chi(Q_\Sigma)$, this completes the proof. \square

3.7. Examples. If $k\Sigma$ is an incidence algebra which is not strongly simply connected, then the conclusion of 3.6 may, or may not, hold true. Let Σ be the poset with quiver



Then $\chi(Q_\Sigma) = 1 - 6 + 8 = 3$, while it easily seen that a set of generators of least cardinality for I_Σ is the set $\{\alpha\gamma - \beta\delta, \alpha\lambda - \beta\mu, \gamma\nu - \lambda\rho, \delta\nu - \mu\rho\}$ which has 4 elements. Note that $k\Sigma$ is simply connected, and even separated and coseparated. Thus, the conclusion of 3.6 does not hold true in this case.

On the other hand, let $\Sigma' = \Sigma \setminus \{x\}$. Then $\chi(Q_{\Sigma'}) = 1 - 5 + 6 = 2$, while $I_{\Sigma'}$ is generated by the set $\{\alpha\gamma - \beta\delta, \alpha\lambda - \beta\mu\}$ which has 2 elements. Note that $k\Sigma'$ is simply connected and separated (but not coseparated). Thus, the conclusion of 3.6 holds true in this case.

4. The Main Results

4.1. Lemma. *Let G be a free group in n generators, and S be a subset of G such that $|S| < n$. Then the normal subgroup H generated by S is a proper subgroup of G .*


PROOF. Let G' be the derived subgroup of G , and assume $G = H$. Then $H/G' = G/G'$ is free abelian of rank n . However, since H/G' is abelian, it is generated by the cosets sG' , with $s \in S$. Hence the rank of H/G' is at most $|S| < n$, and this yields a contradiction. \square

4.2. Before proving the following lemma, we need a simple, but useful observation. Assume $A = kQ_A/I$, where the ideal I is generated by a set of relations of the form $S \cup M$, where M consists of monomials, while S consists of minimal relations. As seen in 2.2, $\pi_1(A) = \pi_1(Q_A)/N$, where N is the normal subgroup of $\pi_1(Q_A)$ generated by all uv^{-1} , with (u, v) a contour in Q_A . It is easily seen that in fact, N is the normal subgroup generated by all uv^{-1} , where (u, v) occurs in a minimal relation belonging to the set S (see [10]).

LEMMA. *Assume that $A = kQ_A/I$ is a schurian simply connected algebra. Let S be a set of minimal relations and M be a set of monomials such that I is generated by $S \cup M$. Then $|S| \geq \chi(Q_A)$.*

PROOF. Since A is schurian, each minimal relation is binomial. Let N be the normal subgroup of $\pi_1(Q_A)$ generated by all uv^{-1} , with u, v are paths occurring in a binomial relation in S . It follows from 2.2 that $\pi_1(A) = \pi_1(Q_A)/N$. Now, the hypothesis implies that $\pi_1(Q_A) = N$. Since $\pi_1(Q_A)$ is free in $\chi(Q_A)$ generators, the result follows from 4.1. \square

4.3. Example. The statement of the lemma is not true if A is not schurian. For instance, the algebra A given by the quiver Q_A




bound by $\alpha_1\beta_1 + \alpha_2\beta_2 + \alpha_3\beta_3 = 0$ is such that $|S| = 1$, but $\chi(Q_A) = 2$.

4.4. Lemma. *Let A be a schurian strongly simply connected algebra. Then:*

- (a) *There exists a unique poset $\Sigma = \Sigma(A)$ such that $Q_\Sigma = Q_A$.*
- (b) *The incidence algebra $k\Sigma$ is strongly simply connected.*

PROOF. (a) It suffices (see, for instance, [11]) to prove that Q_A contains no subquiver of the form



with $\alpha, \beta_1, \dots, \beta_s$ arrows of Q_A . Assume that this is the case, and let $A \cong kQ_A/I$ be an arbitrary presentation of A . Since $\alpha \notin I$, we have that $\beta_1 \cdots \beta_s \notin I$, by 2.3. Since $I \subseteq \text{rad}^2 kQ_A$, there is no relation involving α and $\beta_1 \cdots \beta_s$ and of the form $\lambda\alpha + \mu\beta_1 \cdots \beta_s = 0$ (where $\lambda, \mu \in k$ are non-zero). Consequently, $\dim_k A(x, y) \geq 2$, a contradiction to A being schurian.

(b) Since $Q_\Sigma = Q_A$ and A is schurian strongly simply connected, every irreducible cycle in Q_Σ is an irreducible contour in Q_Σ (this property, indeed, depends on the quiver only). Since any contour in $k\Sigma$ is commutative, we deduce from 2.3 that $k\Sigma$ is strongly simply connected. \square

4.5. One consequence of this lemma is that every schurian strongly simply connected algebra is in fact a quotient of an incidence algebra, a fact already shown in [9](2.7). In fact, we have the following stronger result.


COROLLARY. *An algebra A is schurian and strongly simply connected if and only if there exists a poset Σ such that $k\Sigma$ is strongly simply connected and $A \cong k\Sigma/J$, where J is an ideal of $k\Sigma$ generated by paths which are not entirely contained in irreducible contours.*

PROOF. This follows easily from 4.4 and 2.3. \square

REMARK. If, in particular, $A = k\Sigma/J$ is a quotient of an incidence algebra, with $k\Sigma$ strongly simply connected, then A is strongly simply connected if and only if J is generated

by paths which are not entirely contained in irreducible contours.

4.6. Example. Let A be given by the quiver



bounded by $\alpha\beta = \gamma\delta$, $\delta\nu = \lambda\mu$, $\gamma\lambda = 0$ and $\beta\nu = 0$. Here, $A \cong k\Sigma/J$, where $\Sigma = \Sigma(A)$ and J is generated by the paths $\beta\nu$ and $\gamma\lambda$. These paths are entirely contained in the contour $(\alpha\beta\nu, \gamma\lambda\mu)$. However, this contour is not irreducible. Accordingly, A is schurian strongly simply connected.

4.7. As a first consequence, we have the following characterization of strongly simply connected quotient of an incidence algebras. We recall that the strong simple connectedness of incidence algebras is completely characterized by the absence of full crowns (see [9](3.3)).

THEOREM. *Let $A = k\Sigma/J$ be a quotient of an incidence algebra. Then A is strongly simply connected if and only if A is simply connected and $k\Sigma$ is strongly simply connected.*

PROOF. Since the necessity follows from 4.4, we only need to prove the sufficiency. Assume that A is not strongly simply connected. By 4.5, there exists an irreducible contour (p, q) in Q_A such that $p \in J$ (or, equivalently, $q \in J$). By 3.6, there exists a set $\{\gamma_1, \dots, \gamma_{\chi(Q_A)}\}$ of generators of I_Σ such that $\gamma_1 = p - q$. Let S be the subset

of $\{\gamma_1, \dots, \gamma_{\chi(Q_A)}\}$ corresponding to those contours in Q_A which are non-zero in A . Thus, $|S| < \chi(Q_A)$ since $\gamma_1 \notin S$. Hence, $I_\Sigma + J$ is generated by S and a set $\{m_1, \dots, m_s\}$ of monomials, where $s \geq 2$ and m_1, m_2 are respectively subpaths of p, q . Now, $\pi_1(A) \cong \pi_1(Q_A)/N$, where N is a normal subgroup of $\pi_1(Q_A)$ generated by $|S|$ elements (see the remark in 4.2). Since $|S| < \chi(Q_A)$, the statement of 4.2 yields a contradiction to the simple connectedness of A . \square

4.8. Our second main result characterizes the incidence algebras among the schurian strongly simply connected algebras.

THEOREM. *Let $A = kQ_A/I_A$ be a schurian strongly simply connected algebra, given its normed presentation, and Σ be the associated poset. Then $A \cong k\Sigma$ if and only if I_A has a generating set consisting of exactly $\chi(Q_A)$ elements.*

PROOF. Since the necessity follows at once from 3.6, let us show the sufficiency. Assume that A is schurian and strongly simply connected, and that (Q_A, I_A) is a normed presentation of A . We further assume that I_A has a generating set consisting of exactly $\chi(Q_A)$ relations (such a set is then, by 4.2, of least possible cardinality). Let $\Sigma = \Sigma(A)$ be as in 4.4. It follows from the preceding discussion that $I_A = J + I_\Sigma$, where J is generated by paths in I_A and I_Σ is generated by all possible relations of the form $p - q$, with (p, q) an irreducible contour. Clearly, these two sets of generators are disjoint. Moreover, we get from 3.6 that I_Σ has a generating set consisting of $\chi(Q_\Sigma) = \chi(Q_A)$ elements, all of which are commutativity relations of irreducible contours. This set has as many elements as the given generating set of I_A . Since any generating set of I_A must involve the commutativity of all irreducible contours, we deduce that $J = 0$, that is, $A \cong k\Sigma$. \square

Acknowledgements

The first, second and third authors thank respectively NSERC of Canada, FAPESP and CNPq of Brazil. This work was done during several visits of the authors at the Universidade de São Paulo and the Université de Sherbrooke, they are grateful to each other and to their colleagues in these institutions for the hospitality.

References

- [1] Assem, I.; Castonguay, D.; Marcos, E.N.; Trepode, S. Schurian strongly simply connected algebras. in preparation.
- [2] Assem, I.; Liu, S. Strongly simply connected algebras. *J. Algebra* **1998**, 207, 449-477.
- [3] Assem, I.; de la Peña, J.A. The fundamental groups of a triangular algebra. *Comm. Algebra* **1996** 24, no.1, 187-208.
- [4] Assem, I.; Platzeck, M.I.; Redondo, M.J.; Trepode, S. Simply connected incidence algebras. Preprint (2002), *Discrete Maths*, to appear.
- [5] Assem, I.; Skowroński, A. On some class of simply connected algebras. *Proc. London Math. Soc.* **1988**, (3), 56, 417-450.
- [6] Bardzell, M.J.; Marcos, E.N. $H^1(A)$ and presentations of finite dimensional algebras., in: *Representations of algebras*. Lecture Notes in Pure Applied Math. **2002**, vol. 224, 31-38.
- [7] Bongartz, K.; Gabriel, P. Covering spaces in representation theory. *Invent. Math.* **1981/82**, 65, No. 3, 331-378.

- [8] Bustamante, J.C. On the fundamental group of a schurian algebra. *Comm. Algebra* **2002**, vol. 30, No 11, 5307-5329.
- [9] Dräxler, P. Completely separating algebras. *J. Algebra* **1994**, 165, No.2, 550-565.
- [10] Farkas, D.; Geiss, C.; Green, E.; Marcos, E. Diagonalizable derivations of finite-dimensional algebras I. *Israel Journal of Mathematics* **2000**, 117, 157-183.
- [11] Gatica, M.A.; Redondo, M.J. Hochschild cohomology and fundamental groups of incidence algebras. *Comm. Algebra* **2001**, 29, (5), 2269-2283.
- [12] Martínez-Villa, R.; de la Peña, J.A. The universal cover of a quiver with relations. *J. Pure and Applied Algebra* **1983**, 30, no. 3, 277-292.
- [13] Massey, W.S. *Algebraic topology, an introduction*. Springer-Verlag, Berlin-Heidelberg-New York **1967**.
- [14] Reynaud, E. Algebraic fundamental group and simplicial complexes. *J. Pure Applied Algebra* **2003**, vol.177, no.2, 203-214.
- [15] Rival, I. A fixed point theorem for finite partially ordered sets. *J. Comb. Th. Ser. A* **1976**, 21, no.3, 309-318.
- [16] Skowroński, A. Simply connected algebras and Hochschild cohomologies. in: *Representations of Algebras*. Ottawa, 1992, CMS Conf. Proc. 14, Amer. Math. Soc., Providence, RI, 1993; 431-447,

DÉPARTEMENT DE MATHÉMATIQUES ET D'INFORMATIQUE, FACULTÉ DES SCIENCES, UNIVERSITÉ DE SHERBROOKE, SHERBROOKE, (QUÉBEC) J1K 2R1, CANADA
E-mail address: ibrahim.assem@dm1.usherb.ca

DEPARTAMENTO DE MATEMÁTICA, IME, UNIVERSIDADE DE SÃO PAULO, CAIXA POSTAL 66281, 05315-970, SÃO PAULO, BRAZIL
E-mail address: diane@ime.usp.br

DEPARTAMENTO DE MATEMÁTICA, IME, UNIVERSIDADE DE SÃO PAULO, CAIXA POSTAL 66281, 05315-970, SÃO PAULO, BRAZIL
E-mail address: emmarcos@ime.usp.br

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL DE MAR DEL PLATA, FUNES 3350, 7600, MAR DEL PLATA, ARGENTINA
E-mail address: strepode@mdp.edu.ar

TRABALHOS DO DEPARTAMENTO DE MATEMÁTICA

TRABALHOS DO DEPARTAMENTO DE MATEMÁTICA

TÍTULOS PUBLICADOS

- 2002-01 COELHO, F. U. and LANZILOTA, M. A. On non-semiregular components containing paths from injective to projective modules. 13p.
- 2002-02 COELHO, F. U., LANZILOTTA, M. A. and SAVIOLI, A. M. P. D. On the Hochschild cohomology of algebras with small homological dimensions. 11p.
- 2002-03 COELHO, F. U., HAPPEL, D. and UNGER, L. Tilting up algebras of small homological dimensions. 20p.
- 2002-04 SHESTAKOV, I.P. and UMIRBAEV. U.U. Poisson brackets and two-generated subalgebras of rings of polynomials. 19p.
- 2002-05 SHESTAKOV, I.P. and UMIRBAEV. U.U. The tame and the wild automorphisms of polynomial rings in three variables. 34p.
- 2002-06 ALENCAR, R. and LOURENÇO, M.L. On the Gelbaum-de Lamadrid's result. 16p.
- 2002-07 GRISHKOV, A. Lie algebras with triality. 28p.
- 2002-08 GRISHKOV, A. N. and GUERREIRO, M. Simple classical Lie algebras in characteristic 2 and their gradations, I. 21p.
- 2002-09 MELO, S. T., NEST, R. and SCHROHE, E. K-Theory of Boutet de Monvel's algebra. 8p.
- 2002-10 POJIDAEV, A. P. Enveloping algebras of Filippov algebras. 17p.
- 2002-11 GORODSKI, C. and THORBERGSSON, G. The classification of taut irreducible representations. 47p.
- 2002-12 BORRELLI, V. and GORODSKI, C. Minimal Legendrian submanifolds of S^{2n+1} and absolutely area-minimizing cones. 13p.
- 2002-13 CHALOM, G. and TREPODE, S. Representation type of one point extensions of quasitilted algebras. 16p.
- 2002-14 GORODSKI, C. and THORBERGSSON, G. Variationally complete actions on compact symmetric spaces. 8p.
- 2002-15 GRISHKOV, A. N. and GUERREIRO, M. Simple classical Lie algebras in characteristic 2 and their gradations, II. 15p.
- 2002-16 PEREIRA, Antônio Luiz and PEREIRA, Marcone Corrêa. A Generic Property for the Eigenfunctions of the Laplacian. 28p.

- 2002-17 GALINDO, P., LOURENÇO, M. L. and MORAES, L. A. Polynomials generated by linear operators. 10p.
- 2002-18 GRISHKOV, A. and SIDKI, S. Representing idempotents as a sum of two nilpotents of degree four. 9p.
- 2002-19 ASSEM, I. and COELHO, F. U. Two-sided gluings of tilted algebras. 27p.
- 2002-20 ASSEM, I. and COELHO, F. U. Endomorphism rings of projectives over Laura algebras. 10p.
- 2002-21 CONDORI, L. O. and LOURENÇO, M. L. Continuous homomorphisms between topological algebras of holomorphic germs. 11p.
- 2002-22 MONTES, R. R. and VERDERESI, J. A. A new characterization of the Clifford torus. 5p.
- 2002-23 COELHO, F. U., DE LA PEÑA, J. A. and TREPODE, S. On minimal non-tilted algebras. 27p.
- 2002-24 GRISHKOV, A. N. and ZAVARNITSINE, A. V. Lagrange's theorem for Moufang Loops. 21p
- 2002-25 GORODSKI, C., OLMOS, C. and TOJEIRO, R. Copolarity of isometric actions. 23p.
- 2002-26 MARTIN, Paulo A. The Galois group of $x^n - x^{n-1} - \dots - x - 1$. 18p.
- 2002-27 DOKUCHAEV, M.A. and MILIES, C. P. Isomorphisms of partial group rings. 12p.
- 2002-28 FUTORNY, V. and OVSIENKO, S. An analogue of Kostant theorem for special PBW algebras. 14p.
- 2002-29 CHERNOUSOVA, Zh. T., DOKUCHAEV, M. A., Khibina, M. A., Kirichenko, V. V., Miroshnichenko, S. G. and Zhuravlev, V. N. Tiled orders over discrete valuation rings, finite Markov chains and partially ordered sets. I. 36p.
- 2002-30 FERNÁNDEZ, J. C. G. On commutative power-associative nilalgebras. 10p.
- 2002-31 FERNÁNDEZ, J. C. G. Superalgebras and identities. 11p.
- 2002-32 GRICHKOV, A. N. , GIULIANI, M. L. M. and ZAVARNITSINE, A. V. The maximal subloops of the simple Moufang loop of order 1080. 10p.
- 2002-33 ZAVARNITSINE, A. V. Recognition of the simple groups $L_3(q)$ by element orders. 20p.
- 2002-34 ZUBKOV, A. N. and MARKO, F. When a Schur superalgebra is cellular? 13p.
- 2002-35 COELHO, F. U. and SAVIOLI, A. M. P. D. On shod extensions of algebras. 11p.

- 2003-01 COELHO, F. U. and LANZILOTTA, M. A. Weakly shod algebras. 28p.
- 2003-02 GREEN, E. L., MARCOS, E. and ZHANG, P. Koszul modules and modules with linear presentations. 26p.
- 2003-03 KOSZMIDER, P. Banach spaces of continuous functions with few operators. 31p.
- 2003-04 GORODSKI, C. Polar actions on compact symmetric spaces which admit a totally geodesic principal orbit. 11p.
- 2003-05 PEREIRA, A. L., Generic Hyperbolicity for the equilibria of the one-dimensional parabolic equation $u_t = (a(x)u_x)_x + f(u)$. 19p.
- 2003-06 COELHO, F. U. and PLATZECK, M. I. On the representation dimension of some classes of algebras. 16p.
- 2003-07 CHERNOUSOVA, Zh. T., DOKUCHAEV, M.A., Khibina, M.A., Kirichenko, V.V., MIROSHNICHENKO, S.G., Zhuravlev, V.N. Tiled orders over discrete valuation rings, finite Markov chains and partially ordered sets. II. 43p.
- 2003-08 ARAGONA, J., FERNANDEZ, R. and JURIAANS, S. O. A Discontinuous Colombeau Differential Calculus. 20p.
- 2003-09 OLIVEIRA, L. A. F., PEREIRA, A. L. and PEREIRA, M. C. Continuity of attractors for a reaction-diffusion problem with respect to variation of the domain. 22p.
- 2003-10 CHALOM, G., MARCOS, E., OLIVEIRA, P. Gröbner basis in algebras extended by loops. 10p.
- 2003-11 ASSEM, I., CASTONGUAY, D., MARCOS, E. N. and TREPODE, S. Quotients of incidence algebras and the Euler characteristic. 19p.

Nota: Os títulos publicados nos Relatórios Técnicos dos anos de 1980 a 2001 estão à disposição no Departamento de Matemática do IME-USP.
 Cidade Universitária "Armando de Salles Oliveira"
 Rua do Matão, 1010 - Cidade Universitária
 Caixa Postal 66281 - CEP 05315-970