

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/298196646>

Orogenic structural inheritance and rifted passive margin formation

Conference Paper · April 2016

CITATION

1

READS

73

2 authors:



Claudio Alejandro Salazar Mora
University of São Paulo

21 PUBLICATIONS 97 CITATIONS

[SEE PROFILE](#)



Ritske Huismans
University of Bergen

135 PUBLICATIONS 4,316 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Landscape evolution, sedimentary flux and architecture (Pyrenean system) [View project](#)



Geodynamic numerical modelling of the South Atlantic Rift System: implications for the Neoproterozoic evolution of the Araçuaí Orogen and the São Francisco – Congo cratons [View project](#)

Orogenic structural inheritance and rifted passive margin formation

Claudio A. Salazar Mora (1,2) and Ritske S. Huismans (2)

(1) Geosciences Institute, University of São Paulo, Brazil, (2) Bergen University, Earth Sciences, Bergen, Norway

Structural inheritance is related to mechanical weaknesses in the lithosphere due to previous tectonic events, e.g. rifting, subduction and collision. The North and South Atlantic rifted passive margins that formed during the breakup of Western Gondwana, are parallel to the older Caledonide and the Brasiliano-Pan-African orogenic belts. In the South Atlantic, 'old' mantle lithospheric fabric resulting from crystallographic preferred orientation of olivine is suggested to play a role during rifted margin formation (Tommasi and Vauchez, 2001). Magnetometric and gravimetric mapping of onshore structures in the Camamu and Almada basins suggest that extensional faults are controlled by two different directions of inherited older Brasiliano structures in the upper lithosphere (Ferreira et al., 2009). In the South Atlantic Campos Basin, 3D seismic data indicate that inherited basement structures provide a first order control on basin structure (Fetter, 2009).

Here we investigate the role of structural inheritance on the formation of rifted passive margins with high-resolution 2D thermo-mechanical numerical experiments. The numerical domain is 1200 km long and 600 km deep and represents the lithosphere and the sublithospheric mantle. Model experiments were carried out by creating self-consistent orogenic inheritance where a first phase of orogen formation is followed by extension. We focus in particular on the role of varying amount of orogenic shortening, crustal rheology, contrasting styles of orogen formation on rifted margin style, and the time delay between orogeny and subsequent rifted passive formation. Model results are compared to contrasting structural styles of rifted passive margin formation as observed in the South Atlantic.

Ferreira, T.S., Caixeta, J.M., Lima, F.D., 2009. Basement control in Camamu and Almada rift basins. *Boletim de Geociências da Petrobrás* 17, 69–88.

Fetter, M., 2009. The role of basement tectonic reactivation on the structural evolution of Campos Basin, offshore Brazil: Evidence from 3D seismic analysis and section restoration. *Marine and Petroleum Geology* 26, 873–886.

Tommasi, A., Vauchez, A., 2001. Continental rifting parallel to ancient collisional belts: An effect of the mechanical anisotropy of the lithospheric mantle. *Earth and Planetary Science Letters* 185, 199–210.