

RT-MAT 95-13

On Auslander-Reiten
Components for Quasitilted
Algebras

Flávio U. Coelho
and
Andrzej Skowroński

Abril 1995

ON AUSLANDER-REITEN COMPONENTS FOR QUASITILTED ALGEBRAS

FLÁVIO U. COELHO AND ANDRZEJ SKOWROŃSKI

ABSTRACT. An artin algebra A over a commutative artin ring R is called quasitilted if $\text{gl.dim } A \leq 2$ and for each indecomposable finitely generated A -module M we have $\text{pd } M \leq 1$ or $\text{id } M \leq 1$. In [11] several characterizations of quasitilted algebras were proven. We investigate the structure and homological properties of connected components in the Auslander-Reiten quiver Γ_A of a quasitilted algebra A .

Let A be an artin algebra over a commutative artin ring R , that is, A is an R -algebra which is finitely generated as an R -module. Denote by $\text{ind } A$ the category of indecomposable finitely generated right A -modules, by Γ_A the Auslander-Reiten quiver of A , and by τ_A the Auslander-Reiten translation in Γ_A . Following [10], the algebra A is called tilted if there exists a hereditary artin algebra H and a tilting H -module T such that $A = \text{End}_H(T)$. Recall that a finitely generated H -module T is called tilting if $\text{Ext}_H^1(T, T) = 0$ and there is an exact sequence $0 \rightarrow H_H \rightarrow T_0 \rightarrow T_1 \rightarrow 0$ with T_0 and T_1 in the additive category $\text{add } T$, given by T . The representation theory of tilted algebras is rather well understood. In particular, we know the shape of all connected components of the Auslander-Reiten quivers of tilted algebras (see [8], [12], [13], [17], [18], [19], [20], [27]). It is known that a tilted algebra A is of global dimension at most 2 and no module in $\text{ind } A$ has both the projective and the injective dimension equal to 2. However, these properties do not characterise the tilted algebras. Happel, Reiten and Smalø have shown in [11] that they characterise the class of artin algebras of the form $A = \text{End}(T)$, where T is a tilting object in a hereditary abelian R -category \mathcal{H} , called quasitilted algebras. Besides the tilted algebras, important classes of quasitilted algebras are provided by tubular algebras [19], canonical algebras [14], [19], [21], algebras with separating tubular families of modules [15], [25], and semiregular branch enlargements of such algebras (see [7]). Moreover, it is known that any representation-finite quasitilted algebra is tilted [11]. An important result proven in [11] is the following trisection of the category $\text{ind } A$ of a quasitilted algebra A . Namely, let A be a quasitilted algebra, $\mathcal{C} = \mathcal{C}_A$ be the full subcategory of $\text{ind } A$ formed by the modules whose all successors in $\text{ind } A$ have

1991 *Mathematics Subject Classification.* 16G10, 16G70, 18G20.

Key words and phrases. tilted algebras, complete slices, homological dimensions.

injective dimension at most one, and $\mathcal{D} = \mathcal{D}_A$ be the full subcategory of $\text{ind } A$ formed by the modules whose all predecessors in $\text{ind } A$ have projective dimension at most one. Then we have a trisection

$$\text{ind } A = (\mathcal{D} \setminus \mathcal{C}) \vee (\mathcal{D} \cap \mathcal{C}) \vee (\mathcal{C} \setminus \mathcal{D})$$

such that

$$\text{Hom}_A(\mathcal{D} \cap \mathcal{C}, \mathcal{D} \setminus \mathcal{C}) = 0, \quad \text{Hom}_A(\mathcal{C} \setminus \mathcal{D}, \mathcal{D} \cap \mathcal{C}) = 0, \quad \text{and} \quad \text{Hom}_A(\mathcal{C} \setminus \mathcal{D}, \mathcal{D} \setminus \mathcal{C}) = 0.$$

Moreover, \mathcal{D} (respectively, \mathcal{C}) contains all indecomposable projective (respectively, injective) A -modules.

We investigate here the structure of connected components in the Auslander-Reiten quiver of an arbitrary quasitilted algebra A . Good understanding of the Auslander-Reiten components seems to be the main step in describing the ring structure and representation theory of arbitrary quasitilted algebras (see [15], [24], [25], [26]). We are mainly interested in quasitilted algebras which are not tilted.

In order to state our main results, recall that a (connected) component Γ of Γ_A is called **regular** if Γ contains neither a projective module nor an injective module. Moreover, Γ is called **semiregular** if Γ does not contain both a projective and an injective module. We shall prove the following facts.

Theorem (A). *Let A be a quasitilted artin algebra, and Γ be a component of Γ_A containing an oriented cycle. Then Γ is a semiregular tube.*

We note that a semiregular tube is either regular (that is, of the form $\mathbb{Z}\mathbf{A}_\infty/(\tau^s)$, for some $s \geq 1$) or is obtained from a regular tube by a sequence of ray (or coray) insertions.

Theorem (B). *Let A be a quasitilted algebra, and Γ be a regular component of Γ_A .*

- (i) *If $\Gamma \cap \mathcal{C} \neq \emptyset$, then Γ is contained in \mathcal{C} .*
- (ii) *If $\Gamma \cap \mathcal{D} \neq \emptyset$, then Γ is contained in \mathcal{D} .*

Theorem (C). *Let A be a quasitilted algebra and Γ be a component of Γ_A with infinitely many τ_A -orbits or containing an oriented cycle.*

- (i) *If Γ contains a projective module, then Γ is contained in $\mathcal{D} \setminus \mathcal{C}$.*
- (ii) *If Γ contains an injective module, then Γ is contained in $\mathcal{C} \setminus \mathcal{D}$.*

Theorem (D). *Let A be a quasitilted algebra which is not tilted, and Γ be a component of Γ_A .*

- (i) *If Γ contains a projective module, then Γ is contained in $\mathcal{D} \setminus \mathcal{C}$.*
- (ii) *If Γ contains an injective module, then Γ is contained in $\mathcal{C} \setminus \mathcal{D}$.*

We get also the following immediate consequences of the above theorems.

Corollary (E). *Let A be a quasitilted algebra which is not tilted. Then every component of Γ_A is semiregular.*

Corollary (F). *Let A be a quasitilted algebra which is not tilted. Then every component Γ of Γ_A having a module from $\mathcal{C} \cap \mathcal{D}$ is regular, and hence consists entirely of modules from $\mathcal{C} \cap \mathcal{D}$.*

Further consequences will be discussed in Section 6. This paper is organized as follows. In Section 1 we prove preliminary results on the paths between indecomposable modules over artin algebras, playing a crucial role in our further investigations. In Section 2 we recall some facts on tilted and quasitilted algebras applied in the paper. Sections 3, 4 and 5 are devoted to study the structure of components with oriented cycles, regular components, and nonregular components, respectively, in the Auslander-Reiten quivers of quasitilted algebras. In Section 6 we present some consequences of our main results.

1. PRELIMINARY RESULTS

1.1. Let A be an artin algebra over a commutative artin ring R , that is, A is an R -algebra which is finitely generated as an R -module. Unless otherwise stated all algebras are assumed to be basic and connected. By an A -module it is meant a finitely generated right A -module. We shall denote by $\text{mod } A$ the category of all (finitely generated) A -modules, and by $\text{ind } A$ the full subcategory of $\text{mod } A$ with one representative of each isomorphism class of indecomposable A -modules. Then $\text{rad}(\text{mod } A)$ denotes the Jacobson radical of $\text{mod } A$, that is, the ideal in $\text{mod } A$ generated by all noninvertible morphisms between indecomposable modules in $\text{mod } A$. The infinite radical $\text{rad}^\infty(\text{mod } A)$ of $\text{mod } A$ is the intersection of all powers $\text{rad}^i(\text{mod } A)$, $i \geq 1$, of $\text{rad}(\text{mod } A)$.

1.2. We shall denote by Γ_A the Auslander-Reiten quiver of A , and by $\tau = D\text{Tr}$ and $\tau^- = \text{Tr } D$ the Auslander-Reiten translations in Γ_A . We shall now agree to identify the vertices of Γ_A with the corresponding A -modules in $\text{ind } A$. By a component of Γ_A we mean a connected component of Γ_A . We observe that a morphism between indecomposable modules lying in different components of Γ_A belongs to $\text{rad}^\infty(\text{mod } A)$.

We shall use frequently the fact that, for an A -module X , $\text{pd } X \leq 1$ if and only if $\text{Hom}_A(D(A), \tau X) = 0$ (respectively, $\text{id } X \leq 1$ if and only if $\text{Hom}_A(\tau^- X, A) = 0$) (see [19](p. 74)).

Let Γ be a component of Γ_A . Then, Γ is said to be **regular** if Γ contains neither a projective module nor an injective module, and **semiregular** if Γ does not contain both a projective and an injective module. Also, Γ is said to be **postprojective** (respectively, **preinjective**) if Γ contains no oriented cycles and each module in Γ

belongs to the τ -orbit of a projective (respectively, an injective) module. We shall denote by ${}_{\text{L}}\Gamma$ the left stable part of Γ obtained from Γ by deleting the τ -orbits of projective modules, by ${}_{\text{R}}\Gamma$ the right stable part of Γ obtained from Γ by deleting the τ -orbits of injective modules, and by ${}_{\text{S}}\Gamma$ the stable part of Γ obtained from Γ by deleting the τ -orbits of both the projective and the injective modules.

A module $M \in \text{ind } A$ is called τ -periodic if there exists an $m \geq 1$ such that $\tau^m M \simeq M$. Given $M \in \text{ind } A$, we denote by $\mathcal{O}(M)$ the τ -orbit of M , that is,

$$\mathcal{O}(M) = \{\tau^m M : m \in \mathbb{Z}\}.$$

1.3. Let $M, N \in \text{ind } A$. A path from M to N is given by a sequence of nonzero morphisms

$$M = X_0 \xrightarrow{f_1} X_1 \longrightarrow \cdots \xrightarrow{f_t} X_t = N$$

where, for each i , X_i is an indecomposable module and f_i is in $\text{rad}(\text{mod } A)$. We denote a path from M to N by $M \rightsquigarrow N$. If we have in addition that the morphisms f_i are irreducible, then we shall say that it is a path of irreducible maps. Here, an oriented cycle is a path of irreducible maps from a module to itself. A path of irreducible maps $X_0 \rightarrow X_1 \rightarrow \cdots \rightarrow X_t$ is called sectional if $X_i \not\simeq \tau X_{i+2}$ for each $i = 0, \dots, t-2$.

Given a path $M \rightsquigarrow N$, M is said to be a predecessor of N and N a successor of M . It should be clear the terms predecessor and successor by irreducible maps. Finally, given $M, N \in \text{ind } A$, we denote by $M \text{---} N$ when there is either an irreducible map $M \rightarrow N$ or an irreducible map $N \rightarrow M$. For more details on the Auslander-Reiten theory we refer the reader to [3] and [19].

1.4. We will now prove two lemmas needed later on.

Lemma. *Let A be an artin algebra,*

$$(*) \quad X = X_0 \longrightarrow X_1 \longrightarrow \cdots \longrightarrow X_t = X$$

be an oriented cycle through indecomposable modules, and $r \geq 1$. If $\tau^i X_j \neq 0$ for each $1 \leq i \leq r$ and each $j = 0, \dots, t$, then there exists a path of irreducible maps from X to $\tau^r X$.

Proof. We know, by [4], that the oriented cycle $(*)$ is not sectional. Therefore, there exists an l , $2 \leq l \leq t$, such that $\tau X_l \simeq X_{l-2}$. By hypothesis, one can apply τ to $(*)$ to get

$$(**) \quad \tau X = \tau X_0 \longrightarrow \tau X_1 \longrightarrow \cdots \longrightarrow \tau X_t = \tau X.$$

Observe that the module $\tau X_l \simeq X_{l-2}$ appears in both $(*)$ and $(**)$, and hence there exists a path from X to τX , namely

$$X = X_0 \longrightarrow X_1 \longrightarrow \cdots \longrightarrow X_{l-2} \simeq \tau X_l \longrightarrow \tau X_{l+1} \longrightarrow \cdots \longrightarrow \tau X_t = \tau X.$$

By applying τ and composing the paths, we get the desired result. \square

1.5. The next result extends [22](Lemma 4).

Lemma. *Let A be an artin algebra and denote by n the rank of the Grothendieck group $K_0(A)$ of A . Let Γ be a connected component of Γ_A and Γ' be a connected component of Γ . Assume that Γ' has infinitely many τ -orbits and no oriented cycles. Let M be a module in Γ' such that the length of any walk in Γ from a nonstable module to M is at least $2n$. Then, for each $r \geq 1$, there exists a path*

$$M = X_0 \longrightarrow X_1 \longrightarrow \cdots \longrightarrow X_t = \tau^r M$$

in $\text{mod } A$, with all X_i in Γ .

Proof. It is enough to show that there exists a path in $\text{mod } A$ from M to τM , through modules in Γ and then proceed inductively. By [22](Lemma 4), there is a path

$$M = X'_0 \longrightarrow X'_1 \longrightarrow \cdots \longrightarrow X'_t = M$$

in $\text{mod } A$, with X'_1, \dots, X'_t belonging to Γ . Since Γ has no oriented cycles, then one of the maps in the above path should be in $\text{rad}^\infty(\text{mod } A)$. We can then infer that there exists a path

$$(*) \quad M = Y_0 \xrightarrow{f_0} Y_1 \xrightarrow{f_1} \cdots \longrightarrow Y_r \xrightarrow{f_r} \cdots \longrightarrow Y_{r+n} \xrightarrow{f_{r+n}} Y_{r+n+1} \xrightarrow{f_{r+n+1}} M$$

where the morphisms f_0, \dots, f_{r+n+1} are irreducible maps (this is done by using the lifting properties of almost split sequences). Consider now the path of irreducible maps

$$(**) \quad Y_r \xrightarrow{f_r} Y_{r+1} \longrightarrow \cdots \longrightarrow Y_{r+n+1} \xrightarrow{f_{r+n+1}} M.$$

If $(**)$ is nonsectional, then there exists an $i, r \leq i \leq r+n-1$, such that $Y_i \simeq \tau Y_{i+2}$, and then there exists a path of irreducible maps

$$Y_r \longrightarrow \cdots \longrightarrow Y_i \longrightarrow \tau Y_{i+3} \longrightarrow \cdots \longrightarrow \tau M$$

(observe that the modules in $(**)$ are left stable modules, and hence one can apply τ to them). Therefore, there exists a path from M to τM , passing through modules in Γ , namely

$$M = Y_0 \longrightarrow Y_1 \longrightarrow \cdots \longrightarrow Y_r \longrightarrow \cdots \longrightarrow Y_i \longrightarrow \tau Y_{i+3} \longrightarrow \cdots \longrightarrow \tau M.$$

Suppose now $(**)$ is sectional. Then, by [22](Lemma 2), there exist j and l , $r \leq j, l \leq r+n+1$, and a nonzero map $g \in \text{Hom}_A(Y_j, \tau Y_l) = \text{rad}^\infty(Y_j, \tau Y_l)$. Hence, the following path

$$M = Y_0 \longrightarrow Y_1 \longrightarrow \cdots \longrightarrow Y_j \xrightarrow{g} \tau Y_l \longrightarrow \tau Y_{l+1} \longrightarrow \cdots \longrightarrow \tau M$$

gives the required path from M to τM passing through modules in Γ . Now, since for each $i \geq 1$, $\tau^i M$ is in the conditions of the Lemma, we can iterate the above procedure to get a path from M to each $\tau^r M$, $r \geq 1$, as required. \square

1.6. Corollary. *Let A be an artin algebra and Γ be a regular component of Γ_A with infinitely many τ -orbits. Then, for each $M \in \Gamma$, and each $r \geq 1$, there exists a path in $\text{mod } A$ from M to $\tau^r M$.*

2. QUASITILTED ALGEBRAS

2.1. We shall collect in this section the results on quasitilted algebras needed along the paper. We start recalling some facts on tilted algebras. For details on tilting theory we refer the reader to [10] and [19]. Let H be a hereditary algebra and let T be a tilting H -module, that is, a module such that $\text{Ext}_H^1(T, T) = 0$ and there exists a short exact sequence $0 \rightarrow H \rightarrow T_0 \rightarrow T_1 \rightarrow 0$, where T_0 and T_1 are in $\text{add } T$. The algebra $B = \text{End}_H(T)$ is called a tilted algebra. An important fact on a tilted algebra B is that Γ_B contains a component, called **connecting**, which contains a so-called complete slice Σ which reproduces somehow the structure of the hereditary algebra H . It is well-known that all successors of such Σ have injective dimension at most one, and all predecessors of Σ have projective dimension at most one. Recall that a subquiver Σ in a component Γ of Γ_A is called a **complete slice** if: (a) Σ is sincere; (b) Σ is path closed in $\text{mod } A$, and (c) Σ meets each τ -orbit of Γ exactly once.

Let $B = \text{End}_H(T)$, where T is a tilting module over a representation-infinite hereditary algebra H . If T is a postprojective H -module (equivalently, $\text{rad}^\infty(-, T) = 0$), then the algebra B is called **concealed**. It is a well-established fact that the Auslander-Reiten quiver of a tilted algebra B contains at most two connecting components, and it has exactly two if and only if B is concealed. Also, T is a regular H -module if and only if the connecting component of Γ_B is regular.

2.2. We shall now recall the definition of quasitilted algebras and some results on them. We refer the reader to [11] for the proof of these results.

Definition. An algebra A is said to be **quasitilted** if $\text{gl.dim } A \leq 2$ and for each $X \in \text{ind } A$, either $\text{pd } X \leq 1$ or $\text{id } X \leq 1$.

Tilted algebras are clearly examples of quasitilted algebras. However, as mentioned in the introduction the class of quasitilted algebras is much larger. It has been proven in [11](II.3.6) that representation-finite quasitilted algebras are tilted.

2.3. We mention the next result for later reference.

Theorem. ([11](II.1.14)) *Let A be a quasitilted algebra. Then any path in $\text{ind } A$ starting in an injective module and ending in a projective module has a refinement of irreducible maps and any such a path is sectional.*

2.4. Let A be a quasitilted algebra. An important result is the existence of the following trisection of the category $\text{ind } A$. Let

$$\mathcal{C} = \mathcal{C}_A = \{X \in \text{ind } A : \text{for each } Y \text{ with } X \rightsquigarrow Y, \text{id}Y \leq 1\}$$

$$\mathcal{D} = \mathcal{D}_A = \{X \in \text{ind } A : \text{for each } Y \text{ with } Y \rightsquigarrow X, \text{pd } Y \leq 1\}.$$

This induces a trisection

$$\text{ind } A = (\mathcal{D} \setminus \mathcal{C}) \vee (\mathcal{D} \cap \mathcal{C}) \vee (\mathcal{C} \setminus \mathcal{D})$$

such that

$$\text{Hom}_A(\mathcal{D} \cap \mathcal{C}, \mathcal{D} \setminus \mathcal{C}) = 0, \quad \text{Hom}_A(\mathcal{C} \setminus \mathcal{D}, \mathcal{D} \cap \mathcal{C}) = 0 \quad \text{and} \quad \text{Hom}_A(\mathcal{C} \setminus \mathcal{D}, \mathcal{D} \setminus \mathcal{C}) = 0.$$

Moreover, \mathcal{D} contains all the indecomposable projective modules and it is closed under predecessors, while \mathcal{C} contains all the indecomposable injective modules and it is closed under successors.

2.5. The next result gives a criterion for a quasitilted algebra to be tilted.

Theorem. ([11](II.3.4)) *Let A be a quasitilted algebra. If \mathcal{C} contains a projective module, then A is tilted.*

2.6. Let A be a tilted algebra. Then, clearly, any complete slice in $\text{mod } A$ is contained in $\mathcal{C} \cap \mathcal{D}$. In particular, for tilted algebras $\mathcal{C} \cap \mathcal{D}$ is nonempty. For quasitilted algebras which are not tilted, it is still an open question whether $\mathcal{C} \cap \mathcal{D}$ is nonempty. We shall show (Corollary (F)) that, if $\mathcal{C} \cap \mathcal{D}$ is nonempty for a quasitilted algebra which is not tilted, then $\mathcal{C} \cap \mathcal{D}$ is formed by modules lying in regular components.

3. COMPONENTS WITH ORIENTED CYCLES

3.1. Let A be a quasitilted algebra and Γ be a component of Γ_A containing oriented cycles. We shall show that Γ is in fact a semiregular tube, generalising a result known for tilted algebras (see [12], [13], [17]). The main point in the proof is to show that such a Γ is semiregular. Semiregular components with oriented cycles have been described in [9], [16] and [28], and they are either of the form $\mathbb{Z}\mathbb{A}_\infty/(\tau^m)$, for some $m \geq 1$ (if regular) or obtained from it by a sequence of ray (or coray) insertions.

Theorem (A). *Let A be a quasitilted algebra, and Γ be a component of Γ_A containing an oriented cycle. Then Γ is a semiregular tube.*

Proof. Let Γ be a component of Γ_A containing an oriented cycle. We first observe that Γ is infinite. Indeed, if Γ is finite, then A is representation-finite, and hence by (2.2) it is tilted and $\Gamma (= \Gamma_A)$ is a connecting component. It is well-known that, in this case, Γ has no oriented cycles, a contradiction, and hence Γ is infinite.

If Γ is regular, then by [9] and [28], Γ is a stable tube. If Γ is semiregular but not regular, it follows from [16] that Γ is a semiregular tube.

Suppose then that Γ is nonsemiregular. We first claim that Γ has no τ -periodic modules. Suppose Γ has a τ -periodic module. Since Γ is not regular, we infer, using [2](6.2), that there exists an irreducible map $X \rightarrow Y$, where X is a τ -periodic module, and Y is neither left nor right stable, that is, there are m and m' such that $\tau^m Y$ is a projective module P and $\tau^{m'} Y$ is an injective module I . Therefore, there exists a nonsectional path from I to P , which contradicts our hypothesis that A is quasitilted (2.3). This proves the claim.

Consider now the left and the right stable parts of Γ , respectively, ${}_{\text{L}}\Gamma$ and ${}_{\text{R}}\Gamma$. Since Γ is infinite, either ${}_{\text{L}}\Gamma$ or ${}_{\text{R}}\Gamma$ is nonempty. Suppose ${}_{\text{L}}\Gamma \neq \emptyset$ and let Γ' be a connected component of ${}_{\text{L}}\Gamma$. Clearly, Γ' is infinite because otherwise it would contain a τ -periodic module, contradicting the above claim.

We shall show now that Γ' contains no oriented cycles. Suppose it contains oriented cycles. Observe that then, Γ' contains injective modules, because otherwise it would be a stable tube by [9] and [28], in particular, with τ -periodic modules, a contradiction to the claim. In resume, Γ' is an infinite connected component of ${}_{\text{L}}\Gamma$ with oriented cycles and containing injective modules. Then, by [16](2.3), there exists an infinite sectional path

$$\dots \rightarrow \tau^{2t} X_1 \rightarrow \tau^t X_s \rightarrow \dots \rightarrow \tau^t X_2 \rightarrow \tau^t X_1 \rightarrow X_s \rightarrow \dots \rightarrow X_1$$

with $t > s$ such that $\{X_1, \dots, X_s\}$ is a complete set of representatives of τ -orbits in Γ' . Since Γ' is a component of ${}_{\text{L}}\Gamma$ and Γ is not left stable, there exists an irreducible map $X' \rightarrow X''$ with X'' in the τ -orbit of a projective module and $X' \in \Gamma'$. By applying τ as much as necessary, we have that there exists an irreducible map $X \rightarrow P$ with $X \in \Gamma'$ and P an indecomposable projective module. Since $X \in \Gamma'$, we infer that $\tau^{m'} X \simeq \tau^{m+t} X_j$ for some $1 \leq j \leq s$, and some $m, m' \geq 0$.

It follows from the hypothesis on Γ' that there exists an oriented cycle $(*)$ in Γ' containing an injective module I . By (1.4), there are paths from I to each $\tau^r I$, $r \geq 1$. Observe that I is in the τ -orbit of one of X_1, \dots, X_s , say X_i . Then, we get a path from I to $\tau^{(m+1)t} X_i$. Therefore, there exists a path

$$I \xrightarrow{(*)} \tau^{(m+1)t} X_i \rightarrow \tau^{mt} X_j \rightarrow X \rightarrow P$$

from an injective module to a projective module which is not sectional because $(*)$ is not sectional by [4]. A contradiction to the fact that A is quasitilted. Therefore, none of the components of ${}_{\text{L}}\Gamma$ contains an oriented cycle. Similarly, one can also show that ${}_{\text{R}}\Gamma$ contains no oriented cycles.

However, by hypothesis, Γ contains an oriented cycle

$$Y_0 \rightarrow Y_1 \rightarrow \dots \rightarrow Y_u = Y_0.$$

By above remarks, such an oriented cycle should contain a module which is not left stable and a module which is not right stable. By applying τ^{-1} we can assume that Y_0 is an injective module. Let Y_v and $l \geq 0$ be such that $\tau^l Y_v$ is a projective module,

and assume that $\tau^i Y_j \neq 0$ for each $j \neq v$ and $i \leq l$. By (1.4), there exists a path from Y_0 to $\tau^i Y_0$ and then to $\tau^i Y_v$, which is projective, and this path can be chosen to be nonsectional, which is a contradiction to the fact that A is quasitilted, and the result is proven. \square

4. REGULAR COMPONENTS

4.1. The main aim in this section is to prove Theorem (B) stated in the introduction, which concerns regular components of the Auslander-Reiten quiver of quasitilted algebras. We shall need the following lemma.

Lemma. *Let A be a quasitilted algebra and Γ be a component of Γ_A .*

- (a) *If $\Gamma \cap \mathcal{C} \neq \emptyset$, then each τ -orbit of Γ contains a module from \mathcal{C} .*
- (b) *If $\Gamma \cap \mathcal{D} \neq \emptyset$, then each τ -orbit of Γ contains a module from \mathcal{D} .*

Proof. We shall prove only (a) because the proof of (b) is similar.

(a) Let Γ be a component of Γ_A containing a module from \mathcal{C} . Consider the right stable part $,\Gamma$ of Γ . If $,\Gamma = \emptyset$, or equivalently, if each τ -orbit of Γ contains an injective module, then there is nothing to prove because \mathcal{C} contains all the injective modules. Suppose $,\Gamma \neq \emptyset$ and let Γ' be a connected component of $,\Gamma$. We first claim that Γ' has a module from \mathcal{C} . If Γ has no injective modules then $\Gamma' = ,\Gamma = \Gamma$, and the claim is clear. Suppose that Γ contains an injective module. Then there exists an irreducible map $I \rightarrow X'$, where I is an injective module and $X' \in \Gamma'$. The claim now follows from the fact that $X' \in \mathcal{C}$, because $I \in \mathcal{C}$, and \mathcal{C} is closed under successors.

Let $X \in \Gamma' \cap \mathcal{C}$ and let $Y \in \Gamma'$. We shall show that $\mathcal{O}(Y) \cap \mathcal{C} \neq \emptyset$. Since Γ' is connected, there exists a walk in Γ'

$$X = X_0 - X_1 - \cdots - X_s = Y.$$

Observe that the modules X_i , $i = 1, \dots, s$, are right stable and so one can apply τ^- as much as necessary to them to get a path from X to some $\tau^{-m} Y$, $m \geq 0$. Since \mathcal{C} is closed under successors we get that $\tau^{-m} Y \in \mathcal{C}$, and hence each τ -orbit of $,\Gamma$ has a module from \mathcal{C} . The result now follows from the fact that the τ -orbits which are not in $,\Gamma$ contain an injective module, and hence a module from \mathcal{C} . \square

4.2. We can now prove Theorem (B) from the introduction.

Theorem (B). *Let A be a quasitilted algebra and Γ be a regular component of Γ_A .*

- (a) *If $\Gamma \cap \mathcal{C} \neq \emptyset$, then Γ is contained in \mathcal{C} .*
- (b) *If $\Gamma \cap \mathcal{D} \neq \emptyset$, then Γ is contained in \mathcal{D} .*

Proof. We shall only prove (a) because the proof of (b) is similar.

(a) Let Γ be a regular component containing a module M from \mathcal{C} . If Γ has oriented cycles, then by [9], it is a stable tube and then clearly every module in Γ is a successor of M , therefore belonging to \mathcal{C} (2.4).

Suppose from now on that Γ has no oriented cycles and let $N \in \Gamma$. We shall show that $N \in \mathcal{C}$. Suppose Γ has infinitely many τ -orbits. By (4.1) there exists an $m \in \mathbb{Z}$ such that $\tau^m N \in \mathcal{C}$. By (1.6), there exists a path from $\tau^m N$ to N , and hence N is also in \mathcal{C} . It remains the case when Γ has only finitely many τ -orbits. If $N \notin \mathcal{C}$, then there exists a path

$$N = X_0 \xrightarrow{f_0} X_1 \longrightarrow \dots \xrightarrow{f_{t-1}} X_t = X$$

where $\text{id}X > 1$. It is well-known that then $\text{Hom}_A(\tau X, A) \neq 0$ (see (1.2)). Therefore there exists a path

$$N = X_0 \xrightarrow{f_0} X_1 \longrightarrow \dots \xrightarrow{f_{t-1}} X_t = X \xrightarrow{f_t} X_{t+1} \xrightarrow{f_{t+1}} \tau - X \xrightarrow{f_{t+2}} P$$

where P is an indecomposable projective module, and the morphisms f_t and f_{t+1} are irreducible.

Since Γ is regular, we have that $P \notin \Gamma$ and then at least one of the maps $f_0, f_1, \dots, f_{t-1}, f_{t+2}$ is in $\text{rad}^\infty(\text{mod } A)$. Observe now that if $g : Y \rightarrow Y'$ is a map in $\text{rad}^\infty(\text{mod } A)$, then for each $r \geq 1$, there exist a chain of irreducible maps

$$Y = Y_0 \xrightarrow{g_1} Y_1 \longrightarrow \dots \xrightarrow{g_r} Y_r$$

and a morphism $h_r : Y_r \rightarrow Y'$ such that the composition $h_r g_r \cdots g_1$ is nonzero.

Suppose now that one of f_0, \dots, f_{t-1} is in $\text{rad}^\infty(\text{mod } A)$. By the above and (4.1), we infer that there exists a path from some module in Γ which belongs to \mathcal{C} to X , and so $\text{id}X = 1$, a contradiction. If none of f_0, \dots, f_{t-1} belong to $\text{rad}^\infty(\text{mod } A)$, then $f_{t+2} \in \text{rad}^\infty(\text{mod } A)$. By similar arguments there exists a module $Z \in \Gamma$ such that τZ is in \mathcal{C} and $\text{Hom}_A(Z, P) \neq 0$, or equivalently, $\text{id}\tau Z > 1$, a contradiction. \square

5. NONREGULAR COMPONENTS

5.1. Let A be a quasitilted algebra. We shall now concentrate in the study of nonregular components of Γ_A . In this section we shall prove Theorems (C) and (D) and establish some immediate consequences of them.

Theorem (C). *Let A be a quasitilted algebra and Γ be a component of Γ_A with infinitely many τ -orbits or containing an oriented cycle.*

- (a) *If Γ contains a projective module, then Γ is contained in $\mathcal{D} \setminus \mathcal{C}$.*
- (b) *If Γ contains an injective module, then Γ is contained in $\mathcal{C} \setminus \mathcal{D}$.*

Proof. We shall only prove (a) because the proof of (b) is similar.

(a) Let Γ be a component of Γ_A containing a projective module. Suppose first that Γ has oriented cycles. Then, by (3.1), Γ is a ray tube. Suppose furthermore that there exists a module $M \in \Gamma \cap \mathcal{C}$. Since Γ is a ray tube, any module in Γ which belongs to a cycle is a successor of M , and hence belongs to \mathcal{C} by (2.4). On the other hand, there exists a module X which is a nonprojective summand of the radical of some projective module which belongs to an oriented cycle. By (1.2), we infer that

$\text{id}\tau X > 1$, a contradiction to the fact that τX should be in \mathcal{C} because it is a successor of M . Therefore, $\Gamma \cap \mathcal{C} = \emptyset$, and since $\text{id}A = \mathcal{C} \cup \mathcal{D}$, we get that $\Gamma \subset \mathcal{D} \setminus \mathcal{C}$.

Suppose now Γ has infinitely many τ -orbits but no oriented cycles, and that $\Gamma \cap \mathcal{C} \neq \emptyset$. Therefore, there exists a connected component Γ' of Γ with infinitely many τ -orbits. It follows now from (4.1) that there exists a module $M \in \Gamma' \cap \mathcal{C}$ such that the length of any walk from a nonstable module to M is at least $2n$, where n is the rank of $K_0(A)$. Let

$$M' = X_0 \rightarrow X_1 \rightarrow \cdots \rightarrow X_t = P$$

be a walk in Γ' of minimal length from a module M' in the τ -orbit $\mathcal{O}(M)$ of M to a projective module P . Because of the minimality, all the modules X_0, \dots, X_{t-1} are left stable and then, by applying τ conveniently, there exist an $m \geq 0$ and a path of irreducible maps

$$\tau^m M = Y_0 \rightarrow Y_1 \rightarrow \cdots \rightarrow Y_t = P.$$

Since the modules Y_0, \dots, Y_{t-1} are left stable, we get a path of irreducible maps from $\tau^{m+1}M$ to τY_{t-1} . On the other hand, by (1.5), there exists a path from M to $\tau^{m+1}M$, and then τY_{t-1} is a successor of M , which implies that $\tau Y_{t-1} \in \mathcal{C}$ (2.4). This, however, is a contradiction to the fact that $\text{id}\tau Y_{t-1} > 1$, because $\text{Hom}_A(Y_{t-1}, A) \neq 0$. Therefore $\Gamma \cap \mathcal{C} = \emptyset$ and because $\text{id}A = \mathcal{C} \cup \mathcal{D}$, we have that $\Gamma \subset \mathcal{D} \setminus \mathcal{C}$ as required. \square

5.2. For quasitilted algebras which are not tilted, the above result can be deepened as follows.

Theorem (D). *Let A be a quasitilted algebra which is not tilted, and Γ be a component of Γ_A .*

- (a) *If Γ contains a projective module, then Γ is contained in $\mathcal{D} \setminus \mathcal{C}$.*
- (b) *If Γ contains an injective module, then Γ is contained in $\mathcal{C} \setminus \mathcal{D}$.*

Proof. We shall prove only (a) because the proof of (b) is similar.

(a) Let Γ be a component containing a projective module. If Γ has oriented cycles or infinitely many τ -orbits, then the result follows from (5.1). Suppose then that Γ has no oriented cycles and only finitely many τ -orbits. Therefore, there exists an indecomposable projective module P in Γ such that P has no proper successors in Γ which are also projective modules. Since A is not tilted, $P \notin \mathcal{C}$ (2.5). Therefore, there exists a path

$$P = X_0 \xrightarrow{f_0} X_1 \rightarrow \cdots \xrightarrow{f_{t-1}} X_t = X$$

where $\text{id}X > 1$, or equivalently, $\text{Hom}_A(\tau^{-1}X, A) \neq 0$ (1.2). Hence, there exists a path

$$P = X_0 \xrightarrow{f_0} X_1 \rightarrow \cdots \xrightarrow{f_{t-1}} X_t = X \xrightarrow{f_t} X_{t+1} \xrightarrow{f_{t+1}} \tau^{-1}X \xrightarrow{f_{t+2}} P'$$

where P' is an indecomposable projective module, and f_t and f_{t+1} are irreducible maps. By our hypothesis on P , at least one of the maps $f_0, \dots, f_{t-1}, f_{t+2}$ is in

$\text{rad}^\infty(\text{mod } A)$. Suppose one of f_0, \dots, f_{t-1} is in $\text{rad}^\infty(\text{mod } A)$. Following the considerations made in the proof of Theorem (B) we infer that there exists a path from some module in $\Gamma \cap \mathcal{C}$ to X , a contradiction because $\text{id}X > 1$.

If now $f_0, \dots, f_{t-1} \notin \text{rad}^\infty(\text{mod } A)$, then $f_{t+2} \in \text{rad}^\infty(\text{mod } A)$. Also, by similar considerations of those made in the proof of Theorem (B), we get that there exists a module $Z \in \Gamma$ such that $\tau Z \in \mathcal{C}$ and $\text{Hom}_A(Z, P') \neq 0$, or equivalently, $\text{id}\tau Z > 1$, a contradiction, and this finishes the proof. \square

5.3. We have the following direct consequences of (5.2).

Corollary. *Let A be a quasitilted algebra, and Γ be a component of Γ_A . If Γ is nonsemiregular, then A is tilted and Γ is the (unique) connecting component of Γ_A .*

Note that the above corollary generalises [11](II.3.6) which says that any representation-finite quasitilted algebra is tilted, because clearly the Auslander-Reiten quiver of any representation-finite algebra is nonsemiregular.

5.4 Corollary (E). *Let A be a quasitilted algebra which is not tilted. Then any component of Γ_A is semiregular.*

5.5 Corollary (F). *Let A be a quasitilted algebra which is not tilted. Then, every component Γ of Γ_A having a module from $\mathcal{C} \cap \mathcal{D}$ is regular, and hence consists of modules from $\mathcal{C} \cap \mathcal{D}$.*

6. SOME CONSEQUENCES

6.1. We say that a property holds for almost all modules if it holds for all but finitely many of them. In [1], I. Assem and the first named author have characterized the finite dimensional algebras over algebraically closed fields which satisfy the property that almost all of their indecomposable modules have injective (or projective) dimension at most one. These algebras are called left (respectively, right) glueings of tilted algebras (see [1] for details).

For an artin algebra A such that $\text{id}X \leq 1$ for almost all $X \in \text{ind } A$, it follows from [5], [6] and [23], that Γ_A contains a component Γ containing all the projective modules and such that: (i) almost all of its modules lie in the τ -orbits of projective modules; and (ii) there are at most finitely many modules in Γ belonging to oriented cycles.

6.2. We shall use the above fact to show the following result.

Proposition *Let A be a quasitilted algebra.*

(a) *The following are equivalent:*

- (i) $\text{id}X \leq 1$ for almost all $X \in \text{ind } A$.
- (ii) A is tilted and Γ_A has a postprojective component with a complete slice.
- (iii) \mathcal{C} is cofinite in $\text{ind } A$.

(b) *The following are equivalent:*

- (i) $pdX \leq 1$ for almost all $X \in \text{ind } A$.
- (ii) A is tilted and Γ_A has a preinjective component with a complete slice.
- (iii) \mathcal{D} is cofinite in $\text{ind } A$.

Proof. We shall prove only (a) because the proof of (b) is similar.

(a) (i) \Rightarrow (ii) Suppose $\text{id}X \leq 1$ for almost all $X \in \text{ind } A$. By the above remarks, Γ_A has a component containing all the projective modules and such that almost all of its modules belong to the τ -orbits of projective modules and there are at most finitely many modules in Γ belonging to oriented cycles. Suppose Γ contains an injective module. Then Γ is a nonsemiregular component, and hence, by (5.3), A is tilted and Γ is a connecting component. Clearly, then Γ is postprojective.

If Γ contains no injective modules, then Γ is in fact a postprojective component (see [5](6.7) or [16](2.1)). Clearly, a postprojective component containing all projective modules and no injective modules is indeed connecting and (ii) follows.

(ii) \Rightarrow (iii) By (2.6), all modules which are successors of complete slice belong to \mathcal{C} . Now, if Γ_A has a postprojective component Γ with a complete slice, then almost all modules in $\text{ind } A$ are successors of a complete slice in Γ . This proves (iii).

(iii) \Rightarrow (i) Clear. \square

6.3. Corollary. *Let A be a quasitilted algebra which is not tilted. Then, there are infinitely many indecomposable modules X with $pdX = 2$ and infinitely many indecomposable modules Y with $pdY = 2$.*

6.4. It has been shown independently in [1] and [23] that a representation-infinite algebra is concealed if and only if $pdX \leq 1$ and $\text{id}X \leq 1$ for almost all $X \in \text{ind } A$. The next result is also a direct consequence of (6.2).

Corollary. *The following are equivalent for a representation-infinite artin algebra A :*

- (a) $pdX \leq 1$ and $\text{id}X \leq 1$ for almost all $X \in \text{ind } A$.
- (b) A is concealed.
- (c) A is quasitilted and $\mathcal{C} \cap \mathcal{D}$ is cofinite in $\text{ind } A$.

6.5. We shall also mention the next two results that are direct consequences of the previous sections.

Proposition. *Let A be a quasitilted algebra, and Γ be a component of Γ_A .*

- (a) *If Γ contains a projective module, then $\Gamma \subset \mathcal{D} \setminus \mathcal{C}$ if and only if Γ has no complete slice.*
- (b) *If Γ contains an injective module, then $\Gamma \subset \mathcal{C} \setminus \mathcal{D}$ if and only if Γ has no complete slice.*

Proof. We shall only prove (a) because the proof of (b) is similar.

(a) Let Γ be a component containing a projective module. If $\Gamma \subset \mathcal{D} \setminus \mathcal{C}$ then, clearly, Γ contains no complete slice (2.6).

Suppose now that Γ has no complete slice. If A is not tilted, then by (5.2), $\Gamma \subset \mathcal{D} \setminus \mathcal{C}$. Moreover, if Γ contains oriented cycles or has infinitely many τ -orbits, then by (5.1), $\Gamma \subset \mathcal{D} \setminus \mathcal{C}$. It remains to show the result when A is tilted, and Γ is a component without oriented cycles and with only finitely many τ -orbits. Since by hypothesis, Γ is not a connecting component, we infer that Γ is postprojective and it does not contain injective module. Clearly, $\Gamma \subset \mathcal{D}$. Suppose now that $\Gamma \cap \mathcal{C}$ has a module X .

Observe that Γ does not contain all the projective modules and in fact, since A is connected, there exist indecomposable projective modules $P \in \Gamma$ and $P' \notin \Gamma$ with $\text{Hom}_A(P, P') \neq 0$. Since $\text{Hom}_A(P, P') = \text{rad}^\infty(P, P')$, we infer that for each $t \geq 1$, there exist a path of irreducible maps

$$P = Y_0 \longrightarrow Y_1 \longrightarrow \cdots \longrightarrow Y_t$$

and a nonzero map $f_t \in \text{Hom}(Y_t, P')$. Note that all the successors of X are in \mathcal{C} and hence there are only finitely many modules in Γ which are not in \mathcal{C} . Therefore, there exists t such that $\tau Y_t \in \mathcal{C}$ and $\text{Hom}_A(Y_t, P') \neq 0$, or equivalently $\text{id}\tau Y_t > 1$ (by (1.2)), a contradiction. Therefore, $\Gamma \cap \mathcal{C} = \emptyset$ as required. \square

6.6. Proposition. *The following are equivalent for a quasitilted algebra A :*

- (a) *Each nonregular component is either contained in $\mathcal{D} \setminus \mathcal{C}$ or contained in $\mathcal{C} \setminus \mathcal{D}$.*
- (b) *A is either not tilted or a tilted of the form $A = \text{End}_H(T)$, where T is a regular tilting module over a hereditary algebra H .*

Proof. (a) \Rightarrow (b) Suppose A is tilted. Then Γ_A contains a connecting component Γ . If Γ is nonregular, then by (a), it is contained either in $\mathcal{D} \setminus \mathcal{C}$ or in $\mathcal{C} \setminus \mathcal{D}$, a contradiction to the fact that Γ contains a complete slice lying in $\mathcal{C} \cap \mathcal{D}$ (2.6). Then Γ is a regular and, by (2.1), $A = \text{End}_H(T)$ where T is a regular tilting module over a hereditary algebra H .

(b) \Rightarrow (a) Let Γ be a nonregular component of Γ_A . If A is not tilted, then by (5.2), Γ is either contained in $\mathcal{D} \setminus \mathcal{C}$ or contained in $\mathcal{C} \setminus \mathcal{D}$. If now $A = \text{End}_H(T)$, where T is a regular tilting module over a hereditary algebra H , then Γ should not be the connecting component of Γ_A (2.1), and hence it does not contain complete slice. By [20], Γ is semiregular and by (5.1) Γ is contained in $\mathcal{D} \setminus \mathcal{C}$ in case it has projective modules, or in $\mathcal{C} \setminus \mathcal{D}$ in case it has injective modules. This proves the result. \square

ACKNOWLEDGEMENTS

The both authors gratefully acknowledge support from the Polish Scientific Grant KBN. The first named author also gratefully acknowledges support from FAPESP and CNPq, Brasil, and the hospitality of the Nicholas Copernicus University.

REFERENCES

1. I. Assem, F. U. Coelho, *Glueings of tilted algebras*, J. Pure and Applied Algebra 96(3) (1994) 225-243.
2. M. Auslander, I. Reiten, *Representation theory of artin algebras V*, Comm. Algebra 5 (1977) 519-554.
3. M. Auslander, I. Reiten, S. Smalø, *Representation theory of artin algebras*, Cambridge Studies in Advanced Mathematics 36, Cambridge Univ. Press, 1995.
4. R. Bautista and S. Smalø, *Non-existent cycles*, Comm. Algebra 11 (1983) 1755-1767.
5. F. U. Coelho, *Components of Auslander-Reiten quivers containing only preprojective modules*, J. Algebra 157 (1993) 472-488.
6. F. U. Coelho, *A note on preinjective partial tilting modules*, in: *Representations of Algebras, CMS Conference Proceedings*, Vol. 14 (1994) 109-115.
7. F. U. Coelho, E. N. Marcos, H. A. Merklen, A. Skowroński, *Domestic semiregular branch enlargements of tame concealed algebras*, preprint 1994.
8. V. Dlab and C. M. Ringel, *Indecomposable representations of graphs and algebras*, Memoirs Amer. Math. Soc. 173 (1976).
9. D. Happel, U. Preiser, C. M. Ringel, *Vinberg's characterisation of Dynkin diagrams using sub-additive functions with applications to Dtr-periodic modules*, in: *Representation Theory II*, Springer Lecture Notes Math. 832 (1980) 280-294.
10. D. Happel and C. M. Ringel, *Tilted algebras*, Trans. Amer. Math. Soc. 274 (1982) 399-443.
11. D. Happel, I. Reiten and S. Smalø, *Tilting in abelian categories and quasitilted algebras*, preprint (Trodheim 1994).
12. O. Kerner, *Tilting wild algebras*, J. London Math. Soc. 39 (1989) 29-47.
13. O. Kerner, *Stable components of wild tilted algebra*, J. Algebras 142 (1991) 37-57.
14. H. Lenzing, J. A. de la Peña, *Wild canonical algebras*, Math. Z., to appear.
15. H. Lenzing, J. A. de la Peña, *Algebras with a separating tubular family*, preprint, 1994.
16. S. Liu, *Semi-stable components of an Auslander-Reiten quiver*, Journal London Math. Soc. 47 (1993) 405-416.
17. S. Liu, *The connected components of the Auslander-Reiten quiver of a tilted algebra*, J. Algebra 161 (1993) 505-523.
18. C. M. Ringel, *Finite dimensional hereditary algebras of wild representation type*, Math. Z. 161 (1978) 235-255.
19. C. M. Ringel, *Tame algebras and integral quadratic forms*, Springer Lecture Notes in Math. 1099 (1984).
20. C. M. Ringel, *The regular components of Auslander-Reiten quiver of a tilted algebra*, Chinese Ann. Math. 9B (1988) 1-18.
21. C. M. Ringel, *The canonical algebras*, in: *Topics in Algebra*, Banach Center Publ., 26, Part I (PWN Warsaw 1990) 407-432.
22. A. Skowroński, *Regular Auslander-Reiten components containing directing modules*, Proc. Amer. Math. Soc. 120 (1994) 19-26.

23. A. Skowroński, *Minimal representation-infinite artin algebras*, Math. Proc. of Cambridge Phil. Soc. 116 (1994) 229-243.
24. A. Skowroński, *Cycle-finite algebras*, J. Pure Appl. Algebra, to appear.
25. A. Skowroński, *On omnipresent tubular families of modules*, preprint (Toruń 1994).
26. A. Skowroński, *Module categories with finite short cycles*, in preparation.
27. H. Strauss, *On the perpendicular category of a partial tilting module*, J. Algebra 144 (1991) 43-66.
28. Y. Zhang, *The structure of stable components*, Can. J. Math. 43 (1991) 652-672.

DEPARTAMENTO DE MATEMÁTICA-IME, UNIVERSIDADE DE SÃO PAULO, R. DO MATÃO, 1010
CIDADE UNIVERSITARIA, SÃO PAULO, SP, 05508-900, BRASIL

E-mail address: fucoelho@ime.usp.br

INSTITUTE OF MATHEMATICS, NICHOLAS COPERNICUS UNIVERSITY, CHOPINA 12/18, TORUŃ,
POLAND

E-mail address: skowron@mat.torun.edu.pl

TRABALHOS DO DEPARTAMENTO DE MATEMÁTICA
TÍTULOS PUBLICADOS

- 94-01 BIANCONI, R. A note on the construction of a certain class of Kleinian groups. 9p.
- 94-02 HENTZEL, I.R., JACOBS, D.P., PERESI, L.A. & SVERCHKOV, S.R. Solvability of the ideal of all weight zero elements in Bernstein Algebras. 11p.
- 94-03 CORREA, I. & PERESI, L.A. Bernstein-Jordan Algebras of dimension five, 6p.
- 94-04 ABADIE, B. The range of traces on the K_0 -group of quantum Heisenberg manifolds. 13p.
- 94-05 BIANCONI, R., GUSEVSKII, N. and KLIMENKO, H. Schottky type groups and Kleinian groups acting on S^3 with the limit set a wild Cantor set. 30p.
- 94-06 GORODSKI, C. Minimal Hyperspheres in Rank Two Compact Symmetric Spaces. 20 p.
- 94-07 BARROS, L.G.X. de and WATSON, B. On Modular Loop Algebras of RA2 Loops. 19p.
- 94-08 BARROS, L.G.X. de Loop Rings of I.P. Loops. 12p.
- 94-09 BIANCONI, R. A note on the non-interpretability of a discrete order in an o-minimal (dense) structure. 9p.
- 94-10 GOODAIRE, E.G. and POLCINO MILIES, C. On the loop of units of an alternative loop ring, 10p.
- 94-11 GOODAIRE, E.G. and POLCINO MILIES, C. Finite subloops of units in an alternative loop ring. 8p.
- 94-12 JURIAANS, S.O. Trace Properties of Torsion Units in Group Rings. 7p.
- 94-13 FALBEL, E. and GORODSKI, C. On Contact Sub-Riemannian Symmetric Spaces. 22p.
- 94-14 GORODSKI, C. Minimal Sphere Bundles in Euclidean Spheres. 32p.
- 94-15 COELHO, F.U., MARCOS, E.N., MERKLEN, H.A. and SKOWRONSKI, A. Module categories with infinite radical cube zero. 28p.
- 94-16 CHAVES, R.M.B. On two parameter envelopes of N-Spheres in a real space form. 17p.
- 94-17 BARROS, L.G.X and JURIAANS, S.O. Units in Integral Loop Rings. 13p.
- 94-18 BRASIL, JR. A. Hypersurfaces in $S^4(1)$ with constant mean curvature and constant Gauss-Kronecker curvature. 7p.
- 94-19 COSTA R., NEUBURG, M. and SUAZO, A. Some Results in the Theory of Bernstein Algebras. 19p.
- 94-20 FALBEL E., GORODSKI C. Sub-Riemannian Homogeneous Spaces in Dimensions 3 and 4. 32p.
- 94-21 BARROS, L. G. X. de and POLCINO MILIES, C. Loops Algebras of Code Loops. 13p.
- 94-22 CARRARA, V.L., RUAS, M.A.S. and SAEKI, O. A note on codimension two submanifolds with at most four critical points. 11p.

- 94-23 COELHO F.U., MARCOS E.N., MERKLEN H.A., and SKOWRONSKI, A. Domestic Semiregular Branch Enlargements of Tame Cancealed Algebras. 14p.
- 94-24 DOKUCHAEV M. A., and JURIAANS, O. Finite Subgroups in Integral Group Rings. 20p
- 94-25 FALBEL E., GORODSKI C. and VELOSO J.M. Conformal Sub-Riemannian Geometry in Dimension 3. 12p.
- 95-01 BARROS, L.G.X. de and Juriaans, S.O. Loops whose Loop Algebras are Flexible. 25p.
- 95-02 GUIDORIZZI, H.L. - Jordan canonical form: an elementary proof, 12 p.
- 95-03 CATALAN A., and COSTA R. E-Ideals in Baric Algebras. 14p.
- 95-04 MARTIN, P.A. On the generating function of the decimal expansion of an irracional real numbers. 7p.
- 95-05 COELHO F.U., MARCOS E.N., MERKLEN H.A. and PLATZECK.M.I. Modules of Infinite Projective Dimension over Algebras whose Idempotent Ideals are Projective. 13p.
- 95-06 GUIDORIZZI, H. L. The family of functions $S_{\alpha,k}$ and the Liénard Equation. 22p.
- 95-07 GUIDORIZZI, H. L. On the Existence of Periodic Solution for the Equation $\ddot{x} + \alpha x^{2n+1} \dot{x} + x^{4n+3} = 0$. 5p.
- 95-08 CORTIZO, S.F. Extensões Virtuais. 27p.
- 95-09 CORTIZO, S.F. Cálculo Virtual. 31p.
- 95-10 GUIDORIZZI, H. L. On Periodic Solutions of Systems of the Type $\ddot{x} = H(y)$, $\dot{y} = - \sum_{i=1}^n f_i(x)H_i(y) - g(x)$. 16p.
- 95-11 OLIVA, S. M., PEREIRA, A. L. Attractors for Parabolic Problems with Non linear Boundary Conditions in Fractional Power Spaces. 28p.
- 95-12 CORDARO, P. D. Global hypoellipticity for $\bar{\partial}_b$ on certain compact three dimensional CR manifolds. 11p.
- 95-13 COELHO, F.U. and SKOWRONSKI, A. On Auslander-Reiten Components for Quasitilted Algebras. 16p.

NOTA: Os títulos publicados dos Relatórios Técnicos dos anos de 1980 a 1993 estão à disposição no Departamento de Matemática do IME-USP. Cidade Universitária "Armando de Salles Oliveira" Rua do Matão, 1010 - Butantã Caixa Postal - 66281 (Ag. Cidade de São Paulo) CEP: 05389-970 - São Paulo - Brasil