

HADRONS
2025 Porto Alegre

• • • XVI International Workshop on Hadron Physics

10-14 Mar | Centro Cultural da UFRGS

Invited speakers

Kenji Fukushima
The University of Tokyo

Chun Shen
Wayne State University

Lisheng Geng
Beihang University

Fernanda Steffens
University of Bonn

National Committee

Kanchan Khemchandani (UNIFESP)
Jun Takahashi (UNICAMP)
Letícia Palhares (UERJ)
Luciano Abreu (UFBA)
Ricardo Sonego Farias (UFSM)
Victor Gonçalves (UFPel)
Tiago Nunes (UFSC)

Organizing Committee

Victor Gonçalves (UFPel) - Chair
Gustavo Gil da Silveira (UFRGS)
Magno Machado (UFRGS)
Ricardo Sonego Farias (UFSM)

Registrations open from
10 Aug 2024 to 14 Feb 2025
<https://indico.cern.ch/e/hadrons2025>



Começar (Iniciar) 10 de mar. de 2025 09:00
Fim 14 de mar. de 2025 19:00

America/Sao_Paulo



Centro Cultural - UFRGS
 Jacarandá Auditorium
 Rua Eng. Luiz Englert, 333 - Porto Alegre-RS
[Ir para o mapa](#)



Victor Goncalves



Distribuído por Indico v3.3.7-pre

[Ajuda](#) | [Contato](#) | [Termos e condições](#) | [Encurtador de URL](#) | [Privacidade](#)





ID da Contribuição: 131

Tipos: **Poster**

Studying gauge-space geometry via lattice QCD

quarta-feira, 12 de março de 2025 17:58 (2 minutos)

Quantum chromodynamics (QCD) is the theory that describes the nuclear strong interaction between quarks, mediated by gauge fields carrying color charge, the gluons. QCD has a very important and unsolved problem, which is finding a mechanism to explain why quarks are never seen alone in nature, but only forming bound states, such as protons and neutrons. This constitutes the color confinement problem, which may be studied in the framework of quantum field theory. It is a hard task because of the non-Abelian character of the theory: the gluons themselves interact with each other and, in the low-energy regime, this forbids us to treat the theory perturbatively, as done in quantum electrodynamics (QED).

Consequently, the quantization of non-Abelian gauge fields presents several issues. From the technical point of view, integrating the gauge field over all its possible configurations in the path integral involves the inversion of operators in the Yang-Mills Lagrangian with null eigenvalues, leading to divergences. Using the Faddeev-Popov method, we impose a gauge condition for the gluon field, adding new integration variables in gauge space. The curves that connect physically equivalent fields through gauge transformations in this space are called gauge orbits and, in principle, the resulting formulation solves the divergences by factorization.

Ideally, the gauge-fixing method just described causes a gauge orbit to intersect the region specified by the gauge condition only once. But this is not guaranteed for a general Yang-Mills theory, and hence there are still ambiguities coming from equivalent configurations in gauge space, called Gribov copies. For a gauge transformation, the existence of Gribov copies is directly related to the fact that there are non-trivial eigenstates of the Faddeev-Popov operator with null eigenvalues.

Among the proposed confinement scenarios, the one due to Gribov and Zwanziger associates color confinement to infrared properties of propagators of (gauge-fixed) fields around null eigenvalues of the Faddeev-Popov operator in gauge space. Our main objective is to explore and test the Gribov-Zwanziger confinement scenario, comparing analytical predictions with numerical results from lattice QCD. As a preliminary result of our lattice data, we calculate the static quark-antiquark potential. In addition, we show our results for the Landau gauge fixing procedure using different algorithms and the measurement of the gluon propagator.

Autores: KÜL, João Octavio (Universidade de São Paulo); MENDES, Tereza

Apresentador: KÜL, João Octavio (Universidade de São Paulo)

Classificação da Sessão: Poster session