



PEETERS

EQUALITY IN LINEAR LOGIC

Author(s): Marcelo E. CONIGLIO and Francisco MIRAGLIA

Source: *Logique et Analyse*, Mars - Juin 1996, NOUVELLE SÉRIE, Vol. 39, No. 153/154, CONTEMPORARY BRAZILIAN RESEARCH IN LOGIC: Part I (Mars - Juin 1996), pp. 113-151

Published by: Peeters Publishers

Stable URL: <https://www.jstor.org/stable/44084560>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



Peeters Publishers is collaborating with JSTOR to digitize, preserve and extend access to
Logique et Analyse

JSTOR

EQUALITY IN LINEAR LOGIC

Marcelo E. CONIGLIO and Francisco MIRAGLIA

1. *Quantales*

In this section we introduce the basic definitions and results of the theory of quantales (a good reference is [Ros]). Quantales were introduced by Mulvey ([Mul]) as an algebraic tool for studying representations of non-commutative C^* -algebras. Informally, a quantale is a complete lattice Q equipped with a product distributive over arbitrary sup's. The importance of quantales for Linear Logic is revealed in Yetter's work ([Yet]), who proved that semantics of classical linear logic is given by a class of quantales, named Girard quantales, which coincides with Girard's phase semantics. An analogous result is obtained for a sort of non-commutative linear logic, as well as intuitionistic linear logic without negation, which suggest that the utilisation of the theory of quantales (or even weaker structures, such that $*$ -autonomous posets) might be fruitful in studying the semantic of several variants of linear logic.

As usual, we denote the order in a lattice by \leq , while \vee and \wedge denote the operations of sup and inf, respectively. We write \top for the largest element in a lattice and $\mathbf{0}$ for its smallest element.

Definition 1.1 A quantale is a complete lattice Q with an associative binary operation $\otimes: Q \times Q \longrightarrow Q$, which distributes on the right and on the left of arbitrary sup's, i.e.:

- [Q1] $a \otimes (b \otimes c) = (a \otimes b) \otimes c$, for every $a, b, c \in Q$
- [Q2] $a \otimes (\bigvee_{i \in I} a_i) = \bigvee_{i \in I} (a \otimes a_i)$, $(\bigvee_{i \in I} a_i) \otimes a = \bigvee_{i \in I} (a_i \otimes a)$

A quantale Q is unital if it has an element $\mathbf{1} \in Q$ such that $a \otimes \mathbf{1} = \mathbf{1} \otimes a = a$, for every $a \in Q$. A quantale Q is commutative if $a \otimes b = b \otimes a$, for every $a, b \in Q$.

A morphism of quantales is an operator between quantales which preserves \otimes and arbitrary sup's.

It's easily seen that the above axioms imply that \otimes is increasing in both coordinates, that is

If $a \leq b$ then, $\forall c \in Q$, $a \otimes c \leq b \otimes c$ and $c \otimes a \leq c \otimes b$

We register a classic result:

Proposition 1.2 *The endomorphisms $a \otimes \cdot, \cdot \otimes a : Q \longrightarrow Q$ have right adjoints, denoted by $a \rightarrow_r \cdot$ and $a \rightarrow_l \cdot$, respectively. Thus,*

$$a \otimes c \leq b \text{ iff } c \leq a \rightarrow_r b \quad c \otimes a \leq b \text{ iff } c \leq a \rightarrow_l b$$

and consequently

$$\begin{aligned} a \rightarrow_r b &= \bigvee \{c \in Q : a \otimes c \leq b\} \\ a \rightarrow_l b &= \bigvee \{c \in Q : c \otimes a \leq b\}. \end{aligned}$$

Definition 1.3 *Let Q be a quantale. A map $j : Q \longrightarrow Q$ is said to be a*

a) *quantic nucleus if it satisfies:*

$$\begin{array}{ll} [NQ1] & a \leq b \text{ implies } j(a) \leq j(b) \\ [NQ3] & j(j(a)) = j(a) \end{array} \quad \begin{array}{ll} [NQ2] & a \leq j(a) \\ [NQ4] & j(a) \otimes j(b) \leq j(a \otimes b) \end{array}$$

b) *quantic conucleus if it satisfies:*

$$\begin{array}{ll} [CNQ1] & a \leq b \text{ implies } g(a) \leq g(b) \\ [CNQ3] & g(g(a)) = g(a) \end{array} \quad \begin{array}{ll} [CNQ2] & g(a) \leq a \\ [CNQ4] & g(a) \otimes g(b) \leq g(a \otimes b) \end{array}$$

Quantic nuclei and conuclei are important, because they determine the quotients and subobjects in the category of quantales.

Definition 1.4 *Let Q be a quantale. A subset $S \subseteq Q$ is a subquantale of Q if it is closed under \otimes and arbitrary sup's.*

Proposition 1.5 [Ros]: (a) *If $Q \xrightarrow{j} Q$ is a quantic nucleus, then $Q_j = \{x \in Q : j(x) = x\}$ is a quantale where the operations \otimes^j, \vee^j and \wedge^j in Q_j are given by:*

$$a \otimes^j b = j(a \otimes b) \quad \vee_{i \in I}^j a_i = j(\vee_{i \in I} a_i) \quad \wedge_{i \in I}^j a_i = \wedge_{i \in I} a_i.$$

Moreover, the map $j : Q \longrightarrow Q_j$ given by $a \mapsto j(a)$, is a surjective morphism of quantales. Further, every surjective morphism of quantales can be represented in this form.

(b) If $Q \xrightarrow{g} Q$ is a quantic conucleus, then $Q_g = \{ x \in Q : g(x) = x \}$ is a subquantale where, with notation as in (a), $\bigwedge_{i \in I} a_i = g(\bigwedge_{i \in I} a_i)$. Moreover, every subquantale is of this form, i.e.,

if $S \subseteq Q$ is a subquantale, then there exists a quantic conucleus g in Q such that $S = Q_g$.

Definition 1.6 An element $\perp \in Q$ is dualizing if:

$$(a \rightarrow, \perp) \rightarrow, \perp = a = (a \rightarrow, \perp) \rightarrow, \perp, \text{ for every } a \in Q.$$

An element $s \in Q$ is cyclic if:

$$a \rightarrow, s = a \rightarrow, s, \text{ for every } a \in Q.$$

Proposition 1.7 [Yet]: Let Q be a quantale and s, \perp be elements of Q .

a) $s \in Q$ is cyclic iff for all a_1, \dots, a_n in Q ,

$a_1 \otimes \cdots \otimes a_n \leq s$ implies $a_{\pi(1)} \otimes a_{\pi(2)} \otimes \cdots \otimes a_{\pi(n)} \leq s$,
for all cyclic permutations π of $\{1, \dots, n\}$.

b) If $\perp \in Q$ is dualizing, then Q is unital and we have:

$$\mathbf{1} = \perp \rightarrow, \perp = \perp \rightarrow, \perp.$$

Definition 1.8 A Girard quantale is a quantale which has a cyclic dualizing element \perp . The operator $\cdot \rightarrow, \perp = \cdot \rightarrow, \perp \equiv_{\text{def}} \cdot \rightarrow \perp$ is called linear negation, and we write $a^\perp = a \rightarrow \perp$ (note that $\mathbf{1} = \perp^\perp$, $\perp = \mathbf{1}^\perp$, and $a = a^{\perp\perp}$).

Next proposition is of frequent use when computing in a Girard quantale.

Proposition 1.9 [Ros]: Let Q be a Girard quantale with a cyclic dualizing element \perp and let $a, b \in Q$. Then:

$$\begin{array}{ll} (1) & a \rightarrow, b = (a \otimes b^\perp)^\perp \\ (2) & a \rightarrow, b = (b^\perp \otimes a)^\perp \\ (3) & a \otimes b = (a \rightarrow, b^\perp)^\perp \\ (4) & b \otimes a = (a \rightarrow, b^\perp)^\perp \\ (5) & a \rightarrow, b = b^\perp \rightarrow, a^\perp \\ (6) & a \rightarrow, b = b^\perp \rightarrow, a^\perp \end{array}$$

Proposition 1.10 [Yet]: Let Q be an unital quantale and $s \in Q$ cyclic. Then, $j : Q \longrightarrow Q$, given by $j(a) = (a \rightarrow s) \rightarrow s$ is a quantic nucleus, and $Q_j = \{ a \in Q : j(a) = a \}$ is a Girard quantale, where $\perp = s \in Q_j$.

Example 1.11: The Phase Quantales (Girard)

Let $(M, \cdot, 1)$ be a monoid. We define $A \cdot B = \{ a \cdot b : a \in A, b \in B \}$ $\forall A, B \subseteq M$. Let $\perp \subseteq M$ be such that $a \cdot b \in \perp$ implies $b \cdot a \in \perp$ (for example, \perp can be a semiprime ideal or the complement of a prime ideal of M).

Thus, $\wp(M)$ is a quantale with the product defined above, and with sup's e inf's calculated as unions (\cup) and intersections (\cap). In fact, it's an unital quantale, where $\mathbf{1} = \{ 1 \}$ and \perp is cyclic. Then, by proposition 1.10, $\wp(M)_j$ is a Girard's quantale containing \perp . Their elements are called facts, and we have:

$$A \otimes B = (A \cdot B)^\perp$$

$$\bigvee_{i \in I} A_i = (\bigcup_{i \in I} A_i)^\perp; \text{ in particular, } A \vee B = (A \cup B)^\perp \equiv_{\text{def}} A \oplus B,$$

$$\bigwedge_{i \in I} A_i = \bigcap_{i \in I} A_i; \text{ in particular, } A \wedge B = A \cap B \equiv_{\text{def}} A \& B.$$

The next result tells us that every Girard quantale is of this form.

Proposition 1.12 [Ros]: If Q is a Girard's quantale, then Q is isomorphic to a phase quantale.

We are now going to relate the exponentials treated by Yetter ([Yet]) and Avron ([Avr]) with certain concepts in the theory of quantales.

Definition 1.13 [Yet]: An open modality in a quantale Q is a map $\mu : Q \longrightarrow Q$ satisfying:

$$\begin{array}{ll} [M1] \quad \mu(a) \leq a & [M2] \quad a \leq b \text{ implies } \mu(a) \leq \mu(b) \\ [M3] \quad \mu(\mu(a)) = \mu(a) & [M4] \quad \mu(\mu(a) \otimes \mu(b)) = \mu(a) \otimes \mu(b). \end{array}$$

An open modality μ is said to be

- central if $b \otimes \mu(a) = \mu(a) \otimes b$ for every $a, b \in Q$.
- idempotent if $\mu(a) \otimes \mu(a) = \mu(a)$ for every $a \in Q$.
- weak if Q is unital, $\mu(\mathbf{1}) = \mathbf{1}$, and $\mu(a) \leq \mathbf{1}$ for every $a \in Q$.

Let $M(Q) = \{ \mu : \mu \text{ is an open modality in } Q \}$, partially ordered by pointwise order.

Proposition 1.14 [Yet]: Let Q be a quantale (resp. unital quantale). Then, there exists a unique maximal open modality in Q central (resp. central, idempotent and weak) denoted by c_m (resp. $!_m$) given by:

$$c_m(x) = \bigvee \{ a \in Q : a \leq x, a \in Z(Q) \}$$

$$!_m(x) = \bigvee \{ a \in Q : a \leq x \wedge \mathbf{1}, a = a \otimes a, a \in Z(Q) \},$$

where $Z(Q) = \{ a \in Q : a \otimes b = b \otimes a, \text{ for every } b \in Q \}$.

Definition 1.15 [Avr]: Let Q be an unital quantale. A map $B: Q \longrightarrow Q$ is a modal operation if it satisfies, for all $x, y \in Q$:

$$\begin{array}{ll} [B1] & B(\mathbf{1}) = \mathbf{1} \\ [B3] & B(B(x)) = B(x) \end{array}$$

$$\begin{array}{ll} [B2] & B(x) \leq x \\ [B4] & B(x) \otimes B(y) = B(x \wedge y). \end{array}$$

Proposition 1.16 Let Q be a quantale, and $\mu: Q \longrightarrow Q$ a map.

- (i) μ is an open modality iff μ is a quantic conucleus.
- (ii) If Q is unital, then μ is an open, idempotent and weak modality iff μ is a modal operation.

Proof: (i) We must prove that [M4] is equivalent to [CNQ4]. Let μ be an open modality. Because $\mu(a) \otimes \mu(b) \leq a \otimes b$, we get $\mu(a) \otimes \mu(b) = \mu(\mu(a) \otimes \mu(b)) \leq \mu(a \otimes b)$.

Conversely, if μ is a quantic conucleus, then

$$\begin{aligned} \mu(a) \otimes \mu(b) &= \mu(\mu(a)) \otimes \mu(\mu(b)) \\ &\leq \mu(\mu(a) \otimes \mu(b)) \leq \mu(a) \otimes \mu(b). \end{aligned}$$

(ii) Assume that μ is an open, weak and idempotent modality. Then, it clearly satisfies conditions [B1], [B2], [B3]. Because $\mu(a) \leq a$ and $\mu(b) \leq \mathbf{1}$, we get $\mu(a) \otimes \mu(b) \leq a \otimes \mathbf{1} = a$. Similarly, $\mu(a) \otimes \mu(b) \leq b$ and therefore, $\mu(a) \otimes \mu(b) \leq a \wedge b$. Thus, $\mu(a) \otimes \mu(b) = \mu(\mu(a) \otimes \mu(b)) \leq \mu(a \wedge b)$.

Now, since μ is increasing and $a \wedge b \leq a, b$, then $\mu(a \wedge b) \leq \mu(a), \mu(b)$. Thus,

$$\mu(a \wedge b) = \mu(a \wedge b) \otimes \mu(a \wedge b) \leq \mu(a) \otimes \mu(b),$$

and so μ satisfies [B4].

Conversely, if μ is a modal operation, then [M1] is just [B2]; [M3] is [B3], while [M4] is equivalent to [CNQ4], in the presence of [M1], [M2], [M3], by item (i) above.

Condition [M2] is item (5) of Lemma 4.2 in [Avr] and μ satisfies [CNQ4] by item 8 of Lemma 4.2 in [Avr]. It follows from items 4 and 1 in that same Lemma, that $\mu(x) \leq \mathbf{1}$ for every $x \in Q$. Thus, [B1] yields that μ is weak. \square .

Definition 1.17 A frame (or Complete Heyting algebra) is a quantale where $\otimes = \wedge$.

Definition 1.18 Let Q be a quantale and let \top be the maximum of Q . We say that $x \in Q$ is

- a) right-sided (in Q) if $x \otimes \top \leq x$,
- b) left-sided (in Q) if $\top \otimes x \leq x$,
- c) two-sided (in Q) if it is right and left sided in Q .
- d) idempotent if $x \otimes x = x$.

Proposition 1.19 [Ros]: Let Q be a quantale, and g a quantic conucleus in Q . Are equivalent:

- (a) Q_g is a frame.
- (b) $g(a) \otimes g(b) = g(a \wedge b)$.
- (c) every $x \in Q_g$ is idempotent and two-sided (in Q_g).

Definition 1.20 A map g satisfying the conditions above is called a localic conucleus and Q_g is called a localic subquantale.

Since open, weak idempotent modalities are localic conuclei μ such that $\mu(\mathbf{1}) = \mathbf{1}$, we have

Corollary 1.21 Let Q be an unital quantale, and let μ be an open, weak and idempotent modality. Then, Q_μ is a frame.

Thus, the fixed points of the interpretation of the modality $!$ (of course) lie in a localic subquantale (complete Heyting algebra) of any quantale in which linear logic is interpreted.

2. Linear Calculus with Equality

In this section we discuss the laws for a binary predicate representing equality. The goal is to define a reflexive, symmetric and transitive predicate satisfying the substitution (Leibnitz's) rule for the class of all formulas. We may assume, just as in Classical Logic (CL), that we have substitution for atomic formulas. With this model in mind, we shall define a prototype of a linear calculus with equality, called (LLE_1) . Our formulation will use sequents in Linear Logic (LL). Analogously, we will set down a calculus with equality for the ($MALL$) fragment, i.e., the fragment without exponentials, indicated by (LLE_0) . Starting from the property of substitution for elementary formulas, we prove that $(\cdot = \cdot)$ must be $\leq \mathbf{1}$ and idempotent in $(MALL)$, and open in the general case; in other words, $(\cdot = \cdot)$ must be intuitionistic.

Definition 2.1 A first order linear language with equality \mathbb{L} consists in a countable set of predicate symbols $\wp = \{ P_n : n \in \omega \} \cup \{ = \}$ (where " $=$ " is binary), a countable set of variables $V = \{ v_n : n \in \omega \}$, together with the symbols:

$1, \perp, \top, \mathbf{0}, {}^\perp, \vee, \wedge, \oplus, \&, \otimes, \sqcup, !, ?, .$

Definition 2.2 The formulas of \mathbb{L} , $FOR(\mathbb{L})$, are defined recursively:

- [F1] $1, \perp, \top, \mathbf{0} \in FOR(\mathbb{L})$.
- [F2] If $P \in \wp$ is a predicate of arity n and x_1, \dots, x_n are variables, then $P(x_1, \dots, x_n)$ and $P(x_1, \dots, x_n)^\perp \in FOR(\mathbb{L})$.
- [F3] if $F, G \in FOR(\mathbb{L})$ then $F \otimes G, F \sqcup G, F \& G, F \oplus G \in FOR(\mathbb{L})$.
- [F4] if $F \in FOR(\mathbb{L})$, then $!F, ?F \in FOR(\mathbb{L})$.
- [F5] if $F \in FOR(\mathbb{L})$ and $x \in V$, then $\wedge x.F$ and $\vee x.F \in FOR(\mathbb{L})$.

Every occurrence of a variable x in a formula F is free except in a subformula of the type $\wedge x.G$ or $\vee x.G$ (which are bounded occurrences). We shall write

$ELF(\mathbb{L}) = \{ P(x_1, \dots, x_n) : P \in \wp - \{ = \}, x_i \in V \}$, the set of elementary formulas

and

$ELF^+(\mathbb{L}) = ELF(\mathbb{L}) \cup \{ (x = y) : x, y \in V \}$, the extended set of elementary formulas.

Definition 2.3 The syntactic linear negation is a map $\perp : FOR(\mathbb{L}) \longrightarrow FOR(\mathbb{L})$ given by the usual rules:

- [NL1] $\perp(1) = \perp, \perp(\perp) = 1, \perp(0) = \top, \perp(\top) = \mathbf{0}$
- [NL2] $\perp(P(x_1, \dots, x_n)) = P(x_1, \dots, x_n)^\perp, \perp((x = y)) = (x = y)^\perp$,
 $\perp(P(x_1, \dots, x_n)^\perp) = P(x_1, \dots, x_n), \perp((x = y)^\perp) = (x = y)$
 (here, $P(x_1, \dots, x_n) \in ELF(\mathbb{L})$)
- [NL3] $\perp(F \otimes G) = \perp(F) \sqcup \perp(G), \perp(F \sqcup G) = \perp(F) \otimes \perp(G),$
 $\perp(F \& G) = \perp(F) \oplus \perp(G), \perp(F \oplus G) = \perp(F) \& \perp(G)$
- [NL4] $\perp(!F) = ?\perp(F), \perp(?F) = !\perp(F)$
- [NL5] $\perp(\wedge x.F) = \vee x.\perp(F), \perp(\vee x.F) = \wedge x.\perp(F)$

We write $\perp(F) = F^\perp$; clearly $F = F^{\perp\perp}$ for every $F \in FOR(\mathbb{L})$.

Definition 2.4 Let A a formula; we define $A[y/x]$ to be the formula obtained from A by replacing all bound occurrences of y by z , where z is the first variable (in the natural order of V) not occurring in A , and then replacing all free occurrences of variable x by y .

Definition 2.5 (Girard) The calculus (LL) for first order commutative linear logic is defined by the axioms and rules below (here, A, B denote formulas, and Γ, Δ denote multisets of formulas):

$$[AX1] \quad \vdash A^\perp, A \quad [AX2] \quad \vdash \top, \Gamma \quad [AX3] \quad \vdash \mathbf{1}$$

$$[CUT] \quad \frac{\vdash \Gamma, A \quad \vdash \Delta, A^\perp}{\vdash \Gamma, \Delta}$$

$$[EXCH] \quad \text{If } \Delta \text{ is a permutation of } \Gamma, \text{ then } \frac{\vdash \Gamma}{\vdash \Delta}$$

$$[\&] \quad \frac{\vdash A, \Gamma \quad \vdash B, \Gamma}{\vdash A \& B, \Gamma}$$

$$[\oplus 1] \quad \frac{\vdash A, \Gamma}{\vdash A \oplus B, \Gamma} \quad [\oplus 2] \quad \frac{\vdash B, \Gamma}{\vdash A \oplus B, \Gamma}$$

$$[\perp] \quad \text{If } \Gamma \text{ is not empty, then } \frac{\vdash \Gamma}{\vdash \perp, \Gamma}$$

$$[\otimes] \quad \frac{\vdash A, \Gamma \quad \vdash B, \Delta}{\vdash A \otimes B, \Gamma, \Delta} \quad [\sqcup] \quad \frac{\vdash A, B, \Gamma}{\vdash A \sqcup B, \Gamma}$$

$$[dereliction] \quad \frac{\vdash A, \Gamma}{\vdash ?A, \Gamma} \quad [weakening] \quad \frac{\vdash \Gamma}{\vdash ?A, \Gamma}$$

$$[contraction] \quad \frac{\vdash ?A, ?A, \Gamma}{\vdash ?A, \Gamma} \quad [!] \quad \frac{\vdash A, ?\Gamma}{\vdash !A, ?\Gamma}$$

$$[\vee] \quad \frac{\vdash A[y/x], \Gamma}{\vdash \vee x.A, \Gamma} \quad [\wedge] \quad \text{If } x \text{ is not free in } \Gamma, \text{ then } \frac{\vdash A, \Gamma}{\vdash \wedge x.A, \Gamma}$$

We can consider as defined connectives the linear implication and the linear equivalence, given by:

$$(A \multimap B) \equiv_{\text{def}} (A^\perp \sqcup B) = (A \otimes B^\perp)^\perp$$

$$(A \circ \multimap B) \equiv_{\text{def}} (A \multimap B) \& (B \multimap A).$$

Definition 2.6 The linear calculus with equality (LLE_0) for the (MALL) fragment of (LL) is defined by adding the axioms below to those in the preceding definition:

$$[R=] : \vdash (x = x); \quad [S=] : \vdash (x = y)^\perp, (y = x); \\ [T=] : \vdash (x = y)^\perp, (y = z)^\perp, (x = z).$$

$$[SUBST] : \vdash (x_1 = y_1)^\perp, \dots, (x_n = y_n)^\perp, \\ F(x_1, \dots, x_n)^\perp, F(x_1 \setminus y_1, \dots, x_n \setminus y_n),$$

where $F(x_1, \dots, x_n)$ denotes an elementary formula with (eventually) x_1, \dots, x_n free and $F(x_1 \setminus y_1, \dots, x_n \setminus y_n)$ is obtained from $F(x_1, \dots, x_n)$ by replacing some occurrences of x_i (which are not in the scope of a y_i -quantifier) by y_i .

$$[I=] : \vdash (x = y)^\perp, (x = y) \otimes (x = y) \quad [\leq 1] : \vdash (x = y)^\perp, \mathbf{1}$$

Definition 2.7 The linear calculus with equality (LLE_1) for (LL) is defined by adding, besides the axioms $[R=]$, $[S=]$, $[T=]$, $[SUBST]$ the following rule

$$[!=] : \vdash (x = y)^\perp, !(x = y)$$

Obviously, all the rules (with exception of $[R=]$) could be formulated as linear implications, for example $[T=]$ could be stated as

$$\vdash (x = y) \otimes (y = z) \multimap (x = z)$$

We have as well that (LLE_1) is equivalent to the calculus obtained from (LLE_1) by replacing $[S=]$ and $[!=]$ by the rule:

$$[S!=] \vdash (x = y)^\perp, !(y = x)$$

3. Semantics

In this section we develop interpretations for the calculi described above. It will be necessary to extend the definitions in [Yet] so that the axioms involving equality are verified.

Definition 3.1 An interpretation of a language \mathbb{L} is a triple $\langle \mathcal{A}, \mathcal{V}, |\cdot|_{\mathcal{A}} \rangle$ where

- \mathcal{A} is an algebra for the theory with constants $\mathbf{1}, \perp, \top, \mathbf{0}$;
- a set of unary operations $\{ \perp, !, ? \}$ (the interpretation of the modalities);
- binary operations $\otimes, \sqcup, \&, \oplus$, and infinitary operations \vee and \wedge ;
- $\mathcal{V} \subseteq \mathcal{A}$, the set of valid elements; and
- $|\cdot|_{\mathcal{A}} : ELF^+(\mathbb{L}) \longrightarrow \mathcal{A}$ is a map.

It is straightforward to see that there exists a (unique) extension of $|\cdot|_{\mathcal{A}}$ to $FOR(\mathbb{L})$, also denoted by $|\cdot|_{\mathcal{A}}$. When there is no risk of confusion, we write $|\cdot|$ for $|\cdot|_{\mathcal{A}}$.

Definition 3.2 A semantics for \mathbb{L} is a class of interpretations. A semantics is sound (with respect to a linear calculus P) if:

$\vdash A_1, \dots, A_n$ is provable in P implies $|A_1 \sqcup \dots \sqcup A_n|_{\mathcal{A}} \in \mathcal{V}$ for every $\langle \mathcal{A}, \mathcal{V}, |\cdot|_{\mathcal{A}} \rangle$ in the semantics.

A semantics for \mathbb{L} is complete (with respect to P) if:

$|F|_{\mathcal{A}} \in \mathcal{V}$, for every $\langle \mathcal{A}, \mathcal{V}, |\cdot|_{\mathcal{A}} \rangle$ in the semantics, implies $\vdash F$ is provable in P .

Definition 3.3 An interpretation of quantales for \mathbb{L} is a Girard quantale Q , together with an assignment $! \mapsto \mu \in M(Q)$ and a map $|\cdot|_Q : ELF^+(\mathbb{L}) \longrightarrow Q$, where $\mathcal{V} = \{ a \in Q : a \geq \mathbf{1} \}$ such that

- $\forall a, b \in Q, a \sqcup b \equiv_{\text{def}} (a^\perp \otimes b^\perp)^\perp$ and $?a \equiv_{\text{def}} (\mu(a^\perp))^\perp$;
- We interpret $\wedge x.A$ and $\vee x.A$ as $\wedge_{y \in \mathcal{V}} |A[y/x]|_Q$ and $\vee_{y \in \mathcal{V}} |A[y/x]|_Q$, respectively.

Recall that $M(Q)$ is the lattice of open modalities in Q (Definition 1.13).

Definition 3.4 The semantics of quantales for (commutative) Linear Logic with equality is the class of interpretation of quantales for \mathbb{L} such that:

[S1] Q is commutative. [S2] $\mu \equiv_{\text{def}} !$ is idempotent and weak.

[S3] $|\cdot|_Q : ELF^+(\mathbb{L}) \longrightarrow Q$ satisfies:

(a) For the Calculus (LLE_0):

$$\begin{aligned}
 [=1] : |x = x| &\geq 1; & [=2] : |x = y| \otimes |y = z| &\leq |x = z|; \\
 [=3] : |x = y| &= |y = x|; \\
 [=4] : |x_1 = y_1| \otimes \cdots \otimes |x_n = y_n| &\otimes |F(x_1, \dots, x_n)| \\
 &\leq |F(x_1, \dots, x_n)| \\
 &\quad (\text{same notation as } [SUBST]); \\
 [=5] : |x = y| &\leq |x = y| \otimes |x = y|; & [=6] : |x = y| &\leq 1
 \end{aligned}$$

(b) For the Calculus (LLE₁), conditions [=5] and [=6] are replaced by the stronger:

$$[=7] : |x = y| \leq !|x = y|.$$

The next result shows that it is always possible to define a map $|\cdot|_Q$ satisfying the above conditions.

Proposition 3.5 If $|\cdot|_Q : ELF(\mathbb{L}) \longrightarrow Q$ is a map, then it can be extended to $ELF^+(\mathbb{L})$ satisfying, in the (MALL) case, conditions [=1], ..., [=6] and, in the general case, conditions [=1], ..., [=4], [=7].

Proof: Let's begin by the general case, in which $! \in M(Q)$ is idempotent and weak.

Let $|\cdot|_Q : ELF(\mathbb{L}) \longrightarrow Q$ be a map, and let $x, y \in V$, $x \neq y$. Define $A_{x,y}$ and $B_{x,y}$ as:

$$\begin{aligned}
 A_{x,y} &= \{ F \in ELF(\mathbb{L}) : \text{neither } x \text{ nor } y \text{ occur in } F \} \\
 B_{x,y} &= \{ F \in ELF(\mathbb{L}) : x \text{ or } y \text{ occur in } F \}
 \end{aligned}$$

In $B_{x,y}$ consider the relation:

$$F \sim G \text{ iff } G \text{ is obtained from } F \text{ by replacing some occurrences of } x \text{ by } y \text{ and/or some occurrences of } y \text{ by } x$$

For example, $P(x,z,y) \sim P(x,z,x) \sim P(y,z,y) \sim P(y,z,x)$. Clearly, \sim is an equivalence relation. Now define,

$$T_{x,y} = \bigwedge \{ |F|_Q \rightarrow |G|_Q : F, G \in B_{x,y}, F \sim G \}.$$

Fact 1: If $|\cdot|$ satisfies [=7] and [=3], then it satisfies [=4] iff $|x = y| \leq T_{x,y}$ for $x \neq y$.

To see this, assume that $F, G \in B_{x,y}$, $F \sim G$. Note that there is $H \in ELF(\mathbb{L})$ such that $H = F(x \setminus y)$, $G = H(y \setminus x)$.

Since $!$ is idempotent, [=7] implies [=5] and so, by [=3], [=4] and [=5] we have that $|x = y| \otimes |F| \leq |H|$; thus,

$$|x = y| \otimes |F| \leq |x = y| \otimes (|x = y| \otimes |F|) \leq |x = y| \otimes |H| \leq |G|.$$

Therefore, $|x = y| \leq |F| \rightarrow |G|$, proving that $|x = y| \leq T_{x,y}$.

Conversely, suppose that $|x = y| \leq T_{x,y}$ for $x \neq y$. Then, if $F \in \bigcup_{i=1}^n B_{x_i, y_i}$,

$$|x_n = y_n| \leq |F(x_1, \dots, x_n)| \rightarrow |F(x_1, \dots, x_{n-1}, x_n \setminus y_n)|$$

$$\text{and therefore, } |x_n = y_n| \otimes |F(x_1, \dots, x_n)| \leq |F(x_1, \dots, x_{n-1}, x_n \setminus y_n)|.$$

Analogously,

$$\begin{aligned} |x_{n-1} = y_{n-1}| \otimes (|x_n = y_n| \otimes |F(x_1, \dots, x_n)|) \\ \leq |x_{n-1} = y_{n-1}| \otimes |F(x_1, \dots, x_{n-1}, x_n \setminus y_n)| \\ \leq |F(x_1, \dots, x_{n-2}, x_{n-1}, y_{n-1}, x_n \setminus y_n)|. \end{aligned}$$

We may proceed by induction to get:

$$|x_1 = y_1| \otimes \dots \otimes |x_n = y_n| \otimes |F(x_1, \dots, x_n)| \leq |F(x_1 \setminus y_1, \dots, x_n \setminus y_n)|,$$

as desired.

If $F \in \bigcap_{i=1}^n A_{x_i, y_i}$ then, since [=7] implies [=6], we have:

$$|x_1 = y_1| \otimes \dots \otimes |x_n = y_n| \otimes |F| \leq (\mathbf{1} \otimes \dots \otimes \mathbf{1}) \otimes |F| = |F|,$$

showing that [=4] is valid and proving Fact 1.

By induction, define a map $|\cdot|$ as follows: for $x, y \in V$, $x \neq y$, set

$$\begin{aligned} |x = y|_0 &= T_{x,y}; \\ |x = y|_{n+1} &= \bigwedge_{z \neq x, y} (|x = z|_n \leftrightarrow |y = z|_n) \\ &\quad (\text{$a \leftrightarrow b$ means $(a \rightarrow b) \wedge (b \rightarrow a)$}); \\ |x = y|_\infty &= !(\bigwedge_{n \in \omega} |x = y|_n); \text{ finally, set } |x = x|_\infty = \mathbf{1}. \end{aligned}$$

Fact 2: $|\cdot|_\infty$ satisfies the properties required for the full logic.

In fact: condition [=1] is clear, while [=3] is verified because it's true for $|x = y|_n$, for all $n \in \omega$.

Since $|x = y|_\infty = !|x = y|_\infty$, [=7] is satisfied (the case $x = y$ is valid too, because $!\mathbf{1} = \mathbf{1}$). Observe that, for every $n \geq 0$ and $z \neq x, y$:

$$\begin{aligned} |x = y|_{n+1} \otimes |x = z|_n &\leq |y = z|_n \text{ and therefore,} \\ |x = y|_\infty \otimes |x = z|_\infty &\leq |x = y|_{n+1} \otimes |x = z|_n \leq |y = z|_n \text{ for every } n \geq 0. \end{aligned}$$

Thus,

$$|x = y|_\infty \otimes |x = z|_\infty \leq \bigwedge_{n \in N} |y = z|_n; \text{ now } [=7] \text{ and } [M4] \text{ yield } [=2].$$

Since $|x = y|_\infty \leq |x = y|_0 = T_{x,y}$, Fact 1 guarantees that $|\cdot|_\infty$ satisfies $[=4]$.

For the (MALL) fragment, it's enough to take $! = !_m$ (see Proposition 1.14) in the above computations. \square

Discussion 3.6 Are requirements $[=5], [=6]$ (resp. $[=7]$) too strong ?

The motivation for them is, starting with substitution for elementary formulas, to have substitution for every formula. Obviously, they are sufficient, but indeed, *they are also necessary*. To see this, note that, if $a = |x = y|$, $b = |P(x)|$ and $c = |P(x \setminus y)|$, then we must have that $a \otimes b \leq c$ and $a \otimes c \leq b$, (i.e. $a \leq (b \leftrightarrow c)$) must imply $a \otimes !b \leq !c$ (and, of course, that $a \otimes !c \leq !b$). Since 1 is a formula, we must also have $a = a \otimes 1 \leq 1$. The critical cases are \otimes and $!$. We have

Lemma 3.7 Let Q be a Girard Quantale and let a be an element of Q such that $a \leq 1$. (a) Are equivalent:

- (i) $\forall b, c, d \in Q, a \leq (b \leftrightarrow c), a \leq (d \leftrightarrow e)$
implies $a \leq ((b \otimes d) \leftrightarrow (c \otimes e))$.
- (ii) $a \leq a \otimes a$

(b) Assume that $! \in M(Q)$ is idempotent and weak. Are equivalent:

- (i) $\forall b, c \in Q, a \leq (b \leftrightarrow c) \text{ implies } a \leq (!b \leftrightarrow !c)$.
- (ii) $a = !a$

(c) $\forall b, c \in Q, a \leq (b \leftrightarrow c) \text{ implies } a \leq (b^\perp \leftrightarrow c^\perp)$.

Proof: (a) (i) \Rightarrow (ii): since $a \leq 1$, then $a \leq (1 \leftrightarrow a)$ and so $a \leq ((1 \otimes 1) \leftrightarrow (a \otimes a))$; thus, $a = a \otimes (1 \otimes 1) \leq a \otimes a$.

(i) \Leftarrow (ii): If $a \leq (b \leftrightarrow c), a \leq (d \leftrightarrow e)$, then we have $a \otimes b \leq c$ and $a \otimes d \leq e$, and so $a \otimes (b \otimes d) \leq (a \otimes a) \otimes (b \otimes d) = (a \otimes b) \otimes (a \otimes d) \leq c \otimes e$. Analogously, $a \otimes (c \otimes e) \leq b \otimes d$.

(b) (i) \Rightarrow (ii): since $a \leq 1$ and $!1 = 1$, then $a \leq (1 \leftrightarrow a)$ and therefore $a \leq (1 \leftrightarrow !a)$; thus, $a = a \otimes 1 \leq !a \leq a$.

(i) \Leftarrow (ii): suppose that $a \leq (b \leftrightarrow c)$; thus, $a \otimes b \leq c$ and then $a \otimes !b \leq a \otimes b \leq c$. Thus, by [M4], we have $a \otimes !b = !a \otimes !b = !(a \otimes !b) = !(a \otimes b) \leq !c$. Analogously, we can prove that $a \otimes !c \leq !b$.

(c) Since Q is commutative, then $(x \rightarrow y) = (y^\perp \rightarrow x^\perp)$, by Proposition 1.9. \square

We can now show that we have substitution for every formula:

Proposition 3.8 *Let $|\cdot|: ELF(\mathbb{L}) \longrightarrow Q$ satisfying $[=1] - [=6]$ ($[=1] - [=4]$, $[=7]$ resp.). Then, the extension to $FOR(\mathbb{L})$ verifies:*

$$|x_1 = y_1| \otimes \cdots \otimes |x_n = y_n| \otimes |\phi(x_1, \dots, x_n)| \leq |\phi(x_1 \{ y_1, \dots, x_n \} y_n)|$$

where $\phi(x_1, \dots, x_n) \in FOR(\mathbb{L})$ has (eventually) free occurrences of variables x_1, \dots, x_n , and $\phi(x_1 \{ y_1, \dots, x_n \} y_n)$ is obtained from $\phi(x_1, \dots, x_n)$ by replacing some occurrences of x_i (not in the scope of a y_i -quantifier) by y_i .

Proof. By induction in the complexity of ϕ . As a first step, we have two cases to consider.

- i) $\phi \in ELF^+(\mathbb{L})$.
It's true by $[=2], [=3], [=4]$ and $[=6]$ (or $[=7]$ and ! weak).
- ii) $\phi \in \{ \mathbf{1}, \perp, \top, \mathbf{0} \}$. It's true by $[=6]$ (or $[=7]$ and ! weak).

To proceed with the induction, assume that substitution holds true for every φ with complexity $\leq k$ and let ϕ be a formula with complexity $k + 1$. We have the following cases:

- a) $\phi = \alpha \otimes \beta$.
It's immediate from $[=5]$ (common to both systems) and lemma 3.7.
- b) $\phi = \alpha^\perp$.
Since $|\alpha^\perp| = |\alpha|^\perp$, the conclusion follows from Lemma 3.7.
- c) $\phi = \wedge x. \alpha(x_1, \dots, x_n, x)$. Given $z \in V$, we have:

$$\begin{aligned} \otimes_{i=1}^n |x_i = y_i| \otimes \wedge_{y \in V} |\alpha(x_1, \dots, x_n, x)[y/x]| \\ \leq \otimes_{i=1}^n |x_i = y_i| \otimes |\alpha(x_1, \dots, x_n, x)[z/x]| \\ \leq |\alpha(x_1 \{ y_1, \dots, x_n \} y_n, x)[z/x]|, \text{ and then} \end{aligned}$$

$$\begin{aligned} \otimes_{i=1}^n |x_i = y_i| \otimes \wedge_{y \in V} |\alpha(x_1, \dots, x_n, x)[y/x]| \\ \leq \wedge_{y \in V} |\alpha(x_1 \{ y_1, \dots, x_n \} y_n, x)[y/x]|. \end{aligned}$$

- d) $\phi = \alpha \& \beta$. Similar to c).
- e) (full logic) $\phi = !\alpha$. It follows from $[=7]$ and Lemma 3.7.

Since the other connectives are defined by duality, the proof is complete. \square

Now, we shall extend the results in [Yet] to the calculus with equality.

Theorem 3.9 (Soundness) *The semantics of quantales for commutative Linear Logic with equality is sound with respect to the given calculus.*

Proof: We prove the validity of the new axioms (for the other rules, consult [Yet]). Define $|\vdash A_1, \dots, A_n| = |A_1 \sqcup \dots \sqcup A_n|$. By observing that, for every $a, b \in Q$, $a \leq b$ iff $a^\perp \sqcup b \geq 1$, the validity of each axiom is guaranteed by [=1]–[=6] (resp. [=1]–[=4], [=7]). \square

Theorem 3.10 (Completeness) *The semantic of quantales for commutative Linear Logic with equality is complete with respect to the given calculus.*

Proof: The proof is an extension of the proofs by [Yet] and [Gir]. Let M_1 be the set of finite sequence of formulas in \mathbb{L} ; M_1 is a monoid with the operation of concatenation (and identity the null sequence). Let M be the (commutative) monoid obtained by identifying sequences which are distinct only by a permutation of their elements. Just as in example 1.11, $\wp(M)$ is a commutative quantale and we set $\perp = \{ \Gamma : \vdash \Gamma \text{ is provable} \} \in \wp(M)$.

Since M is commutative, \perp is cyclic and so we can consider the phase quantale $Q = \wp(M)_j$, where $j : \wp(M) \longrightarrow \wp(M)$ is given by $j(A) = (A \rightarrow \perp) \rightarrow \perp$.

Let $Pr : FOR(\mathbb{L}) \longrightarrow \wp(M)$ defined by:

$$Pr(A) = \{ \Gamma : \vdash A, \Gamma \text{ is provable} \}.$$

By theorem 3.4 in [Yet], Pr factores through Q , i.e., $Pr(FOR(\mathbb{L})) = Q$.

Let $|\cdot| = Pr|_{ELF^+(\mathbb{L})}$; clearly, the unique extension of $|\cdot|$ to $FOR(\mathbb{L})$ is Pr , once we have defined in Q the open, weak and idempotent modality $!$ as:

$$!(x) = \bigvee \{ Pr(!A) : Pr(!A) \leq x \}$$

Fact: $!1 = 1$ (where, by definition, $1 = \bigwedge \{ Pr(A) : \vdash A \text{ is provable} \}$).

To see this, Let $\Gamma \in PR(!1)$; thus, $\vdash !1, \Gamma$ (i.e., $\vdash (? \perp)^\perp, \Gamma$) is provable and let $A \in FOR(\mathbb{L})$ be such that $\vdash A$ is provable. Then:

$$\begin{array}{c} \vdash A \\ \hline \begin{array}{c} \vdash A, \perp \\ \hline \vdash A, ?\perp \end{array} \end{array} \quad \vdash (? \perp)^\perp, \Gamma$$

$$\vdash A, \Gamma$$

i.e., $\Gamma \in Pr(A)$, and so $Pr(!1) \leq \bigwedge \{ Pr(A) : \vdash A \text{ is provable} \} = 1$.

Thus, $Pr(!1) \leq \bigvee \{ Pr(!A) : Pr(!A) \leq !1 \} = !1$. In fact, equality holds because $\vdash !1$ is provable, and so $1 \leq Pr(!1) \leq !1 \leq 1$, i.e. $1 = !1$.

We shall now show that $|\cdot|$ satisfies the required properties. Since $\vdash (x = x)$ is provable, $1 \leq Pr((x = x)) = |x = x|$. This verifies [=1].

To prove [=2], let $\Gamma \Delta \in |x = y| \cdot |y = z|$. Then we have:

$$\vdash \frac{\vdash (x = y), \Gamma \quad \vdash (x = y)^\perp, (y = z)^\perp, (x = z)}{\vdash (y = z)^\perp, (x = z), \Gamma \quad \vdash (y = z), \Delta} \vdash (x = z), \Gamma, \Delta$$

i.e., $\Gamma \Delta \in |x = z|$. Thus, $|x = y| \cdot |y = z| \leq |x = z|$ and so

$$|x = y| \otimes |y = z| = (|x = y| \cdot |y = z|)^\perp \perp \leq |x = z|.$$

To prove [=3], let $\Gamma \in |x = y|$. We have:

$$\vdash \frac{\vdash (x = y), \Gamma \quad \vdash (x = y)^\perp, (y = x)}{\vdash (y = x), \Gamma}$$

i.e., $\Gamma \in |y = x|$ and thus $|x = y| \leq |y = x|$

Properties [=4], [=5], [=6], [=7] are verified in a similar way. This shows that we have an interpretation of quantales such that $|A| \geq 1$ iff $\vdash A$ is provable, completing the proof. \square

4. Generalisations of the Calculus

There are alternatives to the treatment of equality given above, using the exponentials of Linear Logic. For example, instead of requiring $(\cdot = \cdot)$ to be open, we could establish that its characteristic properties be valid in the interior of $(\cdot = \cdot)$. Thus, we define the calculus (P_1) by the axioms:

$$\begin{array}{ll} [R!] & \vdash !(x = x) \\ [T!] & \vdash !(x = y) \otimes !(y = z) \end{array} \quad \begin{array}{ll} [S!] & \vdash !(x = y) \multimap !(y = x) \\ & \vdash !(x = z) \multimap !(x = y) \end{array}$$

For every predicate symbol P ,

$$[SUBS!] \vdash !(x_1 = y_1) \otimes \cdots \otimes !(x_n = y_n) \otimes P(x_1, \dots, x_n) \longrightarrow P(x_1 \{ y_1, \dots, x_n \} y_n)$$

Clearly, we have substitution for every formula, and we can say that $!(x = y)$ has the behaviour of a linear equality. An even weaker form of substitution can be defined as follows:

$$\vdash !(x_1 = y_1) \otimes \cdots \otimes !(x_n = y_n) \otimes !P(x_1, \dots, x_n) \longrightarrow !P(x_1 \{ y_1, \dots, x_n \} y_n)$$

or, even

$$\vdash !(x_1 = y_1) \otimes \cdots \otimes !(x_n = y_n) \otimes !P(x_1, \dots, x_n) \longrightarrow ?P(x_1 \{ y_1, \dots, x_n \} y_n).$$

We can modify each axiom combining the modal operators in all possible ways. It is straightforward to verify that (incorporating id , the identity operation on formulas), there are only seven modal operations obtained by successive applications of $\{!, ?\}$, namely, the set




fig. 1

with its natural lattice structure (see figure 1). This is in complete analogy with Kuratowski's problem, in which we can prove that, given a topological space X and $A \subseteq X$, there are only 14 distinct sets that can be obtained from A by combinations of taking complements and closure (in our case, we do not consider modalities of the form $m'(x) = m(x)^\perp$ with $m \in \mathcal{M}$). Starting with this, we can define a general scheme to create a linear calculus with equality.

Definition 4.1 Let $(m_i)_{i=1}^{10}$ be a sequence in \mathcal{M} . A linear calculus with equality $P((m_i))$ is defined by the following axioms:

$$\begin{array}{ll} [R] & \vdash m_1(x = x) & [S] & \vdash m_2(x = y) \multimap m_3(y = x) \\ [T] & \vdash m_4(m_5(x = y) \otimes m_5(y = z)) \multimap m_6(x = z) \end{array}$$

For each predicate symbol P , the axiom

$$[SUBS] \quad \vdash m_7((\otimes_{i=1}^n m_8(x_i = y_i)) \otimes m_9(P(x_1, \dots, x_n))) \multimap m_{10}(P(x_1 \setminus y_1, \dots, x_n \setminus y_n)).$$

Remarks 4.2 (a) If $m_3 \leq m_2$, then [S] implies that, for $m_3 \leq r, s \leq m_2$, $r(x = y)$, $s(x = y)$, $r(y = x)$ and $s(y = x)$ are all equivalent.

(b) If $m_1 = id$ or $m_1 = !$, then $\vdash m(x = x)$ is provable for every $m \in \mathcal{M}$.

(c) If $m_7 = id$, $m_8 \in \{ !, !?, !?!\}$ and $m_9 \geq m_{10}$, then substitution holds for every formula ϕ built up from formulas of the form $m_9(P)$, where $P \in ELF(\mathbb{L})$.

(d) For each $m_i \in \{ !, !?, !?!\}$, we have $m(\otimes_{i=1}^n m_i(A_i)) = m!(\otimes_{i=1}^n m_i(A_i))$.

Examples 4.3 (a) The system (LLE_1) of section 2 is obtained by the assignment

$$m_3 = ! \text{ and } m_i = id, \text{ for every } i \neq 3.$$

(b) If we set

$$m_1 = m_2 = m_3 = m_5 = m_6 = m_8 = ! \text{ and } m_4 = m_7 = m_9 = m_{10} = id,$$

we get the system (P_1) , in which "equality" is the interior of $(\cdot = \cdot)$. This system satisfies substitution for every formula.

(c) If we set

$$m_1 = m_2 = m_5 = m_6 = m_8 = ?, \quad m_4 = m_7 = m_9 = m_{10} = id,$$

and

$$m_3 = !?$$

then we have a system in which "equality" corresponds to the closure of $(\cdot = \cdot)$; from [S] will follow that that $?(\cdot = \cdot)$ is "clopen". This system satisfies the substitution rule for every formula.

(d) By considering

$$m_1 = m_3 = m_6 = m_{10} = ?!, \quad m_2 = m_5 = m_8 = m_9 = !?!$$

and

$$m_4 = m_7 = !? \text{ or } !?!$$

we get a system containing a translation of the classical theory of equality inside (LL), to be studied in next section.

(e) Another translation of classical equality in (LL) can be obtained by considering the assignment

$$m_1 = id, \quad m_2 = m_4 = m_5 = m_7 = m_8 = m_9 = ?$$

and

$$m_3 = m_6 = m_{10} = !.$$

Now, we would like to identify the classes of equivalent calculi. Instead of doing a complete classification of the possible calculi, we present quantale theoretic techniques that are useful in deciding this problem. We start with reflexivity ([R]):

Proposition 4.4 Let Q be a Girard's quantale, and let $x \in Q$. Then:

(a) Are equivalent:

$$(i) \quad ?x \geq \mathbf{1} \quad (ii) \quad !?x \geq \mathbf{1} \quad (iii) \quad ?!?x \geq \mathbf{1}$$

;

(b) *Are equivalent:*

$$(i) \ x \geq 1 \quad (ii) \ !x \geq 1$$

(c) *Are equivalent:*

$$(i) \ ?!x \geq 1 \quad (ii) \ !?x \geq 1$$

Proof: (a): If $?x \geq 1$, then $?x \geq !1 = 1$. If $!x \geq 1$, then $?x \geq ?1 \geq 1$. Finally, $?!x \geq 1$ implies $?x \geq 1$, because $? \geq ?!$.

(b) comes directly from $!1 = 1$, while (c) is consequence of (b). \square

Thus, introducing the notation $r \equiv s$ to denote that calculi obtained with $m_1 = r$ and $m_1 = s$ are equivalent, we have:

$$(a) \ ? \equiv !? \equiv ?! \quad (b) \ id \equiv ! \quad (c) \ ?! \equiv !?$$

and so there are only three non equivalent possibilities for $[R]$.

With respect to axiom $[S]$, consider $\mathcal{M}_1 = \{?, ?, ?, ?\} = \{?m: m \in \mathcal{M}\}$, together with $\mathcal{M}_2 = \{!, ?, ?, ?\} = \{!m: m \in \mathcal{M}\}$. Write $(r,s) \equiv (r',s')$ to denote that calculi obtained with $m_2 = r$, $m_3 = s$ and $m_2 = r'$, $m_3 = s'$ are equivalent. Then we have:

Proposition 4.5 (a) *If $m_3 \in \mathcal{M}_1$, then:*

$$(i) \ (!,m_3) \equiv (?!,m_3) \equiv (!?!,m_3) \quad (ii) \ (!?,m_3) \equiv (?!,m_3) \\ (iii) \ (id,m_3) \equiv (?!,m_3)$$

(b) *If $m_2 \in \mathcal{M}_2$, then:*

$$(i) \ (m_2,?) \equiv (m_2,!?) \equiv (m_2,?!?) \quad (ii) \ (m_2,?) \equiv (m_2,!?) \\ (iii) \ (m_2,id) \equiv (m_2,?)$$

Therefore, there are only 26 non equivalent cases for $[S]$:

- (1): $(!,!?) \equiv (!,?!?) \equiv (!,?) \equiv (!?!,?!?) \equiv (?!,?!?) \equiv (?!,!?) \equiv (!?!,!) \equiv (!?!,?) \equiv (?!,?)$
- (2): $(!?,!?) \equiv (!?,?!?) \equiv (!?,?) \equiv (?!,?!?) \equiv (?!,?)$
- (3): $(!,?) \equiv (!?!,?) \equiv (?!,?) \equiv (!,!) \equiv (!?!,!?)$
- (4): $(id,?) \equiv (?,?) \quad (5): \ (!?,?) \equiv (?!,?) \equiv (!?,!?)$
- (6): $(id,?) \equiv (?!,?) \quad (7): \ (id,?!?) \equiv (?!,?) \quad (8): \ (!,id) \equiv (!,!?)$
- (9): $(!,!) \equiv (!?,id) \quad (10): \ (!?!,!) \equiv (?!,id) \quad (11): \ (id,id) \equiv (!,!?)$
- (12): $(id,!?) \quad (13): \ (id,?) \quad (14): \ (id,?!?) \quad (15): \ (?!,id)$

(16): $(?,!?)$ (17): $(?,!?)$ (18): $(?,!)$ (19): $(?!,id)$
 (20): $(?!,!?)$ (21): $(?!,!?)$ (22): $(?!,!)$ (23): $(?!,id)$
 (24): $(?!,!?)$ (25): $(?!,!?)$ (26): $(?!,!)$

For transitivity ([T]), we shall write $(r,s,t) \equiv (r',s',t')$ when calculi obtained with $m_4 = r$, $m_5 = s$, $m_6 = t$ and $m_4 = r'$, $m_5 = s'$, $m_6 = t'$ are equivalent. Then we have:

Proposition 4.6 If $m_5 \in \mathcal{M}_2$, then:

- (i) If $m_6 \in \mathcal{M}_1$, then $(r,m_5,m_6) \equiv (s,m_5,m_6)$ for every $r,s \in \mathcal{M}$
- (ii) $(r,m_5,?) \equiv (s,m_5,?) \equiv (id,m_5,?) \equiv (!,m_5,?) \equiv (?!,m_5,?) \equiv (!?,m_5,?) \equiv (!?,m_5,!) \equiv (t,m_5,!) \equiv (u,m_5,!)$ for every $r,s,t,u \in \mathcal{M}$.
- (iii) $(r,m_5,!) \equiv (s,m_5,!) \equiv (id,m_5,!) \equiv (!,m_5,!) \equiv (?!,m_5,!) \equiv (!?,m_5,!) \equiv (!?,m_5,!)$ for every $r,s \in \mathcal{M}$.

The results above imply that, for $m_5 \in \mathcal{M}_2$ fixed, we have only 8 possibilities for pairs (m_4, m_6) (in contrast with 26 possible ones), where the numbers refer to the pairs described above.

[1]: (1) \equiv (2) \equiv (4) \equiv (7) \equiv (13)	[2]: (3) \equiv (5) \equiv (6) \equiv (14)
[3]: (8) \equiv (11) \equiv (12)	[4]: (9) \equiv (10)
[5]: (15) \equiv (19) \equiv (23)	[6]: (16) \equiv (20) \equiv (24)
[7]: (17) \equiv (21) \equiv (25)	[8]: (18) \equiv (22) \equiv (26)

With respect to substitution, if we fix $m_8, m_9 \in \mathcal{M}_2$, then there are again, for pairs (m_7, m_{10}) , only the 8 cases above. Thus, these are the non equivalent calculi that can be constructed with the exponentials, satisfying the usual rules of equality. We register that this corresponds to only 8% of the original universe of possible calculi.

5. Interpreting Classical Equality in Linear Logic with Equality

It is well known that the exponentials of linear logic (LL) are important in interpreting intuitionistic logic (IL) and classical logic (CL) inside LL. For each of these logics, we have two translations, one of them based in the fact that every (commutative) Girard's quantale contains a frame (complete Heyting algebra) (corollary 1.21) and a complete Boolean algebra:

$$\mathcal{H} = \{ !x : x \in Q \} \quad \text{and} \quad \mathcal{B} = \{ ?!x : x \in Q \},$$

respectively. Operations and constants in each of these algebras are given by:

(H). *For the frame \mathcal{H} :*

$$\begin{aligned} \mathbf{0}_{\mathcal{H}} &= \mathbf{0}, \top_{\mathcal{H}} = \mathbf{1}, \quad (x \wedge_{\mathcal{H}} y) = (x \otimes y) = !(x \wedge y), \quad (x \vee_{\mathcal{H}} y) = (x \vee y); \\ (x \Rightarrow y) &= !(x \rightarrow y), \quad \neg x = (x \Rightarrow \mathbf{0}_{\mathcal{H}}) = !(x \rightarrow \mathbf{0}), \\ \wedge_{\mathcal{H}} S &= !\wedge S, \quad \vee_{\mathcal{H}} S = \vee S. \end{aligned}$$

(B). *For the complete Boolean algebra \mathcal{B} :*

$$\begin{aligned} \mathbf{0}_{\mathcal{B}} &= \perp, \top_{\mathcal{B}} = ?\mathbf{1}, \quad (x \wedge_{\mathcal{B}} y) = ?(\mathbf{!}x \otimes \mathbf{!}y) = ?!(x \wedge y), \\ (x \vee_{\mathcal{B}} y) &= (x \sqcup y) = (x^\perp \otimes y^\perp)^\perp; \\ (x \Rightarrow y) &= (\mathbf{!}x \rightarrow y), \quad \neg x = (x \Rightarrow \mathbf{0}_{\mathcal{B}}) = (\mathbf{!}x \rightarrow \perp) = ?(x^\perp); \\ \wedge_{\mathcal{B}} S &= ?!\wedge S, \quad \vee_{\mathcal{B}} S = ?\vee \{ !x : x \in S \}. \end{aligned}$$

Thus, we can interpret $A \in FOR(IL)$ as $A^i \in FOR(\mathbb{L})$ by the rules:

$A^i = \mathbf{!}A$ if A is atomic; proceed by induction on complexity using the operations in (H) to define A^i for all intuitionistic formulas (here, $FOR(IL)$ denote the set of intuitionistic formulas).

For classical logic, we have our first translation:

(*) $A^c = ?\mathbf{!}A$ if A is atomic; then proceed by induction on complexity, using the rules in (B).

Thus, if we assume two-hand sequents for (LL) , we have ([Gir]):

$A_1, \dots, A_n \vdash_{IL} A$ is provable in intuitionistic logic
iff $(A_1)^i, \dots, (A_n)^i \vdash A^i$ is provable in LL ,

and

(1st T) : $A_1, \dots, A_n \vdash_{CL} A$ is provable in classical logic iff $!(A_1)^c, \dots, !(A_n)^c \vdash A^c$ is provable in LL .

Another interpretation for classical logic is constructed from polarities for formulas. Thus, given a sequent for classical logic $\Gamma \vdash_{CL} \Delta$, we'll say that occurrences of formulas $A \in \Gamma$ are *positive* (denoted by pA) and occurrences of formulas $B \in \Delta$ are *negative* (denoted by nB). We have the following rules of a second translation (**):

$$\begin{aligned} pA &= A = nA \text{ if } A \text{ is atomic.} \\ p(\neg A) &= (nA)^\perp, \quad n(\neg A) = (pA)^\perp, \\ p(A \vee B) &= p(A) \oplus p(B), \quad n(A \vee B) = \mathbf{!}n(A) \sqcup \mathbf{!}n(B), \\ p(A \wedge B) &= ?p(A) \otimes ?p(B), \quad n(A \wedge B) = n(A) \& n(B), \\ p(A \Rightarrow B) &= n(A)^\perp \oplus p(B), \quad n(A \Rightarrow B) = ?p(A) \multimap \mathbf{!}n(B), \end{aligned}$$

$$p((\forall x)A) = \wedge x.?p(A), n((\forall x)A) = \wedge x.n(A), \\ p((\exists x)A) = \vee x.p(A), n((\exists x)A) = \vee x.!n(A).$$

With this definition, we have ([Gir]):

$$(\ast\ast) \quad A_1, \dots, A_k \vdash_{CL} A \text{ is provable in } (CL) \text{ iff } !n(A_1), \dots, !n(A_k) \vdash ?p(A) \text{ is provable in } (LL).$$

We now turn to the question of determining a linear theory of equality such that, given a classical theory of, its translation into linear logic is defines a linear theory contained in the original linear theory of equality.

For this, we need to relate the deduction of a sequent in (LL) from a set of sequent-axioms and the deduction of a formula from a multiset of hypothesis (in the left-hand side of the sequent), called *external* and *internal* relations of consequence (respectively) by [Avr].

Definition 5.1 Let $A \in FOR(\mathbb{L})$ with exactly x_1, \dots, x_n as free variables. The universal closure of A , denoted by $\wedge A$ is the formula obtained from A by quantifying universally all the free variables of A , i.e., $\wedge A = \wedge x_1. \dots \wedge x_n.A$. Analogously, we define $\vee A$, the existential closure of A .

It is straightforward to prove next result:

Proposition 5.2 Let A_1, \dots, A_n be formulas, $B = !\wedge A_1 \& \dots \& !\wedge A_n$, and Δ a multiset of formulas. Are equivalent:

- (i) $\vdash \Delta$ is provable from the sequent-axioms $\vdash A_1, \dots, \vdash A_n$;
- (ii) there exists $k \geq 0$ such that $\overbrace{B, \dots, B}^{k \text{ times}} \vdash \Delta$ is provable;
- (iii) $!\wedge A_1, \dots, !\wedge A_n \vdash \Delta$ is provable.

Now, assume that \mathcal{A} is the set of axioms (without free variables) of a theory of (CL) in a language without functional symbols. It follows from Proposition 5.2 that if $A_1, \dots, A_n \in \mathcal{A}$ and A is a formula, then the following are equivalent, for the first translation (\ast) of CL into LL :

- (i) $A_1, \dots, A_n \vdash_{CL} A$;
- (ii) $!(A_1)^c, \dots, !(A_n)^c \vdash A^c$ is provable in (LL) ;
- (iii) There is a proof of $\vdash A^c$ from the axioms $\vdash (A_1)^c, \dots, \vdash (A_n)^c$ (recall that the A_i 's have no free variables).

Thus, to each axiom $A \in \mathcal{A}$ of a classical theory, corresponds a sequent-axiom $\vdash A^c$ in (LL) . Similarly, for the second translation $(\ast\ast)$, it can be seen that to each axiom $A \in \mathcal{A}$, there corresponds a sequent-axiom $\vdash n(A)$.

Now assume that \mathcal{A} defines the classical theory of equality. Thus, \mathcal{A} consists in the following axioms:

Now assume that \mathcal{A} defines the classical theory of equality. Thus, \mathcal{A} consists in the following axioms:

$$\begin{aligned}
 & (\forall x)(x = x); \\
 & (\forall x)(\forall y)((x = y) \Rightarrow (y = x)); \\
 & (\forall x)(\forall y)(\forall z)((x = y) \wedge (y = z) \Rightarrow (x = z)); \\
 & (\forall x_1)(\forall y_1) \dots (\forall x_n)(\forall y_n)((x_1 = y_1) \wedge \dots \wedge (x_n = y_n) \wedge P(x_1, \dots, x_n) \\
 & \qquad \qquad \qquad \Rightarrow P(x_1 \setminus y_1, \dots, x_n \setminus y_n))
 \end{aligned}$$

(where P varies over every predicate symbol of arity n of the language).

For the first translation (*), we get a calculus (E_1) with the following axioms:

$$\begin{aligned}
 [R1] \quad & \vdash ?! \wedge x ?!(x = x) \quad [S1] \quad \vdash ?! \wedge x ?! \wedge y (!?!(x = y) \multimap ?!(y = x)) \\
 [T1] \quad & \vdash ?! \wedge x ?! \wedge y ?! \wedge z (!?(!?!(x = y) \otimes !?!(y = z)) \multimap ?!(x = z)) \\
 [SUBST1] \quad & \text{If } P \text{ is a } n\text{-ary predicate symbol of the language} \\
 & \vdash ?! \wedge x_1 ?! \wedge y_1 \dots ?! \wedge x_n ?! \wedge y_n (!?(!?!(x_1 = y_1) \otimes \dots \otimes \\
 & \qquad \qquad \qquad !?!(x_n = y_n) \otimes !?P(x_1, \dots, x_n)) \multimap ?!P(x_1 \setminus y_1, \dots, x_n \setminus y_n))
 \end{aligned}$$

For the second translation (**), we get a calculus (E_2) with the following axioms:

$$\begin{aligned}
 [R2] \quad & \vdash \wedge x (x = x) \quad [S2] \quad \vdash \wedge x \wedge y (?!(x = y) \multimap !(y = x)) \\
 [T2] \quad & \vdash \wedge x \wedge y \wedge z (?!(?!(x = y) \otimes ?!(y = z)) \multimap !(x = z)) \\
 [SUBST2] \quad & \text{If } P \text{ is a } n\text{-ary predicate symbol in the language} \\
 & \vdash \wedge x_1 \wedge y_1 \dots \wedge x_n \wedge y_n (?!(?!(x_1 = y_1) \otimes \dots \otimes \\
 & \qquad \qquad \qquad ?!(x_n = y_n) \otimes ?P(x_1, \dots, x_n)) \multimap !P(x_1 \setminus y_1, \dots, x_n \setminus y_n)).
 \end{aligned}$$

Consider the linear theory (P_1) determined by the axioms:

$$\begin{aligned}
 [LR1] \quad & \vdash ?!(x = x) \quad [LS1] \quad \vdash ?!(x = y) \multimap ?!(y = x) \\
 [LT1] \quad & \vdash !?!(x = y) \otimes !?!(y = z) \multimap ?!(x = z) \\
 [LSUBST1] \quad & \vdash !?!(x_1 = y_1) \otimes \dots \otimes !?!(x_n = y_n) \otimes !?P(x_1, \dots, x_n) \\
 & \multimap ?!P(x_1 \setminus y_1, \dots, x_n \setminus y_n).
 \end{aligned}$$

Clearly, (P_1) contains (E_1) (because every axiom in (E_1) is deducible in (P_1) , using the equivalences of last section) and we have that (P_1) defines a linear theory of equality in the sense of section 4.

Similarly, define the theory (P_2) by the axioms:

$$\begin{aligned}
 [LR2] \quad & \vdash (x = x) \quad [LS2] \quad \vdash ?(x = y) \multimap ! (y = x) \\
 [LT2] \quad & \vdash ?(?(x = y) \otimes ?(y = z)) \multimap ! (x = z) \\
 [LSUBST2] \quad & \vdash ?(?(x_1 = y_1) \otimes \cdots \otimes ?(x_n = y_n)) \otimes ?P(x_1, \dots, x_n) \\
 & \multimap !P(x_1 \{ y_1, \dots, x_n \} y_n).
 \end{aligned}$$

We can reformulate (P_2) as:

$$\begin{aligned}
 [LR2'] \quad & \vdash (x = x) \quad [LS2'] \quad \vdash (x = y) \multimap (y = x) \\
 [LT2'] \quad & \vdash (x = y) \otimes (y = z) \multimap (x = z) \\
 [LSUBST2'] \quad & \vdash ?((x_1 = y_1) \otimes \cdots \otimes (x_n = y_n)) \otimes ?P(x_1, \dots, x_n) \\
 & \multimap !P(x_1 \{ y_1, \dots, x_n \} y_n) \\
 [? \leq !] \quad & \vdash ?(x = y) \multimap !(x = y).
 \end{aligned}$$

It is straightforward to check that (P_2) and (E_2) are equivalents. In fact, (P_2) is the strongest (and therefore more restricted as far as semantics is concerned) of the calculi already defined.

6. Equality and intuitionistic Linear Logic

In this section we shall study how to recover classical logic and intuitionistic logic from a linear calculus without exponentials, as well as analyse another extensions of the relationship between linear logic and quantales. Our basic system is (commutative) first order intuitionistic linear logic, from now on denoted by (LLI) . The appropriate semantics will turn out to be furnished by commutative unital quantales. Similarly, the semantics for non-commutative first order intuitionistic linear shall be proven to be given by unital quantales. From this, it will be seen that, adding appropriate axioms, we can recover intuitionistic logic, classical logic and classical linear logic. Thus, an intuitionistic linear theory of equality provides an intuitionistic theory of equality, a classical theory of equality, and a classical linear theory of equality, simultaneously. The most natural candidate is system (LLE_0) , the simplest already defined.

We define a sequent calculus for (commutative) first-order intuitionistic linear logic without negation, simply by extending the system in [GiLa] and then proving soundness and completeness for this system.

Definition 6.1 The language \mathbb{L}_i for commutative first-order intuitionistic linear logic consist of a countable set of predicate symbols, $\wp = \{ P_n : n \in \omega \}$, a countable set of variables $V = \{ v_n : n \in \omega \}$, the symbols $\otimes, \multimap, \&, \oplus, \vee, \wedge$ and the same rules as in Definition 2.2 for the formation of the set of formulas $FOR(\mathbb{L}_i)$. For $A \in FOR(\mathbb{L}_i)$ and $x, y \in V$, $A[y/x]$ is as in Definition 2.4.

Definition 6.2 The calculus (LLI) for commutative first order intuitionistic linear logic consist in the following rules and axioms (A, B, C denote formulas, and Γ, Δ denote multisets of formulas):

$$\begin{array}{ll}
 \text{[AX1]} \quad A \vdash A & \text{[AX2]} \quad \Gamma \vdash \top \\
 \text{[AX3]} \quad \vdash \mathbf{1} & \text{[AX4]} \quad \Gamma, \mathbf{0} \vdash A \\
 \\
 \text{[CUT]} \quad \frac{\Gamma \vdash A \quad A, \Delta \vdash B}{\Gamma, \Delta \vdash B} & \text{[EXCH]} \quad \frac{\Gamma, A, B, \Delta \vdash C}{\Gamma, B, A, \Delta \vdash C} \\
 \\
 \text{[1L]} \quad \frac{\Gamma \vdash A}{\mathbf{1}, \Gamma \vdash A} & \text{[⊗ R]} \quad \frac{\Gamma \vdash A \quad \Delta \vdash B}{\Gamma, \Delta \vdash A \otimes B} \quad \text{[⊗ L]} \quad \frac{\Gamma, A, B \vdash C}{\Gamma, A \otimes B \vdash C} \\
 \\
 \text{[& R]} \quad \frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \& B} & \text{[& 1]} \quad \frac{\Gamma, A \vdash C}{\Gamma, A \& B \vdash C} \quad \text{[& 2]} \quad \frac{\Gamma, B \vdash C}{\Gamma, A \& B \vdash C} \\
 \\
 \text{[⊕ L]} \quad \frac{\Gamma, A \vdash C \quad \Gamma, B \vdash C}{\Gamma, A \oplus B \vdash C} & \text{[⊕ 1]} \quad \frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B} \quad \text{[⊕ 2]} \quad \frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B} \\
 \\
 \text{[→ R]} \quad \frac{\Gamma, A \vdash B}{\Gamma \vdash A \multimap B} & \text{[→ L]} \quad \frac{\Gamma \vdash A \quad \Delta, B \vdash C}{\Gamma, \Delta, A \multimap B \vdash C} \\
 \\
 \text{[∨ R]} \quad \frac{\Gamma \vdash A[y/x]}{\Gamma \vdash \bigvee x.A} & \text{[∨ L]} \quad \frac{\Gamma, A \vdash B}{\Gamma, \bigvee x.A \vdash B} \text{ if } x \text{ not occurs free in } \Gamma, B \\
 \\
 \text{[∧ L]} \quad \frac{\Gamma, A[y/x] \vdash B}{\Gamma, \bigwedge x.A \vdash B} & \text{[∧ R]} \quad \frac{\Gamma \vdash A}{\Gamma \vdash \bigwedge x.A} \text{ if } x \text{ not occurs free in } \Gamma
 \end{array}$$

Definition 6.3 An interpretation of quantales for (LLI) is a commutative unital quantale $(Q, \vee, *, \mathbf{1})$ and a map $|\cdot|_Q : FOR(\mathbb{L}_i) \longrightarrow Q$ satisfying:

1. $|\mathbf{0}| = \mathbf{0}$, $|\mathbf{1}| = \mathbf{1}$, $|\top| = \top$.
2. $|A \otimes B| = |A| * |B|$, $|A \multimap B| = |A| \rightarrow |B|$.
3. $|A \& B| = |A| \wedge |B|$, $|A \oplus B| = |A| \vee |B|$.
4. $|\bigwedge x.A| = \bigwedge y \in VAR(\mathbb{L}_i) |A[y/x]|$,
 $|\bigvee x.A| = \bigvee y \in VAR(\mathbb{L}_i) |A[y/x]|$.

We say that $A \in FOR(\mathbb{L}_i)$ is valid in Q if $|A|_Q \geq \mathbf{1}$.

A sequent $\Gamma \vdash A$ is valid in Q if $|\Gamma|_Q \leq |A|_Q$, where

$$|\Gamma|_Q = \begin{cases} \mathbf{1} & \text{if } \Gamma = \emptyset \\ \bigotimes_{i=1}^n |A_i|_Q & \text{if } \Gamma = A_1, \dots, A_n \end{cases}$$

Before stating and proving soundness we establish the following simple

Lemma 6.4 Let Q be a quantale. Then, $a \otimes \mathbf{0} = \mathbf{0} \otimes a = \mathbf{0}$ for every $a \in Q$.

Proof. $\mathbf{0} \leq a \rightarrow, \mathbf{0}$ implies $a \otimes \mathbf{0} \leq \mathbf{0}$; $\mathbf{0} \leq a \rightarrow, \mathbf{0}$ implies $\mathbf{0} \otimes a \leq \mathbf{0}$. \square

Theorem 6.5 (Soundness) If $\Gamma \vdash A$ is provable in (LLI), then it is valid in every interpretation of quantales, i.e., $|\Gamma|_Q \leq |A|_Q$ for every unital quantale Q .

Proof. By induction on least length n of a proof of $\Gamma \vdash A$.

Let Q be a quantale. If $n = 1$, [AX1],[AX2] and [AX3] are immediate, while [AX4] is a consequence of Lemma 6.4.

Assume the thesis holds for all sequents with a proof of length $\leq n$ ($n \geq 1$ fixed) and let $\Lambda \vdash D$ be a sequent admitting a proof of minimum length $n + 1$. We discuss the last rule applied in the proof of $\Lambda \vdash D$:

[CUT]: We have $|\Gamma| \leq |A|$ and $|A| * |\Delta| \leq |B|$ by the induction hypothesis; thus, $|\Gamma| * |\Delta| \leq |A| * |\Delta| \leq |B|$.

The passage through the rules [EXCH] and [1L] follow from the fact that Q is commutative and that $\mathbf{1}$ is the unit of Q .

$[\otimes R]$: $|\Gamma| \leq |A|$ and $|\Delta| \leq |B|$ (induction hypothesis) yield, then $|\Gamma| * |\Delta| \leq |A| * |B|$.

$[\otimes L]$: This works by definition of interpretation.

$[\& R]$: By induction, $|\Gamma| \leq |A|$ and $|\Gamma| \leq |B|$, and so $|\Gamma| \leq |A| \wedge |B|$.

$[\& 1]$ and $[\& 2]$: We have $|\Gamma| * |A| \leq |C|$ (induction hypothesis); thus

$$|\Gamma| * (|A| \wedge |B|) \leq |\Gamma| * |A| \leq |C|.$$

The other rule is similar.

$[\oplus L],[\oplus 1],[\oplus 2]$: Same argument as in $\&$, using \vee in place of \wedge .

$[\multimap R]$: Induction yields $|\Gamma| * |A| \leq |B|$; thus, by adjointness $|\Gamma| \leq |A| \rightarrow |B|$.

$[\multimap L]$: The induction hypothesis yields $|\Gamma| \leq |A|$ and $|\Delta| * |B| \leq |C|$. Thus,

$$(|A| \rightarrow |B|) * |\Gamma| * |\Delta| \leq (|A| \rightarrow |B|) * |A| * |\Delta| \leq |B| * |\Delta| \leq |C|.$$

$[\vee R]$: By induction, there is a variable y such that $|\Gamma| \leq |A[y/x]|$; but then

$$|\Gamma| \leq |A[y/x]| \leq \bigvee_y |A[y/x]|.$$

We now state a Fact whose proof is routine:

Fact: If $\Gamma, A \vdash B$ is provable in n steps and x is not free in either Γ or B , then $\Gamma, A[y/x] \vdash B$ is provable in n steps for every $y \in V$.

[$\vee L$]: Suppose that last rule applied was [$\vee L$]. Since x is not free in Γ, B , by induction and the Fact above, we may assume that $|\Gamma| * |A[y/x]| \leq |B|$, for every $y \in V$.

Thus, $|A[y/x]| \leq |\Gamma| \rightarrow |B|$ for every $y \in V$ and so $|\vee x A| = \vee_y |A[y/x]| \leq |\Gamma| \rightarrow |B|$, i.e. $|\Gamma| * |\vee x A| \leq |B|$.

[$\wedge L$]: This is treated just as [$\vee R$], above.

[$\wedge R$]: Suppose that last rule applied was [$\wedge R$]. Since x is not free in Γ , induction and the Fact above allow us to assume that $|\Gamma| \leq |A[y/x]|$, for every $y \in V$. It is then clear that $|\Gamma| \leq \wedge_y |A[y/x]|$, ending the proof. \square

Before we prove completeness, we need a general result about quantales (generalizing an analogous result about Heyting algebras in [Mir]).

Definition 6.6 An autonomous poset is a partially ordered set P with a binary associative operation \otimes such that the endomorphisms $a \otimes \cdot$ and $\cdot \otimes a$ have right adjoints, denoted by $a \rightarrow_r \cdot$ and $a \rightarrow_l \cdot$, respectively.

A [\otimes, \vee]-autonomous lattice L is an autonomous poset where L is a lattice and such that \otimes is distributive for all \vee 's which exists in L , i.e., if $S \subseteq L$ such that exists $\vee S$ in L , then, for every $a \in L$, $\vee s \in S (a \otimes s)$ and $\vee s \in S (s \otimes a)$ exists in L , and we have that $\vee s \in S (a \otimes s) = a \otimes (\vee S)$, $\vee s \in S (s \otimes a) = (\vee S) \otimes a$.

If S, T are subsets of L , define

$$S \cdot T = \{a \otimes b : a \in S \text{ and } b \in T\}.$$

Definition 6.7 Let L be a lattice and $I \subseteq L$ an ideal (i.e., if $x \in I$ and $y \leq x$ then $y \in I$; if $x, y \in I$ then $x \vee y \in I$). We say that I is complete if it satisfies:

if $S \subseteq I$ such that $\vee_L S$ exists, then $\vee_L S \in I$.

Since an arbitrary meet of complete ideals is again a complete ideal,

$CI(L) = \{I \subseteq L : I \text{ is a complete ideal in } L\}$, is a complete lattice ordered by inclusion, and containing, for each $a \in L$,

$$a^\leftarrow = \{x \in L : x \leq a\}.$$

If $a \in L$ and $S \subseteq L$ is a subset of L , define

$$a \rightarrow_r S \equiv_{def} \bigcup_{c \in S} (a \rightarrow_r c)^\leftarrow \quad \text{and} \quad a \rightarrow_l S \equiv_{def} \bigcup_{c \in S} (a \rightarrow_l c)^\leftarrow.$$

Lemma 6.8 Let L be a $[\otimes, \vee]$ -autonomous lattice. Then

- (a) If $a \in L$ and $I \in CI(L)$, then $a \rightarrow_r I$ and $a \rightarrow_l I$ are in $CI(L)$.
- (b) For every $S, T \subseteq L$ and $K \in CI(L)$, we have

$$S \cdot T \subseteq K \text{ iff } S \subseteq \bigcap_{a \in T} (a \rightarrow_l K) \text{ iff } T \subseteq \bigcap_{a \in S} (a \rightarrow_r K).$$

Proof. (a) If $x \in a \rightarrow_r I$, then there is $c \in I$ such that $x \leq a \rightarrow_r c$; thus, if $y \leq x$ then $y \leq a \rightarrow_r c$ and therefore $y \in a \rightarrow_r I$.

If $x, y \in a \rightarrow_r I$, then there are $c, d \in I$ such that $x \leq a \rightarrow_r c$, and $y \leq a \rightarrow_r d$; therefore

$$x \vee y \leq (a \rightarrow_r c) \vee (a \rightarrow_r d) \leq a \rightarrow_r (c \vee d),$$

yields $x \vee y \in a \rightarrow_r I$, because $c \vee d \in I$.

Let $S \subseteq a \rightarrow_r I$ be such that $c = \bigvee_L S$ exists. For every $s \in S$ there is $c_s \in I$ such that $s \leq a \rightarrow_r c_s$, i.e., $a \otimes s \leq c_s$. Since I is an ideal, $a \otimes s \in I$, for all $s \in S$. Then,

$$a \otimes c = a \otimes \bigvee_{s \in S} s = \bigvee_{s \in S} a \otimes s \in I,$$

because I is complete. But this means that $c \in a \rightarrow_r I$. A similar computation will prove that $a \rightarrow_l I \in CI(L)$.

(b) Let $x \in S$ and $a \in T$ and assume that $S \cdot T \subseteq K$; since $x \otimes a = c \in K$, then $x \leq a \rightarrow_l c$ and thus $x \in a \rightarrow_l K$, for every $a \in T$.

Now, suppose that $S \subseteq \bigcap_{a \in T} (a \rightarrow_l K)$, $x \in S$ and $a \in T$; since $x \in a \rightarrow_l K$, there is $c \in K$ such that $x \leq a \rightarrow_l c$, i.e. $x \otimes a \leq c$. Since K is an ideal, we get $x \otimes a \in K$. The other equivalence can be handled similarly. \square

Theorem 6.9 (The completion of a $[\otimes, \vee]$ -autonomous lattice)

Let L be a $[\otimes, \vee]$ -autonomous lattice with $\mathbf{0}, \top$. Then, there exists a quantale Q and an injective map $\phi : L \longrightarrow Q$ such that

1. $\phi(\mathbf{0}) = \mathbf{0}, \phi(\top) = \top$. Moreover, ϕ preserves all \vee 's and \wedge 's existing in L , i.e., if $\bigvee_{i \in I} a_i$ and $\bigwedge_{j \in J} b_j$ exists in L , then

$$\phi(\bigvee_{i \in I} a_i) = \bigvee_{i \in I} \phi(a_i) \text{ and } \phi(\bigwedge_{j \in J} b_j) = \bigwedge_{j \in J} \phi(b_j).$$

2. For all $a, b \in L$,

- (i) $\phi(a \otimes b) = \phi(a) \otimes \phi(b)$
- (ii) $\phi(a \rightarrow_r b) = \phi(a) \rightarrow_r \phi(b)$ and $\phi(a \rightarrow_l b) = \phi(a) \rightarrow_l \phi(b)$.

3. If $a \otimes b = b \otimes a$ for every $a, b \in L$, then Q is commutative.

4. If L is unital, i.e., there is $\mathbf{1} \in L$ such that $a \otimes \mathbf{1} = a = \mathbf{1} \otimes a, \forall a \in L$, then Q is unital and $\phi(\mathbf{1}) = \mathbf{1}_Q$.

5. If in addition, L has a cyclic dualizing element \perp , then Q is a Girard quantale and $\phi(\perp) = \perp$.

Proof: Set $Q = CI(L)$ and define in Q the operation

$$(I) \quad I * J \equiv_{def} \cap \{ K \in Q : K \supseteq I \cdot J \},$$

where $I \cdot J = \{ a \otimes b : a \in I, b \in J \}$.

Clearly $*$ is increasing in both variables. By Lemma 6.8, $\cdot *$ I and $I *$ \cdot have right adjoints $I \rightarrow_r J$ and $I \rightarrow_l J$, given by

$$\cap_{a \in I} (a \rightarrow_r J) \text{ and } \cap_{a \in I} (a \rightarrow_l J),$$

respectively. Thus, $\forall I, J \in Q$,

$$(II) \quad I * J \subseteq K \text{ iff } I \subseteq J \rightarrow_l K \text{ iff } J \subseteq I \rightarrow_r K.$$

Formulas (I) and (II) will be of constant use. To show that $*$ is associative, we need

Fact 1: Let $I, J, K \in Q$. Then

$$\mathcal{A} = \{ H \in Q : H \supseteq (I * J) \cdot K \} = \{ H \in Q : H \supseteq I \cdot (J * K) \} = \mathcal{B}.$$

Proof of Fact 1: We have, using Lemma 6.8, the following sequence of implications

$$\begin{aligned} H \supseteq (I * J) \cdot K &\Rightarrow H \supseteq (I \cdot J) \cdot K = I \cdot (J \cdot K) \\ &\Rightarrow J \cdot K \subseteq I \rightarrow_r H \Rightarrow J * K \subseteq I \rightarrow_r H. \end{aligned}$$

The last term implies $H \supseteq I \cdot (J * K)$ and $\mathcal{A} \subseteq \mathcal{B}$. Similar reasoning yields $\mathcal{B} \subseteq \mathcal{A}$.

It follows directly from the Fact that $*$ is associative. We now must show that $*$ distributes over suprema.

Let $I \in Q$ and $\{I_\alpha\}_{\alpha \in \Lambda} \subseteq Q$; we shall prove that $I * \bigvee I_\alpha = \bigvee (I * I_\alpha)$. Observe that $\bigvee I_\alpha = \cap \{ H \in Q : H \supseteq \bigcup_\alpha I_\alpha \}$, as well as that $\bigvee (I * I_\alpha) \subseteq I * \bigvee I_\alpha$ is always true. Now, we have the following sequence of implications

$$K \supseteq \cup (I^* I_\alpha) \Rightarrow K \supseteq I^* I_\alpha, \forall \alpha \in A \Rightarrow I_\alpha \subseteq I \rightarrow, K, \forall \alpha \in A \\ \Rightarrow \cup I_\alpha \subseteq I \rightarrow, K \Rightarrow \vee I_\alpha \subseteq I \rightarrow, K \Rightarrow I^* \vee I_\alpha \subseteq K,$$

and so $I^* \vee I_\alpha \subseteq \vee (I^* I_\alpha)$. Right distributivity is similar and so Q is indeed a quantale.

Before defining the map ϕ , we state

Fact 2: For $a, b \in Q$

- (i) $a^\leftarrow \rightarrow_r b^\leftarrow = (a \rightarrow_r b)^\leftarrow$ and $a^\leftarrow \rightarrow_l b^\leftarrow = (a \rightarrow_l b)^\leftarrow$.
- (ii) If $S \subseteq L$ is such that $a = \vee S$ exists in L , then $\vee_{s \in S} s^\leftarrow = a^\leftarrow$.
- (iii) If $S \subseteq L$ such that $a = \wedge S$ exists in L , then $a^\leftarrow = \wedge_{s \in S} s^\leftarrow = \cap_{s \in S} s^\leftarrow$.
- (iv) $a^\leftarrow * b^\leftarrow = (a \otimes b)^\leftarrow$.

Proof of Fact 2: (i) Let $K \in Q$ be such that $a^\leftarrow * K \subseteq b^\leftarrow$; for $x \in K$, we have $a \otimes x \leq b$ and so, $x \leq a \rightarrow_r b$. Thus, $K \subseteq (a \rightarrow_r b)^\leftarrow$, proving that $a^\leftarrow \rightarrow_r b^\leftarrow \subseteq (a \rightarrow_r b)^\leftarrow$.

For the reverse inclusion, consider $x \leq a$ and $y \leq a \rightarrow_r b$; then $x \otimes y \leq a \otimes (a \rightarrow_r b) \leq b$ and so $a^\leftarrow \cdot (a \rightarrow_r b)^\leftarrow \subseteq b^\leftarrow$, i.e., $a^\leftarrow * (a \rightarrow_r b)^\leftarrow \subseteq b^\leftarrow$. This last relation implies $(a \rightarrow_r b)^\leftarrow \subseteq a^\leftarrow \rightarrow_r b^\leftarrow$, as needed. For the operation \rightarrow_l , the argument is similar.

(ii) If $\vee S = a$, then $s^\leftarrow \subseteq a^\leftarrow$ for every $s \in S$, and so it is clear that $\vee_{s \in S} s^\leftarrow \subseteq a^\leftarrow$.

If $H \supseteq \cup_{s \in S} s^\leftarrow$, since H is complete, it follows that $a \in H$.

(iii) Is similar to (ii).

(iv) If $x \leq a, y \leq b$ then $x \otimes y \leq a \otimes b$ implies $a^\leftarrow \cdot b^\leftarrow \subseteq (a \otimes b)^\leftarrow$. Thus, $a^\leftarrow * b^\leftarrow \subseteq (a \otimes b)^\leftarrow$.

Now, if $H \supseteq a^\leftarrow \cdot b^\leftarrow$ then $a \otimes b \in H$ and we get $(a \otimes b)^\leftarrow \subseteq H$. By the definition of $*$ (formula (I)), this yields $(a \otimes b)^\leftarrow \subseteq a^\leftarrow * b^\leftarrow$, as desired. This ends the proof of Fact 2.

We now define

$$\phi: L \longrightarrow Q \text{ by } \phi(a) = a^\leftarrow.$$

Clearly ϕ is injective and, by Fact 2.(ii) and (iii), preserves all existing \vee 's and \wedge 's in L . Furthermore, $\phi(\mathbf{0}) = \{ \mathbf{0} \}$ (the smallest complete ideal in L) and $\phi(\mathbf{T}) = L = \mathbf{T}_Q$. The preservation of the other operations is guaranteed by Fact 2.(i) and (iv). This shows that ϕ has the properties in items 1 and 2 of the statement.

It is quite clear from the definition of $*$ (see (I), above) that Q will be commutative if $*$ is commutative in L . Moreover, if L is unital, straightforward computation will show that $\phi(\mathbf{1}) = \mathbf{1}^\leftarrow$ is the unit of Q .

(d) Let $\perp \in L$ be a cyclic dualizing element in L . Clearly $\perp^\leftarrow \in Q$ is cyclic and therefore $Q_j = \{ I \in Q : j(I) = I \}$ is a Girard quantale, where

$j : Q \longrightarrow Q$ given by $j(I) = (I \rightarrow \perp^\leftarrow) \rightarrow \perp^\leftarrow$ (Proposition 1.10). Moreover, $a^\leftarrow \in Q_j$, for every $a \in L$. Thus, the map $j \circ \phi : L \longrightarrow Q_j$ is an embedding of L in a Girard quantale, with all the required properties. \square

We can now prove

Theorem 6.10 (Completeness) *If $|\Gamma|_Q \leq |A|_Q$ for all interpretations in quantales Q , then $\Gamma \vdash A$ is provable in (LLI).*

Proof: Our method will be to show that the Lindenbaum algebra L of (LLI) can be embedded in a commutative unital quantale Q , such that $|\Gamma|_Q \leq |A|_Q$ iff $\Gamma \vdash A$ is provable.

Define in $FOR(\mathbb{L}_i)$ the relation:

$$A \sim B \quad \text{iff} \quad A \vdash B \text{ and } B \vdash A \text{ are provable.}$$

It is easily seen that \sim is an equivalence relation. Its equivalence classes shall be denoted by A/\sim .

Let L be the set of equivalence classes, i.e., $L = \{ A/\sim : A \in FOR(\mathbb{L}_i) \}$. In L , define the relation:

$$A/\sim \leq B/\sim \quad \text{iff} \quad A \vdash B \text{ is provable in (LLI).}$$

By [CUT], \leq is a partial order in L . Moreover, for formulas A, B in LLI, we have

Fact 1: If A, B are formulas in (LLI), then

1. $(A \& B)/\sim = (A/\sim) \wedge (B/\sim)$ and $(A \oplus B)/\sim = (A/\sim) \vee (B/\sim)$.
2. $\bigwedge_{y \in V} (A[y/x]/\sim) = (\bigwedge x. A)/\sim$ and $\bigvee_{y \in V} (A[y/x]/\sim) = (\bigvee x. A)/\sim$.
3. $\mathbf{0} = \mathbf{0}/\sim$ and $\top = \top/\sim$.

Proof of Fact 1: All these equalities can be read off the corresponding rules of the calculus. We do the first one in each of items 1. and 2. in some detail, just naming the rules that should be used for the other cases.

1. From [AX1], [&1] and [&] we get $A \& B \vdash A, B$ and so $(A \& B)/\sim \leq A/\sim$, $(A \& B)/\sim \leq B/\sim$. On the other hand, if $C \vdash A$ and $C \vdash B$, then [&R] yields $C \vdash A \& B$. This means that

$$C/\sim \leq A/\sim, C/\sim \leq B/\sim \Rightarrow C/\sim \leq (A \& B)/\sim,$$

proving that $(A \& B)/\sim = (A/\sim) \wedge (B/\sim)$ in L .

Similarly, [AX1], [\oplus 1], [\oplus 2] and [\oplus L] will yield $(A \oplus B)/\sim = (A/\sim) \vee (B/\sim)$.

2. By [AX1] and $[\wedge L]$, $(\wedge x. A)/_{\sim} \leq A[y/x]/_{\sim}$, for every $y \in V$.

If $B/_{\sim} \leq A[y/x]/_{\sim}$, for every y , let z be a variable not occurring in B . Then, $B/_{\sim} \leq A[z/x]/_{\sim}$ and so $[\wedge R]$ yields $B/_{\sim} \leq (\wedge x. A)/_{\sim}$. This proves that $\bigwedge_{x \in V} (A[y/x]/_{\sim}) = (\wedge x. A)/_{\sim}$.

For the existential quantifier, the reasoning is the same, using [AX1], $[\vee R]$ and $[\vee L]$.

3. This follows directly from [AX2] and [AX4], ending the proof of Fact 1.

Define in L a binary operation $*$ by:

$$(A/_{\sim}) * (B/_{\sim}) \equiv_{\text{def}} (A \otimes B)/_{\sim}.$$

Clearly, $*$ is well defined, is associative, commutative, increasing in both variables and $1/_{\sim}$ is the unit. We have

Fact 2: With the operation $*$ defined above, L is a $[\otimes, \vee]$ autonomous lattice.

Proof of Fact 2: It remains to be shown that $*$ has right adjoints and distributes over the suprema existing in L .

If A, B, C are formulas in (LL) , then

$$(+) \quad A \otimes B \vdash C \text{ is provable} \quad \text{iff} \quad A, B \vdash C \text{ is provable} \\ \text{iff} \quad A \vdash B \multimap C \text{ is provable.}$$

To see this, note that $[\otimes R]$ yields $\frac{A \vdash A \quad B \vdash B}{A, B \vdash A \otimes B}$

Thus if $A \otimes B \vdash C$ is provable, [CUT] implies that $A, B \vdash C$ is provable.

The converse comes directly from $[\otimes L]$ as $\frac{A, B \vdash C}{A \otimes B \vdash C}$.

From $[\multimap R]$ we get $\frac{A, B \vdash C}{A \vdash B \multimap C}$.

To show that the last clause in (+) implies the second, first note that $B, B \multimap C \vdash C$ is provable, because we can use $[\multimap R]$ a $\frac{B \vdash B \quad C \vdash C}{B, B \multimap C \vdash C}$

Thus, the provability of $A \vdash B \multimap C$ and $B, B \multimap C \vdash C$ and [CUT] yield that of $A, B \vdash C$.

It is clear from (+) that $(A \multimap \cdot)_{\sim}$ is the right adjoint to $A_{\sim} * \cdot = \cdot * A_{\sim}$ in L .

To show that $*$ distributes over the sup's in L , let $S \subseteq L$ be such that there is $A_{\sim} \in L$ satisfying $A_{\sim} = \bigvee_L S$. Fix $B_{\sim} \in L$. Since $*$ is increasing, $(B_{\sim})^* (X_{\sim}) \leq (B_{\sim})^* (A_{\sim})$, for every $X_{\sim} \in S$.

Let $F_{\sim} \in L$ such that $(B_{\sim})^* (X_{\sim}) \leq F_{\sim}$, for every $X_{\sim} \in S$. By (+) we have that, for $X_{\sim} \in S$,

$$B \otimes X \vdash F \text{ is provable} \quad \text{iff} \quad X \vdash (B \multimap F) \text{ is provable},$$

$\forall X_{\sim} \in S$; taking sup's, we get $A \vdash (B \multimap F)$ is provable. Thus, $B \otimes A \vdash F$ is provable, and so $(B_{\sim})^* (A_{\sim}) \leq F_{\sim}$. But this means that $\bigvee_{X_{\sim} \in S} ((B_{\sim})^* (X_{\sim}))$ exists in L , and it's equal to $(B_{\sim})^* \bigvee_L S$, ending the proof of Fact 2.

Therefore, L is a commutative $[\otimes, \vee]$ -autonomous lattice and by theorem 6.9 it has a completion $\phi : L \longrightarrow Q$, where Q is an unital commutative quantale. Moreover, ϕ preserves \otimes , \multimap , $\mathbf{0}$, \top , $\mathbf{1}$ and all existing \vee 's and \wedge 's in L . Consequently, if X, Y are formulas in (LL) we have

$$\phi(X_{\sim}) \leq \phi(Y_{\sim}) \quad \text{iff} \quad X_{\sim} \leq Y_{\sim}.$$

It follows from these preservation properties that the map $|\cdot|_Q : FOR(\mathbb{L}_i) \longrightarrow Q$ given by $|A|_Q = \phi(A_{\sim})$ is an interpretation of quantales such that $|\Gamma| \leq |A|$ iff $\Gamma \vdash A$ is provable, as required to complete the proof of the Theorem. \square

Before we give logical applications of the preceding results, we need to investigate when is it that a quantale becomes a frame. From Lemma 6.4 comes:

Corollary 6.11 *Let Q be a quantale. If Q has a largest localic subquantale L , then*

$$L = I(Q) = \{ a \in Q : a^2 = a \}.$$

Proof. We must have $L \subseteq I(Q)$ because every $x \in L$ is idempotent (proposition 1.19).

For $a \in I(Q)$, set $L_a = \{ \mathbf{0}, a \}$. By Lemma 6.4, L_a is closed under \otimes ; it is clearly closed under arbitrary \vee 's, and so L_a is a subquantale of Q . Since a is idempotent, it follows that L_a is localic. Thus, $L_a \subseteq L$, i.e., $a \in L$, and therefore $I(Q) \subseteq L$. \square

The next result is a correction of Theorem 3.4.1 in [Ros], which is false as it stands. In fact, the usual quantale of phases of Linear Logic is a counterexample to that result.

Proposition 6.12 *Let Q be a quantale. Are equivalent:*

- (1) Q has a largest localic subquantale.
- (2) $I(Q)$ is a localic subquantale of Q .
- (3) For all every $a, b \in I(Q)$, $a \otimes b = b \otimes a \leq a \wedge b$ (\wedge in Q or in $I(Q)$).

Proof. (1) \Rightarrow (2) comes from Corollary 6.11, while (2) \Rightarrow (3) is immediate.

(3) \Rightarrow (1): Because $I(Q)$ is commutative, it is closed under \otimes . To show that $I(Q)$ is closed under sup's, let $\{a_i\}_{i \in I} \subseteq I(Q)$. Then:

$$\begin{aligned} (*) \quad \bigvee_{i \in I} a_i &= \bigvee_{i \in I} (a_i \otimes a_i) \leq \bigvee_{i, j \in I} (a_i \otimes a_j) \\ &= (\bigvee_{i \in I} a_i) \otimes (\bigvee_{j \in I} a_j); \\ (***) \quad (\bigvee_{i \in I} a_i) \otimes (\bigvee_{j \in I} a_j) &= \bigvee_{i \in I} (\bigvee_{j \in I} (a_i \otimes a_j)) \leq \bigvee_{i \in I} a_i, \end{aligned}$$

because $a_i \otimes a_j \leq a_i$ for every $j \in I$. From (*) and (**) it follows that $\bigvee_{i \in I} a_i$ is idempotent and so $I(Q)$ is a subquantale of Q .

To show that $I(Q)$ is localic, observe that it follows from 3. that $a \otimes b \leq a$, for every $b \in I(Q)$. Thus, $\forall a \in I(Q)$,

$$a \otimes \bigvee I(Q) = \bigvee_{b \in I(Q)} (a \otimes b) \leq a \text{ and } (\bigvee I(Q)) \otimes a \leq a,$$

which shows that all elements of $I(Q)$ are idempotent and two-sided in $I(Q)$. Now, Proposition 1.19 guarantees that $I(Q)$ is localic. If $L \subseteq Q$ is a localic subquantale then every $x \in L$ is idempotent (Proposition 1.19) and so $L \subseteq I(Q)$. \square

Thus, in a commutative Girard quantale Q , $I(Q) \cap \mathbf{1}^\leftarrow$ is the *largest frame* contained in $\mathbf{1}^\leftarrow$. From the point of view of Logic, we have a privileged localic subquantale, that corresponds to intuitionistic Logic. However, the situation is quite different in the non commutative case: in general, a Girard quantale will not have a largest localic subquantale. If one considers Proposition 1.14, one realizes the importance of commutativity: there is a largest interpretation for $!$ in a Girard's quantale Q , corresponding to the largest among frames L satisfying

$$(i) \quad L \subseteq Z(Q) \quad (ii) \quad L \subseteq \mathbf{1}^\leftarrow = \{x \in Q : x \leq \mathbf{1}\}$$

In fact, $Q_! = I(Q) \cap Z(Q) \cap \mathbf{1}^\leftarrow$, where $Z(Q)$ is the center of Q .

The quantale theoretic setting will be helpful in finding axioms that, when added to various systems of Linear Logic, produce classical intuitionism and classical logic. We have the simple (compare with proposition 1.19),

Lemma 6.13 *Let L be a complete lattice and $* : L \times L \longrightarrow L$ a binary operation on L . Are equivalent:*

- (i) $* = \wedge$.
- (ii) (a) $x * x = x$ for every $x \in L$.
- (b) $x * \top \leq x$, $\top * x \leq x$ for every $x \in L$.
- (c) $*$ is increasing in both variables.

Proof: (i) \Rightarrow (ii) is clear.

(ii) \Rightarrow (i): We have $a * b \leq a * \top \leq a$; $a * b \leq \top * b \leq b$ and therefore, $a * b \leq a \wedge b$.

On the other hand, since $*$ is increasing $a \wedge b = (a \wedge b) * (a \wedge b) \leq a * b$. \square

An analogous (and dual) result holds for the operation \vee . From the preceding results we get

Corollary 6.14 *Let Q be a quantale. Are equivalent:*

- (i) Q is a frame, i.e., $\otimes = \wedge$.
- (ii) every $x \in Q$ is idempotent and two-sided. \square

This Corollary and the completeness Theorem 6.10 yield

Proposition 6.15 (a) *The system (LI) obtained from (LLI) by adding the axioms*

$$\begin{array}{ll} [ID1] & \vdash A \otimes A \multimap A \\ [2S] & \vdash A \otimes \top \multimap A \end{array} \quad \begin{array}{ll} [ID2] & \vdash A \multimap A \otimes A \end{array}$$

determines intuitionistic logic, and therefore if we add to (LI) the axiom

$$[\neg \neg] (A \multimap \mathbf{0}) \multimap \mathbf{0} \vdash A$$

we get classical logic, where $\mathbf{1}$ is equivalent to \top and $\mathbf{1} \multimap \mathbf{0}$ is equivalent to $\mathbf{0}$.

Thus, adding the axioms [ID1],[ID2],[2S] and [R=], [S=], [T=], [SUBST], [$\leq \mathbf{1}$] (see section 2) to (LLI) we obtain a intuitionistic theory of equality,

which becomes first order classical logic with equality upon the addition of $[\neg \neg']$.

(b) The system (MALL1) obtained from (LLI) by adding the constant formula \perp to the language and the axiom

$$[\neg \neg'] (A \multimap \perp) \multimap \perp \vdash A,$$

is equivalent with (MALL) in the following sense:

$$\vdash A \text{ is provable in (MALL) iff } \vdash A \text{ is provable in (MALL1)},$$

where A^\perp and $A \sqcup B$ are interpreted as $A \multimap \perp$ and $((A \multimap \perp) \otimes (B \multimap \perp)) \multimap \perp$, respectively. Thus, adding \perp to the language we have that

$$(LLI) + \text{the axioms } [\neg \neg'], [R=], [S=], [T=], [SUBST], [I=], [\leq 1] \\ \text{is equivalent to } (LLE_0),$$

where (LLE_0) is described in Definition 2.6.

(c) Adding [ID1], [ID2] and [2S] to (MALL1) produces classical logic, where $\mathbf{1}$ is equivalent to \top , and $\mathbf{0}$ is equivalent to \perp . \square

Remark 6.16 The possibility of obtaining (MALL) from (LLI) in part (b) appears in [Dos] proved by a different method: one interprets $A \multimap B$, $A \& B$ and $\wedge x.A$ as $(A \otimes (B \multimap \perp)) \multimap \perp$, $((A \multimap \perp) \oplus (B \multimap \perp)) \multimap \perp$ and $(\vee x.(A \multimap \perp)) \multimap \perp$, respectively, together with the interpretation for \sqcup made in (b).

Since $(a \otimes b) \rightarrow c = a \rightarrow (b \rightarrow c)$ and $b = \mathbf{1} \rightarrow b$, in every unital commutative quantale, it can be shown that his interpretation of linear implication is equivalent to adding $[\neg \neg']$ to (LLI) to obtain (MALL) (translations for $\&$ and \wedge are derived and a consequence of the ones given).

As a matter of fact, we can extend this result to non-commutative intuitionistic linear logic, simply by observing that each rule of (LLI) determines an algebraic property of the Lindenbaum algebra of the calculus. For a general unital quantale, we need two implications, \multimap_l and \multimap_r , to discard the exchange rule [EXCH], as well as to modify the rules in order to describe the properties of each operation. We can define a linear calculus which describes unital quantales, essentially the same as in [Abr].

Definition 6.17 The calculus (NLLI) for noncommutative first-order linear intuitionistic logic consist in the following rules and axioms:

[AX1]	$A \vdash A$	[AX2]	$\Gamma \vdash \top$
[AX3]	$\vdash \mathbf{1}$	[AX4]	$\Gamma, \mathbf{0}, \Delta \vdash A$
[CUT]	$\frac{\Gamma \vdash A \quad \Sigma, A, \Delta \vdash B}{\Sigma, \Gamma, \Delta \vdash B}$	[1L]	$\frac{\Gamma, \Delta \vdash A}{\Gamma, \mathbf{1}, \Delta \vdash A}$
[\otimes R]	$\frac{\Gamma \vdash A \quad \Delta \vdash B}{\Gamma, \Delta \vdash A \otimes B}$	[\otimes L]	$\frac{\Gamma, A, B, \Delta \vdash C}{\Gamma, A \otimes B, \Delta \vdash C}$
[& R]	$\frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \& B}$	[\oplus L]	$\frac{\Gamma, A, \Delta \vdash C \quad \Gamma, B, \Delta \vdash C}{\Gamma, A \oplus B, \Delta \vdash C}$
[& L1]	$\frac{\Gamma, A, \Delta \vdash C}{\Gamma, A \& B, \Delta \vdash C}$	[& L2]	$\frac{\Gamma, B, \Delta \vdash C}{\Gamma, A \& B, \Delta \vdash C}$
[\oplus R1]	$\frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B}$	[\oplus R2]	$\frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B}$
[\multimap_l L]	$\frac{\Gamma \vdash A \quad \Sigma, B, \Delta \vdash C}{\Sigma, A \multimap_r B, \Gamma, \Delta \vdash C}$	[\multimap_r , R]	$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \multimap_r B}$
[\multimap_r , L]	$\frac{\Gamma \vdash A \quad \Sigma, B, \Delta \vdash C}{\Sigma, \Gamma, A \multimap_r B, \Delta \vdash C}$	[\multimap_r , R]	$\frac{A, \Gamma \vdash B}{\Gamma \vdash A \multimap_r B}$
[\vee R]	$\frac{\Gamma \vdash A[y/x]}{\Gamma \vdash \vee x.A}$		
[\vee L]	<i>If x is not free in Γ, Δ, B then</i>		$\frac{\Gamma, A, \Delta \vdash B}{\Gamma, \vee x.A, \Delta \vdash B}$
[\wedge L]	$\frac{\Gamma, A[y/x], \Delta \vdash B}{\Gamma, \wedge x.A, \Delta \vdash B}$		
[\wedge R]	<i>If x is not free in Γ then</i>		$\frac{\Gamma \vdash A}{\Gamma \vdash \wedge x.A}$

With the same method employed in Theorems 6.5 and 6.10, Theorem 6.9 yields that unital quantales are a complete and sound class of models for (NCLL):

Theorem 6.18 (Completeness and Soundness for (NCLL))

A sequent $\Gamma \vdash A$ is provable in [NCLL] iff $\Gamma \vdash A$ is valid in every interpretation of quantales.

Instituto de Matemática e Estatística, Universidade de São Paulo
 Caixa Postal 66.281 - Cep 05315-970, S.Paulo, S.P., Brasil
 e-mail: M.E. Coniglio: coni@ime.usp.br
 F. Miraglia: miraglia@ime.usp.br or miraglia@jussieu.logique.fr

REFERENCES

- [Abr] Abrusci V.M., ‘Non-commutative Intuitionistic Linear Logic’, *Zeitschr. f. math. Logik und Grundlagen d. Math.* 36 (1990), 297-318.
- [Avr] Avron A., ‘The Semantics and Proof Theory of Linear Logic’, *Theoretical Computer Science* 57 (1988), 161-184.
- [BrGu] Brown C. and Gurr D., ‘A Representation Theorem for Quantales’, *Journal of Pure and Applied Algebra* 85 (1993), 27-42.
- [Doš] Došen K., ‘Nonmodal Classical Linear Predicate Logic is a Fragment of Intuitionistic Linear Logic’, *Theoretical Computer Science* 102 (1992), 207-214.
- [Gir] Girard J.Y., ‘Linear Logic’, *Theoretical Computer Science* 50 (1987), 1-102.
- [GiLa] Girard J.Y. and Lafont Y., ‘Linear Logic and Lazy Computation’, *Lectures Notes in Computer Science* 250 (Springer, 1987), 52-66.
- [Mir] Miraglia F., *Sheaves and Logic*, preprint.
- [Ros] Rosenthal K., ‘Quantales and their Applications’, *Pitman Research Notes in Mathematics Series*, vol. 234 (Longman, Harlow, 1990).
- [Yet] Yetter D., ‘Quantales and (Noncommutative) Linear Logic’, *Journal of Symbolic Logic* 55 (1990), 41-64.