

RT-MAT 90-08

Total curvature of foliations

R. Langevin & C. Possani

dezembro 1990

TOTAL CURVATURE OF FOLIATIONS

R. Langevin and C. Possani

Summary :

In this article we study the total curvature of foliations of a manifold perhaps with boundary of constant curvature.

The hyperbolic case is handled by identifying the set of geodesic segments with a quotient of the group of isometries of the hyperbolic space covering the manifold.

Foliations given on the boundary of M are studied using properties of the envelope of the hyperplane field they define on the boundary.

Finally we prove a kinematic formula for a plane foliation and a non-geodesic curve.

R. Langevin
Laboratoire de Topologie
C.N.R.S. DO 755
Université de Bourgogne
Faculté des Sciences Mirande
B.P.133
21004 Dijon - FRANCE

C. Possani
Instituto de Matematica e Estatística
Universidade de São Paulo
C.P. 20570 (agência Iguatemi)
05508 São Paulo S.P.
BRASIL

The total curvature of foliations will be studied with an extensive use of integral geometry. More precisely there is a dictionary between curvature integrals and counting of contact points each line of which we will call an exchange theorem.

It is not a surprise that lower bounds of curvature integrals of a foliation given on the boundary of a domain depend on the integral geometry of the boundary data.

I. EXCHANGE THEOREM

Throughout this paper we'll use the so called "Exchange Theorem". In this paragraph we will state it and make some remarks. For the proof we refer the reader to [B.L.R], or [L].

Let $W \subset \mathbb{R}^{n+1}$ be an open set with $\bar{W}$ compact, $U \subset W$ an open subset of W , and $\mathcal{F}$ a codimension one foliation of W with isolated singularities. If $u \in S^n \subset \mathbb{R}^{n+1}$, let $L(u)$ be the oriented line through the origin of $\mathbb{R}^{n+1}$. For each $t \in L(u)$ let T_t be the hyperplane orthogonal to $L(u)$ passing through t . We define $|\mu|(\mathcal{F}, T_t) \in \mathbb{N} \cup \infty$ to be the number of points of U where the hyperplane T_t is tangent to a leaf of $\mathcal{F}$. This number is finite for almost all pairs $(L(u); t)$.

Let $K(x)$ be the Lipschitz-killing curvature of the leaf through the point x .

With these notations we have :

I.1. Exchange Theorem.

$$(I.1) \quad \int_{S^n} \int_{L(u)} |\mu|(\mathcal{F}, T_t) dt du = \int_U |K|(x) dx$$

(Both sides of the equality may be infinite).

This inequality may, of course, be written :

$$(I.2) \quad 1/2 \int_{S^n} \int_{L(u)} |\mu|(\mathcal{F}, T_t) dt du = \int_U |K|(x) dx$$

When $n = 1$, that is when W is an open set in $\mathbb{R}^2$ with compact closure, the left integral can be understood as an integral over the space $\mathcal{G}$ of non-oriented lines of the plane. This set is endowed with the density m associated to the form $dr d\theta$ where r is the distance of the line T to the origin, and θ the angle between the axis Ox and the line T . For more details see [S].

The exchange theorem becomes :

$$(I.3) \quad \int_U |K|(x) dx = \int_{\mathcal{G}} |\mu|(\mathcal{F}, L) dm(L)$$

Instead of considering contact with lines, we can count contacts between the leaves of $\mathcal{F}$ and line segments of fixed length. Let $\mathcal{G}_d$ be the set of non-oriented line segments in $\mathbb{R}^2$ of length d . We parametrize a segment I of $\mathcal{G}_d$ by coordinates (r, θ, s) where (r, θ) are the coordinates of the line L supporting I and s the coordinate of the center of I in L . This gives a density $m = |dr \wedge d\theta \wedge ds|$ on $\mathcal{G}_d$ which is invariant by the action of the isometries, and therefore another version of the exchange theorem :

$$(I.4) \quad \int_U |K|(x) dx = \frac{1}{d} \int_{\mathcal{G}_d} |\mu|(\mathcal{F}, I) dm(I)$$

We will from now on denote by $K(\mathcal{F})$ the integral $\int_U |K|(x) dx$.

If we consider the analogous situation in a hyperbolic surface M , then, although the set of geodesics of M cannot be given a measure invariant by the isometries, the measure $m = |\cosh r \cdot dr \wedge d\theta \wedge ds|$ on the set $\mathcal{G}_d$ of geodesic segments of length d is invariant by the action of "local isometries" and I.4 still holds, see [L.L1].

II. TOTAL CURVATURE OF FOLIATIONS OF BOUNDED PLANE DOMAINS

In this paragraph we will consider foliations of a given domain $D \subset \mathbb{R}^2$ homeomorphic to the unit disc $D^2 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$, and we suppose that its boundary ∂D is piecewise C^2 .

II.1. Definition : the distance between two points x and y in D is defined by :

$d(x, y) = \inf \{l(\gamma) \mid \gamma : [a, b] \rightarrow D \text{ is a regular curve, } \gamma(a) = x, \gamma(b) = y \text{ and } l(\gamma) \text{ is the length of } \gamma\}$

It is clear that d is a metric in D . In fact, as $D \approx D^2$ there always exists exactly one curve γ joining x and y which satisfies $l(\gamma) = d(x, y)$, such a γ is called a "geodesic" of D .

II.2. Definition : the diameter of D is defined as

$$d = \sup \{d(x, y) \mid x, y \in D\}.$$

Let $\mathcal{F}$ be a foliation by curves of D , tangent to ∂D , with isolated singularities of positive index. $\mathcal{F}$ may not be orientable.

II.3. Theorem : with the above notations we have

$$K(\mathcal{F}) \geq l(\partial D) - 2d$$

This theorem is analogous to one of R. Langevin and G. Levitt that asserts if $\mathcal{F}$ is an orientable foliation of the disc D^2 , tangent to the boundary then

$$K(\mathcal{F}) \geq 2\pi - 4$$

Demonstration :

First of all, let us define the index of a singularity of a non orientable foliation.

The index of a singularity P is an integer which measures the variation over RP^1 of the field of directions of the leaves of $\mathcal{F}$ around a neighbourhood of P . The index we consider is the above integer index divided by 2. If the singularity is orientable, then this index is the usual one.

We are going to show that we can suppose $\mathcal{F}$ has 2 singularities of sunset type, which are of index $\frac{1}{2}$. (fig. 2)




fig. 2

All singularities with the property that the oscilation of its field of directions is uniformly bounded in any circle centered at the singularity, can be substituted by a sunset or a source increasing the total curvature of $\mathcal{F}$ as little as we want.

This can be done by considering in the boundary of the disc D_r , with center at the singular point P and radius r , a homotopy between the angle function determined by $\mathcal{F}$ and that one of the foliations below :





fig. 3

A source can be replaced by two sunsets by the modification :




fig. 4

If the oscillation of the field of directions of $\mathcal{F}$ in the neighbourhood of P is not uniformly bounded then $K(\mathcal{F}) \rightarrow \infty$ in a neighbourhood of P and we are done.

Let P and Q be the two sunsets of $\mathcal{F}$, and γ a geodesic of D joining them. We are going to estimate the number of points of contact of leaves of $\mathcal{F}$ and a line L of the plane. Except for a set of lines of measure zero, each line L meets D in a finite number of segments.

If $\overline{AB}$ is a segment of $L \cap D$ and $\overline{AB} \cap \gamma = \emptyset$ then $\overline{AB}$ divides D into two discs, one of them containing $\{P, Q\}$. In the other disc $\mathcal{F}$ is orientable and certainly there's at least one point of contact between $\overline{AB}$ and leaves of $\mathcal{F}$. (see fig.5)




fig. 5

Let $n(L)$ be the number of segments of $L \cap D$ in which L meets γ , and $c(L)$ the number of segments of $L \cap D$ in which L doesn't meet γ . Then we have

II.4.

$$|\mu|(\mathcal{F}; L) \geq c(L)$$

Cauchy's formula [5] gives us :

$l(c) = \int_{\mathcal{G}} \# \{ \text{segments } L \cap c \} = \frac{1}{2} \int_{\mathcal{G}} \# \{ L \cap c \}$ if c is a piecewise regular closed curve and

$l(c) = \frac{1}{2} \int_{\mathcal{G}} \# \{ L \cap c \}$ if c is a piecewise regular arc.

Then we have

$$l(\partial D) = \int_{\mathcal{G}} (n(L) + c(L)) = \int_{\mathcal{G}} n(L) + \int_{\mathcal{G}} c(L) = \int_{\mathcal{G}} \# \{ L \cap \gamma \} + \int_{\mathcal{G}} c(L)$$

And applying II.4 and the exchange theorem

$$l(\partial D) \leq 2l(\gamma) + \int_{\mathcal{G}} |\mu|(\mathcal{F}; L) = 2l(\gamma) + K(\mathcal{F})$$

And then

$$K(\mathcal{F}) \geq l(\partial D) - 2l(\gamma) \geq l(\partial D) - 2d.$$

with the same techniques we can analyse the case where D is a region of $\mathbb{R}^2$ homeomorphic to the annulus $A = \{(x, y) \in \mathbb{R}^2 \mid 0 < r_1 \leq x^2 + y^2 \leq r_2\}$, with boundary $\partial D = C_1 \cup C_2$, where C_1 and C_2 are piecewise regular closed curves. (fig. 6).

Suppose C_1 is the interior curve of ∂D .

II.5. Definition : If $\mathcal{D}$ is a region in $\mathbb{R}^2$ homeomorphic to the annulus A , with piecewise regular boundary $\partial D = C_1 \cup C_2$, C_1 being the interior curve, we call interior convex envelope of C_1 the curve C'_1 which is the boundary of the convex-hull of C_1 in D , that is the boundary of the intersection of all sets of $\mathbb{R}^2$ that are contained in the interior of C_2 and convex in D . (fig. 6).




fig. 6

We can then prove in an analogous way :

II.6. Theorem : If $D \subset \mathbb{R}^2$ is a region homeomorphic to the annulus A as above, and if $\mathcal{F}$ is a foliation of D , without singularities, tangent to the boundary then

$$K(\mathcal{F}) \geq l(C_2) + l(C_1) - 2l(C'_1)$$

If, in theorem II.3, the curve ∂D is convex then there exists a tight foliation of D . By "tight foliation" we mean a foliation $\mathcal{F}$ that realizes the minimum of $K(\mathcal{F})$. Analogously in theorem II.6, if C_1 and C_2 are convex, $C'_1 = C_1$, $K(\mathcal{F}) \geq l(C_1) - l(C_1)$ then there exists tight foliations. (see fig. 7)




fig. 7

If ∂D , in theorem II.3, or C_i , $i = 1, 2$, in theorem II.6 are not convex then we can show that there do not exist tight foliations of D .

The non existence of tight foliations when the boundary of the domain is not a convex curve follows from the following fact : If $P \in \partial D$ is a point of inflexion of ∂D and a regular point for $\mathcal{F}$ then there will be an open set of lines in the plane with the property that each line in this set has more than one point of contact with the leaves of $\mathcal{F}$, in a neighbourhood of x .

But in this case we can exhibit a sequence of foliations of D , $(\mathcal{F}_n)$ with $K(\mathcal{F}_n) \rightarrow l(\partial D) - 2d$.

We can think of the limit of this sequence of foliations as a foliation having ∂D as critical set (In general, this critical set could be a union of curves), with the "curvature concentrated in ∂D " (see fig. 8) (More details in [P]).




fig. 8

III. FOLIATIONS IN A HYPERBOLIC SURFACE.

So far the only known theorem concerning total curvature of foliations in a hyperbolic surface is due to R. Langevin and G. Levitt [L.L.1] and asserts that if M is a hyperbolic compact surface, without boundary and $\mathcal{F}$ is a foliation of M whose singularities are all isolated and of saddle type then

$$K(\mathcal{F}) \geq (12 \log 2 - 6 \log 3) |\chi(M)|$$

In this paragraph we will prove a theorem for two orthogonal foliations in a compact hyperbolic surface M , without boundary. Our approach will be quite similar to the one the two authors mentioned above used in [L.L.2] to establish an analogous theorem for two orthogonal foliations in the sphere $S^2 \subset \mathbb{R}^3$.

First of all let us recall that the universal covering of a compact hyperbolic surface without boundary is the hyperbolic plane H^2 , and we will think of it as the Poincaré disc.

In hyperbolic geometry the interior angle of a regular n -gon is not determined by n , a priori. Given n , there exist a regular polygon of n sides in H^2 , with interior angle α , for each value of α satisfying

$$0 < \alpha < \frac{n-2}{n} \cdot \Pi$$

For technical reasons, that will be clear later, we need n odd and α equal to $\frac{\Pi}{2}$. The smallest n we can get is 5. Given n and α then the length, l , of the side of the regular n -gon in H^2 becomes determined. For $n = 5$ and $\alpha = \frac{\Pi}{2}$ we have $l \approx 4,275$.

We can now state our theorem :

III.1. Theorem : Let M be a compact, without boundary, hyperbolic (that is, with a metric of constant curvature -1) surface and let $\mathcal{F}$ and $\mathcal{F}^\perp$ be orthogonal foliations of M with isolated singularities. Then

$$K(\mathcal{F}) + K(\mathcal{F}^\perp) \geq \frac{2\Pi^2}{5l} | \chi(M) |,$$

where l is the length of the side of the regular pentagon in H^2 , whose interior angles are $\frac{\Pi}{2}$.

Before proving this we make some definitions and remarks.

III.2. Definition : We let G be space of all orientation preserving isometries of H^2 . We define an equivalence relation in G by

$$g_1 \approx g_2 \Leftrightarrow p \circ g_1 \circ g_2^{-1} = p$$

It's easy to check that $\approx$ is actually an equivalence relation. We will denote $G/\sim$ by $\bar{G}$.

III.3. Définition : As in paragraph I we have :

$\mathcal{G}$ = the space of all geodesics of H^2 , with the canonical measure given by the density $m = \cosh r \, dr \, d\theta$.

$\mathcal{G}_r$ = the space of all geodesic segments of length $r > 0$, with the density $m = \cosh r \, dr \, d\theta \, ds$.

We identify, in H^2 , geodesic segments which project by p over the same geodesic arc in M .

III.4. Definition : In $\mathcal{G}$, we define an equivalence relation by $C_1 \approx C_2 \Leftrightarrow (g \in G, g(C_1) = C_2 \Rightarrow [g] = [Id] \text{ or } [g] = [\theta_A])$ where A is the middle point of C_2 and θ_A is the geodesic symmetry through A).

It's easy to see that $\approx$ is an equivalence relation, and we indicate $\mathcal{G}_r/\sim$ by $\bar{\mathcal{G}}_r$. It's not hard to check that the measure in $\mathcal{G}_r$ induces a measure in $\bar{\mathcal{G}}_r$.

We define now a measure in $\tilde{G}$. Consider a fixed segment C_0 in H^2 , of length r . For each $C \in \mathcal{G}$, we have two isometries $g_i \in G, i = 1, 2$ with the property that $g_i(C_0) = C$. (They satisfy $g_1 = \theta_A \circ g_2$). So G is a two-fold covering of $\mathcal{G}$, and we can consider in G a measure induced by $\mathcal{G}$. This measure induces, through the canonical projection, a measure in $\tilde{G}$.

Remark : The measures in $\mathcal{G}$, and in $\tilde{G}$ could have been obtained by parametrizing the geodesic segments by the position in H^2 of it's middle point and by the direction of the tangent line at this point. The above construction allows us to think of an equivalence class $[g]$ as a geodesic segment in M . We return now to our theorem.

Proof of theorem III.1.

Let $\mathcal{F}$ be a foliation of M and $\mathcal{F}^\perp$ be the orthogonal foliation. $\mathcal{F}$ and $\mathcal{F}^\perp$ induce by $p: H^2 \rightarrow M$, foliations in H^2 , and we will also denote them by $\mathcal{F}$ and $\mathcal{F}^\perp$. As p is locally an isometry and considering the identifications we made above we have

$$K(\mathcal{F}) = \frac{1}{r} \int_{\mathcal{G}} |\mu|(\mathcal{F}; [C])$$

where $[C]$ is the equivalence class of a segment C of length r , and $|\mu|(\mathcal{F}; [C])$ is the number of points of contact between leaves of $\mathcal{F}$ in H^2 and a representative of the class $[C]$.

We can also consider $K(\mathcal{F})$ as an integral on the space $\tilde{G}$:

$$K(\mathcal{F}) = \frac{1}{2l} \int_{\tilde{G}} |\mu|(\mathcal{F}; [g](C)) =$$

where C is fixed, with $l(C) = l$, $|\mu|(\mathcal{F}; [g](C))$ is the number of contact points between $\mathcal{F}$ in H^2 and $g(C), g \in [g]$.

The factor $\frac{1}{2}$ appears in this equality since each point of contact between leaves of $\mathcal{F}$, and a segment C will be counted twice in $\tilde{G}$.

We can fix C , and let $\tilde{G}$ act on $\mathcal{F}$ to obtain

$$K(\mathcal{F}) = \frac{1}{2l} \int_{\tilde{G}} |\mu|([g]\mathcal{F}; C)$$

where $|\mu|([g]\mathcal{F}; C)$ is the number of points of contact between leaves of $g(\mathcal{F}), g \in [g]$ and C .

From now on we will fix P , a regular pentagon in H^2 whose interior angles are $\frac{\pi}{2}$ and count contacts between its sides and leaves of $g(\mathcal{F})$. We have

$$K(\mathcal{F}) = \frac{1}{10l} \int_{\tilde{G}} |\mu|([g]\mathcal{F}; P)$$

III.5. Lemma : $|\mu|([g]\mathcal{F}; P) + |\mu|([g]\mathcal{F}; P) \geq 1$.

Proof : Here we need a odd number of sides and interior angles $\frac{\pi}{2}$.

This idea is due to R. Conelly, after [L.L.2]. Except for a set of isometries of zero measure, at the vertices v_i of P we have two cases to consider :

i) The leaf trough v_i , L_i is locally contained in the exterior of P . We call it "exterior leaf".

ii) v_i divides L_i into two half-leaves, one of them locally contained in the interior of P , the other contained in the exterior of P . We call it "interior leaf". (see fig. 9)

As we have 5 vertices there are two consecutives vertices in which the leaves of $\mathcal{F}$ are both exterior or both interior (see fig. 9).




fig. 9

If we have two consecutive vertices, say v_1, v_2 , for which $\mathcal{F}$ has interior leaves, or for which $\mathcal{F}$ has exterior leaves, then there exists a point v on the edge $v_1 v_2$, through which the leaf L_v of $\mathcal{F}$ is either tangent or perpendicular to the side $v_1 v_2$. If the leaf is tangent $|\mu|(\mathcal{F}, P) \geq 1$, if it is orthogonal $\mu(\mathcal{F}^\perp, P) \geq 1$. (see fig. 10).




fig 10

This concludes the proof of Lemma III.5 and we return to the proof of the theorem.

We now have :

$$\begin{aligned} K(\mathcal{F}) + K(\mathcal{F}^\perp) &= \frac{1}{10l} \int_G |\mu| (|g|\mathcal{F}; P) + \frac{1}{10l} \int_G |\mu| (|g|\mathcal{F}^\perp; P) \\ &= \frac{1}{10l} \int_G [|\mu| (|g|\mathcal{F}; P) + |\mu| (|g|\mathcal{F}^\perp; P)] \end{aligned}$$

and then

$$K(\mathcal{F}) + K(\mathcal{F}^\perp) \geq \frac{1}{10l} \int_G 1 = \frac{1}{10l} m(\tilde{G}).$$

The theorem then follows from the two following lemmas :

II.6. lemma : The area of a fundamental domain of M is $2\Pi \cdot |\chi(M)|$

Proof : It's an immediate consequence of the Gauss-Bonnet theorem applied either to M or to a fundamental domain F_g .

II.7. Lemma : $m(\tilde{G}) = 4\Pi^2 \cdot |\chi(M)|$.

According to the above remark, the measure in $\tilde{G}$ can be viewed as the measure over segments of fixed length in H^2 (or in M). The middle point of a segment varies in F_g (or in M) whose area is $2\Pi \cdot |\chi(M)|$. The direction of a segment varies in a segment $[0; \Pi]$. Then the measure of all segments of fixed length is $2\Pi^2 \cdot |\chi(M)|$, and so $m(\tilde{G}) = 4\Pi^2 \cdot |\chi(M)|$.

IV. FOLIATIONS AND ENVELOPES

In this paragraph we turn our attention to the following question : suppose we are given in S^n a n -plane field, $\mathcal{P} : S^n \rightarrow \mathcal{A}_{n,n+1}$. We want to calculate a lower bound for $K(\mathcal{F})$, where $\mathcal{F}$ is a orientable codimension one foliation of the unit ball B^n , with isolated singularities, which extends $\mathcal{P}$, that is, if $x \in S^n$ such that

$$\mathcal{P}(x) = T_x \mathcal{F}_x.$$

First of all note that such a foliation may not exist. As it will be clear in the following, even in this case our theorem make sense, as a result over n -plane fields that extend $\mathcal{P}$.

We will suppose that the envelope determined by $\mathcal{P}$ is a herisson with support function $h : S^n \rightarrow \mathbb{R}$. By this we mean that H is the solution of the system :

$$\begin{cases} \langle x; z \rangle &= h(z) \\ \langle x; dz \rangle &= dh(z) \end{cases}, \quad z \in S^n$$

The symetric part of the support function h , is

$$h^*(u) = \frac{1}{2}(h(u) + h(-u)), u \in S^n$$

h^* doesn't depend of the origin in $\mathbb{R}^n$.

We would like to remark that the notion of an herisson was introduced by R. Langevin, G. Levitt, and H. Rosenberg in [L.L.R], and this paper is the reference to the reader who wants to have more details on it.

We are now able to state our theorem.

IV.1. Theorem : Let $\mathcal{P} : S^n \rightarrow \mathcal{A}_{n,n+1}$ be a n -plane field given along S^n , and $\mathcal{F}$ be a orientable codimension one foliation of the unit ball $B^n \subset \mathbb{R}^{n+1}$ which has isolated singularities and extends $\mathcal{P}$. Suppose the envelope determined by $\mathcal{P}$ is a herisson with support function h .

Then :

$$K(\mathcal{F}) \geq \int_{S^n} |h^*(u)| du$$

Proof : with the notations of the first paragraph, we have

$$K(\mathcal{F}) = \frac{1}{2} \int_{S^n} \int_{L(u)} |\mu| (\mathcal{F}; T_i) dt du$$

We will prove that

$$\text{IV.2.} \quad \frac{1}{2} \int_{L(u)} |\mu|(\mathcal{F}; T_i) dt \geq |h^*(u)|, \forall u \in S^n$$

and this will give us the inequality of theorem IV.1.

As $\mathcal{H}$ is a herisson, the orthogonal projection $P_{L(u)}|_{\mathcal{H}} : \mathcal{H} \rightarrow L(u)$ has, for almost all $u \in S^n$, exactly two critical points x_1, x_2 . Let's suppose $h(u) = P_{L(u)}(x_1)$, and say that $t_i = P_{L(u)}(x_i), i = 1, 2$, and let $y_i \in S^n$ be the points in S^n where $\rho(y_i) = T_i, i = 1, 2$. (see fig. 11)




fig. 11

We are going to show that for almost all $t \in [t_2, t_1]$ there exists at least one point of contact between T_t and leaves of $\mathcal{F}$. This proves IV.2 above.

As $\mathcal{F}$ is orientable we can define in B^n a kind of "Gauss map" $\gamma_{\mathcal{F}}$ which assigns to each regular point, x , of $\mathcal{F}$, the unitary vector $n(x)$ normal, in x , to $\mathcal{F}_x$. The vector $n(x)$ is determined by the orientation of $\mathcal{F}$.

This function $\gamma_{\mathcal{F}}$ is continuous in $B\text{-}Sing(\mathcal{F})$. We indicate by $\tilde{\gamma}_{\mathcal{F}}$, it's restriction to S^n , and due to the fact that $\mathcal{H}$ is a herisson $\tilde{\gamma}_{\mathcal{F}}$ is injective and hence is a homeomorphism over its image in S^n .

With u fixed in S^n , let $D_t = B^n \cap T_t$ and $S_t = \partial D_t$. If $t \in [t_2, t_1]$ and $T_t \cap Sing(\mathcal{F}) = \emptyset$ then either u or $-u$ belongs to $\gamma_{\mathcal{F}}(D_t)$, as we shall now show.

In fact, either $u \in \tilde{\gamma}_{\mathcal{F}}(S_{t_1})$ or $-u \in \tilde{\gamma}_{\mathcal{F}}(S_{t_1})$.

$P_{L(t_1)}^{-1}([t_2; t_1]) \cap S^n$ is a cylinder whose image by $\tilde{\gamma}_f$ is a cylinder in S^n with boundary $\tilde{\gamma}_f(S_{t_1}) \cup \tilde{\gamma}_f(S_{t_2})$.

$\tilde{\gamma}_f(S_{t_i})$ is an hypersurface in S^n that separates S^n into two hemispheres, one of them containing u , the other containing $-u$.

$\tilde{\gamma}_f(D_i)$ must contain one of the two hemispheres determined by $\tilde{\gamma}_f(S_{t_i})$, then $\gamma_f(D_i)$ must contains u or $-u$.

But this means that there is $x \in$ such that $n(x) = u$ or $n(x) = -u$, and then

$$|\mu|(\mathcal{F}; T_i) \geq 1.$$

This gives us inequality IV.2., and we have finished with theorem IV.1.

Remark 1 : when $n = 1$, the integral $\int_{S^1} |h'(u)| du$ is called the "absolute length" of $\mathcal{H}$. [L.L.R].

Remark 2 : If $\mathcal{F}$ is a foliation of B^{n+1} such that S^n is a leaf, then we have :

$$\mathcal{H} = S^n; h'(u) = 2, \forall u \in S^n, \text{ and } K(\mathcal{F}) \geq \frac{1}{2} \int_{S^1} 2 du = \text{area}(S^1) = \frac{2\pi \frac{n+1}{2}}{\Gamma\left(\frac{n+1}{2}\right)}$$

where Γ is the gamma function.

V. EXCHANGE THEOREM WITH PLANE CURVES

In this paragraph we generalize the Exchange Theorem by "counting contacts" with a regular curve of $\mathbb{R}^2$, instead of line segments. This seems to be very natural, after the constructions in paragraph III, where we used a pentagon to estimate $K(\mathcal{F})$.

We believe it will be possible to improve these ideas in higher dimensions. For example, if $\mathcal{F}$ is a foliation by surfaces of an open set of $\mathbb{R}^3$, we can "count contacts" of leaves of $\mathcal{F}$, with a fixed surface of $\mathbb{R}^3$ which will roll over the leaves by the action of the isometries of $\mathbb{R}^3$, and obtain integral formulas relating the total curvature of $\mathcal{F}$, the principal curvatures of its leaves and the principal curvatures of the surface.

In this paragraph W will be an open set in $\mathbb{R}^2$, with $\bar{W}$ compact, $\mathcal{F}$ will be a foliation of W , with isolated singularities and C will be a regular curve fixed in $\mathbb{R}^2$, parametrized by arc length.

We define $\varphi : W \times C \rightarrow G$ by $\varphi(x; y; s) = g \Leftrightarrow g$ is an isometry of $\mathbb{R}^2$ wich sends $s \in C$ to $(x, y) \in W$ in such a way that $\mathcal{F}_{(x, y)}$ and $g(C)$ are tangent in $(x; y)$ and have the same orientation.

Lemma : The jacobian of the map φ satisfies :

$$|\text{Jac } \varphi_{(x, y, s)}| = |K_{\mathcal{F}}(x, y) - K_g(s)|$$

where $K_f(x, y)$ is the curvature of the leaf $\mathcal{F}_{(x, y)}$ at the point (x, y) and $K_g(s)$ is the curvature of $g(C)$ at s , both considered with sign.

We recall that the space G of all orientation preserving isometries of $\mathbb{R}^2$ is the cylinder $\mathbb{R}^2 \times S^1$, and that in G we have a canonical measure given by the density

$$m = dx dy d\theta$$

Given $(x, y, s) \in W \times C$ there are exactly two isometries $g_i, i = 1, 2$ which send s to (x, y) in such a way that the leaf $\mathcal{F}_{(x, y)}$ is tangent in (x, y) to the curve $g_i(C), i = 1, 2$. These isometries differ by a rotation of angle Π . (see fig. 12).




fig. 12

Proof of the Lemma : We consider two orthogonal basis :

- $[e_1, e_2, C'(s)]$ in the tangent space to $W \times C$ in (x, y, s) where $[e_1, e_2]$ is the canonical basis of $\mathbb{R}^2$.

- $[e_1, e_2, f]$ is the tangent space to $\tilde{G} = \mathbb{R}^2 \times S^1$ where $[e_1, e_2]$ is the canonical basis of $\mathbb{R}^2$ and f is the unitary vector tangent to the family of isometry classes in $\tilde{G}$ which roll C over $\mathcal{F}_{(x, y)}$ without sliding in the positive sense in $\mathcal{F}_{(x, y)}$. When C rolls over $\mathcal{F}_{(x, y)}$ without sliding then the point of contact between $g(C)$ and $\mathcal{F}$ has, at each moment, velocity of translation equal to zero and component of rotation equal to $K_f - K_g$. Then :

$$\begin{aligned} |\text{Jac } \varphi_{(x, y, s)}| &= \left| \det \begin{bmatrix} 1 & 0 & * \\ 0 & 1 & * \\ 0 & 0 & K_f(x, y) - K_g(s) \end{bmatrix} \right| = \\ &= |K_f(x, y) - K_g(s)| \end{aligned}$$

The exchange theorem with curves follows directly from this lemma :

Theorem : If W is an open set in $\mathbb{R}^2$, with $\bar{W}$ compact, $\mathcal{F}$ is a foliation of W by curves with isolated singularities, and C is a regular C in $\mathbb{R}^2$, parametrized by arc length, then :

$$\frac{1}{2} \int_G |\mu|(\mathcal{F}; g(C)) \, dm(g) = \int_{W \times C} |K_{\mathcal{F}}(x, y) - K_g(s)| \, dx \, dy \, ds,$$

Where : $- |\mu|(\mathcal{F}; g(C))$ is the number of contact points between leaves of $\mathcal{F}$ and $g(C)$.

- $K_{\mathcal{F}}(x, y)$ denotes the curvature of the leaf of $\mathcal{F}$ through (x, y) at (x, y) .

- $K_g(s)$ denotes the curvature of $g(C)$ at s .

(both curvatures take orientations into account).

Proof :

$$\frac{1}{2} \int_G |\mu|(\mathcal{F}; g(C)) \, dm(g) = \int_{W \times C} |K_{\mathcal{F}}(x, y) - K_g(s)| \, dx \, dy \, ds$$

Remark : An analogous result is valid in any surface of constant curvature.

Example : We consider $W =]0, 1[\times]0, 1[\subset \mathbb{R}^2$, $\mathcal{F}$ the foliation of W by horizontal lines and $C = S(r) = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = r^2\}$ the circle of radius r in $\mathbb{R}^2$, $0 < r < \frac{1}{2}$, parametrized by arc length.

Then $K_{\mathcal{F}(x, y)} = 0$ and $|K_g(s)| = \frac{1}{r}$, and

$$\int_{W \times C} |K_{\mathcal{F}(x, y)} - K_g(s)| = \text{area}(W) \cdot \int_C \frac{1}{r} = 1 \cdot \int_0^{2\pi} \frac{1}{r} \cdot r \cdot d\theta = 2\pi$$

Each isometry g of $\mathbb{R}^2$ is determined by the image of the origin, or in other words by the centre of $g(C)$, and by an angle $\theta \in S^1$ (its component of rotation).

If the centre of $g(C)$ lies in the rectangle $]0, 1[\times]r, 1 - r[$ then $|\mu|(\mathcal{F}; g(C)) = 2$, if the centre lies in $]0, 1[\times [-r, r] \cup]0, 1[\times [1 - r, 1 + r]$ then $|\mu|(\mathcal{F}; g(C)) = 1$, and $|\mu|(\mathcal{F}; g(C)) = 0$ otherwise (see fig. 13).

It follows then that

$$\int_G |\mu|(\mathcal{F}; g(C)) = 2\pi[2 \cdot (1 - 2r) + 2r + 2r] = 4\pi$$




fig. 13

REFERENCES

[B.L.R] F. Brito, R. Langevin, H. Rosenberg : "Intégrales de courbures sur des variétés feuilletées". J. Diff. Geom. 16 (1981), 19-50.

[L] R. Langevin : "Feuilletages Tendus". Bull. Soc. Math. France, 107 (1979) 271-281.

[L.L.1] R. Langevin, G. Levitt : "Courbure totale des feuilletages des surfaces". Comment. Math. Helvetici 57 (1982) 175-195.

[L.L.2] R. Langevin, G. Levitt : "Sur la courbure totale des feuilletages des surfaces a bord". Bol. Soc. Bras. Mat. 16 (1985) 1-13.

[L.L.R] R. Langevin, G. Levitt, H. Rosenberg : "Hérissons et multihérissons (enveloppes paramétrées par leur application de Gauss)". Banach Center Publ. 20 (1988) 245-253.

[P] C. Possani : "Curvatura total de Folheações" Thesis IME.USP. 1989.

[S] L.A. Santalo : "Integral Geometry and Geometric, Probabilty". Encyclopedia of Mathematics and its Applications. Addison-Wesley Pub. Co. 1976.

TRABALHOS DO DEPARTAMENTO DE MATEMATICA

TITULOS PUBLICADOS

- 80-01 PLETCH, A. Local freeness of profinite groups. 10 p.
- 80-02 PLETCH, A. Strong completeness in profinite groups. 8 p.
- 80-03 CARNIELLI, W.A. & ALCANTARA, L.P. de Transfinite induction on ordinal configurations. 22 p.
- 80-04 JONES RODRIGUES, A.R. Integral representations of cyclic p-groups. 13 p.
- 80-05 CORRADA, M. & ALCANTARA, L.P. de Notes on many-sorted systems. 25 p.
- 80-06 POLCINO MILIES, F.C. & SEHGAL, S.K. FC-elements in a group ring. 10 p.
- 80-07 CHEN, C.C. On the Ricci condition and minimal surfaces with constantly curved Gauss map. 10 p.
- 80-08 CHEN, C.C. Total curvature and topological structure of complete minimal surfaces. 21 p.
- 80-09 CHEN, C.C. On the image of the generalized Gauss map of a complete minimal surface in R^4 . 8 p.
- 81-10 JONES RODRIGUEZ, A.R. Units of ZC_p . 7 p.
- 81-11 KOTAS, J. & COSTA, N.C.A. da Problems of model and discursive logics. 35 p.
- 81-12 BRITO, F.B. & GONÇALVES, D.L. Algebras não associativas, sistemas diferenciais polinomiais homogêneos e classes características. 7 p.
- 81-13 POLCINO MILIES, F.C. Group rings whose torsion units form a subgroup II. Iv. (não paginado).
- 81-14 CHEN, C.C. An elementary proof of Calabi's theorems on holomorphic curves. 5 p.
- 81-15 COSTA, N.C.A. da & ALVES, E.H. Relations between paraconsistent logic and many-valued logic. 8 p.
- 81-16 CASTILLA, M.S.A.C. On Przymusiński's theorem. 6 p.
- 81-17 CHEN, C.C. & GOES, C.C. Degenerate minimal surfaces in R^4 . 21 p.
- 81-18 CASTILLA, M.S.A.C. Imagens inversas de algumas aplicações fechadas. 11 p.
- 81-19 ARAGONA VALLEJO, A.J. & EXEL FILHO, R. An infinite dimensional version of Hartogs' extension theorem. 9 p.
- 81-20 GONÇALVES, J.Z. Groups rings with solvable unit groups. 15p.
- 81-21 CARNIELLI, W.A. & ALCANTARA, L.P. de Paraconsistent algebras. 16 p.
- 81-22 GONÇALVES, D.L. Nilpotent actions. 10 p.
- 81-23 COELHO, S.P. Group rings with units of bounded exponent over the center. 25 p.
- 81-24 PARMENTER, M.M. & POLCINO MILIES, F.C. A note on isomorphic group rings. 4 p.
- 81-25 MERKLEN GOLDSCHMIDT, H.A. Hereditary algebras with maximum spectra are of finite type. 10 p.
- 81-26 POLCINO MILIES, F.C. Units of group rings: a short survey. 32 p.

- 84-76 GONÇALVES, J.Z. Normal and subnormal subgroups in the group of units of a group rings. 13 p.
- 85-77 ARAGONA VALLEJO, A.J. On existence theorems for the ∂ -operator on generalized differential forms. 13 p.
- 85-78 POLCINO MILIES, C.; RITTER, J. & SEHGAL, S.K. On a conjecture of Zassenhaus on torsion units in integral group rings II. 14 p.
- 85-79 JONES RODRIGUEZ, A.R. & MICHLER, G.O. On the structure of the integral Green ring of a cyclic group of order p^2 . The Jacobson radical of the integral Green ring of a cyclic group of order p^2 . 26 p.
- 85-80 VELOSO, J.M.M. & VERDERESI, J.A. Three dimensional Cauchy-Riemann manifolds. 19 p.
- 85-81 PERESI, L.A. On baric algebras with prescribed automorphisms II. 18 p.
- 85-82 KNUDSEN, C.A. O impasse aritmo-geométrico e a evolução do conceito de número na Grécia antiga. 43p.
- 85-83 VELOSO, J.M.M. & VERDERESI, J.A. La géométrie, le problème d'équivalence et la classification des CR-varietés homogenes en dimension 3. 30 p.
- 85-84 GONÇALVES, J.Z. Integral group rings whose group is solvable, an elementary proof. 11 p.
- 85-85 LUCIANO, O.O. Nebuleuses infinitesimélement fibrées. 5 p.
- 85-86 ASPERTI, A.C. & DAJCZER, M. Conformally flat Riemannian manifolds as hypersurfaces of the lighth cone. 3 p.
- 85-87 BELOQUI, J.A. A quasi-transversal Hopf bifurcations. 11 p.
- 85-88 POLCINO MILIES, F.C. & RAPHAEL, D.M. A note on derivations with power central values in prime rings. 7 p.
- 85-89 POLCINO MILIES, F.C. Torsion units in group rings and a conjecture of H.J.Zassenhaus. 14 p.
- 86-90 LOURENÇO, M.L. Riemann domains over (DFC) spaces. 32 p.
- 86-91 ARAGONA VALLEJO, A.J. & FERNANDES, J.C.D. The Hartogs extension theorem for holomorphic generalized functions. 9 p.
- 86-92 CARRARA ZANETIC, V.L. Extensions of immersions in dimension two. 27 p.
- 86-93 PERESI, L.A. The derivation algebra of gametic and zygotic algebras for linked loci. 25 p.
- 86-94 COELHO, S.P. A note on central idempotents in group ring. 5 p.
- 86-95 PERESI, L.A. On derivations of baric algebras with prescribed automorphisms. 21 p.
- 86-96 COELHO, F.U. A generalization of a theorem of Todorov on preprojectives partitions. 18 p.
- 86-97 ASPERTI, A.C. A note on the minimal immersions of the two-sphere. 11 p.
- 86-98 COELHO, S.P. & POLCINO MILIES, F.C. A note on central idempotents in group rings II. 8 p.

- 86-99 EXEL FILHO, R. Hankel matrices over right ordered amenable groups. 18 p.

NOVA SERIE

- 86-01 GOODAIRE, E.G. & POLCINO MILIES, F.C. Isomorphisms of integral alternative loop rings. 11 p.
- 86-02 WALCZAK, P.G. Foliations which admit the most mean curvature functions. 11 p.
- 86-03 OLIVEIRA, M.E.G.G. Minimal Klein bottles with one end in R_3 and R_4 . 12 p.
- 86-04 MICALI, A. & VILLAMAYOR, O.E. Homologie de Hochschild de certaines algebres de groupes. 11 p.
- 86-05 OLIVEIRA, M.E.G.G. Minimal Klein bottles in R_3 with finite total curvature. 9 p.
- 86-06 CARRARA ZANETIC, V.L. Classification of stable maps between 2-manifolds with given singular set image. 22 p.
- 87-01 BRITO, F.G.B. & WALCZAK, P.G. Total curvature of orthogonal vector fields on three-manifolds. 4 p.
- 87-02 BRITO, F.G.B. & LEITE, M.L. A remark on rotational hypersurfaces of S_n . 13 p.
- 87-03 GONÇALVES, J., RITTER, J. & SEHGAL, S. Subnormal subgroups in $U(ZG)$. 13 p.
- 87-04 ARAGONA VALLEJO, A.J. & COLOMBEAU, J.F. The interpolation theorem for holomorphic generalized functions. 12 p.
- 87-05 ALMEIDA, S.C. de & BRITO, F.G.B. Immersed hypersurfaces of a space form with distinct principal curvatures. 9 p.
- 87-06 ASPERTI, A.C. Generic minimal surfaces. 21 p.
- 87-07 GOODAIRE, E.G. & POLCINO MILIES, F.C. Torsion units in alternative group rings. 17 p.
- 87-08 MERKLEN GOLDSCHMIDT, H.A. Algebras which are equivalent to a hereditary algebra modulo preprojectives II. 27 p.
- 87-09 REYNOL FILHO, A. P-localization of some classes of groups. 30 p.
- 88-01 CARLSON, J.F. & JONES, A. An exponential property of lattices over group rings. 22 p.
- 88-02 CARLSON, J.F. & JONES, A. Wild categories of periodic modules. 6 p.
- 88-03 SALLUM, E.M. The nonwandering set of flows on a Reeb foliation. 13 p.
- 88-04 ALMEIDA, R. Cohomologie des suites d'Atiyah. 14 p.
- 88-05 GUZZO JR., H. Alguns teoremas de caracterização para álgebras alternativas à direita. 16 p.
- 88-06 HARLE, C.E. Subvariedades isoparamétricas homogêneas. 5 p.

- 81-27 CHEN, C.C. & GACKSTATTER, F. Elliptic and hyper-elliptic functions and complete minimal surfaces with handles. 14 p.
- 81-28 POLCINO MILIES, F.C. A glance at the early history of group rings. 22 p.
- 81-29 FERRER SANTOS, W.R. Reductive actions of algebraic groups on affine varieties. 52 p.
- 81-30 COSTA, N.C.A. da The philosophical import of paraconsistent logic. 26 p.
- 81-31 GONÇALVES, D.L. Generalized classes of groups, spaces c-nilpotent and "the Hurewicz theorem". 30 p.
- 81-32 COSTA, N.C.A. da & MORTENSEN, Chris. Notes on the theory of variable binding term operators. 18 p.
- 81-33 MERKLEN GOLDSCHMIDT, H.A. Homogenes 1-hereditary algebras with maximum spectra. 32 p.
- 81-34 PERESI, L.A. A note on semiprime generalized alternative algebras. 10 p.
- 81-35 MIRAGLIA NETO, F. On the preservation of elementary equivalence and embedding by filtered powers and structures of stable continuous functions. 9 p.
- 81-36 FIGUEIREDO, G.V.R. Catastrophe theory: some global theory a full proof. 91 p.
- 82-37 COSTA, R.C.F. On the derivations of gametic algebras. 17 p.
- 82-38 FIGUEIREDO, G.V.R. A shorter proof of the Thom-Zeeman global theorem for catastrophes of cod ≤ 5 . 7 p.
- 82-39 VELOSO, J.M.M. Lie equations and Lie algebras: the intransitive case. 97 p.
- 82-40 GOES, C.C. Some results about minimal immersions having flat normal bundle. 37 p.
- 82-41 FERRER SANTOS, W.R. Cohomology of comodules II. 15 p.
- 82-42 SOUZA, V.H.G. Classification of closed sets and diffeos of one-dimensional manifolds. 15 p.
- 82-43 GOES, C.C. The stability of minimal cones of codimension greater than one in R^n . 27 p.
- 82-44 PERESI, L.A. On automorphisms of gametic algebras. 27 p.
- 82-45 POLCINO MILIES, F.C. & SEHGAL, S.K. Torsion units in integral group rings of metacyclic groups. 18 p.
- 82-46 GONÇALVES, J.Z. Free subgroups of units in group rings. 8 p.
- 82-47 VELOSO, J.M.M. New classes of intransitive simple Lie pseudogroups. 8 p.
- 82-48 CHEN, C.C. The generalized curvature ellipses and minimal surfaces. 10 p.
- 82-49 COSTA, R.C.F. On the derivation algebra of zygotical algebras for polyploidy with multiple alleles. 24 p.
- 83-50 GONÇALVES, J.Z. Free subgroups in the group of units of group rings over algebraic integers. 3 p.

- 83-51 MANDEL, A. & GONÇALVES, J.Z. Free k -triples in linear groups. 7 p.
- 83-52 BRITO, F.G.B. A remark on closed minimal hypersurfaces of S^4 with second fundamental form of constant length. 12 p.
- 83-53 KIIHL, J.C.S. U-structures and sphere bundles. 8 p.
- 83-54 COSTA, R.C.F. On genetic algebras with prescribed derivations. 23 p.
- 83-55 SALVITTI, R. Integrabilidade das distribuições dadas por subálgebras de Lie de codimensão finita no $gh(n, C)$. 4 p.
- 83-56 MANDEL, A. & GONÇALVES, J.Z. Construction of open sets of free k -Tuples of matrices. 18 p.
- 83-57 BRITO, F.G.B. A remark on minimal foliations of codimension two. 24 p.
- 83-58 GONÇALVES, J.Z. Free groups in subnormal subgroups and the residual nilpotence of the group of units of group rings. 9 p.
- 83-59 BELOQUI, J.A. Modulus of stability for vector fields on 3-manifolds. 40 p.
- 83-60 GONÇALVES, J.Z. Some groups not subnormal in the group of units of its integral group ring. 8 p.
- 84-61 GOES, C.C. & SIMOES, P.A.Q. Imersões mínimas nos espaços hiperbólicos. 15 p.
- 84-62 GIANBRUNO, A.; MISSO, P. & POLCINO MILIES, F.C. Derivations with invertible values in rings with involution. 12 p.
- 84-63 FERRER SANTOS, W.R. A note on affine quotients. 6 p.
- 84-64 GONÇALVES, J.Z. Free-subgroups and the residual nilpotence of the group of units of modular and p -adic group rings. 12 p.
- 84-65 GONÇALVES, D.L. Fixed points of S^1 -fibrations. 18 p.
- 84-66 RODRIGUES, A.A.M. Contact and equivalence of submanifolds of homogenous spaces. 15 p.
- 84-67 LOURENÇO, M.L. A projective limit representation of (DFC)-spaces with the approximation property. 20 p.
- 84-68 FORNARI, S. Total absolute curvature of surfaces with boundary. 25 p.
- 84-69 BRITO, F.G.B. & WALCZAK, P.G. Totally geodesic foliations with integral normal bundles. 6 p.
- 84-70 LANGEVIN, R. & POSSANI, C. Quase-folhações e integrais de curvatura no plano. 26 p.
- 84-71 OLIVEIRA, M.E.G.G. de Non-orientable minimal surfaces in RN . 41 p.
- 84-72 PERESI, L.A. On baric algebras with prescribed automorphisms. 42 p.
- 84-73 MIRAGLIA NETO, F. & ROCHA FILHO, G.C. The measurability of Riemann integrable-function with values in Banach spaces and applications. 27 p.
- 84-74 MERKLEN GOLDSCHMIDT, H.A. Artin algebras which are equivalent to a hereditary algebra modulo pre-projectives. 38 p.
- 84-75 GOES, C.C. & SIMOES, P.A.Q. The generalized Gauss map of minimal surfaces in H^3 and H^4 . 16 p.

- 88-07 LANGEVIN, R. Vers une classification des difféomorphismes Morse-Smale d'une surface. 18 p.
- 88-08 ARAGONA VALLEJO, A. J. & BIAGIONI, H. A. An intrinsic definition of the Colombeau algebra of generalized functions. 48 p.
- 88-09 BORSARI, H.D. A Cohomological characterization of reductive algebraic groups. 20 p.
- 88-10 COELHO, S.P. & POLCINO MILIES, F.C. Finite conjugacy in group rings. 20 p.
- 88-11 FERNANDEZ, R. Characterization of the dual of an Orlicz Space. 19 p.
- 89-01 GONÇALVES, J.Z. & MANDEL, A. Embedding the Free Group Ring into Formal Series. 4 p.
- 89-02 REYNOLDS, A. Nilpotent spaces: some inequalities on nilpotency degrees. 17 p.
- 89-03 GOUVEA, F.Q. Deforming Galois Representations: Controlling the Conductor. 22 p.
- 89-04 CORDARD, P. & TREVES, F. Homology & Cohomology in Hypo-Analytic Structures of the Hypersurfaces Type. 108p.
- 89-05 ARAGONA VALLEJO, A. J. & VILLARREAL ALVARADO, F. Colombeau's theory and shock Waves in a problem of Hydrodynamics. 48 p.
- 90-01 MANDEL, A. & GONÇALVES, J.Z. Semigroup laws and free semigroups in the group of units of group rings. 11 p.
- 90-02 LINTZ, R.G. Generation of Quantum-fields by non-deterministic fields. 32p.
- 90-03 LINTZ, R.G. A critical study on the foundations of Geometry. 41p.
- 90-04 LINTZ, R.G. The theory of manifolds from the non-deterministic point of view. 32p.
- 90-05 GUIDORIZZI, H.L. Oscillating and Periodic solutions of Liénard equations. 18p.
- 90-06 BIANCONI, R. Model Completeness Results for Elliptic and Abelian Functions. 21p.
- 90-07 LEAL, G. & MILIES, C.P. Isomorphic Group (and Loop) Algebras. 21p.
- 90-08 LANGEVIN, R. & POSSANI, C. Total curvature of foliations. 18p.