

Pacific Journal of Mathematics

**LIGHT GROUPS OF ISOMORPHISMS OF BANACH SPACES
AND INVARIANT LUR RENORMINGS**

LEANDRO ANTUNES, VALENTIN FERENCZI,
SOPHIE GRIVAUX AND CHRISTIAN ROSENDAL

LIGHT GROUPS OF ISOMORPHISMS OF BANACH SPACES AND INVARIANT LUR RENORMINGS

LEANDRO ANTUNES, VALENTIN FERENCZI,
SOPHIE GRIVAUX AND CHRISTIAN ROSENDAL

Megrelishvili (2001) defines *light groups* of isomorphisms of a Banach space as the groups on which the weak and strong operator topologies coincide, and proves that every bounded group of isomorphisms of Banach spaces with the point of continuity property (PCP) is light. We investigate this concept for isomorphism groups G of classical Banach spaces X without the PCP, specially isometry groups, and relate it to the existence of G -invariant LUR or strictly convex renormings of X .

1. Introduction

The general objective of this note is to determine conditions on a bounded group of isomorphisms of Banach spaces that ensure the existence of a locally uniformly rotund (LUR) renorming invariant under the action of this group. In particular, we are interested in this context in the notion of lightness for such groups.

Light groups. A frequent problem in functional analysis is to determine under which conditions weak convergence and norm convergence coincide. For example, it is well known that conditions of convexity of the norm of a Banach space ensure that weak and strong convergence are equivalent on its unit sphere. The corresponding problem for isomorphisms of Banach spaces (or more generally of locally convex spaces) was studied by Megrelishvili [2001] in the context of group representations, using the concept of fragmentability.

Given a (real) Banach space X , we denote by $L(X)$ the set of bounded linear operators on X , and by $GL(X)$ the group of bounded isomorphisms of X . We

Antunes was fully supported by UTFPR (Process 23064.004102/2015-40). Ferenczi was supported by CNPq, grant 303034/2015-7 and Fapesp, grants 2013/11390-4 and 2016/25574-8. Grivaux was supported in part by the project FRONT of the French National Research Agency (ANR-17-CE40-0021) and by the Labex CEMPI (ANR-11-LABX-0007-01). Rosendal was supported by the NSF (DMS #1464974).

MSC2010: 22F50, 46B03.

Keywords: groups of isomorphisms of Banach spaces, isometry groups, LUR renormings, renormings of Banach spaces, light groups, invariant renormings.

also denote by $\text{Isom}(X)$ the group of surjective linear isometries of X . If G is a subgroup of $\text{GL}(X)$, we write $G \leq \text{GL}(X)$. Recall that given a Banach space X , the strong operator topology on $L(X)$ is the topology of pointwise convergence, i.e., the initial topology generated by the family of functions $f_x : L(X) \rightarrow X$, $x \in X$, given by $f_x(T) = Tx$, $T \in L(X)$, and the weak operator topology on $L(X)$ is generated by the family of functions $f_{x,x^*} : L(X) \rightarrow \mathbb{R}$, $x \in X$, $x^* \in X^*$, given by $f_{x,x^*}(T) = x^*(Tx)$, $T \in L(X)$.

Definition 1.1 [Megrelishvili 2001]. A group $G \leq \text{GL}(X)$ of isomorphisms on a Banach space X is *light* if the weak operator topology (WOT) and the strong operator topology (SOT) coincide on G .

Observe that since the two operator topologies are independent of the specific choice of norm on X , the same holds for lightness of G .

Well-known examples of light groups are the groups $U(H)$ of unitary operators on Hilbert spaces H . However, the main result of [Megrelishvili 2001] concerning light groups indicates that we have a similar phenomenon in a more general context. Recall that a Banach space X has the point of continuity property (PCP) if for every norm-closed nonempty bounded subset C of X , the identity on C has a point of continuity from the weak to the norm topology:

Theorem 1.2 [Megrelishvili 2001]. *If X is a Banach space with the point of continuity property (PCP) and if $G \leq \text{GL}(X)$ is bounded in norm, then G is light.*

In particular, if X has the Radon–Nikodym property (RNP) (e.g., if X is reflexive or is a separable dual space), then every bounded subgroup of $\text{GL}(X)$ is light. For example, the isometry group of ℓ_1 , $\text{Isom}(\ell_1)$, is light.

We note here that in the literature (and indeed in [Megrelishvili 2001]) PCP sometimes appears as the formally weaker condition “*every weakly-closed nonempty bounded subset has a weak-to-norm point of continuity for the identity*”. However, as was pointed out to us by G. Godefroy, if X satisfies this definition and F is norm-closed and bounded, then any point of continuity of the weak closure \bar{F}^w belongs to F , so the two definitions are equivalent. In fact, if $x \in \bar{F}^w$ is a weak-to-norm point of continuity for the identity, there exists a net $(x_\alpha)_{\alpha \in I} \subset F$ such that $x_\alpha \xrightarrow{w} x$. Hence, $x_\alpha \xrightarrow{\|\cdot\|} x$ and, since F is norm-closed, $x \in F$.

Bounded nonlight groups. A natural question that arises from Megrelishvili’s result is to investigate in which respect his result is optimal, and whether “smallness” assumptions on G or weaker assumptions than the PCP on X could imply that G is light. We show (Theorem 4.6) that any separable space containing an isomorphic copy of c_0 admits a bounded cyclic group of isomorphisms which is not light. This shows that we cannot really expect further general results in this direction.

Megrelishvili gives the group $\text{Isom}(C([-1, 1]^2))$ as an example of a nonlight group. His proof uses a construction of Helmer [1980] of a separately continuous group action on $[-1, 1]^2$ that is not jointly continuous, and the equivalence of pointwise compactness and weak compactness of bounded subsets of $C(K)$. This leads to the following question:

Question 1.3. For which compact sets K is the isometry group $\text{Isom}(C(K))$ not light?

We first prove (Proposition 4.3) that the isometry group of c , the space of real convergent sequences, is not light. Neither is the isometry group of $C(\{0, 1\}^{\mathbb{N}})$ (Corollary 5.6). On the other hand, as a consequence of Theorem 5.9, we show that the isometry group of $C[0, 1]$ is light, while those of the spaces $C([0, 1]^n)$, $n \geq 2$, are not light. These constructions simplify the initial example of Megrelishvili.

Light groups and LUR renormings. In another direction, we study the relation between light groups and the existence of LUR renormings invariant under the action of the group. Recall that a norm $\|\cdot\|$ on X is *rotund* or *strictly convex* if whenever the vectors x, y belong to the unit sphere S_X of X and $\|x + y\| = 2$, $x = y$. It is *locally uniformly convex* (LUC) or *locally uniformly rotund* (LUR) at a vector $x_0 \in X$ if whenever $(x_n)_{n \in \mathbb{N}}$ is a sequence of vectors of X such that $\lim \|x_n\| = \|x_0\|$ and $\lim \|x_0 + x_n\| = 2\|x_0\|$, $\lim \|x_n - x_0\| = 0$. Another equivalent definition (in fact, Lovaglia's original definition) is the following: the norm is LUR at a vector $x_0 \in S_X$ if for every $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon, x_0) > 0$ such that

$$\frac{\|x + y\|}{2} \leq 1 - \delta \quad \text{whenever } \|x - y\| \geq \varepsilon \text{ and } \|y\| = 1.$$

The norm is said to be LUR in X if it is LUR at every point $x_0 \neq 0$ of X (or, equivalently, of S_X). The property of the dual norm $\|\cdot\|_*$ on X^* being strictly convex or LUR is closely related to the differentiability of the norm $\|\cdot\|$ on X , in the senses of Fréchet and Gateaux, respectively. All (real) separable Banach spaces admit an equivalent LUR renorming. For this and much more on renormings of Banach spaces, see [Deville et al. 1993].

A fundamental result in the study of LUR renorming is the following theorem:

Theorem 1.4 [Lancien 1993]. *If X is a separable Banach space with the RNP, X admits an isometry invariant LUR renorming.*

If $G \leq \text{GL}(X, \|\cdot\|)$ is a bounded group of isomorphisms on X , the norm on X defined by

$$|||x||| = \sup_{g \in G} \|gx\|, \quad x \in X,$$

is a G -invariant renorming of X . In other words, $G \leq \text{Isom}(X, |||\cdot|||)$. So a consequence of Theorem 1.4 is that whenever X is a separable space with the RNP

and G is a bounded group of isomorphisms on X , there exists a G -invariant LUR renorming of X . The existence of G -invariant LUR renormings for general groups of isomorphisms G was first investigated by Ferenczi and Rosendal [2013]. They studied problems of maximal symmetry in Banach spaces, analyzing the structure of subgroups of $\text{GL}(X)$ when X is a separable reflexive Banach space. An example of a super-reflexive space with no maximal bounded group of isomorphisms was also exhibited in [Ferenczi and Rosendal 2013].

The relation between light groups and G -invariant LUR renormings is given by Theorem 2.3. We observe that if a Banach space X admits a G -invariant LUR renorming, then G is light. In fact, this is true even if the norm is LUR only on a dense subset of S_X . We also show that the converse assertion is false: although the isometry group of $C[0, 1]$ is light, $C[0, 1]$ admits no strictly convex isometry invariant renorming (Corollary 5.12). This link between the existence of a G -invariant LUR renorming and the lightness of G is a natural one: if X is a Banach space with an LUR norm $\|\cdot\|$, the weak topology and the norm topology coincide on the unit sphere of $(X, \|\cdot\|)$.

Light groups and distinguished families. Ferenczi and Galego [2010] investigated groups that may be seen as the group of isometries of a Banach space under some renorming. Among other results, they proved that if X is a separable Banach space and G is a finite group of isomorphisms of X such that $-\text{Id} \in G$, X admits an equivalent norm $\|\cdot\|$ such that $G = \text{Isom}(X, \|\cdot\|)$. They also proved that if X is a separable Banach space with LUR norm $\|\cdot\|$ and if G is an infinite countable bounded isometry group of X such that $-\text{Id} \in G$ and such that G admits a point $x \in X$ with $\inf_{g \neq \text{Id}} \|gx - x\| > 0$, then $G = \text{Isom}(X, \|\cdot\|)$ for some equivalent norm $\|\cdot\|$ on X . In [Ferenczi and Rosendal 2011], a point x satisfying the condition

$$\inf_{g \neq \text{Id}} \|gx - x\| > 0$$

is called a *distinguished point* of X for the group G .

Ferenczi and Rosendal [2011] generalized results of [Ferenczi and Galego 2010] to certain uncountable Polish groups, and also defined the concept of a *distinguished family* for the action of a group G on a Banach space X . It is clear that if G is an isometry group with a distinguished point, G is SOT-discrete. However, the following question remained open: if G is an isomorphism group of X which is SOT-discrete, should X have a distinguished point for G ? In Proposition 6.1 we see that the answer to this question is negative, and we give an example of an infinite countable group of isomorphisms G of c_0 which is SOT-discrete but does not admit a distinguished point for G . In addition, this group is also not light.

Light groups on quasinormed spaces. Although Megrelishvili has defined the concept of light group only for groups of isomorphisms on locally convex spaces,

we can extend the definition to quasinormed spaces, even if they are not locally convex. We finish this article by investigating whether the isometry groups of the quasinormed spaces ℓ_p and $L_p[0, 1]$, $0 < p < 1$, are light.

2. LUR renormings and light groups

Let G be a bounded group of isomorphisms on a Banach space $(X, \|\cdot\|)$. In this section we are interested in the existence of a G -invariant LUR renorming of X , i.e., in the existence of an equivalent norm $|||\cdot|||$ on X which is both invariant under the action of G and is LUR; or in the existence of a G -invariant *dense LUR renorming*, meaning a renorming which is invariant under the action of G and is LUR on a dense subset of S_X . When $G = \text{Isom}(X, \|\cdot\|)$ we shall talk about *isometry invariant* renormings. Our first result is the following:

Proposition 2.1. *Let X be a Banach space and let $G \leq \text{GL}(X)$. If G is SOT-compact and if X admits an LUR renorming, X admits a G -invariant LUR renorming.*

Proof. Suppose that $\|\cdot\|$ is an equivalent LUR norm on X . The formula

$$|||x||| = \sup_{T \in G} \|Tx\|, \quad x \in X,$$

defines a G -invariant LUR renorming of X . Indeed, suppose that x_n and x are vectors of X such that $|||x_n||| = |||x||| = 1$ for every $n \in \mathbb{N}$ and $\lim |||x_n + x||| = 2$. Then we can find elements T_n , $n \in \mathbb{N}$, of G such that $\lim \|T_n x_n + T_n x\| = 2$. By SOT-compactness of G we can assume without loss of generality that T_n tends SOT to some element T of G , from which it follows that $\|T_n x_n + T x\|$ converges to 2. Since $\|T_n x_n\| \leq |||x_n||| = 1$ for every $n \in \mathbb{N}$ and $\|T x\| \leq |||x||| = 1$, we deduce that $\|T x\| = 1$ and that $\|T_n x_n\|$ converges to 1. In particular, if we set, for every $n \in \mathbb{N}$, $y_n = T_n x_n / \|T_n x_n\|$, then y_n belongs to the unit sphere of $(X, \|\cdot\|)$ and

$$\|y_n + T x\| = \frac{1}{\|T_n x_n\|} \|T_n x_n + \|T_n x_n\| T x\| \rightarrow 2.$$

By the LUR property of $\|\cdot\|$ at the point $T x$, we deduce that y_n converges to $T x$. So $T_n x_n$ converges to $T x$. So $|||x_n - x||| = |||T_n x_n - T_n x|||$ converges to 0 since both $T_n x_n$ and $T_n x$ converge to $T x$. This shows that $|||\cdot|||$ is LUR. \square

It is also worth mentioning that every SOT-compact group of isomorphisms is light:

Proposition 2.2. *Let $G \leq \text{GL}(X)$ be a group of isomorphisms of a Banach space X . If G is SOT-compact, then G is light.*

Proof. The assumption implies that G is also WOT compact, since the WOT is weaker than the SOT. However, the WOT is also Hausdorff, and so the two topologies must agree on G . In other words, G is light. \square

Ferenczi and Rosendal [2011] investigated LUR renormings in the context of transitivity of norms. Recall that a norm $\|\cdot\|$ on X is called *transitive* if the orbit $\mathcal{O}(x)$ of every point $x \in S_X$ under the action of the isometry group $\text{Isom}(X)$ is the whole sphere S_X . If for every $x \in X$ the orbit $\mathcal{O}(x)$ is dense in S_X , we say that $\|\cdot\|$ is *almost transitive*, and if the closed convex hull of $\mathcal{O}(x)$ is the unit ball B_X , we say that $\|\cdot\|$ is *convex transitive*. Ferenczi and Rosendal proved that if an almost transitive norm on a Banach space is LUR at some point of the unit sphere, it is uniformly convex. They also proved that if a convex transitive norm on a Banach space is LUR on a dense subset of the unit sphere, it is almost transitive and uniformly convex.

In the next theorem, we exhibit a relation between the existence of LUR renormings and light groups.

Theorem 2.3. *Let $G \leq \text{GL}(X)$ be a group of isomorphisms of a Banach space X . If X admits a G -invariant renorming which is LUR on a dense subset of S_X , then G is light.*

Proof. Let $\|\cdot\|$ be a G -invariant renorming of X which is LUR on a dense subset of S_X . Let (T_α) be a net of elements of G which converges WOT to the identity operator Id on X , and assume that T_α does not converge SOT to Id . Let $x \in S_X$ be such that $T_\alpha x$ does not converge to x . Without loss of generality, we can suppose that the norm is LUR at x , and that there exists $\delta > 0$ such that, for every α , $\|T_\alpha x - x\| \geq \delta$. Since $\|T_\alpha x\| = \|x\| = 1$ for every α , the LUR property forbids having $\lim \|T_\alpha x + x\| = 2$. So we can assume that there exists $\varepsilon > 0$ such that $\|T_\alpha x + x\| \leq 2 - \varepsilon$ for every α .

Let $\phi \in X^*$ with $\|\phi\| = 1$ be such that $\phi(x) = 1$. Since T_α converges WOT to Id , $\phi(T_\alpha x) \rightarrow 1$. On the other hand,

$$\begin{aligned} 2 - \delta &\geq \|T_\alpha x + x\| \\ &= \max_{\psi \in X^*, \|\psi\|=1} |\psi(T_\alpha x + x)| \geq |\phi(T_\alpha x) + 1| \quad \text{for every } \alpha, \end{aligned}$$

which contradicts the WOT convergence of T_α to Id . □

Remark 2.4. In fact, the proof of Theorem 2.3 gives us a formally stronger result: if X admits a G -invariant renorming which is LUR on a dense subset of S_X then G is *orbitwise light*. Megrelshvili [2003] defines a group $G \leq \text{GL}(X)$ as orbitwise light (or orbitwise Kadec) if for every $x \in X$ the orbit $\mathcal{O}(x) = \{Tx ; T \in G\}$ is a set on which the weak and the strong topologies coincide. It is readily seen that if G is orbitwise light, then it is light, but whether the converse holds is still an open question.

3. Light groups and distinguished points

As recalled in the introduction, Lancien [1993] proved that if X is separable with the RNP, X admits an isometry invariant LUR renorming. Although separable spaces always admit LUR renormings, the generalization of Lancien's result to all separable spaces is false. For example, if $X = C([-1, 1]^2)$ and $G = \text{Isom}(X)$ then, since G is not light [Megrelishvili 2001], by Theorem 2.3 there is no equivalent G -invariant (not even dense) LUR renorming. Another example mentioned in [Ferenczi and Rosendal 2011] is the case where $X = L_1[0, 1]$ and $G = \text{Isom}(L_1[0, 1])$. In this case there is not even a strictly convex G -invariant renorming.

Here we discuss conditions which clarify the relations between the two properties of a group $G \leq \text{GL}(X)$ being light and X having a G -invariant LUR renorming, in the case when G is SOT-discrete. The following notion was defined in [Ferenczi and Rosendal 2011].

Definition 3.1. Let X be a Banach space, let $G \leq \text{GL}(X)$ be a bounded group of isomorphisms of X , and let $x \in X$. We say that x is *distinguished* for G (or for the action of G on X) if

$$\inf_{T \neq \text{Id}} \|Tx - x\| > 0.$$

If $\{x_1, \dots, x_n\}$ is a finite family of vectors of X , then it is distinguished for G if

$$\inf_{T \neq \text{Id}} \max_{1 \leq i \leq n} \|Tx_i - x_i\| > 0,$$

or, equivalently, if the n -tuple (x_1, \dots, x_n) is distinguished for the canonical action of G on X^n .

This notion does not depend on the choice of an equivalent norm on X . Note also that G is SOT-discrete exactly when it admits a distinguished finite family of vectors. We also have, considering the adjoint action of G on X^* :

Lemma 3.2. Assume that $G \leq \text{GL}(X)$ is light. If G acts as an SOT-discrete group on X , then G acts as an SOT-discrete group on X^* .

Proof. Define $\psi : G \rightarrow \text{GL}(X^*)$ by setting $\psi(T)(x^*) = x^* \circ T^{-1}$ for every $T \in G$ and $x^* \in X^*$. We want to show that $\psi(G)$ is an SOT-discrete subgroup of $\text{GL}(X^*)$. It suffices to show the existence of $\varepsilon > 0$ and $x_1^*, \dots, x_n^* \in S_{X^*}$ such that the only element T of G such that $\|\psi(T)(x_i^*) - x_i^*\| < \varepsilon$ for every $1 \leq i \leq n$ is the identity operator Id_X on X . Since G is light and acts as an SOT-discrete group on X , it is WOT-discrete. So there exist $\varepsilon > 0$, $x_1, \dots, x_m \in S_X$ and $x_1^*, \dots, x_n^* \in S_{X^*}$ such that the only element T of G such that $|x_i^*(T^{-1}x_j - x_j)| < \varepsilon$ for every $1 \leq i \leq n$ and $1 \leq j \leq m$ is $T = \text{Id}_X$. The conclusion follows immediately. \square

Lemma 3.3. Let X be a Banach space, let G be a bounded subgroup of $\text{GL}(X)$, and let $\{x_1, \dots, x_n\}$ be a distinguished family of vectors for the action of G on X .

Let $\|\cdot\|$ be a G -invariant norm on X which is LUR at x_i for every $1 \leq i \leq n$. For any functional $x_i^* \in S_{X^*, \|\cdot\|}^*$ such that $x_i^*(x_i) = \|x_i\|$ for every $1 \leq i \leq n$, the family $\{x_1^*, \dots, x_n^*\}$ is distinguished for the action of G on X^* .

Proof. Assume $\|x_i\| = 1$ for every $1 \leq i \leq n$. Let $\alpha = \inf_{T \neq \text{Id}_X} \max_{1 \leq i \leq n} \|Tx_i - x_i\| > 0$. For every $T \neq \text{Id}_X$, choose $1 \leq i \leq n$ such that $\|T^{-1}x_i - x_i\| \geq \alpha$. By the LUR property of the norm at x_i , there exists $\varepsilon > 0$ depending on α but not on i such that $\|T^{-1}x_i + x_i\| \leq 2 - \varepsilon$. So $x_i^*(T^{-1}x_i) \leq 1 - \varepsilon$. From this it follows, using the notation introduced in the proof of Lemma 3.2, that $\psi(T)(x_i^*)(x_i) - x_i^*(x_i) \leq -\varepsilon$, so that $\|\psi(T)(x_i^*) - x_i^*\| \geq \varepsilon$. This being true for every $T \neq \text{Id}_X$, $\{x_1^*, \dots, x_n^*\}$ is distinguished for the action of G on X^* . \square

As a direct corollary, we obtain:

Corollary 3.4. *Let X be a Banach space, let $G \leq \text{GL}(X)$ be SOT-discrete, and assume that X admits a G -invariant dense LUR renorming. If there exists a distinguished family of cardinality n for the action of G on X , there also exists a distinguished family of cardinality n for the action of G on X^* .*

4. Bounded groups which are not light

Isometry groups are especially relevant to our study. We introduce the following definition:

Definition 4.1. A Banach space X is *light* if $\text{Isom}(X)$ is a light subgroup of $\text{GL}(X)$.

Observe that since the isometry group of a Banach space $(X, \|\cdot\|)$ is not invariant by equivalent renorming, the notion of lightness for a Banach space depends very much on the choice of the norm. In our terminology, Megrelishvili [2001] proved that all spaces with the PCP are light but that $C([0, 1]^2)$ is not light. Also, we have the following example:

Example 4.2. The space c_0 is light.

In fact, every isometry T of c_0 (endowed with the usual supremum norm) has the form

$$T((x_k)_{k \in \mathbb{N}}) = (\varepsilon_k x_{\sigma(k)})_{k \in \mathbb{N}}, \quad (x_k)_{k \in \mathbb{N}} \in c_0,$$

where $(\varepsilon_k)_{k \in \mathbb{N}} \in \{-1, 1\}^{\mathbb{N}}$ and σ is a permutation of \mathbb{N} . For $i \in \mathbb{N}$, denote by φ_i the i -th coordinate functional on c_0 . Let (T_α) be a net in $\text{Isom}(c_0)$, such that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$. Write each T_α as

$$T_\alpha((x_k)_{k \in \mathbb{N}}) = (\varepsilon_{\alpha, k} x_{\sigma_\alpha(k)})_{k \in \mathbb{N}}, \quad (x_k)_{k \in \mathbb{N}} \in c_0,$$

with $(\varepsilon_{\alpha, k})_{k \in \mathbb{N}} \in \{-1, 1\}^{\mathbb{N}}$ and σ_α is a permutation of \mathbb{N} . We have for every $x \in c_0$ and every $i \in \mathbb{N}$,

$$|\varphi_i(T_\alpha x) - \varphi_i(x)| = |\varepsilon_{\alpha, i} x_{\sigma_\alpha(i)} - x_i| \rightarrow 0.$$

Since this holds for every x belonging to the space c_{00} of finitely supported sequences, we must eventually have $\sigma_\alpha(i) = i$ and $\varepsilon_{\alpha,i} = 1$. Hence $\|T_\alpha x - x\| \rightarrow 0$ for every $x \in c_{00}$, and by density of c_{00} in c_0 , $\|T_\alpha x - x\| \rightarrow 0$ for every $x \in c_0$.

Another proof of [Example 4.2](#) is based on the observation that c_0 admits a particular LUR renorming, namely the Day's renorming given by

$$\|x\|_D = \sup \left\{ \left(\sum_{k=1}^n \frac{x_{\sigma(k)}^2}{4^k} \right)^{\frac{1}{2}} \right\}, \quad x \in c_0,$$

where the supremum is taken over all $n \in \mathbb{N}$ and all permutations σ of \mathbb{N} (see [\[Deville et al. 1993, p. 69\]](#)). Since this renorming is isometry invariant, it follows from [Theorem 2.3](#) that c_0 is light.

Note that Day's renorming is actually defined on ℓ_∞ , and therefore on the space c of convergent real sequences. In view of [Proposition 4.3](#), it may be amusing to observe that Day's renorming is not strictly convex on c (not even on a dense subset of S_c). In fact, it is not strictly convex at the point $(1, 1, \dots)$ since for every $x = (x_k)_{k \in \mathbb{N}} \in c$ such that $\|x\|_\infty = 1$ and $|x_k| = 1$ for infinitely many indices k , we have $\|x\|_D = \|(1, 1, \dots)\|_D$.

We now provide an elementary example of a space which is not light.

Proposition 4.3. *There exists a subgroup $G \leq \text{Isom}(c)$ which has a distinguished point, but whose dual action on ℓ_1 is not SOT-discrete. In particular the space c is not light.*

Proof. Define G as the subgroup of isometries T of c of the form

$$T((x_k)_{k \in \mathbb{N}}) = (\varepsilon_k x_k)_{k \in \mathbb{N}}, \quad (x_k)_{k \in \mathbb{N}} \in c,$$

where the sequence $(\varepsilon_k)_k \in \{-1, 1\}^\mathbb{N}$ is eventually constant. We easily see that $(1, 1, \dots)$ is a distinguished point for G . On the other hand, the dual space of c identifies isomorphically with ℓ_1 , where $\varphi = (y_k)_{k \in \mathbb{N}} \in \ell_1$ acts on an element $x = (x_k)_{k \in \mathbb{N}} \in c$ by the formula

$$\varphi(x) = y_1 \lim_{k \rightarrow \infty} x_k + \sum_{k=2}^{\infty} y_k x_{k-1}.$$

For every $n \in \mathbb{N}$, define the operator $T_n \in G$ by setting, for every $(x_k)_{k \in \mathbb{N}} \in c$,

$$T_n(x_1, x_2, \dots, x_{n-1}, x_n, x_{n+1}, \dots) = (x_1, x_2, \dots, x_{n-1}, -x_n, x_{n+1}, \dots).$$

Obviously $T_n \xrightarrow{\text{SOT}} \text{Id}$, but for every $n \in \mathbb{N}$ and every $x \in c$ we have

$$\varphi(T_n(x)) = y_1 \lim_{k \rightarrow \infty} (T_n(x))_k + \sum_{k=2}^{\infty} y_k (T_n(x))_{k-1} = \left(y_1 \lim_{k \rightarrow \infty} x_k + \sum_{k=2}^{\infty} y_k x_{k-1} \right) - 2y_n x_n$$

which tends to $\varphi(x)$ as n tends to infinity. Hence $T_n \xrightarrow{\text{WOT}} \text{Id}$ and G is not light, which implies that $\text{Isom}(c)$ itself is not light. Actually the inequality $|(T_n^* \varphi - \varphi)(x)| = 2|y_n x_n| \leq 2|y_n| \|x\|$, $x \in c$, $\varphi \in \ell_1$, implies that T_n^* tends SOT to Id , so the dual action of G on ℓ_1 is not SOT-discrete. \square

Remark 4.4. Note that the nonlight subgroup G of $\text{Isom}(c)$ constructed in the proof of Proposition 4.3 has the property that all its elements belong to the group $\text{Isom}_f(c)$ of isometries which are finite rank perturbations of the identity.

We observe the following relation between groups acting on a space and on a complemented subspace.

Lemma 4.5. *Assume Y embeds complementably in X . If every bounded group of isomorphisms on X is light, then every bounded group of isomorphisms on Y is light.*

Proof. Let Z be a closed subspace of X such that $X \simeq Y \oplus Z$. Let $G \leq GL(Y)$ be a bounded subgroup and for each $T \in G$, consider the operator $\tilde{T} = T \oplus \text{Id}_Z \in GL(X)$. These operators form a bounded subgroup \tilde{G} of $GL(X)$ which is therefore light.

Let $(T_\alpha)_{\alpha \in I}$ be a net in G such that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}_Y$. Then $\tilde{T}_\alpha \xrightarrow{\text{WOT}} \text{Id}_X$, and since \tilde{G} is light, $\tilde{T}_\alpha \xrightarrow{\text{SOT}} \text{Id}_X$. Since for every $y \in Y$,

$$\|T_\alpha(y) - y\|_Y = \|\tilde{T}_\alpha(y, 0) - (y, 0)\|_X \rightarrow 0,$$

it follows that $T_\alpha \xrightarrow{\text{SOT}} \text{Id}_Y$. \square

Assume that X is separable and that $G \leq GL(X)$ is a bounded group of isomorphisms on X . As we have seen, if X has the RNP or if G is SOT-compact, then X admits a G -invariant LUR-renorming. It is natural to wonder whether the assumption on G may be weakened somewhat and, in particular, whether a similar result holds true for cyclic groups G . We show that it is not the case.

Theorem 4.6. *Let X be a separable Banach space containing an isomorphic copy of c_0 . Then $GL(X)$ contains a WOT-indiscrete bounded cyclic subgroup G with a distinguished point in X . In particular, G is not light.*

Proof. Consider the space $c(\mathbb{R}^2)$ of convergent sequences in the euclidean space \mathbb{R}^2 with the supremum norm. We define an isometry T of $c(\mathbb{R}^2)$ by setting

$$T((x_n)_{n \in \mathbb{N}}) = (R_n x_n)_{n \in \mathbb{N}} \quad \text{for every } x = (x_n)_{n \in \mathbb{N}} \in c(\mathbb{R}^2),$$

where, for every $n \in \mathbb{N}$,

$$R_n = \begin{pmatrix} \cos\left(\frac{2\pi}{n}\right) & -\sin\left(\frac{2\pi}{n}\right) \\ \sin\left(\frac{2\pi}{n}\right) & \cos\left(\frac{2\pi}{n}\right) \end{pmatrix}$$

is the rotation of \mathbb{R}^2 of angle $2\pi/n$. Observe that, since $\lim_n 2\pi/n = 0$, we have

$$\lim_n T((x_n)_{n \in \mathbb{N}}) = \lim_n (R_n x_n)_{n \in \mathbb{N}} = \lim_n (x_n)_{n \in \mathbb{N}} \quad \text{for every } x = (x_n)_{n \in \mathbb{N}} \in c(\mathbb{R}^2).$$

As also $R_n^{k!} = \text{Id}_{\mathbb{R}^2}$ whenever $k \geq n$, we deduce that $T^{k!} \xrightarrow{\text{WOT}} \text{Id}$. So the cyclic subgroup $\langle T \rangle$ of $\text{GL}(c(\mathbb{R}^2))$ generated by T is indiscrete in the WOT.

On the other hand, if we define $x = (x_n)_{n \in \mathbb{N}} \in c(\mathbb{R}^2)$ by setting $x_n = (1, 0)$ for every $n \in \mathbb{N}$, we find that, for every $k \in \mathbb{N}$,

$$\|T^k x - x\|_{c(\mathbb{R}^2)} \geq \|R_{2k}^{k!} x_{2k} - x_{2k}\|_2 = \|(-1, 0) - (1, 0)\|_2 = 2.$$

So x is a distinguished point for $\langle T \rangle$.

Observe that $c(\mathbb{R}^2) \simeq c \oplus c \simeq c_0 \oplus c_0 \simeq c_0$, so T can be seen as an automorphism of c_0 . Also, if X is a separable Banach space containing c_0 , then c_0 is complemented in X by Sobczyk's theorem, i.e., X can be written as $X = c_0 \oplus Y$ for some subspace Y of X . Then [Lemma 4.5](#) applies. Actually the group G generated by $S = T \oplus \text{Id}$ on X is not light, since $S^{k!} \xrightarrow{\text{WOT}} \text{Id}$, while G has a distinguished point in X . \square

Remark 4.7. It follows from [Theorem 4.6](#) that any separable Banach space X containing an isomorphic copy of c_0 admits a renorming $\|\cdot\|$ such that $(X, \|\cdot\|)$ is not light.

We finish this section with the following observation:

Lemma 4.8. *Suppose G is an abelian group acting by isometries on a metric space (X, d) without isolated points, and inducing a dense orbit $G \cdot x$ for some element $x \in X$. Then, for every $\varepsilon > 0$, there exists $g \in G \setminus \{1\}$ such that $\sup_{z \in X} d(gz, z) < \varepsilon$.*

Proof. Indeed, since X has no isolated points and the orbit $G \cdot x$ is dense, we may pick $g \in G$ so that $0 < d(gx, x) < \varepsilon$. For any y in $G \cdot x$, written $y = hx$ for $h \in G$, we have

$$d(gy, y) = d(ghx, hx) = d(hgx, hx) = d(gx, x) < \varepsilon.$$

The result follows by density. \square

As a particular instance, note that if G is an SOT-discrete group of isometries of a Banach space X of dimension > 1 with a dense orbit on S_X , then G cannot be abelian.

5. LUR and strictly convex isometry invariant renormings

[Theorem 2.3](#) leads to the following question:

Question 5.1. Does there exist a light Banach space X which admits no isometry invariant LUR renorming?

It was observed in [Ferenczi and Rosendal 2011] that $X = L_1[0, 1]$ does not admit any isometry invariant dense LUR renorming. In fact, since the norm of $L_1[0, 1]$ is almost transitive and is not strictly convex, any equivalent renorming is just a multiple of the original norm, so it is not strictly convex either, and hence is not LUR. Thus $L_1[0, 1]$ could be a natural example of a light Banach space which admits no isometry invariant LUR renorming. However, this is not the case:

Proposition 5.2. *The space $L_1[0, 1]$ is not light.*

Proof. For every $n \in \mathbb{N}$, define $\varphi_n : [0, 1] \rightarrow [0, 1]$ by setting

$$\varphi_n(t) = t + \frac{1 - \cos(2^n \pi t)}{2^n \pi}, \quad t \in [0, 1],$$

and $T_n : L_1[0, 1] \rightarrow L_1[0, 1]$ by

$$T_n(f)(t) = \varphi'_n(t) f(\varphi_n(t)), \quad f \in L_1[0, 1], \quad t \in [0, 1].$$

Note that φ_n is a differentiable bijection from $[0, 1]$ into itself. So T_n is a surjective linear isometry of $L_1[0, 1]$. Moreover, $T_n \xrightarrow{\text{SOT}} \text{Id}$, since for $f \equiv 1$ we have

$$\|T_n(1) - 1\|_1 = \|\sin(2^n \pi x)\|_1 = \frac{2}{\pi} \quad \text{for every } n \in \mathbb{N}.$$

On the other hand, $T_n \xrightarrow{\text{WOT}} \text{Id}$. To prove this, we need to check that

$$\int_0^1 T_n(f)(t)g(t) dt \rightarrow \int_0^1 f(t)g(t) dt \quad \text{for every } f \in L_1[0, 1] \text{ and } g \in L_\infty[0, 1].$$

By the linearity of T_n and the density of step functions in $L_1[0, 1]$, it is sufficient to consider the case where f is the indicator function of a segment $I_{m,k} = [\frac{2k}{2^m}, \frac{2(k+1)}{2^m}]$, where $m \geq 1$ and $0 \leq k \leq 2^{m-1} - 1$. In this case the function φ_n is a bijection from $I_{m,k}$ into itself for every $n \geq m$. Thus $f \circ \varphi_n = \varphi_n$, and

$$\begin{aligned} \int_0^1 T_n(f)(t)g(t) dt &= \int_0^1 \varphi'_n(t) f(\varphi_n(t))g(t) dt = \int_0^1 \varphi'_n(t) f(t)g(t) dt \\ &= \int_0^1 f(t)g(t) dt + \int_0^1 \sin(2^n \pi t) f(t)g(t) dt. \end{aligned}$$

The result then follows from the Riemann–Lebesgue lemma. \square

Remark 5.3. Another space of which it is well known that it does not admit any LUR renorming is ℓ_∞ . Indeed ℓ_∞ does not admit any equivalent norm with the Kadec–Klee property ([Deville et al. 1993, Chapter 2, Theorem 7.10]), while every LUR norm satisfies the Kadec–Klee property ([Deville et al. 1993, Chapter 2, Proposition 1.4]). However, ℓ_∞ does admit a strictly convex renorming (see [Diestel 1975, p. 120]). We note here that it does not admit any isometry invariant strictly convex renorming. To see this, consider the points $x = (1, 1, 0, 1, 0, 1, 0, \dots)$ and

$y = (-1, 1, 0, 1, 0, 1, 0, \dots)$. Setting $z = (x + y)/2 = (0, 1, 0, 1, 0, 1, 0, \dots)$, it is readily seen that there exist two isometries T and S of ℓ_∞ such that $Tx = y$ and $Sx = z$. So, for any isometry invariant renorming $\|\cdot\|$ of ℓ_∞ we have $\|x\| = \|y\| = \|z\|$, and therefore $\|\cdot\|$ cannot be strictly convex.

Proposition 5.4. *The space ℓ_∞ is not light.*

Proof. Consider the sequence of isometries $T_n : \ell_\infty \rightarrow \ell_\infty$, $n \in \mathbb{N}$, defined by

$$T_n(x_1, \dots, x_{n-1}, x_n, x_{n+1}, \dots) = (x_1, \dots, x_{n-1}, -x_n, x_{n+1}, \dots), \quad x = (x_k)_{k \in \mathbb{N}} \in \ell_\infty.$$

Notice that $T_n \not\xrightarrow{\text{SOT}} \text{Id}$, since $\|T_n(1, 1, \dots) - (1, 1, \dots)\|_\infty = 2$ for every $n \in \mathbb{N}$. On the other hand, observe that $T_n \xrightarrow{\text{WOT}} \text{Id}$. Indeed, if $(e_j)_{j \in \mathbb{N}}$ denotes the canonical basis of ℓ_∞ , the sequence $(\beta(e_j))_{j \in \mathbb{N}}$ belongs to ℓ_1 for every $\beta \in \ell_\infty^*$. In particular, $\beta(e_j) \rightarrow 0$. Thus $\beta(T_n x - x) = -2x_n \beta(e_n) \rightarrow 0$ for every $x \in \ell_\infty$ and $\beta \in \ell_\infty^*$, showing that $T_n \xrightarrow{\text{WOT}} \text{Id}$. \square

A similar proof allows us to construct many examples of $C(K)$ -spaces which are not light.

Theorem 5.5. *Let K be a compact space with infinitely many connected components. Then $C(K)$ is not light.*

Proof. We claim that, under the assumption of [Theorem 5.5](#), there exists a sequence $(N_n)_{n \in \mathbb{N}}$ of disjoint clopen subsets of K . Indeed, choose two points x_1 and y_1 of K which belong to two different connected components of K . Since the connected component of a point x of K is the intersection of all the clopen subsets of K containing x , there exists a clopen subset K_1 of K such that $x_1 \in K_1$ and $y_1 \in L_1 := K \setminus K_1$. The two sets K_1 and L_1 are compact, and one of them, say K_1 , has infinitely many connected components. We set then $N_1 = L_1$, and repeat the argument starting from the compact set K_1 . This yields a sequence $(N_n)_{n \in \mathbb{N}}$ of disjoint clopen subsets of K .

For each integer $n \in \mathbb{N}$, define $T_n \in \text{Isom}(C(K))$ by setting, for every $f \in C(K)$ and every $x \in K$,

$$T_n(f)(x) = \begin{cases} -f(x) & \text{if } x \in N_n, \\ f(x) & \text{otherwise.} \end{cases}$$

If χ_n denotes the indicator function of the set N_n , we have $T_n(f) = f(1 - 2\chi_n)$ for every $f \in C(K)$. Applying this to the constant function $f \equiv 1$, we have $\|T_n(f) - f\|_\infty = 2$ for every $n \in \mathbb{N}$, so that $T_n \not\xrightarrow{\text{SOT}} \text{Id}$. On the other hand, the same kind of argument as in [Proposition 5.4](#) shows that $T_n \xrightarrow{\text{WOT}} \text{Id}$. Indeed, we have $\Phi(T_n f - f) = -2\Phi(f\chi_n)$ for every functional $\Phi \in C(K)^*$ and every $f \in C(K)$. For every sequence $(\alpha_n)_{n \in \mathbb{N}} \in c_0$, the series $\sum_{n \in \mathbb{N}} \alpha_n f \chi_n$ converges in $C(K)$, so that the series $\sum_{n \in \mathbb{N}} \alpha_n \Phi(f\chi_n)$ converges. It follows that $\Phi(f\chi_n) \rightarrow 0$ as $n \rightarrow +\infty$, which proves our claim. \square

As a direct consequence of [Theorem 5.5](#) we retrieve the result, proved in [Proposition 4.3](#) above, that the space c of convergent sequences is not light. Also, we immediately deduce that the space of continuous functions on the Cantor space is not light.

Corollary 5.6. *The space $C(\{0, 1\}^{\mathbb{N}})$ is not light.*

In view of the results above, combined with the known fact that the space $C([0, 1]^2)$ is not light, it may seem reasonable to conjecture that none of the spaces $C(K)$, where K is any infinite compact space, is light. However, our next result shows that this is not the case.

Theorem 5.7. *Let K be an infinite compact connected space. Then $C(K)$ is light if and only if the topologies of pointwise and uniform convergence coincide on the group $\text{Homeo}(K)$ of homeomorphisms of K .*

Proof. Suppose first that the topologies of pointwise and uniform convergence coincide on $\text{Homeo}(K)$. Let $(T_\alpha)_{\alpha \in I}$ be a net of isometries of $C(K)$ such that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$. By the Banach–Stone theorem and the connectedness of K , each isometry T_α of $C(K)$ has the form

$$T_\alpha(f) = \varepsilon_\alpha f \circ \varphi_\alpha \quad \text{for every } f \in C(K),$$

where $\varepsilon_\alpha \in \{-1, 1\}$ and $\varphi_\alpha \in \text{Homeo}(K)$. Since $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$, $\varepsilon_\alpha \rightarrow 1$, so we can suppose without loss of generality that $\varepsilon_\alpha = 1$ for every $\alpha \in I$. Moreover, the fact that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$ also implies that the net $(\varphi_\alpha)_{\alpha \in I}$ converges pointwise to the identity function id_K on K . Our assumption then implies that $(\varphi_\alpha)_{\alpha \in I}$ converges uniformly to id_K on K , from which it easily follows that $T_\alpha \xrightarrow{\text{SOT}} \text{Id}$. Thus $C(K)$ is light.

Conversely, suppose that $C(K)$ is light. Let $(\varphi_\alpha)_{\alpha \in I}$ be a net of elements of $\text{Homeo}(K)$ which converges pointwise to $\varphi \in \text{Homeo}(K)$. Consider the isometries T_α and T of $C(K)$ defined by

$$T_\alpha(f) = f \circ \varphi_\alpha \quad \text{and} \quad T(f) = f \circ \varphi \quad \text{for every } f \in C(K).$$

Then $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$. Since $C(K)$ is light, $T_\alpha \xrightarrow{\text{SOT}} \text{Id}$ and thus $(\varphi_\alpha)_{\alpha \in I}$ converges to φ uniformly on K . \square

Remark 5.8. [Theorem 5.7](#) characterizes the lightness of $C(K)$ for infinite connected compact spaces K . We may naturally wonder whether the connectedness assumption is really necessary. It is indeed the case: there exist compact spaces K with infinitely many connected components which are *rigid* in the sense that their homeomorphism group $\text{Homeo}(K)$ is trivial (it consists uniquely of the identity map on K). The existence of such compacta is proved in [\[de Groot and Wille 1958\]](#) (see the remark on p. 443 at the end of Section 2). By [Theorem 5.5](#), $C(K)$ is not light for such a compact K , but the topologies of pointwise and uniform convergence obviously coincide on $\text{Homeo}(K)$.

Birkhoff [1934] studied various topologies on so-called “transformation spaces”, in particular on the groups of homeomorphisms of topological spaces. He introduced the notions of A -, B - and C -convergence of sequences of homeomorphisms on a given space X corresponding respectively to pointwise convergence, continuous convergence, and continuous convergence in both directions. Since on compact spaces continuous convergence and uniform convergence coincide, Theorem 5.7 can be rephrased, using Birkhoff’s language, as saying that for compact connected spaces K , $C(K)$ is light if and only if A - and B -convergence coincide on $\text{Homeo}(K)$. Now, it is observed in [Birkhoff 1934, Theorem 18] that A -convergence implies B - and C -convergence for homeomorphisms of (disjoint or not) finite unions of segments of the real line (this is essentially the content of Dini’s second convergence theorem), while if K contains an n -dimensional region with $n \geq 2$ (i.e., an open set homeomorphic to an open subset of \mathbb{R}^n), A -convergence implies neither B - nor C -convergence for homeomorphisms of K ([Birkhoff 1934, Theorem 19]). In more modern language, under this assumption there exists a sequence $(\varphi_n)_{n \in \mathbb{N}}$ of homeomorphisms of K such that φ_n converges pointwise but not uniformly on K to the identity function on K . Combined with Theorem 5.7 above, this yields:

Theorem 5.9. *Let K be an infinite compact connected space.*

- (a) *If K is homeomorphic to a finite union of segments of \mathbb{R} , $C(K)$ is light.*
- (b) *If K contains an n -dimensional region for some $n \geq 2$, $C(K)$ is not light.*

For instance, the space $C[0, 1]$ is light, while spaces $C([0, 1]^n)$, $n \geq 2$, are not light. We thus retrieve in a natural way the original example of Megrelishvili of a nonlight space.

Theorem 5.9 allows us to answer Question 5.1 in the negative. Although $C[0, 1]$ is light, it does not admit any isometry invariant LUR renorming. In fact, $C[0, 1]$ does not admit any isometry invariant strictly convex renorming. In order to prove this, we need the following lemma:

Lemma 5.10. *Let $f \in C[0, 1]$ be such that there exists an interval $[a, b] \subset [0, 1]$, $a < b$, on which f is strictly monotone. Then there exists $g \in C[0, 1]$ with the following three properties:*

- (a) $\|f\|_\infty = \|g\|_\infty = \|(f + g)/2\|_\infty$;
- (b) $\|f - g\|_\infty > 0$;
- (c) *There exist two homeomorphisms φ and ψ of $[0, 1]$ such that $g = f \circ \varphi$ and $(f + g)/2 = f \circ \psi$.*

Proof. Let $0 \leq a < b \leq 1$ be such that f is strictly monotone on $[a, b]$. Without loss of generality, suppose f is strictly increasing on $[a, b]$. Let $\xi : [a, b] \rightarrow [f(a), f(b)]$

be an increasing homeomorphism such that $\xi \not\equiv f|_{[a,b]}$. Define $g \in C[0, 1]$ and a homeomorphism $\varphi : [0, 1] \rightarrow [0, 1]$ by

$$g(x) = \begin{cases} \xi(x) & \text{if } x \in [a, b], \\ f(x) & \text{otherwise,} \end{cases} \quad \varphi(x) = \begin{cases} f^{-1}(\xi(x)) & \text{if } x \in [a, b], \\ x & \text{otherwise.} \end{cases}$$

Then, $g = f \circ \varphi$, $\|g\|_\infty = \|f\|_\infty = \|(f + g)/2\|_\infty$ and $\|f - g\|_\infty > 0$. Moreover, $f \circ \psi = (f + g)/2$, where $\psi : [0, 1] \rightarrow [0, 1]$ is the homeomorphism defined by

$$\psi(x) = \begin{cases} f^{-1}\left(\frac{\xi(x) + f(x)}{2}\right) & \text{if } x \in [a, b], \\ x & \text{otherwise.} \end{cases} \quad \square$$

Proposition 5.11. *Let $\|\cdot\|$ be an isometry invariant renorming of $C[0, 1]$. Then there exists a dense subset of $C[0, 1]$ where $\|\cdot\|$ is not strictly convex.*

Proof. Let $f \in C[0, 1]$ be a nonconstant and affine function, and take g, φ and ψ as in Lemma 5.10. Since $f \mapsto f \circ \varphi$ and $f \mapsto f \circ \psi$ define surjective linear isometries of $C[0, 1]$,

$$\|g\| = \|f \circ \varphi\| = \|f\| = \|f \circ \psi\| = \left\| \frac{f+g}{2} \right\|.$$

So $\|\cdot\|$ is not strictly convex at the point f . The result then follows from the fact that the set of piecewise linear functions is dense in $C[0, 1]$. \square

Combining Theorem 5.9 and Proposition 5.11, we obtain:

Corollary 5.12. *The space $C[0, 1]$ is light, but does not admit any isometry invariant LUR renorming.*

Remark 5.13. Using the same arguments as in the proofs of Proposition 5.11, Theorem 5.7, and Theorem 5.9 one can prove that $C_0(\mathbb{R})$ is light, but does not admit a strictly convex isometry invariant renorming either.

Remark 5.14. The examples presented in this section show that there is no general relation between closed subspaces and their respective isometry groups, in terms of being light, apart from Lemma 4.5. In fact:

- (1) c_0 is a closed subspace of c , c_0 is light, but c is not;
- (2) c is isometrically isomorphic to a closed subspace of $C[0, 1]$, c is not light but $C[0, 1]$ is light.

Corollary 5.12 gives us a positive answer to Question 5.1. On the other hand, Remark 2.4 suggests the following new question:

Question 5.15. Does there exist a Banach space X and an orbitwise light group $G \leq GL(X)$ such that X admits no G -invariant LUR renorming?

The next proposition shows that the isometry group of $C[0, 1]$ also gives a positive answer to [Question 5.15](#):

Proposition 5.16. *The group $\text{Isom}(C[0, 1])$ is orbitwise light.*

Proof. Let $f \in C[0, 1]$ and let $(g_\alpha)_{\alpha \in I}$ be a net in the orbit $\mathcal{O}(f)$ of f under the action of the group $\text{Isom}(C[0, 1])$ such that g_α converges weakly to $g \in \mathcal{O}(f)$. By the Banach–Stone theorem, there exist homeomorphisms $\varphi, \varphi_\alpha \in \text{Hom}([0, 1])$ and $\varepsilon, \varepsilon_\alpha \in \{-1, 1\}$ such that $g = \varepsilon \cdot f \circ \varphi$ and $g_\alpha = \varepsilon_\alpha \cdot f \circ \varphi_\alpha$. Since g_α converges weakly to g (hence, pointwise), we can assume that the φ_α are increasing homeomorphisms, $\varepsilon = \varepsilon_\alpha = 1$ for every $\alpha \in I$ and $g = f$.

Suppose by contradiction that $f \circ \varphi_\alpha$ does not converge uniformly to f . Then we can assume that there exists $\varepsilon > 0$ and for every $\alpha \in I$ there exists $x_\alpha \in [0, 1]$ such that $|f(\varphi_\alpha(x_\alpha)) - f(x_\alpha)| > 2\varepsilon$. We also can assume that $x_\alpha \rightarrow x \in [0, 1]$ and $x_\alpha \leq x$ for every α . Then by the continuity of f at the point x ,

$$|f(\varphi_\alpha(x_\alpha)) - f(x)| > \varepsilon.$$

Let $\delta > 0$ be such that $|x - y| < \delta \implies |f(x) - f(y)| < \frac{\varepsilon}{8}$. Then $\varphi_\alpha(x_\alpha) \notin (x - \delta, x + \delta)$ for every α , and $\varphi_\alpha(x_\alpha) < x - \delta$ for infinitely many indices $\alpha \in I$, or $\varphi_\alpha(x_\alpha) > x + \delta$ for infinitely many indices $\alpha \in I$. Without loss of generality, we may assume that

$$\varphi_\alpha(x_\alpha) < x - \delta, \quad \text{for every } \alpha \in I.$$

We also may assume that

$$x - \delta < x_\alpha \leq x, \quad \text{for every } \alpha \in I$$

(the cases $\varphi_\alpha(x_\alpha) > x + \delta$ and/or $x < x_\alpha < x + \delta$ for every $\alpha \in I$ are similar).

Let $\alpha_1 \in I$ and let

$$y_{1,1} = \varphi_{\alpha_1}(x_{\alpha_1}).$$

We claim that for every $n \geq 2$, there exists a finite sequence in $[0, 1]$,

$$y_{n,1} < y_{n,2} < \cdots < y_{n,2n-1} < x - \delta,$$

such that

$$|f(y_{n,2k+1}) - f(x)| > \varepsilon - \frac{\varepsilon}{8} - \frac{\varepsilon}{2^{k+4}} \left(\sum_{j=0}^{n-k-2} \frac{1}{2^j} \right) > \frac{3\varepsilon}{4} \quad \text{for } k = 0, 1, \dots, n-1$$

and

$$|f(y_{n,2k}) - f(x)| < \frac{\varepsilon}{8} + \frac{\varepsilon}{2^{k+3}} \left(\sum_{j=0}^{n-k-1} \frac{1}{2^j} \right) < \frac{\varepsilon}{4} \quad \text{for } k = 1, 2, \dots, n-1.$$

Notice that the existence of such a sequence for every $n \geq 2$ contradicts the uniform continuity of f on $[0, 1]$. Hence it suffices to prove this claim in order to complete the proof of [Proposition 5.16](#).

We prove this claim by induction. Since $f \circ \varphi_\alpha$ converges pointwise to f and $x_\alpha \rightarrow x$, we can take $\alpha_2 \succcurlyeq \alpha_1$ such that $x_{\alpha_1} < x_{\alpha_2} < x$, $|f(\varphi_\beta(y_{1,1})) - f(y_{1,1})| < \varepsilon/16$ and $|f(\varphi_\beta(x_{\alpha_1})) - f(x_{\alpha_1})| < \varepsilon/16$ for every $\beta \succcurlyeq \alpha_2$. Let

$$y_{2,1} = \varphi_{\alpha_2}(y_{1,1}), \quad y_{2,2} = \varphi_{\alpha_2}(x_{\alpha_1}) \quad \text{and} \quad y_{2,3} = \varphi_{\alpha_2}(x_{\alpha_2}).$$

Since φ_{α_2} is an increasing homeomorphism and $y_{1,1} < x - \delta < x_{\alpha_1} < x_{\alpha_2}$, we have $y_{2,1} < y_{2,2} < y_{2,3}$ and $y_{2,3} = \varphi_{\alpha_2}(x_{\alpha_2}) < x - \delta$. Moreover,

$$|f(y_{2,1}) - f(x)| > \varepsilon - \frac{\varepsilon}{8} - \frac{\varepsilon}{16}, \quad |f(y_{2,2}) - f(x)| < \frac{\varepsilon}{8} + \frac{\varepsilon}{16}, \quad |f(y_{2,3}) - f(x)| > \varepsilon - \frac{\varepsilon}{8},$$

which proves the inequalities for $n = 2$.

Suppose the inequalities hold for n . Let $\alpha_{n+1} \succcurlyeq \alpha_n$ such that $x_{\alpha_n} < x_{\alpha_{n+1}} < x$, $|f(\varphi_\beta(y_{n,r})) - f(y_{n,r})| < \varepsilon/2^{n+3}$ and $|f(\varphi_\beta(x_{\alpha_n})) - f(x_{\alpha_n})| < \varepsilon/2^{n+3}$ for every $r = 1, 2, \dots, 2n - 1$ and every $\beta \succcurlyeq \alpha_2$. Let

$$y_{n+1,r} = \varphi_{\alpha_{n+1}}(y_{n,r}) \quad \text{for } r = 1, \dots, 2n - 1,$$

and let

$$y_{n+1,2n} = \varphi_{\alpha_{n+1}}(x_{\alpha_n}) \quad \text{and} \quad y_{n+1,2n+1} = \varphi_{\alpha_{n+1}}(x_{\alpha_{n+1}}).$$

It follows that

$$|f(y_{n+1,2k+1}) - f(x)| > \varepsilon - \frac{\varepsilon}{8} - \frac{\varepsilon}{2^{k+4}} \left(\sum_{j=0}^{n-k-1} \frac{1}{2^j} \right) \quad \text{for } k = 0, 1, \dots, n,$$

and

$$|f(y_{n+1,2k}) - f(x)| < \frac{\varepsilon}{8} + \frac{\varepsilon}{2^{k+3}} \left(\sum_{j=0}^{n-k} \frac{1}{2^j} \right) \quad \text{for } k = 1, 2, \dots, n.$$

Since $\varphi_{\alpha_{n+1}}$ is an increasing homeomorphism, and $y_{n,1} < y_{n,2} < \dots < y_{n,2n-1} < x - \delta < x_{\alpha_n} < x_{\alpha_{n+1}}$, we have $y_{n+1,1} < y_{n+1,2} < \dots < y_{n+1,2n+1} = \varphi_{\alpha_{n+1}}(x_{\alpha_{n+1}}) < x - \delta$, which proves the claim. \square

6. An example of a group with a discrete orbit but no distinguished point

In this section we solve a problem of [Ferenczi and Rosendal 2011], mentioned in the introduction, by exhibiting an SOT-discrete group of isomorphisms of c_0 which admits no distinguished point. More generally, we show the following:

Proposition 6.1. *For any integer $r \geq 2$, there exists a bounded infinite SOT-discrete group of isomorphisms of c_0 of the form $\text{Id} + F$, $F \in L(c_0)$ of finite rank, admitting a distinguished family of cardinality r , but none of cardinality $r - 1$.*

Proof. Since $c_0 \simeq \ell_1^r \oplus_\infty c_0$, it is enough to define the group G as an infinite bounded SOT-discrete group of isomorphisms on $\ell_1^r \oplus_\infty c_0$.

Let $(e_n)_{n \in \mathbb{N}}$ be the canonical basis of c_0 , and let $(U_n)_{n \in \mathbb{N}}$ be the sequence of isometries of c_0 defined by setting, for every $n, m \in \mathbb{N}$, $U_n(e_n) = -e_n$ and $U_n(e_m) = e_m$ whenever $m \neq n$. Let $(\phi_n)_{n \in \mathbb{N}}$ be dense in the unit sphere of ℓ_∞^r , and define the rank-one operator $R_n : \ell_1^r \rightarrow c_0$ by $R_n(x) = \phi_n(x)e_n$, $x \in \ell_1$. We then define an operator T_n on $\ell_1^r \oplus_\infty c_0$ in matrix form as

$$T_n = \begin{pmatrix} \text{Id} & 0 \\ R_n & U_n \end{pmatrix}.$$

It is readily checked that $T_n^2 = \text{Id}$ for every $n \in \mathbb{N}$ and that for all distinct integers n_1, \dots, n_k ,

$$T_{n_1} \dots T_{n_k} = \begin{pmatrix} \text{Id} & 0 \\ R_{n_1} + \dots + R_{n_k} & U_{n_1} \dots U_{n_k} \end{pmatrix}.$$

Therefore the group G generated by the operators T_n is abelian. Furthermore, since for every $x \in \ell_1^r$

$$\|(R_{n_1} + \dots + R_{n_k})x\| = \|\phi_{n_1}(x)e_{n_1} + \dots + \phi_{n_k}(x)e_{n_k}\| \leq \max_i \|\phi_{n_i}(x)\| \cdot \|x\|,$$

it follows that $\|T_{n_1} + \dots + T_{n_k}\| \leq 2$; thus G is a bounded subgroup of $\text{GL}(\ell_1^r \oplus_\infty c_0)$.

We claim that no family $\{x_1, \dots, x_{r-1}\}$ of $\ell_1^r \oplus c_0$ is distinguished for G . Indeed, writing each vector x_i as (y_i, z_i) with $y_i \in \ell_1^r$ and $z_i \in c_0$, we note that $U_n z_i \rightarrow z_i$ for every $1 \leq i \leq r-1$. Since the vectors y_1, \dots, y_{r-1} generate a subspace of dimension strictly less than r of ℓ_1^r , there exists a norm 1 functional $\phi \in \ell_\infty^r$ such that $\phi(y_i) = 0$ for every $1 \leq i \leq r-1$. Let $D \subset \mathbb{N}$ be such that $\phi_n \rightarrow \phi$ in ℓ_∞^r as n tends to infinity along D . Then $R_n(y_i) \rightarrow 0$ as n tends to infinity along D , and therefore $T_n(x_i) \rightarrow x_i$ as n tends to infinity along D for every $1 \leq i \leq r-1$. So the family $\{x_1, \dots, x_{r-1}\}$ is not distinguished for G .

On the other hand, if we denote by (f_1, \dots, f_r) the canonical basis of ℓ_1^r , then the family $\{f_1 \oplus 0, \dots, f_r \oplus 0\}$ is distinguished for G . To check this, note that for any operator $T \in \text{GL}(\ell_1^r \oplus_\infty c_0)$ of the form

$$T = \begin{pmatrix} \text{Id} & 0 \\ \sum_{k \in F} R_k & U \end{pmatrix},$$

where F is a nonempty subset of \mathbb{N} , and U is an isometry of c_0 , we have

$$\|T(f_s \oplus 0) - f_s \oplus 0\| = \max_{k \in F} |\phi_k(f_s)| \quad \text{for every } 1 \leq s \leq r.$$

Since, for each $k \in F$, ϕ_k is normalized in ℓ_∞^r , $|\phi_k(f_s)| \geq 1$ for at least one index s . It follows that

$$\max_{1 \leq s \leq r} \|T(f_s \oplus 0) - f_s \oplus 0\| \geq 1,$$

and so

$$\inf_{T \in G, T \neq \text{Id}} \left\{ \max_{1 \leq s \leq r} \|T(f_s \oplus 0) - f_s \oplus 0\| \right\} \geq 1.$$

Hence $\{f_1, \dots, f_r\}$ is a distinguished family for G . □

We immediately deduce the following:

Corollary 6.2. *The group of isomorphisms of c_0 which is constructed in the proof of Proposition 6.1 is not light.*

Proof. For every $x \in \ell_1^r$, the sequence $(R_n(x))_{n \in \mathbb{N}}$ tends weakly to 0 in c_0 . We also know that the sequence $(U_n)_{n \in \mathbb{N}}$ tends WOT to Id. Therefore $(T_n)_{n \in \mathbb{N}}$ also tends WOT to Id. On the other hand, we have for every $x \in \ell_1^r$ and every $n \in \mathbb{N}$ that

$$\|T_n(x \oplus 0) - x \oplus 0\| = \|R_n(x)\| = \|\phi_n(x)\|.$$

By the density of the sequence $(\phi_n)_{n \in \mathbb{N}}$ in the unit sphere of ℓ_∞^r , this implies that the sequence $(T_n(x \oplus 0))_{n \in \mathbb{N}}$ does not tend to x in norm, and thus $(T_n)_{n \in \mathbb{N}}$ does not tend SOT to Id. □

We have thus proved:

Corollary 6.3. *There exists a bounded group G of isomorphisms of c_0 which is infinite, not light, SOT-discrete, and does not admit a distinguished point.*

Proof. Take $r = 2$ in Proposition 6.1. □

7. Quasinormed spaces

Although Megrelishvili has defined the concept of light group of isomorphisms only for locally convex spaces, we can extend the definition to quasinormed spaces, even if these spaces are not locally convex. One could ask if there is a general answer for the isometry groups of nonlocally convex spaces, in terms of being light. The spaces ℓ_p and $L_p[0, 1]$, $0 < p < 1$, are examples that give a negative answer to this question.

Recall that for $0 < p < 1$, $(L_p[0, 1])^* = \{0\}$, i.e., the only linear continuous functional $f : L_p[0, 1] \rightarrow \mathbb{R}$ is the constant function $f \equiv 0$ (see [Kalton et al. 1984, p. 18]). Considering the sequence $(T_n)_{n \in \mathbb{N}}$ constantly equal to $-\text{Id}$, we observe that $T_n \not\xrightarrow{\text{SOT}} \text{Id}$ while $T_n \xrightarrow{\text{WOT}} \text{Id}$. So $L_p[0, 1]$ is trivially nonlight for every $0 < p < 1$. On the other hand, we have:

Proposition 7.1. *For $0 < p < 1$, the space ℓ_p is light.*

Proof. Let $0 < p < 1$, and let $(T_\alpha)_{\alpha \in I}$ be a net in $\text{Isom}(\ell_p)$ such that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$. Each T_α acts on vectors $(x_n)_{n \in \mathbb{N}} \in \ell_p$ as $T_\alpha((x_n)_{n \in \mathbb{N}}) = (\varepsilon_n^{(\alpha)} x_{\sigma_\alpha(n)})_{n \in \mathbb{N}}$, where σ_α is a permutation of \mathbb{N} and $(\varepsilon_n^{(\alpha)})_{n \in \mathbb{N}}$ is a sequence of elements of $\{-1, 1\}$ (the proof of this statement is similar to the case where $p > 1$ and $p \neq 2$, which can be found in [Banach 1932, p. 178]). Assume, by contradiction, that $T_\alpha \not\xrightarrow{\text{SOT}} \text{Id}$.

Then there exist $x \in \ell_p$, $\varepsilon > 0$ and an infinite sequence $(\alpha_i)_{i \in \mathbb{N}}$ of indices in I such that $\|T_{\alpha_i}x - x\|_p^p > \varepsilon$ for every $i \in \mathbb{N}$. Since $x \in \ell_p$, there exists $N \in \mathbb{N}$ such that $\sum_{k=N+1}^{\infty} |x_k|^p < \varepsilon/2$. The dual space of ℓ_p identifies isomorphically with ℓ_{∞} , where $\Phi = (y_k)_{k \in \mathbb{N}}$ acts on an element $x = (x_k)_{k \in \mathbb{N}} \in \ell_p$ by the formula $\Phi(x) = \sum_{k \in \mathbb{N}} y_k x_k$ (see [Kalton et al. 1984, p. 21]). Considering for $1 \leq j \leq N$ the functionals Φ_j identified with the vectors of the canonical basis $e_j \in \ell_{\infty}$, as well as the vectors $e_k \in \ell_p$ for $1 \leq k \leq N$, we obtain by the WOT convergence of T_{α} to Id that

$$\Phi_j(T_{\alpha}(e_k)) - \Phi_j(e_k) = \varepsilon_j^{(\alpha)} \delta_{\sigma_{\alpha}(k),j} - \delta_{k,j} \rightarrow 0,$$

where $\delta_{k,j} = 1$ if $k = j$ and $\delta_{k,j} = 0$ if $k \neq j$. In particular, $\varepsilon_k^{(\alpha)} \delta_{\sigma_{\alpha}(k),k} \rightarrow 1$ for every $1 \leq k \leq N$. So we may assume that the permutations σ_{α} fix the first N integers and that $\varepsilon_k^{(\alpha)} = 1$ for every $1 \leq k \leq N$. Hence we have for every $i \in \mathbb{N}$

$$\sum_{k=N+1}^{\infty} |(T_{\alpha_i}(x))_k|^p = \sum_{k=N+1}^{\infty} |x_k|^p < \frac{\varepsilon}{2}.$$

However, taking $z = (z_k)_{k \in \mathbb{N}} \in \ell_p$ defined by $z_k = 0$ if $1 \leq k \leq N$ and $z_k = 1$ if $k > N$, we have $\|z\|_p^p = \sum_{k=N+1}^{\infty} |x_k|^p < \varepsilon/2$ and

$$\|T_{\alpha_i}x - x\|_p^p = \|T_{\alpha_i}z - z\|_p^p \leq \|T_{\alpha_i}z\|_p^p + \|z\|_p^p = \varepsilon$$

for every $i \in \mathbb{N}$, which is a contradiction. □

We finish the paper with a few related questions and comments.

8. Questions and comments

Our first question concerns renormings of the space c . Since c is not light, as proved in Proposition 4.3, it does not admit any isometry invariant LUR renorming. But it may still admit an isometry invariant strictly convex renorming.

Question 8.1. Does c admit an isometry invariant strictly convex renorming?

We have observed in Section 4 that if the isometry group $\text{Isom}(X)$ of a Banach space X of dimension > 1 acts almost transitively on S_X and is SOT-discrete, it is not abelian.

Question 8.2. Suppose X is a separable Banach space of dimension > 1 and $G \leq \text{Isom}(X)$ is an SOT-discrete amenable subgroup. Can G have a dense orbit on S_X ?

We have seen in Corollary 6.3 that there exists a bounded group G of isomorphisms of c_0 which is infinite, not light, SOT-discrete, and does not admit a distinguished point. We may wonder about the role of the space c_0 in this construction. For example, we can ask:

Question 8.3. Does there exist a reflexive space X with an SOT-discrete bounded group $G \leq GL(X)$ that does not admit a distinguished point?

Of course such a group G , if it exists, must be light, as all reflexive spaces are light. Noting that the example of [Proposition 4.3](#) is a group of finite rank perturbations of the identity on the space c_0 , a question in the same vein is:

Question 8.4. Does there exist a reflexive space X with an SOT-discrete infinite bounded group $G \leq GL(X)$ such that all elements of G are finite rank perturbations of the identity?

This question is relevant to [\[Ferenczi and Rosendal 2013\]](#), where isometry groups on complex, reflexive, separable, hereditarily indecomposable spaces are studied. A negative answer would imply that all isometry groups on such spaces act almost trivially, i.e., there would exist an isometry invariant decomposition $F \oplus H$ of the space where F is finite dimensional and all elements of the group act as multiples of the identity on H , [\[Ferenczi and Rosendal 2013, Theorem 6.9\]](#).

Another natural space which could be investigated in this context is the universal space of Gurarii, whose isometry group possesses a very rich structure (see [\[Gurarii 1966\]](#) for its definition and [\[Garbulińska and Kubiś 2011\]](#) for a recent survey).

Question 8.5. Is the isometry group of the Gurarii space light?

Finally, whether the converse to Megrelishvili's result holds remains an open question:

Question 8.6. Does a Banach space X have the PCP if and only if all bounded subgroups of $GL(X)$ are light?

The answer is positive when X has an unconditional basis; this follows from [Theorem 4.6](#), the fact that an unconditional basis whose span does not contain c_0 must be boundedly complete, and the fact that separable dual spaces have the RNP and therefore the PCP.

Acknowledgement

The authors would like to thank the anonymous referee for the corrections and many valuable suggestions which helped to improve the manuscript.

References

- [Banach 1932] S. Banach, *Théorie des opérations linéaires*, Monografie Mat. **1**, Seminar. Mat. Uniw. Warszaw., 1932. [Zbl](#)
- [Birkhoff 1934] G. Birkhoff, “The topology of transformation-sets”, *Ann. of Math.* (2) **35**:4 (1934), 861–875. [MR](#) [Zbl](#)
- [Dewille et al. 1993] R. Dewille, G. Godefroy, and V. Zizler, *Smoothness and renormings in Banach spaces*, Pitman Mono. Surveys Pure Appl. Math. **64**, Longman, Harlow, UK, 1993. [MR](#) [Zbl](#)

- [Diestel 1975] J. Diestel, *Geometry of Banach spaces: selected topics*, Lecture Notes in Math. **485**, Springer, 1975. [MR](#) [Zbl](#)
- [Ferenczi and Galego 2010] V. Ferenczi and E. M. Galego, “Countable groups of isometries on Banach spaces”, *Trans. Amer. Math. Soc.* **362**:8 (2010), 4385–4431. [MR](#) [Zbl](#)
- [Ferenczi and Rosendal 2011] V. Ferenczi and C. Rosendal, “Displaying Polish groups on separable Banach spaces”, *Extracta Math.* **26**:2 (2011), 195–233. [MR](#) [Zbl](#) [arXiv](#)
- [Ferenczi and Rosendal 2013] V. Ferenczi and C. Rosendal, “On isometry groups and maximal symmetry”, *Duke Math. J.* **162**:10 (2013), 1771–1831. [MR](#) [Zbl](#)
- [Garbulińska and Kubiś 2011] J. Garbulińska and W. Kubiś, “Remarks on Gurarii spaces”, *Extracta Math.* **26**:2 (2011), 235–269. [MR](#) [Zbl](#)
- [de Groot and Wille 1958] J. de Groot and R. J. Wille, “Rigid continua and topological group-pictures”, *Arch. Math.* **9** (1958), 441–446. [MR](#) [Zbl](#)
- [Gurarii 1966] V. I. Gurarii, “Spaces of universal placement, isotropic spaces and a problem of Mazur on rotations of Banach spaces”, *Sibirsk. Mat. Z.* **7**:5 (1966), 1002–1013. In Russian; translated in *Sib. Math. J.* **7**:5 (1966), 799–807. [MR](#)
- [Helmer 1980] D. Helmer, “Joint continuity of affine semigroup actions”, *Semigroup Forum* **21**:2-3 (1980), 153–165. [MR](#) [Zbl](#)
- [Kalton et al. 1984] N. J. Kalton, N. T. Peck, and J. W. Roberts, *An F -space sampler*, London Math. Soc. Lecture Note Series **89**, Cambridge Univ. Press, 1984. [MR](#) [Zbl](#)
- [Lancien 1993] G. Lancien, “Dentability indices and locally uniformly convex renormings”, *Rocky Mountain J. Math.* **23**:2 (1993), 635–647. [MR](#) [Zbl](#)
- [Megrelishvili 2001] M. G. Megrelishvili, “Operator topologies and reflexive representability”, pp. 197–208 in *Nuclear groups and Lie groups* (Madrid, 1999), edited by E. Martín Peinador and J. Núñez García, Res. Exp. Math. **24**, Heldermann, Lemgo, Germany, 2001. [MR](#) [Zbl](#)
- [Megrelishvili 2003] M. Megrelishvili, “Fragmentability and representations of flows”, *Topology Proc.* **27**:2 (2003), 497–544. [MR](#) [Zbl](#)

Received December 21, 2017. Revised May 25, 2018.

LEANDRO ANTUNES
 DEPARTAMENTO DE MATEMÁTICA
 UNIVERSIDADE TECNOLÓGICA FEDERAL DO PARANÁ
 CAMPUS TOLEDO
 TOLEDO, PR
 BRAZIL
leandroantunes@utfpr.edu.br

VALENTIN FERENCZI
 DEPARTAMENTO DE MATEMÁTICA
 UNIVERSIDADE DE SÃO PAULO
 INSTITUTO DE MATEMÁTICA E ESTATÍSTICA
 SÃO PAULO, SP
 BRAZIL
ferenczi@ime.usp.br

SOPHIE GRIVAUX

CNRS, UNIV. LILLE, UMR 8524 - LABORATOIRE PAUL PAINLEVÉ

FRANCE

sophie.grivaux@univ-lille.fr

CHRISTIAN ROSENDAL

DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE

UNIVERSITY OF ILLINOIS AT CHICAGO

CHICAGO, IL

UNITED STATES

rosendal.math@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$665/year for print and electronic.
Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 301 No. 1 July 2019

Multiplicity upon restriction to the derived subgroup	1
JEFFREY D. ADLER and DIPENDRA PRASAD	
Unknotting number and Khovanov homology	15
AKRAM ALISHAHI	
Light groups of isomorphisms of Banach spaces and invariant LUR renormings	31
LEANDRO ANTUNES, VALENTIN FERENCZI, SOPHIE GRIVAUX and CHRISTIAN ROSENDAL	
Some uniform estimates for scalar curvature type equations	55
SAMY SKANDER BAHOURA	
Complemented copies of $c_0(\tau)$ in tensor products of $L_p[0, 1]$	67
VINÍCIUS MORELLI CORTES, ELÓI MEDINA GALEGO and CHRISTIAN SAMUEL	
On the volume bound in the Dvoretzky–Rogers lemma	89
FERENC FODOR, MÁRTON NASZÓDI and TAMÁS ZARNÓCZ	
Lifting of Elliptic curves	101
SANOLI GUN and V. KUMAR MURTY	
Loxodromics for the cyclic splitting complex and their centralizers	107
RADHIKA GUPTA and DERRICK WIGGLESWORTH	
Lie 2-algebroids and matched pairs of 2-representations: a geometric approach	143
MADELEINE JOTZ LEAN	
Algorithmic homeomorphism of 3-manifolds as a corollary of geometrization	189
GREG KUPERBERG	
Harish-Chandra modules for divergence zero vector fields on a torus	243
ZHIQIANG LI, SHAOBIN TAN and QING WANG	
Weighted estimates for rough singular integrals with applications to angular integrability	267
FENG LIU and DASHAN FAN	
∞ -tilting theory	297
LEONID POSITSELSKI and JAN ŠTOVÍČEK	
The “quantum” Turán problem for operator systems	335
NIK WEAVER	
BCOV torsion and degenerations of Calabi–Yau manifolds	351
WEI XIA	
Classification of gradient expanding and steady Ricci solitons	371
FEI YANG, SHOUWEN FANG and LIANGDI ZHANG	