

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/270761061>

Black wattle insect pests currently in Brazil

Conference Paper · December 2014

CITATIONS

0

READS

319

3 authors, including:



Maria Angélica Ono

Universidade Federal de Lavras (UFLA)

1 PUBLICATION 0 CITATIONS

[SEE PROFILE](#)



Wesley Godoy

University of São Paulo

139 PUBLICATIONS 1,520 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Ecological modelling of the larval dispersal of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) on Bt cotton: implications for evolution and resistance management [View project](#)



Pollination services provided to agricultural crops by wild and managed insects [View project](#)



Global Advanced Research Journal of Agricultural Science (ISSN: 2315-5094) Vol. 3(12) pp. 409-414, December, 2014 Special Anniversary Review Issue.

Available online <http://garj.org/garjas/index.htm>

Copyright © 2014 Global Advanced Research Journals

Review

Black wattle insect pests currently in Brazil

Maria Angélica Ono, Elisângela Novais Lopes Ferreira, Wesley Augusto Conde Godoy

Departamento de Entomologia e Acarologia, Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo, Av. Pádua Dias, 11, 13418-900 Piracicaba – São Paulo, Brazil.

Accepted 12 December, 2014

This review treats the main insect pests associated with black wattle in Brazil, as an attempt to generate information useful for pest-management programs in plantations of *Acacia mearnsii*. This is the first compilation of insect species associated with black wattle in Brazil. Essential data about biology, behaviour, and evidence of infestation are also presented to improve the detection of pests and prevent invasions of new areas. Occurrences of pests in black wattle are described, and the damage caused by these insects to the plantations is also discussed. The main black wattle insect pest in Brazil is *Oncideres impluviata* (Coleoptera, Cerambycidae), which has been recorded annually in plantations in the state of Rio Grande do Sul.

Keywords: *Acacia mearnsii* pests, black wattle pest status

INTRODUCTION

Acacia mearnsii De Wild (black wattle) is a small tree or a large shrub native to south-eastern Australia (Nair 2007). The black wattle has high adaptive ability, as evidenced by its presence in several regions of the world (Endress 1994; Yazaki 1997; Seigler 2002). Besides Australia, the black wattle also occurs in South Africa; North, Central and South America; many countries of Asia, particularly in India, Indonesia, and China; and several countries of Europe. It is widely grown by companies and small farmers for commercial purposes (Nair 2007; Searle 1997; Stein and Tonietto 1997; <http://www.cabi.org/isc/datasheet/2326>).

The trees are grown on a large scale because the bark is a good source of tannin, which is essential for leather tanning (Sherry 1971; Schneider 1999; Boland 2006; Nair 2007). Tannin can also be used as anti-corrosive agent

and in water treatment procedures (Mangrich *et al.* 2014). Black wattle is also a source of gum, usually employed in the pharmaceutical and food industries (Searle 1997); and is used as an ornamental plant, to control erosion, as a nitrogen-fixer for soil replenishment, and as raw material for paper (Sherry 1971; Kannegiesser 1990; Lorenzi *et al.* 2003; Caldeira *et al.* 2004).

A. mearnsii can also be used to produce charcoal (Oliveira 1968; Caldeira *et al.* 2003; Riegel *et al.* 2008), and due to its low lignin content, it has been used to produce cellulose (Freddo *et al.* 1999). Mixing *A. mearnsii* with *Pinus elliottii* Engelm and *Eucalyptus grandis* (Hill, 1862) can improve the structure of particleboard (Hillig *et al.* 2002). The plant has also been used in erosion control and reforestation projects (Schneider *et al.* 2005). In Brazil, *A. mearnsii* is economically and socially important in the southern Brazilian state of Rio Grande do Sul, because large numbers of small growers cultivate the tree under contract with companies (Stein and Tonietto 1997).

*Corresponding Author's Email: wacgodoy@usp.br

Among the most common phytosanitation problems for black wattle are: root rot, a disease caused by the fungus *Phytophthora* (Santos *et al.* 2001); and the insect species *Adeloneivaia subangulata* (Herrich-Schäffer, 1855) (Lepidoptera: Saturniidae); *Platypus sulcatus* (Chapuis, 1865) (Coleoptera: Platypodidae); and especially *Oncideres impluviata* (Germar, 1824) (Coleoptera: Cerambycidae), the longhorn beetle (Amante *et al.* 1976; Bressan and Santos 1985; Santana and Santos 2001). In Brazil, *O. impluviata* is currently the main pest of *A. mearnsii* (Amante *et al.* 1976; Baucke 1958; Stein and Tonietto 1997). In this review we present the first compilation of the main black wattle insect pests, and also potential pests of this plant occurring in Brazil.

Acacia spp. in the world and in Brazil

The genus *Acacia* contains about 1300 species (Seigler 2002; Endress 1994), of which more than 800 are endemic to Australia (Yazaki 1997), predominantly in tropical and subtropical areas (Dillon and Dillon 1945; Rice 1986). *A. mearnsii* is the third most cultivated forest species in Brazil along with rubber, teak and araucaria trees, behind *Eucalyptus* spp. and *Pinus* spp. (Abraf 2013). The first seeds were brought from South Africa in 1928 to Rio Grande do Sul (Oliveira 1968; Fleig 1993; Stein and Tonietto 1997) and since 1930 acacia growing has become an important agricultural activity, allowing the country to reach self-sufficiency in tannin production (Schneider and Tonini 2003). Black wattle plantations in Brazil are currently estimated to cover 140,000 to 200,000 hectares (Higa and Resende 1994; Stein and Tonietto 1997; Ildes 2002, 2007; Nair 2007; Attias *et al.* 2013). Many small farmers currently cultivate *A. mearnsii* in Brazil (Attias *et al.* 2013), comprising 60% of the black wattle plantations in the country (Tonietto and Stein 1997). The raw wood produced is exported as wood chips to Japan for the manufacture of cellulose (Lisboa 2009).

The main insect pests of Acacia spp. outside Brazil

Nair (2007) provided a valuable overview of pests of *A. mearnsii* occurring in Australia, South Africa, Kenya, China, India, and included two citations for Brazil. Previous surveys in South Africa also indicated the existence of a significant diversity of phytophagous insects associated with *Acacia* spp. (Hepburn 1966; Swain and Prinsloo 1986), belonging to the orders Coleoptera (46.1%), Lepidoptera (35.7%), Hemiptera (11.8%), Isoptera (3.6%), Psocoptera (1.4%), Orthoptera (0.9%) and Thysanoptera (0.5%). Govender (2007) recorded 14 groups of pests responsible for 60% of the mortality of *A. mearnsii* seedlings; members of the order Coleoptera, including the

family Scarabaeidae and the subfamilies Rutelinae and Melolonthinae, were the dominant taxonomic group.

The main insect pests of A. mearnsii in Brazil

In Brazil, although cultivation of *A. mearnsii* has increased (Abraf 2013), knowledge of the insect fauna associated with the different phenological stages of the plant is still sparse. Comparing the study by Menschoy and Baucke (1966) of insects associated with *Acacia* spp., with the findings of Govender (2007) and the records documented in Brazil (Amante *et al.* 1976; Bressan and Santos 1985; Pedrosa-Macedo *et al.* 1993; Tarragó and Costa 1990; Oliveira *et al.* 2006) suggests that the diversity at the order level present in Brazil is similar to other countries, including the predominance of coleopterans.

Insects are certainly among the main phytosanitation problems in black wattle plantations in Brazil. Their role as pests is directly related to their population density and the size of the cultivated area (Stein and Tonietto 1997). Insects attack the entire plant, ingesting both the bark and leaf tissues, sucking the sap, or attacking the wood, seeds and roots (Searle 1997).

The insects that are most frequently associated with black wattle belong to the orders Coleoptera, Lepidoptera, Hymenoptera and Hemiptera. In Brazil particularly, it is very common to find branches that have been severed by beetles (Amante *et al.* 1976), which significantly alter the tree architecture and can cause significant economic damage (Calderón-Cortés *et al.* 2011). The leaves are generally destroyed by defoliating caterpillars (Lepidoptera) or beetles (Bressan and Santos 1985; Pedrosa-Macedo *et al.* 1993; Tarragó and Costa 1990; Oliveira *et al.* 2006). Ants are important consumers of leaves and seeds (Costa and Link 2014).

Order Coleoptera

Coleopterans are economically important because many species are associated with forests (BertiFilho 1981), seriously damaging the trees in a variety of ways (Pedrosa-Macedo *et al.* 1993; Oliveira *et al.* 2006; Oliveira and Costa 2009). This is the most important order of black wattle insect pests. Below we present records of the presence and damage caused to black wattle by beetles, including comments about the type of injury caused by the insects and also some evidence of their presence in the trees.

Stator limbatus (Horn 1873) is a species of the family Chrysomelidae, which has been associated with black wattle seed predation (Oliveira and Costa 2009). These beetles lay eggs in seeds, their larvae feed on them, and hatching adults make round holes, damaging the seeds. The wood borer *Platypus sulcatus* (Platypodidae) has been

documented in black wattle as building transverse and longitudinal galleries in the trees and consequently weakening the plant support, as well as facilitating the entry of pathogenic microorganisms that cause several diseases (Pedrosa-Macedo *et al.* 1993). Some typical signs of this infestation are the presence of gum and sawdust (Santana and Santos 2001). The occurrence of gummosis caused by the fungus *Phytophthora* sp. is associated with weakening of the tree, as a consequence of attacks by *P. sulcatus* (Santana and Santos 2001).

Two beetle species of the family Scarabaeidae, *Macrodactylus suturalis* (Mannerheim, 1829) (Melolonthinae) and *Rutela lineola* (Linnaeus, 1767) (Rutelinae), have also been observed attacking leaves and inflorescences of *A. mearnsii*. On the other hand, *M. suturalis* has also been observed as a black wattle pollen disperser (Gomes Costa 1943; Oliveira *et al.* 2006; Alves and Marins-Corder 2009).

The beetle *Phaops thunbergi* (Dalman, 1823) (Curculionidae) has been recorded defoliating the *A. mearnsii* follicular system in commercial plantations (Oliveira *et al.* 2006). The species feeds preferentially on young leaves, significantly reducing the leaf area with serious consequences for the photosynthetic activity as well as the development of the plant. Severe attacks on seedlings will kill the plants (Oliveira *et al.* 2006).

The occurrence of members of Scolytinae (Curculionidae) in Brazilian *A. mearnsii* plantations has been recorded by using ethanol-baited flight intercept traps. Machado *et al.* (2014) recorded 25 species of Scolytinae, of which the most abundant were *Hypothenemus eruditus* (Westwood 1836), *Xyleborinus saxeseni* (Ratzeburg 1837), *Hypothenemus* sp., *Microcorthylus quadridens* (Wood 2007) and *Corthylus pharax* (Schedl 1976). A previous study conducted by Murrai (2005) found 37 species, and the main genera were *Ambrosiodmus*, *Hypothenemus*, *Monarthrum*, *Xyleborinus*, *Xyleborus* and *Xylosandrus*. However, these records were obtained using only traps.

One study has reported specimens of Scolytinae collected from branches and trunks, although the species has not yet been identified (Lemes *et al.* 2013). The direct association between beetles and branches is essential for studies emphasizing phytosanitation aspects in *A. mearnsii*, since members of Scolytinae may transmit pathogens, as documented for the fungus *Ceratocystis fimbriata* Ell. & Halst, the agent of mango wilt, which is disseminated by the species *Hypocryphalus mangiferae* Stebbing 1914 (Batista 2010; Ferreira *et al.* 2010). This has particular relevance because *C. fimbriata* has been identified as the fungus causing the similar disease in black wattle in Brazil (Ribeiro *et al.* 1988; Santos 2004).

The family Cerambycidae is the most important taxonomic group for black wattle in Brazil. This family has

more than 25,000 species described in the Neotropical region, with 1,550 genera, and nine of them, which are twig-girdlers, have been recorded in Brazil (Costa Lima 1955; Baucke 1958; Dillon and Dillon 1946; Amante *et al.* 1976; Monné 2005; Galileo and Martins 2006). Members of the genus *Oncideres* are the most frequent and abundant in *A. mearnsii* (Monné 2002), occurring throughout Brazil (Link *et al.* 1994; Witeck Neto and Link 1997; Coutinho *et al.* 1998). Usually, adult females cut branches and lay eggs inside them, changing the entire plant architecture and causing serious economic damage (Amante *et al.* 1976; Calderón-Cortés 1997).

Several twig-girdler species are considered to be potential pests of blackwattle in Brazil (Galileo and Martins 2006). Link *et al.* (1984) listed about 35 plant species capable of hosting *Oncideres saga* (Dalman, 1823), including black wattle (Magistrali *et al.* 2008). Three other twig-girdler species, *O. dejeani* (Thomson, 1868), *O. ocularis* (Thomson, 1868) and *O. impluviata*, have been recorded as injuring black wattle and reaching other Brazilian regions; they have been recorded mainly in Rio Grande do Sul (Baucke 1958; Amante *et al.* 1976; Galileo and Martins 2006; Seffrin *et al.* 2006), probably because this state has the largest area planted to black wattle in the country, with a high concentration of monocultures.

O. impluviata is native to South America, and its host species include members of Mimosaceae, Caesalpinaceae and Aquifoliaceae (Galileo and Martins 2006). This is the most important pest of black wattle currently in Brazil, and the available records of damage indicate its negative effect on the plants, causing heavy damage to the plantations (Cabi 2005). This species attacks black wattles of all ages, but prefers older plants, which have thicker branches (Costa and Marques 1988; Link and Costa 1993). Attack by *O. impluviata* may kill trees that are less than four years old. Older trees may survive but their branches tend to bifurcate, signaling the presence of *O. impluviata* (Amante *et al.* 1976).

O. impluviata females ring the branches of black wattle, cutting them before laying their eggs. This behavior interrupts the sap flow, damaging the tree (Amante *et al.* 1976). The larvae feed on the dry wood, boring into it and building galleries inside the branch (Amante *et al.* 1976). The insect spends about a year in this stage, and in Brazil the adults generally emerge between November and January, when they then infest new trees (Baucke 1958; Amante *et al.* 1976). Control of *O. impluviata* involves manually collecting and burning the branches (Amante *et al.* 1976; Kirch 1983).

Order Lepidoptera

Two species of this order have been cited as the main black wattle pests, *Adeloneivaia subangulata* (Lepidoptera:

Saturniidae) and *Thyrinitea arnobia* (Stoll, 1782) (Lepidoptera, Geometridae). Newly hatched caterpillars of *A. subangulata* scrape the leaf blade, and from the third instar start to eat leaves and apical buds. *T. arnobia* has been recorded as completely defoliating tree tops (Tarragó and Costa 1990). This species is also considered to be the main defoliator of *Eucalyptus* spp., and has been cited as the cause of frequent outbreaks in these plantations (Zanuncio *et al.* 2000).

Other orders

Little information is available concerning other orders of insects with species that damage *A. mearnsii* in Brazil. This section summarizes the information on different orders, which is scattered through the literature. In Hymenoptera, leaf-cutting ants are the most important taxonomic group affecting black wattle plantations. Leaf-cutting ants are currently considered to be the main Brazilian forestry pest, because of their habit of totally eating the leaf blade, causing heavy damage to trees (Della Lucia 2011). In *A. mearnsii* plantations, the leaf-cutting ant most frequently recorded is *Acromyrmex* spp. (Costa *et al.* 2011). Although leafhoppers are often observed on black wattle, mainly on nursery plants, the only record in South America is of *Aethalion reticulatum* (Linnaeus, 1767) (Hemiptera: Aethalionidae) on *Acacia* spp. (Arce de Hamity. 2003).

REMARKS

Based on the benefits that *A. mearnsii* provides to Brazil, especially the southern region, more extensive studies of the insect diversity in black wattles are needed in order to determine the best way to deal with the insect diversity, not only pests, but also natural enemies and beneficial species. This review describes the current status of black wattle insect pests and their significant impacts on the plantations. We hope that new information about insect diversity associated with *A. mearnsii* will soon be available to help in the implementation of alternative pest-management programs, especially the use of natural enemies, to attenuate the negative impacts of pests and make the Brazilian forestry sector sustainable.

ACKNOWLEDGEMENTS

Works by MAO and ENLF were supported by a postgraduate fellowship respectively from CAPES and CNPq.

REFERENCES

Abraf. associação brasileira de produtores de florestas plantadas (2013). Anuário estatístico da ABRAF: Ano base 2012. 130.

- Alves EMS, Marins Corder MP (2009). Biologia reprodutiva de *Acacia mearnsii* De Wild. (Fabaceae) IV: visitantes florais. Rev. Árvore 33: 443-450.
- Amante E, Berlato MA, Gessinger GI (1976). Biologia do "serrador" da acácia-negra, *Oncideres impluviata* (Germar, 1824) (Coleoptera: Cerambycidae) no Rio Grande do Sul. I Etologia. Agron.Sulriograndense 12: 3-56.
- Arce de Hamity MC, Neder de Román LE, Zamar ML (2003). Aspectos bioecológicos de *Aethalion reticulatum* (Linnaeus, 1767) (Hemiptera: Aethalionidae) espécie prejudicial em árvores y arbustos de laPrepunajujeña. Jujuy-Argentina. Idesia21:31-39.
- Attias N, Siqueira MF, Bergallo HG (2013). Acácias australianas no Brasil: histórico, formas de uso e potencial de invasão. Biodiv.Bras. 3: 74-96. At:http://sistemasdeproducao.cnptia.embrapa.br/FontesHTML/Manga/CultivodaMangueira_2ed/doencas.htm - Accessed on: 3 Dec. 2014.
- Batista DC (2010). Cultivo da mangueira. Embrapa Semiárido, Sistemas de Produção, 2 - 2ª edição, ISSN 1807-0027 Online VersionAug/2010.
- Baucke O (1958). Biologia e controle do "serrador da acácia negra" *Oncideres impluviata* (Germ., 1842) (Cerambycidae - Lamiinae - Onciderini). Tese (Doutorado) – Escola de Agronomia "Eliseu Maciel", Universidade Federal de Pelotas, RS. 67.
- Berti Filho E (1981). Insetos associados a plantações de espécies do gênero *Eucalyptus* nos estados da Bahia, Espírito Santo, Mato Grosso do Sul, Minas Gerais e São Paulo. Tese (Doutorado). Escola Superior de Agricultura "Luiz de Queiroz" - ESALQ, Piracicaba, SP. 176.
- Boland DJ, Brooker MIH, Chippendale GM, Hall N, Hyland BPM, Johnston RD, Kleinig DA, Turner JD (2006). Forest trees of Australia. 5th ed. CSIRO Publishing, Collingwood. 768.
- Bressan DA, Santos HR (1985). Avaliação da densidade, viabilidade e do parasitismo em pupas de *Adeloneivaia subangulata* (Herrich-Schäffer, 1855) Travassos, 1940 (Lep. Attacidae) em povoamentos de *Acacia mearnsii* (de Wild.). R. Flor. 15: 38-42. Available at: <http://ojs.c3sl.ufpr.br/ojs2/index.php/floresta/article/viewArticle/6360>. Accessed on: 2 Dec. 2014.
- CABI (Commonwealth Agricultural Bureau International) (2005). Forestry Compendium, 2005 edn., (CD version). Wallingford, UK: CAB International.
- Caldeira MVW, Rondon Neto RM, Schumacher MV (2003). Conteúdo e exportação de micronutrientes em Acácia negra (*Acacia mearnsii* De Wil.) procedência batemansbay (Austrália). R. Árvore 27:9-14.
- Caldeira MVW, Rondon Neto RM, Schumacher MV (2004). Eficiência do uso de micronutrientes e sódio em três procedências de Acácia negra (*Acacia mearnsii* De Wild.). R. Árvore. 28: 39-47.
- Calderón-Cortés N, Quesada M, Escalera-Vasquez LH (2011). Insect as stem engineers: Interactions mediated by the Twig-Girdler *Oncideres albomarginata* chamela enhance arthropod diversity. PLOS ONE 6: e19083.
- Costa EC, D'Avila M, Cantarelli EB, Murari AB (2011). Entomologia florestal. Editora UFSM, Santa Maria. 247.
- Costa EC, Marques EM (1988). Aspectos etológicos de *Oncideres impluviata* (Germar, 1824) em bracinga. Rev. Ci. Rurais 18:219-228.
- Costa Lima AM (1955). Insetos do Brasil. 9º Tomo (Coleópteros). Ministério de Agricultura, Escola Nacional de Agronomia, Rio de Janeiro, Série Didática. 121-124.
- Costa EC, Link D (2014). Alguns problemas entomológicos em essências florestais no Rio Grande do Sul. Available at: <http://ainfo.cnptia.embrapa.br/digital/bitstream/item/102081/1/AlgunsProblemas.pdf> Accessed on: 2 Dec. 2014.
- Coutinho CL, Carvalho AG, Oliveira ES, Veiga BGA (1998). *Oncideres saga* (Dalman,1823) (Coleoptera, Cerambycidae) e a arborização urbana em Seropédica, RJ. Floresta e Ambiente 5:50-54.
- Della Lucia TMC (2011). Formigas cortadeiras, da bioecologia ao manejo. Ed. UFV, Viçosa. 421.
- Dillon LS, Dillon ES (1946). The tribe Onciderini (Coleoptera: Cerambycidae). Part II. scientific publications. Reading Public Museum and Art Gallery, Scientific Publications 6: 189-413.
- Endress PK (1994). Diversity and evolutionary biology of tropical flowers. Cambridge University Press, Cambridge. 284.

- Ferreira EM, Harrington TC, Thorpe DJ, Alfenas AC (2010). Genetic diversity and interfertility among highly differentiated populations of *Ceratocystis fimbriata* in Brazil. *Plant Path.* 59:721-735.
- Fleig FD, Schneider PR, Brum ET (1993). Análise econômica dos sistemas de produção com acácia negra (*Acacia mearnsii* De Wild.) no Rio Grande do Sul. *Ciênc. Flor.* 3: 203-240.
- Freddo A, Foelkel CEB, Frizzo SMB, Silva MCM (1999). Elementos minerais em madeiras de eucalipto e Acácia Negra e sua influência na indústria de celulose Kraft branqueada. *Ciênc. Flor.* 9: 193-209.
- Galileo MHM, Martins UR (2006). Cerambycidae (Coleoptera, Insecta) do Parque Copesul de Proteção Ambiental, Triunfo, Rio Grande do Sul. Museu de Ciências Naturais da Fund. Zoobotânica do RS, Porto Alegre. 314.
- Gomes Costa R (1943). Pragas das plantas cultivadas no Rio Grande do Sul. *Rev. Agric.* 1241-1242.
- Govender P. (2007). Status of seedling establishment pests of *Acacia mearnsii* De Wild. (Mimosaceae) in South Africa. *S. Afr. J. Sci.* 103:141-147.
- Hepburn GA (1966). A revised list of wattle insects and spiders of southern Africa. In: WRI Annual Research Report, Wattle Research Institute, University of Natal 67:83.
- Higa AR, Resende MDV (1994). Breeding *Acacia mearnsii* in Southern Brazil. In: Australian Tree Species Research in China, Zhabgzhou. Canberra: ACIAR. 158-160.
- HilligÉ, Haselein CR, Santini EJ (2002). Propriedades mecânicas e chapas aglomeradas estruturas fabricadas com madeiras de Pinus, Eucalipto e Acácia negra. *Ciênc. Flor.* 12:59-70.
- Ildis (2002). International Legume Database and Information Service. University of Southampton, UK. Available at: <http://www.ildis.org/database/>.
- Ildis (2007). International Legume Database and Information Service. Reading, UK: School of Plant Sciences, University of Reading. Available at: <http://www.ildis.org/>. Accessed on Dec. 04.
- Kannegiesser SU (1990). Apuntes sobre algunas acacias australianas. 1.- *Acaciamearnsii* De Wild. *Ciênc. Invest. Flor.* 2:198-212.
- Kirch E (1983). Estudo dos inimigos naturais de *Oncideres impluviata* (Germar, 1824) em *Mimosa scabrella* Benth. Dissertação (Mestrado), UFPR, Curitiba. 65.
- Lemes PG, Anjos N, Jorge IR (2013). Bioecology of *Oncideres ocularis* Thomson (Coleoptera: Cerambycidae) on *Acacia mangium* Willd. (Fabaceae). *J. Kansas Entomol. Soc.* 86:307-317.
- Link D, Costa EC (1993). Aspectos da biologia de *Oncideres impluviata* e de *O. captiosa* em Santa Maria, RS (Coleoptera: Cerambycidae). *Ciênc. Flor.* 3: 77-84.
- Link D, Costa EC, Alvarez Filho A, Carvalho S, Tarragó MFS (1984). Serrador: levantamento das espécies, épocas de ocorrência e especificidade hospedeira. 2. *Oncideres* spp. e plantas hospedeiras. In: Congresso Florestal Estadual. Anais. 2: 244-254.
- Link D, Costa EC, Thum AB (1994). Binomia comparada dos serradores, *Oncideres saga* (Dalman, 1823) e *Oncideres dejeani* (Thomson, 1868) (Coleoptera: Cerambycidae) em *Parapiptadenia rigida*. *Ciênc. Flor.* 4: 137-144.
- Lisboa RS (2009). Estratégias de suprimento e governança no setor florestal: um estudo de caso múltiplo na indústria de base florestal do Rio Grande do Sul. Tese (Mestrado). UFSM, Santa Maria. 128.
- Lorenzi H, Souza HM, Torres MAV, Bacher L (2003). Árvores Exóticas no Brasil: Madeiras, ornamentais e aromáticas. Nova Odessa: Instituto Plantarum de Estudos da Flora. 352.
- Machado LM, Costa EC, Magistrali IC, Boscardin J, Machado DN, Garlet J (2014). Escolitíneos associados a uma população de acácia-negra (*Acacia mearnsii* De Wild.). *R. Biotemas* 27:57-63.
- Magistrali IC, Magistrali PR, Nadai J, Goellner AA (2008). Parâmetros biológicos de *Oncideres saga* (Dalman, 1823) (Coleoptera: Cerambycidae) em *Acacia mearnsii* De Wild. *R. Trópica - Ciênc. Agr. Biol.* 2: 1-10.
- Mangrich AS, Doumer ME, Mallmannn AS, Wolf CR (2014). Química verde no tratamento de águas: Uso de coagulante derivado de tanino de *Acaciamearnsii*. *R. Virt. Quím.* 6:2-15.
- Menschoy AB, Baucke O (1966). Segunda relação das pragas das plantas cultivadas no Rio Grande do Sul. *Pesq. Agropec. Bras.* 1: 17-46.
- Monné MA (2002). Catalogue of the Neotropical Cerambycidae (Coleoptera) with known host plant. Part V. *Pub. Av. Mus. Nac.* 96: 1-72.
- Monné MA (2005). Catalogue of the Cerambycidae (Coleoptera) of the Neotropical Region. Part II. Subfamily Lamiinae. *Zootaxa* 1023:1-759.
- Murari AB (2005). Levantamento populacional de Scolytidae (Coleoptera) em povoamento de acácia-negra. Dissertação (Mestrado). UFSM, Santa Maria. 63.
- Nair KSS (2007). Tropical Forest Insect Pests: Ecology, Impact, and Management. Cambridge: Cambridge University Press. 404.
- Oliveira HA (1968). Acácia-negra e tanino no Rio Grande do Sul. Canoas: La Salle. 2:121.
- Oliveira LS, Costa EC (2009). Predação de sementes de *Acacia mearnsii* De Wild. (Fabaceae, Mimosoideae). *Biotemas* 22: 39-44.
- Oliveira LS, Costa EC, Cantarelli EB, Perrando ER, Pacheco DDP (2006). Ocorrência de *Phaop sthungeri* (Coleoptera: Curculionidae) em *Acacia mearnsii* De Wild. *Ciênc. Rural* 36: 971-972.
- Pedrosa-Macedo JH, Berti Filho E, Santos HR, Costa EC, Marques EN, Peres Filho O, Mueller JA, Fava HHP, Rocha MP, Pitrowski V, Nascimento EN, Silva LKF (1993). Pragas florestais do sul do Brasil. Manual de Pragas em Florestas. IPEF/SLF. 2:111.
- Ribeiro IJA, Fumikoito M, Paradela Filho O, Castro JL (1988). Gomose da acácia-negra causada por *Ceratocystis fimbriata* Ell. & Halst. *Bragantia* 47:71-74.
- Rice ME (1986). Winter mortality of *Oncideres pustulatus* Le Conte (Coleoptera: Cerambycidae) larvae induced by freezing temperatures. *J. of the Kansas Entomological Society* 5: 423-427.
- Riegel I, Moura ABD, Morisso FDP, Mello FS (2008). Análise termogravimétrica da pirólise da Acácia negra (*Acacia mearnsii* De Wild.) cultivada no Rio Grande do Sul, Brasil. *R. Árvore* 32:533-543.
- Santana DLQ, Santos AF (2001). Ocorrência de *Platypussulcatusem* Acácia-negra (*Acacia mearnsii*). *Bol. Pesq. Flor., Colombo* 42:153-158.
- Santos AF (2004). Murcha de *Ceratocystis* em Acácia-Negra. Comunicado técnico 155, ISSN 1517-5030, Colombo, PR, December, 2004. Available at: http://www.cnpf.embrapa.br/publica/comuntec/edicoes/com_tec125.pdf
- Santos AF, Grigoletti Júnior A, Auer CG, Santana DLQ (2001). O complexo gomose da acácia-negra. Embrapa, Circular Técnica 44, ISSN 1517-5278. Available at: <http://www.cnpf.embrapa.br/publica/circtec/edicoes/circ-tec44.pdf>
- Accessed on: 2 Dec. 2014.
- Schneider PR, Desten G, Brilla, Mainard GL (1999). Determinação da produção de casca em acácia negra. *Ciênc. Flor.* 1:64-75.
- Schneider PR, Finger CAG, Sobrinho VG, Schneider PSP (2005). Determinação indireta do estoque de biomassa e carbono em povoamentos de Acácia-negra (*Acaciamearnsii* De Wild.). *Ciênc. Flor.* 15: 391-402.
- Schneider PR, Tonini H (2003). Utilização de variáveis Dummy em equações de volume para *Acacia mearnsii* De Wild. *Ciênc. Flor.* 13:121-129.
- Searle SD (1997). *Acacia mearnsii* De Wild. (Black Wattle) in Australia. In: Brown AG, Ho CK. (Eds). *BlackWattleand Its Utilisation*. Canberra: Rural Ind. Res. Develop. Corpor. 2-13.
- Seffrin RCAS, Costa EC, Couto MRM, Lopes SJ (2006). Medidas morfométricas de fêmeas e machos de *Oncideres dejeani* Thompson, 1868 (Coleoptera: Cerambycidae). *Ciênc. Rural.* 36:1313-1316.
- Seigler DS (2002). Economic potential from Western Australian *Acacia* species: secondary plant products. *Cons. Sci. West. Aust.* 4: 109-116.
- Sherry SP (1971). The Black Wattle (*Acacia mearnsii*). Pietermaritzburg: University of Natal. 402.
- Stein PP, Toniello L (1997). Black Wattle Silviculture in Brazil. In Brown AG, Ho CK. (Eds). *Black Wattle and Its Utilisation*. Canberra: Rural Industries Res. Develop. Corp. 2-13.

- Swain VM, Prinsloo GL (1986). A List of the Phytophagous Insects and Mites on Forest Trees and Shrubs in South Africa. Department of Agricultural Technical Services, Pretoria. 66:1-91.
- Tarragó MFS, Costa EC (1990). Ocorrência de *Thyriniteina arnobia* (Stoll, 1782) (Lepidoptera: Geometridae), em acácia negra no Rio Grande do Sul. Rev. Ciênc. Rurais 20:219-221.
- Tonietto L, Stein PP (1997). Silvicultura da acácia negra (*Acacia mearnsii* De Wild.) no Brasil. Flor. Estat. 4: 11-16.
- Witeck Neto L, Link D (1997). Cerambycidae associados à Lauraceae, na região central do Rio Grande do Sul, Brasil. Ciênc. Flor.7: 33-39.
- Yazaki Y (1997). *Acacia stroyi*: a potential tannin-producing species. Aust.For. J.60:24-28.
- Zanuncio JC, Zanuncio TV, Lopes ET, Ramalho FS (2000). Temporal variation of Lepidoptera collected in an Eucalyptus plantation in the state of Goiás, Brazil. J. Of Zoology 50:435-443.