

ANAIS



VI SIMPÓSIO BRASILEIRO DE
METALOGENIA
17 A 20/8/2025 - SALVADOR - BA

**A Metalogenia para o
Desenvolvimento do Setor Mineral
Brasileiro.**



UFBA
Universidade
Federal da Bahia



ADIMB
Agência para o Desenvolvimento e
Inovação do Setor Mineral Brasileiro



ANAIS DO VI SIMPÓSIO BRASILEIRO DE METALOGENIA: A Metalogenia para o Desenvolvimento do Setor Mineral Brasileiro

Salvador, 17 a 20 de agosto de 2025

Organizadores:

Comissão Organizadora do VI Simpósio Brasileiro de Metalogenia

ADIMB – Agência para o desenvolvimento Tecnológico

Salvador, Bahia, Brasil

2025



Ficha Catalográfica

Dados Internacionais de Catalogação na Publicação (CIP) (Câmara Brasileira do Livro, SP, Brasil)

Simpósio Brasileiro de Metalogenia
(6. : 2025 : Salvador, BA)
Anais do VI Simpósio Brasileiro de Metalogenia
[livro eletrônico] : a metalogenia para o
desenvolvimento do setor mineral brasileiro /
organização Poliana Iara de Freitas Toledo, Simone
Cerqueira Pereira Cruz. -- Salvador, BA : ADIMB,
2025.

PDF

Vários autores.
ISBN 978-65-988953-0-3

1. Geologia - Brasil 2. Minerais - Brasil
I. Toledo, Poliana Iara de Freitas. II. Cruz, Simone
Cerqueira Pereira. III. Título.

25-304917.0

CDD-549

Índices para catálogo sistemático:

1. Rochas e minerais : Mineralogia 549

Eliane de Freitas Leite - Bibliotecária - CRB 8/8415



3D GEOPHYSICAL MODELING FOR THE EXPLORATION OF IOCG-TYPE DEPOSITS IN THE AQUIRI REGION, CARAJÁS MINERAL PROVINCE, BRAZIL

Serêjo, G.¹; Monteiro, L.V.¹; Marco, C.J.¹, Louro, V.A.², De Oliveira, S.B.¹; Dentith, M.³

¹ Universidade de São Paulo, São Paulo, São Paulo, Brazil, gabriela.serejo.oliveira@usp.br

² CSIRO, Perth, Western Australia, Australia.

³ University of Western Australia, Perth, Western Australia, Australia.

The Aquiri Region, located in the western portion of the Carajás Domain (Carajás Mineral Province), represents an emerging frontier for mineral exploration in Brazil. Similar to the eastern Carajás Domain — which hosts world-class iron oxide-copper-gold (IOCG) deposits — the Aquiri Region contains several promising targets currently under exploration by VALE S. A. , including the AQW2 target. This particular target comprises two prominent magnetic anomalies, referred to as the Deep and Infill targets. Both targets are predominantly hosted within an Archaean-age gabbroic body attributed to the Cateté Suite, structurally controlled by a NW-SE shear zone. The Infill target exhibits characteristics consistent with an iron oxide-copper-gold (IOCG) system. Hydrothermal alteration is zoned and progresses through distinct assemblages: sodic → potassic-sodic → calcic-ferric → ferric-calcic → ferric (Cu-Au). The copper-gold mineralization is associated with the later-stage ferric alteration and is hosted within a breccia system exhibiting a predominant E–W structural orientation. The mineralization consists primarily of chalcopyrite and shows a direct spatial association with magnetite. Geophysical signatures are notably similar to those of IOCG deposits worldwide, characterized by strong and juxtaposed magnetic and gravity anomalies, as well as uranium and potassium enrichment, intermediate resistivity (up to 180 Ohm. m) and high chargeability (up to 20 mV/V). In contrast, the Deep target presents a greater exploration challenge due to the presence of a broad magnetic anomaly that has been poorly tested to date. Limited drilling has revealed hydrothermal alteration patterns similar to those observed at the Infill target, including sodic → potassic-sodic → calcic-ferric assemblages. However, significant ferric alteration associated with magnetite, typically linked to copper-gold mineralization, has not been identified to date. In order to investigate the source of this broad magnetic anomaly, 3D magnetotelluric (MT) and DC-resistivity/induced polarization (DC-IP) models were calculated. The 3D MT resistivity model revealed a large, deep-seated conductivity anomaly extending from approximately 400 to 800 m in depth, dipping eastward, with resistivity values around 10 Ohm·m. In addition, the 3D resistivity and chargeability models from the DC-IP survey identified a significant shallower anomaly, up to 2 km in length, characterized by resistivity values reaching 150 Ohm. m and chargeability values up to 10 mV/V. Both anomalies exhibit geophysical signatures consistent with those of the Infill target, suggesting the presence of a potential new mineralizing system with IOCG deposit characteristics in depth.

Acknowledgements: Special thanks to VALE S.A., the University of São Paulo, CAPES, the University of Western Australia, the Centre for Exploration Targeting, UBC, Terra Resources, and Moombarriga Geoscience for their valuable support and collaboration.