

RT-MAT 2004-24

**ENGEL SUBGROUPS OF TRIANGULAR
MATRICES OVER LOCAL RINGS**

**M. Dokuchaev,
V. Kirichenko and
C. Polcino Milies**

Setembro 2004

Esta é uma publicação preliminar ("preprint").

Engel subgroups of triangular matrices over local rings⁰

M. Dokuchaev¹, V. Kirichenko² and C. Polcino Milies³

^{1,3} Instituto de Matemática e Estatística
Universidade de São Paulo
Caixa Postal 66281, São Paulo, SP
05315-970 – Brazil

²Faculty of Mechanics and Mathematics,
Kiev National Taras Shevchenko University,
Vladimirska Str., 64
01033 Kiev, Ukraine

Mathematics Subject Classifications (2000): Primary 20H25; Secondary 20H20, 20F18, 20F45.

Key words and phrases: triangular matrix group, nilpotent subgroup, Engel subgroup.

Abstract

We consider Engel subgroups of the group $T(n, R)$ of upper triangular matrices over a local ring R which satisfies a weak commutativity condition. If, in addition, R is artinian then we give a complete description of the maximal Engel subgroups of $T(n, R)$ up to conjugacy. These subgroups turn out to be nilpotent and we study their nilpotency class.

1 Introduction

The group $UT(n, K)$ of the upper-unitriangular $n \times n$ -matrices over a field K is a classical example of a nilpotent group which appears naturally in various situations in algebra and geometry. Its direct product with the group of scalar matrices $\{\alpha I \mid 0 \neq \alpha \in K\}$, which we shall denote by $N(n, K)$, gives

⁰This work was partially supported by CNPq and FAPESP of Brazil.

a maximal nilpotent subgroup of the group $T(n, K)$ of the invertible upper-triangular $n \times n$ -matrices over K . It is even maximal Engel. It is easily seen that the group of invertible diagonal matrices, which is obviously abelian, is also maximal Engel. What are the other maximal Engel subgroups in $T(n, K)$? Observe that the group of diagonal matrices is the direct product of n copies of $N(1, K) \cong K^*$.

The main result in this paper (Theorem 3.1) implies that an arbitrary maximal Engel subgroup of $T(n, K)$, up to conjugacy in the general linear group $GL(n, K)$, is a direct product of the form $N(n_1, K) \times N(n_2, K) \times \dots \times N(n_s, K)$ where $n = n_1 + \dots + n_s$. As a consequence, it follows that it is actually nilpotent. Many interesting results about nilpotent linear groups appear in [1], [3] and [4].

We shall use the following notation: given a ring R we write $e_{ij}(r)$, $r \in R$, for the elementary matrix, whose (i, j) th entry is r and all other entries are 0; $T(n, R)$ denotes the group of invertible upper-triangular $n \times n$ -matrices over R , while $UT(n, R)$ stands for the group of upper unitriangular $n \times n$ -matrices, that is the subgroup of $T(n, R)$, generated by the upper-triangular transvections $I + e_{ij}(r)$, $i < j$, $r \in R$. We shall also write

$$g = (g_{ij})_{i \leq j} = \begin{pmatrix} g_{11} & g_{12} & \dots & g_{1n} \\ 0 & g_{22} & \dots & g_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & g_{nn} \end{pmatrix}.$$

In Section 2 some preliminary results are given. In Section 3 we prove that an Engel subgroup of upper-triangular matrices over a commutative local ring R is conjugate, in $UT(n, R)$, to a group G with the following property:

$$\begin{aligned} &\text{If for a pair of indices } i < j \text{ there is an element } g' = (g'_{ij})_{i \leq j} \\ &\text{in } G \text{ such that } g'_{ii} - g'_{jj} \in \mathcal{U}(R) \text{ then } g_{ij} = 0 \text{ for all } g \in G. \end{aligned} \quad (1)$$

Furthermore, with the additional assumption that R is artinian, we describe, up to conjugacy, the maximal Engel subgroups of $T(n, R)$. Moreover, in the latter result we are able to substitute the commutativity condition on a local ring R with maximal ideal \mathcal{M} by the weaker assumption:

$$R/\mathcal{M} \text{ is commutative} \quad \text{and} \quad \mathcal{M}/\mathcal{M}^2 \subseteq 3(R/\mathcal{M}^2), \quad (2)$$

where $\mathfrak{Z}(R/\mathcal{M}^2)$ denotes the centre of R/\mathcal{M}^2 .

Notice that, for arbitrary $r \in R$ and $m_1, m_2 \in \mathcal{M}$ we have that

$$\begin{aligned} [r, m_1 m_2]_{\text{Lie}} &= r m_1 m_2 - m_1 m_2 r = r m_1 m_2 - m_1 (r m_2 + [m_2, r]_{\text{Lie}}) \\ &= (r m_1 - m_1 r) m_2 - m_1 [m_2, r]_{\text{Lie}} \in \mathcal{M}^3. \end{aligned}$$

Thus $\mathcal{M}^2/\mathcal{M}^3 \subseteq \mathfrak{Z}(R/\mathcal{M}^3)$ and an inductive argument readily shows that condition (2) is actually equivalent to the following.

$$\mathcal{M}^k/\mathcal{M}^{k+1} \subseteq \mathfrak{Z}(R/\mathcal{M}^{k+1}), \quad \forall k \geq 0. \quad (3)$$

A non-commutative example of a local ring satisfying condition (2) is given by the group algebra of a finite p -group G over a field K of characteristic p . In fact, it is well known that this is a local ring whose radical is $\Delta(G)$, the augmentation ideal of KG (see [2, p. 27]). Moreover, in any group ring KG we have that

$$\Delta(G)^k/\Delta(G)^{k+1} \subseteq \mathfrak{Z}(KG/\Delta(G)^{k+1}).$$

2 Preliminary results

Lemma 2.1. *Suppose that R is either commutative or an artinian local ring satisfying condition (2). Given an upper-triangular $n \times n$ -matrix $a = (a_{ij})_{i \leq j}$ over R , there is an element $t \in UT(n, R)$ such that for each $1 \leq i < j \leq n$ with $a_{ii} - a_{jj} \in \mathcal{U}(R)$ the (i, j) th entry of $t^{-1}at$ is 0.*

Proof. For a transvection $t = I + e_{ij}(r)$, $i < j$, $r \in R$, we see that $t^{-1}at$ can be obtained from a by adding to the j th column the i th multiplied by r from the right and subtracting from the i th row the j th one multiplied by r from the left. Thus the (i, j) th entry of $t^{-1}at$ is $a_{ij} + a_{ii}r - ra_{jj}$ and the diagonal entries do not change.

Suppose first that R is commutative. Then taking $r = -a_{ij}(a_{ii} - a_{jj})^{-1}$ we see that the (i, j) th entry of $t^{-1}at$ becomes 0. We want to do this for each (i, j) with $a_{ii} - a_{jj} \in \mathcal{U}(R)$ preserving the zeros obtained in previous steps. It is easily seen that this happens if we work parallel to the main diagonal as follows: $(1, 2) \rightarrow (2, 3) \rightarrow \dots \rightarrow (n-1, n) \rightarrow (1, 3) \rightarrow (2, 4) \rightarrow \dots \rightarrow (n-2, n) \rightarrow \dots \rightarrow (1, n)$, omitting, of course, those places (i, j) for which $a_{ii} - a_{jj} \notin \mathcal{U}(R)$. Thus we obtain an element $t \in UT(n, R)$ such that $t^{-1}at$ possesses the desired property.

Suppose now that R is an artinian local ring with maximal ideal \mathcal{M} satisfying (2). Then R/\mathcal{M} is a field and it follows, by the commutative case, that there is $t_1 \in UT(n, R)$ such that the desired property holds for $t_1^{-1}at_1$ modulo \mathcal{M} . Assume by induction that we found already an element $t_k \in UT(n, R)$ such that each (i, j) th entry $a_{ij}^{(k)}$ of $a^{(k)} = t_k^{-1}at_k$ lies in \mathcal{M}^k for every $1 \leq i \leq j \leq n$ with $a_{ii} - a_{jj} \in \mathcal{U}(R)$. Because the element $r = -a_{ij}^{(k)}(a_{ii} - a_{jj})^{-1} \in \mathcal{M}^k$ is central modulo \mathcal{M}^{k+1} , we see that r satisfies $a_{ij}^{(k)} + a_{ii}r - ra_{jj} \equiv 0 \pmod{\mathcal{M}^{k+1}}$. Hence the (i, j) th entry of $(I + e_{ij}(r))^{-1}a^{(k)}(I + e_{ij}(r))$ is contained in \mathcal{M}^{k+1} . Working parallel to the main diagonal as we did in the commutative case we come to an element $t_{k+1} \in UT(n, R)$ such that the (i, j) th entry of $t_{k+1}^{-1}at_{k+1}$ belongs to \mathcal{M}^{k+1} for all $1 \leq i \leq j \leq n$ with $a_{ii} - a_{jj} \in \mathcal{U}(R)$. Since R is artinian, $\mathcal{M}^s = 0$ for some $s \geq 0$ and consequently $t_s^{-1}at_s$ satisfies the required property. \square

Remark 2.2. Observe that conjugation by a transvection t as above does not change the diagonal entries of a .

Lemma 2.3. *Suppose that R is either commutative or a local ring with condition (2) such that $\cap_{l=0}^{\infty} \mathcal{M}^l = 0$. Suppose further that an Engel group G of upper-triangular matrices over R contains an element $g = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ in which*

$$a = \begin{pmatrix} a_1 & & * \\ & \ddots & \\ 0 & & a_k \end{pmatrix}, \quad b = \begin{pmatrix} b_1 & & * \\ & \ddots & \\ 0 & & b_m \end{pmatrix} \quad \text{and } a_i - b_j \in \mathcal{U}(R)$$

for each $1 \leq i \leq k$, $1 \leq j \leq m$. Then any $x \in G$ is of the form

$$x = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \quad \text{with } \alpha = \begin{pmatrix} \alpha_1 & & * \\ & \ddots & \\ 0 & & \alpha_k \end{pmatrix}, \quad \text{and } \beta = \begin{pmatrix} \beta_1 & & * \\ & \ddots & \\ 0 & & \beta_m \end{pmatrix}.$$

Proof. Write $x = \begin{pmatrix} \alpha & \gamma \\ 0 & \beta \end{pmatrix}$ in which α and β are as above and $\gamma = (\gamma_{ij})$ is a $k \times m$ -matrix. We need to show that $\gamma = 0$.

We have that

$$x^{-1} = \begin{pmatrix} \alpha^{-1} & -\alpha^{-1}\gamma\beta^{-1} \\ 0 & \beta^{-1} \end{pmatrix}$$

and

$$x' = [x, g] = \begin{pmatrix} \alpha^{-1}a^{-1}\alpha a & \alpha^{-1}a^{-1}\gamma b - \alpha^{-1}\gamma\beta^{-1}b^{-1}\beta b \\ 0 & \beta^{-1}b^{-1}\beta b \end{pmatrix}.$$

Compute first $\alpha^{-1}a^{-1}\gamma b$. It equals

$$\begin{pmatrix} \alpha_1^{-1}a_1^{-1} & & & * \\ & \alpha_2^{-1}a_2^{-1} & & \\ & & \ddots & \\ 0 & & & \alpha_k^{-1}a_k^{-1} \end{pmatrix} \begin{pmatrix} \gamma_{11} & \gamma_{12} & \dots & \gamma_{1m} \\ \gamma_{21} & \gamma_{22} & \dots & \gamma_{2m} \\ \dots & \dots & \dots & \dots \\ \gamma_{k1} & \gamma_{k2} & \dots & \gamma_{km} \end{pmatrix} \begin{pmatrix} b_1 & & & * \\ & b_2 & & \\ & & \ddots & \\ 0 & & & b_m \end{pmatrix},$$

which belongs to

$$\begin{pmatrix} \alpha_1^{-1}a_1^{-1}\gamma_{11} + \sum_{i \geq 2} R\gamma_{i1} & \alpha_1^{-1}a_1^{-1}\gamma_{12} + \sum_{i \geq 2} R\gamma_{i2} & \dots & \alpha_1^{-1}a_1^{-1}\gamma_{1m} + \sum_{i \geq 2} R\gamma_{im} \\ \alpha_2^{-1}a_2^{-1}\gamma_{21} + \sum_{i \geq 3} R\gamma_{i1} & \alpha_2^{-1}a_2^{-1}\gamma_{22} + \sum_{i \geq 3} R\gamma_{i2} & \dots & \alpha_2^{-1}a_2^{-1}\gamma_{2m} + \sum_{i \geq 3} R\gamma_{im} \\ \dots & \dots & \dots & \dots \\ \alpha_k^{-1}a_k^{-1}\gamma_{k1} & \alpha_k^{-1}a_k^{-1}\gamma_{k2} & \dots & \alpha_k^{-1}a_k^{-1}\gamma_{km} \end{pmatrix} \times$$

$$\times \begin{pmatrix} b_1 & & & * \\ & b_2 & & \\ & & \ddots & \\ 0 & & & b_m \end{pmatrix}.$$

Similarly, $\alpha^{-1}\gamma\beta^{-1}b^{-1}\beta b$ is contained in

$$\begin{pmatrix} \alpha_1^{-1}\gamma_{11} + \sum_{i \geq 2} R\gamma_{i1} & \alpha_1^{-1}\gamma_{12} + \sum_{i \geq 2} R\gamma_{i2} & \dots & \alpha_1^{-1}\gamma_{1m} + \sum_{i \geq 2} R\gamma_{im} \\ \alpha_2^{-1}\gamma_{21} + \sum_{i \geq 3} R\gamma_{i1} & \alpha_2^{-1}\gamma_{22} + \sum_{i \geq 3} R\gamma_{i2} & \dots & \alpha_2^{-1}\gamma_{2m} + \sum_{i \geq 3} R\gamma_{im} \\ \dots & \dots & \dots & \dots \\ \alpha_k^{-1}\gamma_{k1} & \alpha_k^{-1}\gamma_{k2} & \dots & \alpha_k^{-1}\gamma_{km} \end{pmatrix} \times \beta^{-1}b^{-1}\beta b.$$

Suppose now that R is commuative. Then

$$\beta^{-1}b^{-1}\beta b = \begin{pmatrix} 1 & & * \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{pmatrix}.$$

We see that the $(k, 1)$ th entry of $\gamma^{(l)} = \alpha^{-1}a^{-1}\gamma b - \alpha^{-1}\gamma\beta^{-1}b^{-1}\beta b$ is $\alpha_k^{-1}\gamma_{k1}(a_k^{-1}b_1 - 1)$.

$$\text{Write } x^{(l)} = [x, \underbrace{g, \dots, g}_l] = \begin{pmatrix} \alpha^{(l)} & \gamma^{(l)} \\ 0 & \beta^{(l)} \end{pmatrix}, \gamma^{(l)} = \begin{pmatrix} \gamma_{11}^{(l)} & \gamma_{12}^{(l)} & \dots & \gamma_{1m}^{(l)} \\ \dots & \dots & \dots & \dots \\ \gamma_{k1}^{(l)} & \gamma_{k2}^{(l)} & \dots & \gamma_{km}^{(l)} \end{pmatrix}.$$

Since R is commutative,

$$\alpha^{(l)} = \begin{pmatrix} 1 & & * \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{pmatrix} \text{ and } \beta^{(l)} = \begin{pmatrix} 1 & & * \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{pmatrix}.$$

The above calculations with $x = x^{(l-1)}$ tell us that $\gamma_{k1}^{(l)} = \alpha_k^{-1}\gamma_{k1}(a_k^{-1}b_1 - 1)^l$ for each $l \geq 1$. Since α_k^{-1} and $a_k^{-1}b_1 - 1$ are units in R and G is Engel, it follows that γ_{k1} must be 0. Hence $\gamma_{k1}^{(l)} = 0$ for all $l \geq 1$. Then

$$\gamma_{k-1,1}^{(1)} = \alpha_{k-1}^{-1}\gamma_{k-1,1}(a_{k-1}^{-1}b_1 - 1)$$

and, more generally,

$$\gamma_{k-1,1}^{(l)} = \alpha_{k-1}^{-1}\gamma_{k-1,1}(a_{k-1}^{-1}b_1 - 1)^l, \quad l \geq 1.$$

We derive again from the Engel property of G that $\gamma_{k-1,1} = 0$ and, consequently, $\gamma_{k-1,1}^{(l)} = 0$ for all $l \geq 1$. Going up this way by the first column we conclude that $\gamma_{11}^{(l)} = \gamma_{21}^{(l)} = \dots = \gamma_{k1}^{(l)} = 0$ for each $l \geq 1$. Suppose by induction that the first s columns of $\gamma^{(l)}$ are all zero for each $l \geq 1$. Then $\gamma_{k,s+1}^{(l)} = \alpha_k^{-1}\gamma_{k,s+1}(a_k^{-1}b_{s+1} - 1)^l$. As above, this yields $\gamma_{k,s+1}^{(l)} = 0$ for every $l \geq 1$. It is easily seen that we can go up by the $(s+1)$ st column of γ as we did for the first one and conclude that it is also zero. Hence the $(s+1)$ st column of $\gamma^{(l)}$ is zero for each $l \geq 1$. It follows by induction that γ is the zero $k \times m$ -matrix.

Next we adjust our proof for the case when R is a local ring with $\cap_{r=0}^{\infty} \mathcal{M}^r = 0$ and which satisfies (2). We keep the above notation. Because R/\mathcal{M} is a

field, it follows by the commutative case that all entries of γ are in \mathcal{M} . Suppose by induction we know already that each entry of γ is in \mathcal{M}^r . Thus every γ_{ij} is central modulo \mathcal{M}^{r+1} . Since working modulo \mathcal{M} we have

$$\beta^{-1}b^{-1}\beta b \equiv \begin{pmatrix} 1 & & * \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{pmatrix},$$

it follows that $\gamma_{k1}^{(1)} \equiv \alpha_k^{-1}\gamma_{k1}(a_k^{-1}b_1 - 1) \bmod \mathcal{M}^{r+1}$. Furthermore,

$$\alpha^{(l)} \equiv \begin{pmatrix} 1 & & * \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{pmatrix} \bmod \mathcal{M}, \beta^{(l)} \equiv \begin{pmatrix} 1 & & * \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{pmatrix} \bmod \mathcal{M}$$

and hence $\gamma_{k1}^{(l)} \equiv \alpha_k^{-1}\gamma_{k1}(a_k^{-1}b_1 - 1)^l \bmod \mathcal{M}^{r+1}$ for each $l \geq 1$. The Engel property of G implies that $\gamma_{k1}^{(l)} \in \mathcal{M}^{r+1}$ for all $l \geq 1$. The rest of the argument goes as in the commutative case and shows that all entries of γ are in \mathcal{M}^{r+1} . Hence, by induction, they are contained in all powers of \mathcal{M} and we conclude from $\cap_{r=0}^{\infty} \mathcal{M}^r = 0$ that γ is the zero matrix. □

Corollary 2.4. *Let R be as in Lemma 2.3. Suppose further that an Engel group G of upper-triangular matrices over R contains an element $g = (g_{ij})_{i \leq j}$ such that, for some index i , we have $g_{i,i} - g_{i+1,i+1} \in \mathcal{U}(R)$ and $g_{i,i+1} = 0$. Then $x_{i,i+1} = 0$ for all $x = (x_{ij})_{i \leq j} \in G$.*

Proof. Notice that the set

$$\left\{ \begin{pmatrix} x_{i,i} & x_{i,i+1} \\ 0 & x_{i+1,i+1} \end{pmatrix} : x \in G \right\}$$

is an Engel group of upper-triangular matrices over R . By the previous lemma, we readily obtain that $x_{i,i+1} = 0$ for all $x \in G$. □

3 The Engel subgroups of upper-triangular matrices

We recall that, for a group G of upper-triangular matrices over R , property (1) is as follows:

If for a pair of indices $i < j$ there is an element $g' = (g'_{ij})_{i \leq j}$ in G such that $g'_{ii} - g'_{jj} \in \mathcal{U}(R)$ then $g_{ij} = 0$ for all $g \in G$.

Theorem 3.1. *Let G be an Engel subgroup of upper triangular $n \times n$ -matrices over a local ring R . Suppose further that R is either commutative or artinian with condition (2). Then G is conjugate in $UT(n, R)$ to a subgroup with property (1).*

Proof. For simplicity we shall say that an upper-triangular matrix $g = (g_{ij})_{i \leq j}$ with entries in R satisfies property (x) if

$$g_{ii} - g_{jj} \in \mathcal{U}(R) \Rightarrow g_{ij} = 0.$$

Let \mathcal{M} be the (unique) maximal ideal of R and $g = (g_{ij})_{i \leq j}$ be an arbitrary fixed element of G . By Lemma 2.1 we may assume, up to conjugacy in $UT(n, R)$, that g satisfies property (x). Suppose that, for some index i , the diagonal entries g_{ii} and $g_{i+1, i+1}$ of g are not congruent modulo \mathcal{M} . By property (x) we have that $g_{i, i+1} = 0$. Let h be the matrix obtained from the identity $n \times n$ -matrix by interchanging its i th and $(i+1)$ st rows:

$$h = \begin{pmatrix} I & & \\ & 0 & 1 \\ & 1 & 0 \\ & & I \end{pmatrix}, \quad (4)$$

It is readily seen that conjugation by h transposes the i th and $(i+1)$ st rows and also the i th and $(i+1)$ st columns. Since $g_{i, i} - g_{i+1, i+1} \in \mathcal{U}(R)$, $g_{i, i+1} = 0$ and G is Engel, we know from Corollary 2.4, that $x_{i, i+1} = 0$ for all $x \in G$. Hence, all elements of G remain triangular under conjugation by h . Moreover, $h^{-1}gh$ still has property (x) and its (i, i) th and $(i+1, i+1)$ st diagonal entries are permuted. Applying this type of conjugation several times if necessary, we see that G is conjugate by an element $\tilde{h} \in GL(n, R)$ to a group \tilde{G} where g becomes

$$\tilde{g} = \text{diag}(a, b) = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, \quad (5)$$

in which a and b are upper-triangular matrices, the diagonal entries of a are pairwise congruent modulo \mathcal{M} and no diagonal entry of a is congruent modulo \mathcal{M} to a diagonal entry of b . The element \tilde{h} is a product of permutation matrices of the form (4). By Lemma 2.3, any x of \tilde{G} is of the form $x = \text{diag}(\alpha_x, \beta_x)$ where α_x and β_x have the same sizes as a and b respectively. Let $\tilde{G}_1 = \langle \alpha_x : x \in \tilde{G} \rangle$, $\tilde{G}_2 = \langle \beta_x : x \in \tilde{G} \rangle$. Since \tilde{G}_1 and \tilde{G}_2 are Engel groups of upper-triangular matrices of smaller sizes, induction on n implies that there exists $t_1 \in UT(n_1, R)$ and $t_2 \in UT(n_2, R)$, where n_1 and n_2 are the sizes of a and b respectively, such that both $t_1^{-1}\tilde{G}_1t_1$ and $t_2^{-1}\tilde{G}_2t_2$ satisfy (1). Then, taking $t = \text{diag}(t_1, t_2)$, the group $t^{-1}\tilde{G}t$ obviously verifies (1). Now, let

$$\tilde{\tilde{G}} = \tilde{h}t^{-1}\tilde{G}t\tilde{h}^{-1} = (\tilde{h}t\tilde{h}^{-1})^{-1}\tilde{G}(\tilde{h}t\tilde{h}^{-1}).$$

Because conjugation by \tilde{h} preserves property (1) we see that $\tilde{\tilde{G}}$ is upper-triangular and satisfies (1). Hence we will be done if we show that $\tilde{h}t\tilde{h}^{-1} \in UT(n, R)$.

Without loss of generality we may suppose that t_1 and t_2 are transvections. Write $\hat{t}_1 = \text{diag}(t_1, I)$, $\hat{t}_2 = \text{diag}(I, t_2) \in UT(n, R)$ where the I 's are identity matrices of appropriate sizes. Then evidently $t = \hat{t}_1\hat{t}_2$. Write also $\tilde{h} = h_{\tau_1}h_{\tau_2} \dots h_{\tau_m}$, where $h_{\tau_1}, h_{\tau_2}, \dots$ are permutation matrices of the form (4) and τ_1, τ_2, \dots denote the corresponding permutations (in fact, transpositions) of indices. Observe, in particular, that τ_m is the transposition $(n_1, n_1 + 1)$, since the indices which are involved in a are $1, \dots, n_1$. Now, notice that when we conjugate an upper-triangular transvection t' , say, by one of the permutation matrices h_τ involved in \tilde{h} we obtain again an upper-triangular transvection unless τ interchanges the two indices which determine t' . However, we can avoid this in our case, as shown below.

We remark that conjugation by h_{τ_m} induces a permutation of indices after which we can distinguish two sets which form a partition of the set of all indices: the set \mathfrak{a}_m of those indices coming from $\{1, \dots, n_1\}$, the set of indices of a , and the set \mathfrak{b} of those coming from the set of indices of b . Moreover $h_{\tau_m}\hat{t}_1h_{\tau_m}^{-1}$ is an upper-triangular transvection whose indices are both in \mathfrak{a}_m . In a similar way, $h_{\tau_m}\hat{t}_2h_{\tau_m}^{-1}$ is an upper-triangular transvection whose indices are both in \mathfrak{b}_m .

Next, since conjugation by $h_{\tau_{m-1}}$ permutes one index from \mathfrak{a}_m with one from \mathfrak{b}_m , we again obtain two disjoint sets of indices \mathfrak{a}_{m-1} and \mathfrak{b}_{m-1} , as above, and such that $h_{\tau_{m-1}}h_{\tau_m}\hat{t}_1h_{\tau_m}^{-1}h_{\tau_{m-1}}^{-1}$ and $h_{\tau_{m-1}}h_{\tau_m}\hat{t}_2h_{\tau_m}^{-1}h_{\tau_{m-1}}^{-1}$ are upper-triangular transvections whose indices are in \mathfrak{a}_{m-1} and \mathfrak{b}_{m-1} respectively.

Inductively, we obtain that $h\hat{t}_1h^{-1}$ and $h\hat{t}_2h^{-1}$ are both upper triangular transvections and the result follows. \square

Denote by $N(n, R)$ the group of all invertible upper-triangular $n \times n$ -matrices $g = (g_{ij})_{i \leq j}$ with entries in a local ring R such that $g_{ii} \equiv g_{jj} \pmod{\mathcal{M}}$ for all $1 \leq i, j \leq n$.

Theorem 3.2. *Let R be a local artinian ring with (unique) maximal ideal \mathcal{M} which satisfies condition (2). For any fixed decomposition $n = n_1 + \dots + n_s$ the group*

$$G_{n_1, \dots, n_s} = \begin{pmatrix} N(n_1, R) & & & 0 \\ & N(n_2, R) & & \\ & & \ddots & \\ 0 & & & N(n_s, R) \end{pmatrix},$$

is nilpotent and is a maximal Engel subgroup of $T(n, R)$. Moreover, every maximal Engel subgroup of $T(n, R)$ is conjugate in $GL(n, R)$ to one of the groups G_{n_1, \dots, n_s} .

Proof. Notice that, by Theorem 3.1, every Engel subgroup $G \subseteq T(n, R)$ is conjugate to a group satisfying property (1). Conjugating by permutation matrices of the form (4) we can determine a decomposition $n = n_1 + \dots + n_s$ such that G is conjugate in $GL(n, R)$ to a subgroup of G_{n_1, \dots, n_s} . Thus, it remains to show that G_{n_1, \dots, n_s} is nilpotent.

Evidently, $G_{n_1, \dots, n_s} \simeq N(n_1, R) \times N(n_2, R) \times \dots \times N(n_s, R)$ and hence it suffices to show that $G = N(n, R)$ is nilpotent ($n \geq 1$).

Let m be the nilpotency index of \mathcal{M} , that is $\mathcal{M}^m = 0$ and $\mathcal{M}^{m-1} \neq 0$. Observe that (2) implies that the $(k+1)$ st term $\Gamma_{k+1}(\mathcal{U}(R))$ of the lower central series of $\mathcal{U}(R)$ is contained in $1 + \mathcal{M}^k$, $k \geq 1$, and thus $\mathcal{U}(R)$ is nilpotent of class at most m . Write

$$\mathcal{N}^{(1)} = \begin{pmatrix} 0 & R & R & R & \dots \\ 0 & 0 & R & R & \dots \\ 0 & 0 & 0 & R & \dots \\ 0 & 0 & 0 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}, \quad \mathcal{N}^{(2)} = \begin{pmatrix} 0 & \mathcal{M} & R & R & \dots \\ 0 & 0 & \mathcal{M} & R & \dots \\ 0 & 0 & 0 & \mathcal{M} & \dots \\ 0 & 0 & 0 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix},$$

$$\mathcal{N}^{(3)} = \begin{pmatrix} 0 & \mathcal{M}^2 & \mathcal{M} & R & \dots \\ 0 & 0 & \mathcal{M}^2 & \mathcal{M} & \dots \\ 0 & 0 & 0 & \mathcal{M}^2 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}, \quad \mathcal{N}^{(4)} = \begin{pmatrix} 0 & \mathcal{M}^3 & \mathcal{M}^2 & \mathcal{M} & \dots \\ 0 & 0 & \mathcal{M}^3 & \mathcal{M}^2 & \dots \\ 0 & 0 & 0 & \mathcal{M}^3 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix},$$

etc. As \mathcal{M} is a nilpotent ideal of index m , the last non-zero ideal will be

$$\mathcal{N}^{(n+m-2)} = \begin{pmatrix} 0 & 0 & \dots & 0 & \mathcal{M}^{m-1} \\ 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}.$$

Denote by D the subgroup of diagonal elements of G . We claim that

$$\begin{aligned} G \supset \Gamma_2(D) + \mathcal{N}^{(1)} \supset \Gamma_3(D) + \mathcal{N}^{(2)} \supset \dots \supset \Gamma_m(D) + \mathcal{N}^{(m-1)} \supset I + \mathcal{N}^{(m)} \supset \dots \\ \supset I + \mathcal{N}^{(n+m-2)} \supset I \end{aligned} \quad (6)$$

is a central series of G . Indeed, given a diagonal matrix $d = \text{diag}(g_{11}, \dots, g_{nn}) \in G$ and a transvection $t = I + e_{ij}(x)$ with $x \in \mathcal{M}^l$ we have that

$$[d, t] = I + e_{ij}(g_{ii}^{-1}(g_{ii}x - xg_{jj})) \in I + e_{ij}(\mathcal{M}^{l+1}), \quad (7)$$

as $g_{ii} - g_{jj} \in \mathcal{M}$. Moreover, for any two transvections with $i \neq j'$, one has

$$[I + e_{ij}(x), I + e_{i'j'}(y)] = I + \delta_{j,i'} e_{ij'}(xy), \quad (8)$$

where $\delta_{j,i'}$ is the Kronecker delta. Notice that, if $j \geq j'$ then the right-hand side of (8) is equal to I .

Since G is generated by $UT(n, R)$ and D , it follows in view of (7) and (8) that $[G, I + \mathcal{N}^{(k)}] \subseteq I + \mathcal{N}^{(k+1)}$ for all $k \geq 1$. Moreover, for any $h = \text{diag}(h_{11}, \dots, h_{nn}) \in \Gamma_{k+1}(D)$ one has $h_{ii} \in \Gamma_{k+1}(\mathcal{U}(R)) \subseteq 1 + \mathcal{M}^k$, $i = 1, \dots, n$. Hence taking $t = I + e_{ij}(x)$ with $x \in R$ we see by (7) that $[t, d] \in I + e_{ij}(\mathcal{M}^k)$ and, consequently, $[UT(n, R), \Gamma_{k+1}(D)] \subseteq I + \mathcal{N}^{(k+1)}$. This yields that (6) is a central series and $G = N(n, R)$ is nilpotent. \square

Let R be as in Theorem 3.2 and let τ be a partition $\{1, \dots, n\} = \tau_1 \cup \dots \cup \tau_s$ of $\{1, \dots, n\}$ into a disjoint union of subsets. Denote by G_τ the group of all matrices $(g_{ij})_{i \leq j} \in T(n, R)$ such that

- (i) if $i, j \in \tau_k$ for some $k \in \{1, \dots, s\}$ then $g_{ii} \equiv g_{jj} \pmod{\mathcal{M}}$;
- (ii) $g_{ij} = 0$ for all $i \in \tau_k$ and $j \in \tau_{k'}$ with $k \neq k'$.

Corollary 3.3. *The groups G_τ , with τ running over all partitions of $\{1, \dots, n\}$ into disjoint union of subsets, are nilpotent, pairwise non-conjugate in $T(n, R)$ and every maximal Engel subgroup of $T(n, R)$ is conjugate to one of them. Moreover, all the elements in the conjugacy class of G_τ are obtained conjugating G_τ by the elements of $UT(n, R)$.*

Proof. By Theorem 3.1 every Engel subgroup of $T(n, R)$ satisfies property (1) and hence, up to conjugacy in $UT(n, R)$, is a subgroup of G_τ for some partition τ . It follows from the proof of Theorem 3.1 that G_τ is conjugate in $GL(n, R)$ to G_{n_1, \dots, n_s} where n_i is the number of elements in τ_i ($i = 1, \dots, s$). Hence G_τ is maximal Engel and nilpotent in view of Theorem 3.2. \square

From Theorem 3.2 we readily obtain the following.

Corollary 3.4. *The groups $G_{n_1, n_2, \dots}$ with $n = n_1 + n_2 + \dots$ running over all decompositions such that $n_1 \geq n_2 \geq \dots$, form a full set of representatives of the isomorphism classes of the maximal Engel subgroups of $T(n, R)$.*

The length of the central series (6) of the group $G = N(n, R)$ is $n + m - 1$, so we wonder whether or not it is the nilpotency class of G . We examine this by analyzing the lower central series of $N(n, R)$. It was seen above that $[G, I + \mathcal{N}^{(k)}] \subseteq I + \mathcal{N}^{(k+1)}$. We have that

$$[D, I + e_{ij}(\mathcal{M}^l)] = I + e_{ij}(\mathcal{M}^{(l+1)}) \quad (9)$$

for each $l \geq 0$. In fact, take $I + e_{ij}(z)$ with $z \in \mathcal{M}^{l+1}$. Write $z = \sum xy$ where $x \in \mathcal{M}^l, y \in \mathcal{M}$. Then $d = \text{diag}(1, \dots, 1, 1 - y, 1, \dots, 1) \in D$ with $1 - y$ placed in the j th position. By (7) we have that $[d, I + e_{ij}(x)] = I + e_{ij}(xy)$ and thus $I + e_{ij}(z) = \prod I + e_{ij}(xy) \in [D, I + e_{ij}(\mathcal{M}^l)]$ which means that $[D, I + e_{ij}(\mathcal{M}^l)] \supseteq I + e_{ij}(\mathcal{M}^{l+1})$, proving (9).

Using formulas (8) and (9), it follows that the commutator subgroup of $N(n, R)$ is

$$[D(I + \mathcal{N}^{(1)}), D(I + \mathcal{N}^{(1)})] = \Gamma_2(D)(I + \mathcal{N}^{(2)}).$$

Next for $g = \text{diag}(g_{11}, \dots, g_{nn}) \in \Gamma_2(D)$ observe that $g_{ii} - g_{jj} \in \mathcal{M}^2$ for any $1 \leq i, j \leq n$. Indeed, it will suffice to prove our claim in the case when g is a generator of $\Gamma_2(D)$ and thus $g_{ii} = [h_{ii}, f_{ii}]$ and $g_{jj} = [h_{jj}, f_{jj}]$ for some $h_{ii}, f_{ii}, h_{jj}, f_{jj} \in \mathcal{U}(R)$. We know that $h_{jj} = h_{ii}(1+x)$ and $f_{jj} = f_{ii}(1+y)$ with $x, y \in \mathcal{M}$. Then

$$g_{jj} = [h_{ii}(1+x), f_{ii}(1+y)] = ([h_{ii}, 1+y][h_{ii}, f_{ii}]^{(1+y)})^{(1+x)}[1+x, 1+y][1+x, f_{ii}]^{(1+y)}.$$

Since \mathcal{M} is central modulo \mathcal{M}^2 it follows that $g_{jj} \in [h_{ii}, f_{ii}](1 + \mathcal{M}^2)$ and consequently $g_{ii} - g_{jj} \in \mathcal{M}^2$ as desired.

We see now that if $g = \text{diag}(g_{11}, \dots, g_{nn}) \in \Gamma_2(D)$ then for any $I + e_{ij}(z) \in I + \mathcal{N}^{(1)}$ we have by (7) that

$$[I + e_{ij}(z), g] \subseteq I + e_{ij}(\mathcal{M}^2) \subseteq I + \mathcal{N}^{(3)},$$

so that

$$[I + \mathcal{N}^{(1)}, \Gamma_2(D)] \subseteq I + \mathcal{N}^{(3)}$$

and thus, in view of formula (9), the third term of the lower central series of $N(n, R)$ is

$$[D(I + \mathcal{N}^{(1)}), \Gamma_2(D)(I + \mathcal{N}^{(2)})] = \Gamma_3(D)(I + \mathcal{N}^{(3)}).$$

Is it true in general that $g_{ii} - g_{jj} \in \mathcal{M}^k$ for any $1 \leq i, j \leq n$ if $g = \text{diag}(g_{11}, \dots, g_{nn}) \in \Gamma_k(D)$ and $k \geq 1$? If we assume this for $k-1$, we may write $g_{ii} = [h_{ii}, f_{ii}]$, $g_{jj} = [h_{jj}, f_{jj}]$, $h_{jj} = h_{ii}(1+x)$, $f_{jj} = f_{ii}(1+y)$ with $h_{ii} \in \Gamma_{k-1}(\mathcal{U}(R))$, $f_{ii} \in \mathcal{U}(R)$, $x \in \mathcal{M}^{k-1}$ and $y \in \mathcal{M}$. Then, since clearly

$$[1+x, 1+y][1+x, f_{ii}]^{(1+y)} \in 1 + \mathcal{M}^k,$$

the above commutator calculation shows that $g_{jj} \in g_{ii}(1 + \mathcal{M}^k)$ if and only if $[h_{ii}, 1+y][h_{ii}, f_{ii}]^{(1+y)} \in g_{ii}(1 + \mathcal{M}^k)$. Since $[h_{ii}, f_{ii}] = g_{ii} \in 1 + \mathcal{M}^{k-1}$, we have $[g_{ii}, 1+y] \in 1 + \mathcal{M}^k$ and thus

$$g_{ii} - g_{jj} \in \mathcal{M}^k \iff [h_{ii}, 1+y] \in 1 + \mathcal{M}^k.$$

We conclude from the above argument that if $[\Gamma_{l-1}(\mathcal{U}(R)), 1 + \mathcal{M}] \subseteq 1 + \mathcal{M}^l$ for all $2 \leq l \leq k$ then for each such l and any $g = \text{diag}(g_{11}, \dots, g_{nn}) \in \Gamma_l(D)$ one has $g_{ii} - g_{jj} \in \mathcal{M}^l$ for all $1 \leq i, j \leq n$. Applying (7) again we see that $[I + \mathcal{N}^{(1)}, \Gamma_l(D)] \subseteq I + \mathcal{N}^{(l+1)}$ which yields that the $(l+1)$ st term of the lower central series of $N(n, R)$ is

$$[D(I + \mathcal{N}^{(l)}), \Gamma_l(D)(I + \mathcal{N}^{(l)})] = \Gamma_{l+1}(D)(I + \mathcal{N}^{(l+1)}), \quad (10)$$

$2 \leq l \leq k$.

Thus we have shown that if $[\Gamma_{k-1}(\mathcal{U}(R)), 1 + \mathcal{M}] \subseteq 1 + \mathcal{M}^k$ does hold for all k then all terms of the lower central series of $N(n, R)$ are determined by (10) and the nilpotency class of $N(n, R)$ is $n + m - 2$. This obviously happens if $\Gamma_2(\mathcal{U}(R)) \subseteq 1 + \mathcal{M}^2$ as, by induction, $[\Gamma_{k-1}(\mathcal{U}(R)), 1 + \mathcal{M}] \subseteq [1 + \mathcal{M}^{k-1}, 1 + \mathcal{M}] \subseteq 1 + \mathcal{M}^k$ for all k .

On the other hand, this is true also if $[\mathcal{M}, \mathcal{M}]_{\text{Lie}} \subseteq \mathcal{M}^3$. Indeed, induction shows that $[\mathcal{M}^{k-2}, \mathcal{M}]_{\text{Lie}} \subseteq \mathcal{M}^k$. Since $\Gamma_{k-1}(\mathcal{U}(R)) \subseteq 1 + \mathcal{M}^{k-2}$, it follows again that $[1 + \mathcal{M}^{k-1}, 1 + \mathcal{M}] \subseteq 1 + \mathcal{M}^k$ for all k .

Suppose that $[\Gamma_{k-1}(\mathcal{U}(R)), 1 + \mathcal{M}] \subseteq 1 + \mathcal{M}^k$ does not hold for all k and let k_0 be the least integer with $[\Gamma_{k_0}(\mathcal{U}(R)), 1 + \mathcal{M}] \not\subseteq 1 + \mathcal{M}^{k_0+1}$. Evidently $2 \leq k_0 \leq m - 1$. It follows from the above consideration that if $h \in \Gamma_{k_0}(\mathcal{U}(R))$ and $y \in \mathcal{M}$ are such that $[h, 1 + y] \notin 1 + \mathcal{M}^{k_0+1}$, then taking $h' = \text{diag}(h, \dots, h)$ and $f = \text{diag}(1, 1 + y, 1, \dots)$ we see that $g = [h', f] = \text{diag}(1, [h, 1 + y], 1, \dots, 1) \in \Gamma_{k_0+1}(D)$ and by (7),

$$[g, I + e_{12}(1)] = I + e_{12}(v), \text{ with } v = 1 - [h, 1 + y] \in \mathcal{M}^{k_0} \setminus \mathcal{M}^{k_0+1}.$$

Thus $\tilde{\mathcal{Z}} = [I + \mathcal{N}^{(1)}, \Gamma_{k_0+1}(D)] \not\subseteq I + \mathcal{N}^{(k_0+2)}$. Since $\Gamma_{k_0+1}(\mathcal{U}(R)) \subseteq 1 + \mathcal{M}^{k_0}$ one has that $\tilde{\mathcal{Z}} \subseteq I + \mathcal{N}^{(k_0+1)}$. Then the $(k_0 + 2)$ nd term of the lower central series of $N(n, R)$ is

$$\begin{aligned} [D(I + \mathcal{N}^{(1)}), \Gamma_{k_0+1}(D)(I + \mathcal{N}^{(k_0+1)})] &= \Gamma_{k_0+2}(D)\tilde{\mathcal{Z}}(I + \mathcal{N}^{(k_0+2)}) \\ &\supsetneq \Gamma_{k_0+2}(D)(I + \mathcal{N}^{(k_0+2)}). \end{aligned}$$

Suppose that $k_0 = m - 1$. If $n = 2$, then $k_0 + 2 = m - 1 + 2 = n + m - 1$ and the above calculated term $\Gamma_{k_0+2}(N(n, R))$ is the last non-identity term of the lower central series of $N(n, R)$. If $n > 2$, taking an element from $\tilde{\mathcal{Z}}$ which is not contained in $I + \mathcal{N}^{(k_0+2)}$ we see using (8) that $[I + \mathcal{N}^{(1)}, \tilde{\mathcal{Z}}]$ has a nonidentity element from $I + e_{13}(\mathcal{M}^{m-1})$. Since this element is not contained in $I + \mathcal{N}^{(k_0+3)}$, the $(k_0 + 3)$ rd term of the lower central series of $N(n, R)$ strictly contains $\Gamma_{k_0+3}(D)(I + \mathcal{N}^{(k_0+3)})$. We see by induction that the $(n + m - 1)$ st term of the lower central series of $N(n, R)$ contains a non-identity element from $I + e_{1n}(\mathcal{M}^{m-1})$. We have shown thus that if $k_0 = m - 1$ then the nilpotency class of $N(n, R)$ is $n + m - 1$.

If $k_0 < m - 1$, the nilpotency class of $N(n, R)$ still depends on the properties of R . However, for all $l \geq 2$ the l th term of the lower central series of $N(n, R)$ contains $\Gamma_l(D)(I + \mathcal{N}^{(l)})$ and is contained in $\Gamma_l(D)(I + \mathcal{N}^{(l-1)})$ so that the nilpotency class is either $n + m - 2$ or $n + m - 1$. Assuming a rather natural condition on R we can guarantee that it will be $n + m - 1$. The condition is

$$v \in \mathcal{M}, v\mathcal{M} \subseteq \mathcal{M}^s \Rightarrow v \in \mathcal{M}^{s-1}, \text{ for all } s \geq 2. \quad (11)$$

Assuming (11) and taking the above considered element $v = 1 - [h, 1 + y]$ we see that there exists $z \in \mathcal{M}$ with $vz \notin \mathcal{M}^{k_0+2}$. Then by (7) one has $[\text{diag}(1, 1 - z, 1, \dots, 1), I + e_{12}(v)] = I + e_{12}(vz)$ and the $k_0 + 3$ rd term of the lower central series of $N(n, R)$ strictly contains $\Gamma_{k_0+3}(D)(I + \mathcal{N}^{(k_0+3)})$. Using (11) in this way, we produce by induction an element $I + e_{12}(w)$ of the $(m+1)$ st term of the lower central series of $N(n, R)$ which is not contained in $I + \mathcal{N}^{(m+1)}$. Then applying (8) several times, as in case $k_0 = m - 1$, we obtain in the $n + m - 1$ st term of the lower central series of $N(n, R)$ a nonidentity element from $I + e_{1n}(\mathcal{M}^{m-1})$. This means that the nilpotency class is in fact $n + m - 1$.

We have obtained the following result.

Proposition 3.5. *Let R be as in Theorem 3.2 and $n \geq 2$.*

(i) *If $[\Gamma_{k-1}(\mathcal{U}(R)), 1 + \mathcal{M}] \subseteq 1 + \mathcal{M}^k$ for all k then the lower central series of $N(n, R)$ is*

$$G \supset \Gamma_2(D) + \mathcal{N}^{(2)} \supset \Gamma_3(D) + \mathcal{N}^{(3)} \supset \dots \supset \Gamma_m(D) + \mathcal{N}^{(m)} \supset I + \mathcal{N}^{(m+1)} \supset \dots \supset I + \mathcal{N}^{(n+m-2)} \supset I,$$

and the nilpotency class of $N(n, R)$ is $n + m - 2$. This happens if either $\Gamma_2(\mathcal{U}(R)) \subseteq 1 + \mathcal{M}^2$ or if $[\mathcal{M}, \mathcal{M}]_{\text{Lie}} \subseteq \mathcal{M}^3$.

(ii) *If k_0 is the least integer with $[\Gamma_{k_0}(\mathcal{U}(R)), 1 + \mathcal{M}] \not\subseteq 1 + \mathcal{M}^{k_0+1}$ then $2 \leq k_0 \leq m - 1$, the first $k_0 + 1$ terms of the lower central series of $N(n, R)$ are as in item (i) and for each $k_0 + 2 \leq l \leq n + m - 1$ the l th term $\Gamma_l(N(n, R))$ of the lower central series of $N(n, R)$ satisfies*

$$\Gamma_l(D)(I + \mathcal{N}^{(l)}) \subseteq \Gamma_l(N(n, R)) \subseteq \Gamma_l(D)(I + \mathcal{N}^{(l-1)}).$$

In particular, the nilpotency class of $N(n, R)$ is either $n + m - 2$ or $n + m - 1$. The latter occurs whenever R satisfies (11) or $k_0 = m - 1$.

The next corollary immediately follows from item (i) of Proposition 3.5.

Corollary 3.6. *If R is a commutative local artinian ring and $n \geq 2$ then $N(n, R)$ is nilpotent of class $n + m - 2$.*

On the other hand, since $2 \leq k_0 \leq m - 1$, item (ii) of Proposition 3.5 does not occur if $m = 2$. Thus we immediately have:

Corollary 3.7. *Let R be as in Theorem 3.2 and $n \geq 2$. If $\mathcal{M} \neq 0$ and $\mathcal{M}^2 = 0$ then $N(n, R)$ is nilpotent of class $n + m - 2 = n$.*

If R is a K -algebra satisfying the conditions of Theorem 3.2 and $R = K \oplus \mathcal{M}$ as vector spaces, then $\Gamma_2(\mathcal{U}(R)) = \Gamma_2(1 + \mathcal{M}) \subseteq 1 + \mathcal{M}^2$, so that R verifies item (i) of Proposition 3.5. An example of such a ring is given by the group algebra KG of a finite p -group G over a field K of characteristic p . Indeed, it was observed already in the Introduction that KG is local with $\mathcal{M} = \Delta(G)$ which satisfies (2). Moreover, evidently $KG = K \oplus \Delta(G)$. Hence we obtain the following:

Corollary 3.8. *Let R be as in Theorem 3.2 and $n \geq 2$. If R is a K -algebra and $R = K \oplus \mathcal{M}$ as vector spaces, then $N(n, R)$ is nilpotent of class $n + m - 2$. In particular, this occurs if $R = KG$, the group algebra of a finite p -group G over a field K of characteristic p .*

Acknowledgments

The first author is grateful to Prof. Arnaldo Mandel for useful conversations.

References

- [1] M. Shirvani, B. A. F. Wehrfritz, *Skew Linear Groups*, Cambridge Univ. Press, Cambridge, 1986.
- [2] S. K. Sehgal, *Topics in Group Rings*, Marcel Dekker, New York, 1978.
- [3] D. A. Suprunenko, *Matrix groups*, Translations of mathematical monographs, 45, AMS, Providence, R. I., 1976.
- [4] B. A. F. Wehrfritz, *Infinite Linear Groups*, Springer, New York, 1973.

TRABALHOS DO DEPARTAMENTO DE MATEMÁTICA

TÍTULOS PUBLICADOS

- 2003-01 COELHO, F.U. and LANZILOTTA, M.A. Weakly shod algebras. 28p.
- 2003-02 GREEN, E.L., MARCOS, E. and ZHANG, P. Koszul modules and modules with linear presentations. 26p.
- 2003-03 KOSZMIDER, P. Banach spaces of continuous functions with few operators. 31p.
- 2003-04 GORODSKI, C. Polar actions on compact symmetric spaces which admit a totally geodesic principal orbit. 11p.
- 2003-05 PEREIRA, A.L. Generic Hyperbolicity for the equilibria of the one-dimensional parabolic equation $u_t = (a(x)u_x)_x + f(u)$. 19p.
- 2003-06 COELHO, F.U. and PLATZECK, M.I. On the representation dimension of some classes of algebras. 16p.
- 2003-07 CHERNOUSOVA, Zh. T., DOKUCHAEV, M.A., Khibina, M.A., Kirichenko, V.V., MIROSHNICHENKO, S.G., Zhuravlev, V.N. Tiled orders over discrete valuation rings, finite Markov chains and partially ordered sets. II. 43p.
- 2003-08 ARAGONA, J., FERNANDEZ, R. and JURIAANS, S.O. A Discontinuous Colombeau Differential Calculus. 20p.
- 2003-09 OLIVEIRA, L.A.F., PEREIRA, A.L. and PEREIRA, M.C. Continuity of attractors for a reaction-diffusion problem with respect to variation of the domain. 22p.
- 2003-10 CHALOM, G., MARCOS, E., OLIVEIRA, P. Gröbner basis in algebras extended by loops. 10p.
- 2003-11 ASSEM, I., CASTONGUAY, D., MARCOS, E.N. and TREPODE, S. Quotients of incidence algebras and the Euler characteristic. 19p.
- 2003-12 KOSZMIDER, P. A space $C(K)$ where all non-trivial complemented subspaces have big densities. 17p.
- 2003-13 ZAVARNITSINE, A.V. Weights of the irreducible $SL_3(q)$ -modules in defining characteristic. 12p.
- 2003-14 MARCOS, E. N. and MARTÍNEZ-VILLA, R. The odd part of a N-Koszul algebra. 7p.
- 2003-15 FERREIRA, V.O., MURAKAMI, L.S.I. and PAQUES, A. A Hopf-Galois correspondence for free algebras. 12p.
- 2003-16 KOSZMIDER, P. On decompositions of Banach spaces of continuous functions on Mrówka's spaces. 10p.

- 2003-17 GREEN, E.L., MARCOS, E.N., MARTÍNEZ-VILLA, R. and ZHANG, P. D-Koszul Algebras. 26p.
- 2003-18 TAPIA, G. A. and BARBANTI, L. Um esquema de aproximação para equações de evolução. 20p.
- 2003-19 ASPERTI, A. C. and VILHENA, J. A. Björling problem for maximal surfaces in the Lorentz-Minkowski 4-dimensional space. 18p.
- 2003-20 GOODAIRE, E. G. and MILIES, C. P. Symmetric units in alternative loop rings. 9p.
- 2003-21 ALVARES, E. R. and COELHO, F. U. On translation quivers with weak sections. 10p.
- 2003-22 ALVARES, E.R. and COELHO, F.U. Embeddings of non-semiregular translation quivers in quivers of type ZA . 23p.
- 2003-23 BALCERZAK, M., BARTOSZEWICZ, A. and KOSZMIDER, P. On Marczewski-Burstin representable algebras. 6p.
- 2003-24 DOKUCHAEV, M. and ZHUKAVETS, N. On finite degree partial representations of groups. 24p.
- 2003-25 GORODSKI, C. and PODESTÀ, F. Homogeneity rank of real representations of compact Lie groups. 13p.
- 2003-26 CASTONGUAY, D. Derived-tame blowing-up of tree algebras. 20p.
- 2003-27 GOODAIRE, E.G. and MILIES, C. P. When is a unit loop f -unitary? 18p.
- 2003-28 MARCOS, E.N., MARTÍNEZ-VILLA, R. and MARTINS, M.I.R. Hochschild Cohomology of skew group rings and invariants. 16p.
- 2003-29 CIBILIS, C. and MARCOS, E.N. Skew category, Galois covering and smash product of a category over a ring. 21p.
- 2004-01 ASSEM, I., COELHO, F.U., LANZILOTTA, M., SMITH, D. and TREPODE, SONIA Algebras determined by their left and right parts. 34p.
- 2004-02 FUTORNY, V., MOLEV, A. and OVSIENKO, S. Harish-Chandra Modules for Yangians. 29p.
- 2004-03 COX, B. L. and FUTORNY, V. Intermediate Wakimoto modules for affine $sl(n+1, C)$. 35p.
- 2004-04 GRISHKOV, A. N. and ZAVARNITSINE, A. V. Maximal subloops of simple Moufang loops. 43p.
- 2004-05 GREEN, E.L. and MARCOS, E. δ -Koszul Algebras. 15p.
- 2004-06 GORODSKI, C. Taut Representations of compact simple lie groups. 16p.
- 2004-07 ASPERTI, A.C. and VALÉRIO, B.C. Ruled helicoidal surfaces in a 3-dimensional space form. 9p.
- 2004-08 ASSEM, I., CASTONGUAY, D., MARCOS, E.N. and TREPODE, S. Strongly simply connected schurian algebras and multiplicative bases. 26p.

- 2004-09 RODRIGUES, A.A.M., MIRANDA FILHO, R.C. and SOUZA, E.G. Definability and Invariance in First Order Structures. 15p.
- 2004-10 GIAMBRUNO, A. and MILIES, C. P. Free groups and involutions in the unit group of a group algebra. 5p.
- 2004-11 RAO, S. ESWARA and FUTORNY, V. Classification of integrable modules for affine lie superalgebras. 20p.
- 2004-12 GRISHKOV, A. N. and ZAVARNITSINE, A. V. Groups with triality. 22p.
- 2004-13 BUSTAMANTE, J. C. and CASTONGUAY, D. Fundamental groups and presentations of algebras. 10p.
- 2004-14 FERNANDES, S.M. and MARCOS, E.N. On the Fundamental Derivation of a Finite Dimensional Algebra. 15p.
- 2004-15 CARRIÓN, H.; GALINDO, P. and LOURENÇO, M.L. Banach spaces whose bounded sets are bounding in the bidual. 10p.
- 2004-16 KOSZMIDER, P. Projections in weakly compactly generated Banach spaces and Chang's conjecture. 15p.
- 2004-17 KOSZMIDER, P. On a problem of Rolewicz about Banach spaces that admit support sets. 19p.
- 2004-18 GOODAIRE, E.G., MILIES, C.P. and PARMENTER, M.M. Central units in alternative loop rings. 8p.
- 2004-19 GIAMBRUNO, A. and MILIES, C. P. Free groups and involutions in the unit group of a group algebra. 5p.
- 2004-20 CARRIÓN, H. Entire Functions on l_1 . 16p.
- 2004-21 SHESTAKOV, I. and ZHUKAVETS, N. Universal multiplicative envelope of free Malcev superalgebra on one odd generator. 28p.
- 2004-22 ASSEM, I., BUSTAMANTE, J.C., CASTONGUAY, D. and NOVOA, C. A note on the fundamental group of a one-point extension. 6p.
- 2004-23 BAHTURIN, Y. A., SHESTAKOV, I. P. and ZAICEV, M. V. Gradings on Simple Jordan and Lie Algebras. 25p.
- 2004-24 DOKUCHAEV, M., KIRICHENKO, V. and MILIES, C. P. Engel subgroups of triangular matrices over local rings. 16p.

Nota: Os títulos publicados nos Relatórios Técnicos dos anos de 1980 a 2002 estão à disposição no Departamento de Matemática do IME-USP.
Cidade Universitária "Armando de Salles Oliveira"
Rua do Matão, 1010 - Cidade Universitária
Caixa Postal 66281 - CEP 05315-970