

A novel semantic theory of the assembly rules of interaction networks

Marco A. R. Mello^{§1}, Sharlene E. Santana², Pierre-Michel Forget³, Cristina A. Kita^{1,4}, Nastaran Lotfi⁵, Isabel C. S. Machado⁶, Renata L. Muylaert⁷, Francisco A. Rodrigues⁸, Tanja M. Straka⁹ & Carsten F. Dormann¹⁰

1. Department of Ecology, Institute of Biosciences, Universidade de São Paulo. Rua do Matão, 321/Travessa 14, 05508-090 São Paulo, Brazil. ORCID: <https://orcid.org/0000-0002-9098-9427>. E-mail: marmello@usp.br

2. Department of Biology and Burke Museum of Natural History and Culture, University of Washington, 4303 Memorial Way Northeast, 98195 Seattle, USA. ORCID: <https://orcid.org/0000-0001-6463-3569>. E-mail: ssantana@uw.edu.

3. MECADEV-ECOTROP, Département Adaptations du Vivant, Muséum National d'Histoire Naturelle, Bâtiment 50, 45 rue Buffon, 75005 Paris, France. <https://orcid.org/0000-0002-9252-974X>. E-mail: pierre-michel.forget@mnhn.fr

4. Programa de Pós-Graduação em Ecologia, Instituto de Biociências, Universidade de São Paulo. Rua do Matão, 321/Travessa 14, 05508-090 São Paulo, Brazil. ORCID: <https://orcid.org/0000-0002-4079-2677>. E-mail: c.akemikita@gmail.com

5. Department of Neurology, University of Texas Health Science Center at Houston, Houston, TX 77030. ORCID: <https://orcid.org/0000-0001-7258-6117>. E-mail: Nastaran.lotfi@uth.tmc.edu

6. Programa de Pós-graduação em Biologia Vegetal, Universidade Federal de Pernambuco, Av. Prof. Moraes Rego s/n, Cidade Universitária, 50670-901, Recife-PE, Brazil. ORCID: <https://orcid.org/0000-0001-5015-2393>. E-mail: isabel.machado@ufpe.br

7. Sydney School of Veterinary Science, Faculty of Science, The University of Sydney, Sydney, 2050 New South Wales, Australia. ORCID: <https://orcid.org/0000-0002-6466-6210>. E-mail: renata.muylaert@sydney.edu.au

8. Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo, São Carlos, São Paulo, Brazil. ORCID: <https://orcid.org/0000-0002-0145-5571>. E-mail: francisco@icmc.usp.br

9. Institute for Biology, Freie Universität Berlin, Königin-Luise-Str. 1-3, 14195 Berlin, Germany. ORCID: <https://orcid.org/0000-0003-4118-4056>. E-mail: tanja.straka@fu-berlin.de

10. Department of Biometry and Environmental System Analysis, Albert-Ludwigs-Universität Freiburg. Tennenbacher Straße 4, 79106 Freiburg, Germany. ORCID: <https://orcid.org/0000-0002-9835-1794>. E-mail: carsten.dormann@biom.uni-freiburg.de

§ Corresponding author.

* Authors from the 3rd to the penultimate position contributed equally and so are listed in alphabetical order of last name.

45

46 **Abstract**

47 In the web of life, every interaction between species tells a story of cooperation,
48 conflict, or chance. For centuries, ecologists have charted these stories to better
49 understand phenomena such as pollination and disease. We have been using the lens of
50 network science to distill them into topological patterns such as nestedness or
51 modularity. Yet, like in the old buddhist parable of the blind monks and the elephant,
52 these separate views have often missed the bigger picture. In this Perspective, we
53 review the history of topological studies of interaction networks, challenge the “black-
54 and-white” paradigm, and point out a way forward: the Integrative Theory of Interaction
55 Networks (ITIN). This theory unveils the logic of how interaction networks assemble
56 through sequential, rule-bound processes shaped mainly by resource dissimilarity and
57 interaction type. We show that compound topologies are likely the norm in function-
58 oriented, taxonomically-inclusive, well-sampled networks. ITIN is deductive,
59 predictive, and testable, mapping from first principles to real-world systems, thus
60 helping us grasp “the whole elephant”. As ecologists confront the polycrisis, from
61 irreparable biodiversity loss to global health threats, ITIN offers a coherent framework
62 to understand, predict, and restore the threads of the web of life.

63

64 **Introduction**

65 Ecological interactions between individual organisms of different species (i.e., species
66 interactions) have drawn the attention of scientists, philosophers, and stakeholders for
67 millennia ¹. This broad family of phenomena ranges from mutualistic (both sides gain
68 fitness) to antagonistic (one or both sides lose fitness), and result in crop pollination,
69 plant protection, seed dispersal, parasitism, and disease, among many other outcomes ².

70 Species interactions have been studied from multiple perspectives and using numerous
71 approaches ³. One approach that has become mainstream is network science, an
72 interdisciplinary field that provides not only tools for data analysis but also concepts
73 that may be integrated across scientific disciplines to obtain new insights ⁴. The study of
74 interaction networks as we know it began in the late 19th century (or even earlier),
75 through a classical effort to understand complex relationships in lake food webs ⁵. Only
76 more recently, between the late 1980s and early 2000s, the focus slowly shifted from
77 food webs to mutualistic networks ⁶. The intriguing structural patterns of interaction
78 networks, also known as topologies, has become a fashionable research topic in itself,
79 irrespective of the effect they may, or may not, have on the community and ecosystem
80 dynamics ⁷. This focus, although being an important first descriptive effort, seems to
81 have led to worrisome epistemological biases, such as confounding the ecological
82 processes with the network patterns they generate ⁸.

83 Biases in topological studies can be compared to the ancient Buddhist parable of the
84 blind monks and the elephant ([Figure 1](#)). Once upon a time, a group of blind monks
85 heard that a huge, strange animal had arrived near the temple, and, curious to know
86 what it was like, decided to examine it by touch. Each monk felt a different part of the
87 elephant and drew a different conclusion. To one monk, the trunk felt like a snake. To
88 another, the ear felt like a fan. To others, a leg felt like a tree trunk, the belly like a wall,
89 the tail like a rope, and a tusk like a spear. Each monk described the elephant based on
90 their separate experience, illustrating how partial perspectives can lead to incomplete or
91 conflicting understandings of a greater truth.




Figure 1: The blind monks and the elephant. As in the old Buddhist parable, we have been analyzing only network parts but drawing conclusions for the whole. Currently, mutualistic networks are generally assumed to be nested and antagonistic networks, modular. Gradient networks have been seldom considered, usually being associated with parasitism. Much fewer studies discussed compound topologies, although these have been drawing more interest lately. The image was modified from the original *ukiyo-e* print illustration by Itcho Hanabusa (Wikimedia Commons, public domain).

Not so different from those monks, most ecological studies have focused on interactions in only one context, ignoring that sites with different abundances will yield different networks and observed interaction preferences⁸. Moreover, by focusing on isolated interaction types, while trying to draw conclusions for all types, those studies ignore how interactions create complex systems with emergent properties⁹. Recent advances, however, show that networks require a more integrative approach to be fully understood¹⁰. Integration is especially needed for understanding not only the topological patterns observed in nature, but also how they emerge as a result of ecological and evolutionary processes. In other words, the assembly rules of interaction networks.

In this Perspective, we aim to demonstrate how different views about network topologies and their assembly rules could be integrated, in order to help refocus our research efforts. First, we synthesize the history of network topology studies. Second, we show how the Integrative Hypothesis of Specialization (IHS) provides a comprehensive approach to see the whole elephant, in addition to shedding light on the processes behind the patterns. Third, we point to new directions, as after a series of developments, empirical tests, and additional synthesis, IHS is evolving into a semantic theory, the Integrative Theory of Interaction Networks (ITIN), which we first introduce.

Archetypal topologies

A seminal study ahead of its time in the mid-2000s proposed a taxonomy of network topologies¹¹. From its perspective, there would be four archetypal topologies ([Figure 2](#)) that capture the essence of most real-world systems (but see an alternative taxonomy¹²).



125 Figure 2: The four archetypal topologies. The conceptual matrices presented here were
126 modified from those originally proposed in a seminal paper ¹¹. (A) A nested topology
127 resembles a Russian doll: nodes with fewer links are connected to a subset of the links
128 captured by the most connected nodes, sequentially, for the rows and columns separately.
129 (B) The modular topology looks like the filter bubbles observed in social media, as the
130 network is divided into cohesive subgroups of nodes (i.e., modules), more densely
131 connected to one another than to other subgroups. (C) In a gradient topology, as in a chess
132 board, there is little to no overlap in the links made by different nodes, and there are also
133 much fewer links than in a modular network, leading to the formation of truly
134 disconnected components. (D) A compound topology combines a modular structure in

the network, with a nested structure within its modules, resulting in “Russian dolls trapped inside filter bubbles”, and tends to be larger than a modular network.

The first archetype, nested networks (quantified as nestedness), emerges when the links made by species with fewer interactions represent a subset of those made by species with more interactions, sequentially. The second archetype, modular networks (quantified as modularity), represents systems that consist of cohesive subgroups of species (i.e., modules), which are densely interconnected but have fewer interactions with other subgroups. Gradient networks are the third and least studied archetype, usually observed when sampling in steep environmental gradients, leading to few links and almost no overlap in the interactions made by different species, therefore forming several small components (i.e., totally disconnected subgroups). Finally, the fourth archetype, compound networks, combines modularity in the network and nestedness within its modules.

Lately, different research groups have independently proposed topologies similar to the compound archetype, with slight variations in conceptualization and operationalization^{13–15}. Their structure would in general resemble a polycentric heterarchy¹⁶, almost equivalent to a dissortative network¹⁷. That is, in such a network there are differences in influence between nodes, with one or a few nodes dominating each module and playing the role of module hubs.

Despite the multiple options provided by the four archetypal topologies, another seminal study published a little earlier proposed the universality of nestedness in mutualistic networks¹⁸. By comparing food webs, pollination, and seed dispersal systems, the study invoked co-evolutionary arguments to conclude that mutualisms would form highly cohesive, nested networks in which a core of generalist species would drive the dynamics and maybe the evolution of the entire local community. In contrast, antagonisms would be characteristically modular, as traditionally assumed¹⁹. The almost overlooked gradient topology, which was not considered then and still remains poorly studied²⁰, would be found only in the most specialized systems with high interaction intimacy²¹ or along steep gradients with substantial turnover in species²².

Since nestedness was advanced, it became the dominant view in the study of interaction networks, especially mutualistic ones. Most studies have been either trying to validate a nested topology for the respective system of interest or aiming to link nestedness to the stability of the entire community, sometimes resorting to circular reasoning²³, despite how controversial this view has always been²⁴. Nestedness has been pointed out as an artifact²⁵ or a simple byproduct of abundance²⁶, and not a pattern derived from a clearly defined process²⁷. In addition, nestedness is also frequently found in antagonisms²⁸. A striking critique emerged from a study that revealed conceptual flaws in early nestedness indices and algorithms²⁹. The high nestedness scores reported for many systems were artificially inflated, and, as a more accurate index was proposed, it became clear that nestedness was indeed highly variable and lower on average than previously measured in many empirical networks³⁰, and even lower than expected by chance from differences in abundance³¹.

What should have been the final blow to the universality of nestedness came from a naturalistic perspective. Different studies started to point out that the mutualistic interactions sampled at a local scale and low completeness levels, included in popular databases, only represent fragments, not entire networks^{32–34}. This deserves attention considering that ecological functions, such as pollination, and its derived ecosystem services, such as crop pollination, are not delivered by single taxa, but by guilds composed of several taxa acting together at a local scale.

Therefore, variations in the definition of nestedness, conceptual limitations in its proxies, and a narrow taxonomic focus have raised questions regarding its universality. Nonetheless, nestedness remains a central concept in the study of mutualistic networks and continues to provide a benchmark for assessing the structure of mutualistic interactions across pristine, degraded, and restored environments³⁵. More recently, however, studies started adopting conceptually refined approaches that offer fresh perspectives on network topology³⁶.

A way forward

Given the challenges presented above, a new theoretical perspective was needed to make sense of patterns and processes. The Integrative Hypothesis of Specialization (IHS) was proposed as a way forward, originally presented as a graphical model³⁷, and

subsequently mathematically formalized as an algorithmic model ³⁸. IHS was built as a minimalist mechanism that integrates sequential processes, which themselves represent the mechanistic assembly rules of interaction networks.

In short, IHS proposes that interaction constraints of geographic, morphological, physiological, or behavioral nature would lead to the formation of layers and modules in interaction networks. Within these modules, the specialisation-niche breadth trade-off predominates once the constraints are overcome. Consequently, we can deduce from IHS that the assembly rules of interaction networks include a threshold of influence of different ecological and evolutionary processes at a given hierarchical level (entire network, layers, modules, motifs, and dyads), which determines which archetypal topology predominates at that level.

To evaluate the four archetypal topologies with as much generality as possible, IHS assumes that all interaction types can be viewed as pure consumer models, similarly to other network models ^{15,39,40}, but aiming at higher generality and parsimony. We know that most interactions are more complex than this, but this simplification of reality has proven to be surprisingly efficient in deducing topology. IHS is based on three first principles that apply to evolving consumer-resource relationships in general ⁴¹: (i) each type of resource has a set of characteristics that affect its use, and the types of resources may be more or less similar to each other in these characteristics; (ii) a consumer's mutation that increases its ability to use a given resource tends to improve its efficiency on similar resources, but worsen its on different resources, generating clusters of usable and non-usable resources; and (iii) a consumer's ability to use each resource in a given cluster at a given time is the result of its previous adaptations and maladaptations (i.e. mutations are small). Based on this, the free parameters of IHS are: (a) consumer species richness, (b) resource species richness, (c) method used to generate the initial innate performance matrix (innate method), (d) maximum dissimilarity between two resource species (maximum dissimilarity), and (e) number of resource species clusters (number of clusters).

The structure of IHS leads to the logical conclusion that the topology of a given interaction network depends mainly on the dissimilarity between the resources contained within it, from the point of view of consumer adaptations ⁴¹. In turn, this conclusion leads to different secondary deductions, for example: the more dissimilar the resources, say in traits, the more likely a modular topology emerges. Assuming that

most resources in nature tend to be organized in clusters ⁴², very large networks with high heterogeneity would tend to have a compound topology: modular across clusters, but nested within. In turn, small and homogeneous networks, which would actually represent clusters, would tend to be nested. At the extreme of this continuum, small networks with very high resource dissimilarity, probably resulting from very high interaction intimacy or steep environmental gradients, would be organized as a gradient. These deductions are valid mainly for interactions studied at the community level, although further expansions of IHS to interactions between individuals have already been made ⁴³. Although many studies propose that coevolution is an important driver of interaction networks ⁴⁴, IHS assumes that their structure can emerge without *mutual* adaptation between consumers and resources. This does not mean that coevolution is not important for understanding species interactions in general. Rather, within the scope of IHS, coevolution is not *necessary* to explain the assembly rules that generate the four archetypal topologies.

A novel theory

Starting from the realization that IHS already has all the characteristics of a semantic theory ^{45–47}, all that remains is to formalize it as: the Integrative Theory of Interaction Networks (or ITIN, for short).

To build a novel semantic theory of the assembly rules of interaction networks, it is essential to understand the concepts and assumptions that guide this approach. Models of this type are constructed theoretically and do not directly depend on real-world data, which allows for a more flexible and adaptable approach to the conceptualization of the expected patterns in natural phenomena. The development of these models involves the application of a mathematical or logical formalism to deduce the models from the axioms, allowing a meaningful comparison with the observed phenomena. This crucial step has already been taken in building the algorithmic model used to make the proof of concept of IHS ⁴¹.

Since its proof of concept, IHS has undergone a series of empirical tests of several predictions deduced from its central axiom. IHS proved to be capable of explaining the multiplicity of network topologies in real-world networks of pathogen infection ³⁷, parasitism ⁴⁸, seed dispersal ⁴⁹, pollination ⁵⁰, resin collection ⁵¹, burrow sharing ⁵²,

predation ⁵³, resource partitioning ⁵⁴, multilayered interactions ⁵⁵, and food crop management ⁵⁶, among many other examples, widely extending its original domain of validity.

Later studies also demonstrated that compound and modular topologies are actually much more common than previously thought ¹⁴, as expected from theory. Most interaction networks, in the end, resemble filter bubbles filled with Russian dolls, and a good example of this kind of compound topology was observed in a system formed by small mammals and their ectoparasitic fleas ⁴⁸. So, if modular and compound topologies are the most common structures observed in well-sampled, taxonomically comprehensive interaction networks studied at different spatial scales, from local to continental, what about nested and gradient topologies? Explaining the logical relationship between archetypal topologies is the next step in the development of our novel theory (Figure 3).




Figure 3: A roadmap to the archetypal topologies of interaction networks according to the Integrative Theory of Interaction Networks (ITIN). (1) As an example (see section Three archetypal stories), two species arrive at an open area after a disturbance, such as a forest fire, and begin to interact. (2) Ecological succession follows, bringing more

species and interactions, forming a shifting topology that may resemble a random network. (3) A critical point is reached, a phase transition, after which the network may or may not blossom, giving rise to different stable topologies. (4) When resource dissimilarity follows a tree-like, hierarchical structure, the network becomes fractal, resembling a Cantor matrix. (5) If network growth is halted by external factors, such as hunting, its topology remains undefined. (6) When resource dissimilarity is discontinuous and the network continues to grow, a compound topology emerges, with modularity across scales and nestedness within modules. (7) Nested topologies, therefore, represent the internal structure of modules, not the whole mature network. (8) In mutualistic systems (White Queen Hypothesis), resources strengthen associations with better consumers, leading to modular or compound topologies. (9) In antagonistic systems (Red Queen Hypothesis), escape dynamics promote specialization, resulting in gradient topologies.

Conceptual examples might help understand our rationale ([Box 1](#)).

Box 1: Three archetypal stories

The Integrative Theory of Interaction Networks (ITIN) is a description of how real-world networks develop over time. The following “stories” exemplify possible paths towards an observed network topology.

The niche sequence along a steep gradient story

Imagine several plant species from the same family, growing along a mountain’s altitudinal gradient. Evolved from the same ancestor, they still share most of their niche characteristics, but differ, say, in frost tolerance. Each species has thus a position along the growth-frost tolerance gradient. A set of herbivore species adapted to feeding on this family will feed on all species. The network therefore will be fully connected. Over time, slow-growing, high-altitude species will invest more in chemical defense, as each herbivore attack leads to tissue loss, which they cannot readily compensate for by fast growth given that they experience a short growing season. In response to the increased chemical anti-herbivore defense, some herbivores will start avoiding these species, while others will adapt and even specialize on them. The resulting topology will depend

on the degree of separation: if almost every herbivore becomes a specialist, we end up with a modular network. If most herbivores exhibit some lower preference for the species a bit higher up and lower down the gradient, we end up with a gradient topology.

The abundance story

Most species are rare⁵⁷. The causes are multifaceted, from neutral ecological drift to competitive hierarchies⁵⁸ or statistical inevitability of random processes⁵⁹. Either way, the evolutionary pressures to evade an abundant predator are much stronger than those to evade a rare one. Antelopes run (almost) fast enough to evade a cheetah, not only fast enough to outrun the much slower jackal. As a consequence, all antelopes face predation by common predators, and common antelopes face all predators, simply as a matter of probability. We can compute this as the matrix product of the abundance vectors of antelopes and predators, normalised to sum to 1. The resulting network is nested. Interestingly, two common predators may compete for common prey, leading to specialisation *for the most common prey* only. This will be visible as anti-nestedness for the most common prey, but still a strong nestedness signal overall. This makes “nestedness” an awkward concept: the observed competition signal leads to a *reduced* nestedness, while the overall topology is still highly nested.

The introduced species story

The effect of adding a new species to a network is likely to be extremely case-specific. Imagine a pollination network to which a non-native plant species is added⁶⁰. If that species is similar in traits to species already present in the network, we can expect it to “slot in”, maintaining whatever topology existed before. However, if the plant’s attractive traits vastly exceed those of the existing species (e.g., flowers that are more nectar rich), it may actually withdraw flower visitors from other plants. The network topology is dependent on what happens next. If the pollinators keep their visitation pattern, but prefer the new plant species, then the network becomes a bit more nested. If, however, the new plant leads to a strong re-alignment of trait preferences, it may lead to network fragmentation, and a more modular structure. The point here is that the topology, in concert with the relevant traits, will in the end help us to understand what happened.

341

342 ITIN's mechanism sets the thresholds that define when one topology ends and another
343 begins. In the simplest and least realistic scenario, considering only a tree-like
344 distribution of resource dissimilarity, such as in a dendrogram, with progressive
345 dissimilarity between hierarchical levels, the fundamental topology of interaction
346 networks should be fractal, like in a Cantor matrix ⁶¹. Nevertheless, this should not be
347 expected in the real world, as virtually all trait distributions are discontinuous ⁴².

348 Considering even more realistic scenarios, sometimes network assembly is halted by a
349 process, such as overhunting, that prevents the addition of new nodes or the
350 establishment of links between them. In this case, the network might not reach an
351 archetypal topology, remaining frozen in an undefined transitional topology.

352 If the network continues to grow, it might develop to a point where it becomes
353 compound. First, because not all consumers are empirically able to use all resources ⁶²,
354 this leads to the formation of layers and modules. Second, because after constraints are
355 met, there is more freedom of association within a module. Consequently, neutral
356 processes leading to differences in abundance ⁶³ or universal network assembly
357 processes such as preferential attachment ⁶⁴ should lead to a nested structure within the
358 modules. Nested topologies, therefore, would represent pieces and not entire networks.

359 Two processes could further shape interaction networks into the remaining archetypal
360 topologies (Figure 4). On one hand, if the interaction type is antagonistic, according to
361 the Red Queen Hypothesis ⁶⁵ the resources would tend to escape the consumers,
362 reducing niche overlap and even the total number of links observed, maybe even
363 resulting in the extinction of some consumers, leading to a gradient topology. On the
364 other hand, if the interaction type that defines the links is mutualistic, then the resources
365 would tend to strengthen their association with the most advantageous consumers, while
366 escaping the worst consumers, which could be called a "White Queen Hypothesis". This
367 would lead to a truly modular topology, with smaller modules that are not internally
368 nested, because freedom of association is still limited after the main constraints are
369 resolved.

370




Figure 4: A novel semantic theory of the assembly rules of interaction networks. As a development of the Integrative Hypothesis of Specialization (IHS), the Integrative Theory of Interaction Networks (ITIN) has been expanded by the addition of two first principles and one free parameter. This enhances its ability to explain which processes determine where one archetypal topology ends and another begins. See also Figure 3.

Therefore, in addition to the first principles considered in the original IHS, ITIN also includes the following assumptions: (iv) if resource dissimilarities are distributed discontinuously, there are clusters of resources and consumers linked within the network; and (v) if the interaction is mutualistic, associations between consumers and resources tend to be strengthened, but if the interaction is antagonistic, resources tend to escape the consumers. Consequently, ITIN includes an additional free parameter: (f) interaction type.

From description to explanation

As discussed in the previous sections, the study of interaction networks has been guided by a multitude of metrics. Connectance, modularity, nestedness, specialization: each captures a piece of the puzzle. Yet, as in the parable of the blind monks and the elephant, these fragmented perspectives did not converge into the bigger picture. The result has been a proliferation of metrics that quantify patterns but do not explain


mechanistically how they arise or what they mean for ecological functions and ecosystem services ⁸.

Without a mechanism that ties pattern to process, our ability to predict and intervene remains limited. Conservation biologists struggle to anticipate how seed dispersal networks will respond to human disturbances ⁶⁶. Agroecologists seek to design sustainable crops but are left with taxon-biased network fragments that overlook the functional guilds actually delivering vital ecosystem services ⁶⁷. Epidemiologists face difficulties estimating the risk of pathogen spillover without understanding the assembly rules that govern host-pathogen networks ⁶⁸. The black-and-white paradigm of network topology has constrained both scientific inference and practical action, leaving us only with numbers yet still in search for meaning.

ITIN provides this much needed meaning. By rooting network topologies in first principles and free parameters mappable into operational variables, it shifts the focus from static descriptors to dynamic assembly rules. It explains not only what structures we observe, but why they emerge and where the thresholds lie between one topology and another.

Practical applications

As an empirically tested semantic theory, ITIN might be used to help solve problems. Not only theoretical puzzles, but also practical challenges within the frameworks provided by the UN's Sustainable Development Goals ⁶⁹ and the One Health Initiative ⁷⁰ ([Figure 5](#)).



413

414 Figure 5: Conceptual framework illustrating how network studies can help translate
 415 knowledge about ecosystem services and diseases into solutions to real-world problems.
 416 On the left, pollinators contribute to ecosystem services such as crop pollination,
 417 supporting food production. On the right, animal hosts and pathogens represent the
 418 ecological interactions underlying disease emergence. By using network science,
 419 including the new framework provided by ITIN, both research areas can be aligned with
 420 the global agendas of Sustainable Development Goals and One Health.

421

422 The central driver of network topologies according to ITIN is resource dissimilarity. In
 423 order to develop practical solutions, it is of paramount importance to sample entire local
 424 networks responsible for a given ecosystem service of interest. A function-oriented
 425 modelling of interaction networks and the ecosystem services they deliver requires
 426 collaboration beyond taxonomic boundaries⁹. For example, if the aim is to restore or
 427 improve pollination services delivered to a given crop, all functional groups (e.g., buzz
 428 pollinators and oil-collectors) that compose the local guild should be considered⁷¹.

After replacing the black-and-white paradigm and focusing on functions and services, another step would be to carefully consider what kind of phenomenon is actually being represented as a network. In other words, what is flowing from node to node: matter, energy, or information ¹⁰? A network approach may help us better predict how pathogens flow from one host to the other, leading to spillover or outbreaks ⁷². Such a network approach could be guided by ITIN adapted to transient flows ⁷³. This area of application highlights the need to invest in initiatives targeted at biosurveillance and big data integration ⁷⁴.

There are critical interfaces across crops and diseases as well ⁶⁸, uniting the ecological and social layers of real-world networks ⁷⁵. It is vital to better understand how the relationships between humans, wildlife, crops, and livestock determine economic and health gains and losses ⁷⁶. ITIN can be also helpful in this human-domestic-wildlife interface, considering the huge resource dissimilarity observed in extensively changed social-ecological systems ⁷⁷.

Moving from the entire network to its layers and modules, another practical application of ITIN is the use of centrality metrics as proxies for the concept of keystone species ⁷⁸. Within this family of metrics, some have been pointed out as very useful not only for identifying the most important species responsible for maintaining different ecosystem services ⁷¹, but also to help select species to improve agroecosystems and crops in general ⁷⁹, and identify pathogen superspreaders ⁸⁰. Now it is also possible to assess the topology of multilayer networks ⁸¹, as well as the multilayer centrality of a species in such a network ⁸², especially considering thresholds in resource dissimilarity that create constraints for consumers between and within layers.

Testing for archetypal topologies

Traditionally, ecologists have used different topological indices of avoidance ⁸³, connectance ⁸⁴, complementary specialization ⁸⁵, nestedness ⁸⁶, or modularity ⁸⁷, just to give a few examples, to measure the structure of empirical or synthetical networks. Most commonly, the next step is to test whether the observed index is significantly different from expected by chance ³¹. However, neither a topological score nor a P-value alone can mechanistically explain the structure of empirical networks if the analysis is not guided by a hypothesis-oriented, strong-inference framework ⁸⁸.

Attempts to provide such a framework have been recently made in the case of nestedness⁸⁹ and compound topologies⁴⁸. Similar hypothesis-driven efforts are still lacking for the other two archetypal topologies, modular and gradient, as is an information-theoretical way to compare the fits of these four archetypal topologies to the data. Together, these advances and gaps signal that we need to work further on conceptualization and operationalization to better understand network topologies and, most importantly, their assembly rules, so we can improve our mechanistic understanding of interaction networks.

To help fill these gaps, we provide an analytical workflow built as an R tutorial to test for network topologies, using the best solutions currently available ([Supplementary Code 1](#)). Our workflow is based on five main axioms that create a framework to interpret the results of each test. First, a network's topology is a matter of degree. Second, no single metric is able to tell the whole story of a network's topology. Third, null model analysis should be used to test hypothesis-derived predictions. Fourth, networks can have mixed topologies. Fifth, weighted interaction networks have more statistical power.

Concluding remarks

Grounded in both logic and empirical evidence, the Integrative Theory of Interaction Networks brings together a coherent set of mechanistic hypotheses from which testable predictions can be drawn. It aims for generality while focusing on the assembly rules that give rise to the four archetypal topologies observed in interaction networks. Because it maps directly onto operational variables, it offers a practical framework for studying real-world systems. Over time, our field is slowly moving from a black-and-white paradigm toward a more balanced view of network assembly, something akin to a Buddhist "middle way." Like the elephant whose parts we have long examined, the whole continues to reveal more than the sum of its parts. The moment calls for widening our view, deepening our understanding of species interactions, and applying that knowledge to the urgent challenges set by the Sustainable Development Goals. There is still much to learn, and we can only move forward together.

Acknowledgements

The Integrative Theory of Interaction Networks is the outcome of exciting teamwork carried out since 2007. We are deeply grateful to all colleagues who have taken this journey together with us, by either coauthoring our previous papers or contributing with insightful discussions, especially Ana Crestani, Boris Krasnov, Cullen Geiselman, Elisângela Bezerra, Érika Braga, Fernanda Costa, Gabriel Félix, Gustavo Graciolli, Jochen Fründ, Judith Bronstein, Marcelo Nogueira, Marco Tschapka, Nico Blüthgen, Rafael Pinheiro, Richard Stevens, and Robert Poulin. TMS also acknowledges Jonathan Jeschke and the CURT network for the insightful and inspiring discussions on social-ecological networks, which significantly shaped her thinking. Our study is dedicated to the memory of Elisabeth Kalko, our dear friend and mentor, who started this research program together with us many years ago, paving the way for our theory. Her legacy will live forever in our hearts and minds. MARM was supported by grants, fellowships, and scholarships given to him and his team by the Alexander von Humboldt Foundation (AvH, 1134644), São Paulo Research Foundation (FAPESP, 2023/03083-6, 2023/02881-6, and 2023/17728-9), National Council for Scientific and Technological Development (CNPq, 305204/2024-6), and Consulate General of France in São Paulo. RLM is supported by an Australian Research Council Australian Laureate Fellowship (FL240100037) funded by the Australian Government. PMF is supported by the Muséum National d'Histoire Naturelle, Centre National de la Recherche Scientifique, and Office Français de la Biodiversité (OFB). CAK thanks the Coordination for the Improvement of Higher Education Personnel (CAPES, 8888.802356/2023-00), Graduate School in Ecology of the University of São Paulo (PPGE/IB-USP), and São Paulo Research Foundation (FAPESP, 2023/17728-9) for the Ph.D. scholarships. NL thanks the National Institutes of Health for a grant (NIH, R01AG074226). TMS thanks the CURT network funded by the Deutsche Forschungsgemeinschaft (DFG; STR 1714/2-1). ICSM thanks the National Council for Scientific and Technological Development (CNPq, 310508/2019-3). CFD acknowledges funding by the DFG Research Unit 5207 "Reassembly". Last, but not least, we thank FAPESP, CNPq, CAPES, and German Academic Exchange Service (DAAD) for the scholarships and fellowships granted to our students and postdocs.

Author contributions

MARM has since 2007 coordinated the research program that led to the novel theory. MARM, SES, and CFD discussed the final synthesis of the theory, wrote the first draft of the manuscript and the supplementary material, and coordinated the writing. CAK and RLM worked on the literature review and figures. MARM, CFD, and RLM worked on the supplement. All other authors contributed by discussing the ideas that form the corpus of ITIN and reviewing drafts of the manuscript and supplement.

Competing interests

None to declare.

Disclosure of generative AI-use

We used ChatGPT (OpenAI, <https://chatgpt.com>) in combination with GitHub Copilot (GitHub, <https://github.com/features/copilot>) to optimize the code provided in the supplement.

Supplementary materials

Supplementary Code 1: We have prepared a repository on GitHub with an analytical workflow that includes data, code, and tutorials aimed at helping assess the topology of empirical and synthetical interaction networks, within the framework proposed by the Integrative Theory of Interaction Networks: https://github.com/marmello77/new_theory.

References

1. Ings, T. C. & Hawes, J. E. The History of Ecological Networks. in *Ecological Networks in the Tropics* (eds. Dáttilo, W. & Rico-Gray, V.) 15–28 (Springer International Publishing, Cham, 2018).
2. Hervías-Parejo, S. *et al.* On the structure of species-function participation in

- 544 multilayer ecological networks. *Nat. Commun.* **15**, 8910 (2024).
- 545 3. Bianco, G., Manning, P. & Schleuning, M. A quantitative framework for
546 identifying the role of individual species in Nature’s Contributions to People. *Ecol.*
547 *Lett.* **27**, (2024).
- 548 4. Barabasi, A.-L. *Network Science*. (Cambridge University Press, Cambridge,
549 England, 2016).
- 550 5. Camerano, L. On the equilibrium of living beings by means of reciprocal
551 destruction. *Atti della Reale Accademia delle Scienze di Torino* **15**, 393–414
552 (1880).
- 553 6. Bascompte, J. & Jordano, P. *Mutualistic Networks*. (Princeton University Press,
554 Princeton, NJ, 2014).
- 555 7. Guimarães, P. R. The Structure of Ecological Networks Across Levels of
556 Organization. *Annu. Rev. Ecol. Evol. Syst.* **51**, 433–460 (2020).
- 557 8. Blüthgen, N. & Staab, M. A Critical Evaluation of Network Approaches for
558 Studying Species Interactions. *Annu. Rev. Ecol. Evol. Syst.* (2024)
559 doi:10.1146/annurev-ecolsys-102722-021904.
- 560 9. Brimacombe, C., Bodner, K., Michalska-Smith, M. & Fortin, M.-J. Shortcomings
561 of reusing species interaction networks created by different sets of researchers.
562 *PLoS Biol.* (2023) doi:10.1371/journal.pbio.3002068.
- 563 10. Brose, U. *et al.* Embedding information flows within ecological networks. *Nat.*
564 *Ecol. Evol.* **9**, 547–558 (4 2025).
- 565 11. Lewinsohn, T. M., Inácio Prado, P., Jordano, P., Bascompte, J. & M. Olesen, J.
566 Structure in plant-animal interaction assemblages. *Oikos* **113**, 174–184 (2006).
- 567 12. Ulrich, W. *et al.* A comprehensive framework for the study of species co-
568 occurrences, nestedness and turnover. *Oikos* **126**, 1607–1616 (2017).

- 569 13. Palazzi, M. J., Borge-Holthoefer, J., Tessone, C. J. & Solé-Ribalta, A. Macro- and
570 mesoscale pattern interdependencies in complex networks. *J. R. Soc. Interface* **16**,
571 20190553–20190553 (2019).
- 572 14. Lampo, A., Palazzi, M. J., Borge-Holthoefer, J. & Solé-Ribalta, A. Hybrid
573 structural arrangements mediate stability and feasibility in mutualistic networks.
574 (2021).
- 575 15. Flores, C. O., Valverde, S. & Weitz, J. S. Multi-scale structure and geographic
576 drivers of cross-infection within marine bacteria and phages. *ISME J.* **7**, 520–532
577 (2013).
- 578 16. Cumming, G. S. Heterarchies: Reconciling Networks and Hierarchies. *Trends*
579 *Ecol. Evol.* **31**, 622–632 (2016).
- 580 17. Newman, M. E. J. Assortative mixing in networks. *Phys. Rev. Lett.* **89**, 208701–
581 208701 (2002).
- 582 18. Bascompte, J., Jordano, P., Melián, C. J. & Olesen, J. M. The nested assembly of
583 plant–animal mutualistic networks. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 9383–9387
584 (2003).
- 585 19. Pimm, S. L. *Food Webs*. (The University of Chicago Press, Chicago, IL, 2002).
- 586 20. Fonseca, C. R. & Ganade, G. Asymmetries, Compartments and Null Interactions in
587 an Amazonian Ant-Plant Community. *J. Anim. Ecol.* **65**, 339–339 (1996).
- 588 21. Guimarães, P. R. *et al.* Interaction Intimacy Affects Structure and Coevolutionary
589 Dynamics in Mutualistic Networks. *Curr. Biol.* **17**, 1797–1803 (2007).
- 590 22. Tylianakis, J. M. & Morris, R. J. Ecological Networks Across Environmental
591 Gradients. *Annu. Rev. Ecol. Evol. Syst.* **48**, 25–48 (2017).
- 592 23. Strona, G. Nestedness and Ecological Network Stability. in 137–147 (2022).
- 593 24. Dormann, C. F. The rise, and possible fall, of network ecology. in *Defining*

- 594 *Agroecology: A Festschrift for Teja Tscharntke* (eds. Dormann, C. F. et al.) 143–
595 159 (Tredition, Hamburg, 2023).
- 596 25. Staniczenko, P. P. A., Kopp, J. C. & Allesina, S. The ghost of nestedness in
597 ecological networks. *Nat. Commun.* **4**, 1391–1391 (2013).
- 598 26. Krishna, A., Guimaraes, P. R., Jordano, P. & Bascompte, J. A neutral-niche theory
599 of nestedness in mutualistic networks. *Oikos* **117**, 1609–1618 (2008).
- 600 27. James, A., Pitchford, J. W. & Plank, M. J. Disentangling nestedness from models
601 of ecological complexity. *Nature* **487**, 227–230 (2012).
- 602 28. Fortuna, M. A. *et al.* Nestedness versus modularity in ecological networks: two
603 sides of the same coin? *J. Anim. Ecol.* **79**, 811–817 (2010).
- 604 29. Almeida-Neto, M., Guimaraes, P. R. & Lewinsohn, T. M. On nestedness analyses:
605 rethinking matrix temperature and anti-nestedness. *Oikos* **116**, 716–722 (2007).
- 606 30. Almeida-Neto, M., Guimarães, P., Guimarães, P. R., Loyola, R. D. & Ulrich, W. A
607 consistent metric for nestedness analysis in ecological systems: reconciling concept
608 and measurement. *Oikos* **117**, 1227–1239 (2008).
- 609 31. Dormann, C. F., Frund, J., Bluthgen, N. & Gruber, B. Indices, Graphs and Null
610 Models: Analyzing Bipartite Ecological Networks. *Open Ecol. J.* **2**, 7–24 (2009).
- 611 32. Bezerra, E. L. S., Machado, I. C. & Mello, M. A. R. Pollination networks of oil-
612 flowers: a tiny world within the smallest of all worlds. *J. Anim. Ecol.* **78**, 1096–
613 1101 (2009).
- 614 33. Mello, M. A. R. *et al.* The modularity of seed dispersal: differences in structure
615 and robustness between bat– and bird–fruit networks. *Oecologia* **167**, 131–131
616 (2011).
- 617 34. Donatti, C. I. *et al.* Analysis of a hyper-diverse seed dispersal network: modularity
618 and underlying mechanisms. *Ecol. Lett.* **14**, 773–781 (2011).

- 619 35. Windsor, F. M. *et al.* Network science: Applications for sustainable
620 agroecosystems and food security. *Perspect. Ecol. Conserv.* **20**, 79–90 (2022).
- 621 36. Dansereau, G. *et al.* Overcoming the disconnect between species interaction
622 networks and biodiversity conservation. *Trends Ecol. Evol.* (07 2025)
623 doi:10.1016/j.tree.2025.05.010.
- 624 37. Pinheiro, R. B. P. *et al.* Trade-offs and resource breadth processes as drivers of
625 performance and specificity in a host–parasite system: a new integrative
626 hypothesis. *International Journal for Parasitology* **46**, 115–121 (2016).
- 627 38. Pinheiro, R. B. P., Felix, G. M. F., Dormann, C. F. & Mello, M. A. R. A new
628 model explaining the origin of different topologies in interaction networks.
629 *Ecology* **100**, e02796–e02796 (2019).
- 630 39. Martinez, N. D. Allometric Trophic Networks From Individuals to Socio-
631 Ecosystems: Consumer–Resource Theory of the Ecological Elephant in the Room.
632 *Front. Ecol. Evol.* **8**, (2020).
- 633 40. Case, T. J. *An Illustrated Guide to Theoretical Ecology*. (Oxford University Press,
634 London, England, 1999).
- 635 41. Mello, M. A. R. & Dormann, C. F. We need to talk about the topology of
636 interaction networks. *Oecol. Australis* **29**, 190–195 (2025).
- 637 42. Clune, J., Mouret, J.-B. & Lipson, H. The evolutionary origins of modularity.
638 *Proc. Biol. Sci.* **280**, 20122863–20122863 (2013).
- 639 43. Crestani, A. C., Mello, M. A. R. & Cazetta, E. Interindividual variations in plant
640 and fruit traits affect the structure of a plant-frugivore network. *Acta Oecol.*
641 *(Montrouge)* **95**, 120–127 (2019).
- 642 44. Guimarães, P. R., Pires, M. M., Jordano, P., Bascompte, J. & Thompson, J. N.
643 Indirect effects drive coevolution in mutualistic networks. *Nature* **550**, 511–514

(2017).

45. Travassos-Britto, B., Pardini, R., El-Hani, C. N. & Prado, P. I. Towards a pragmatic view of theories in ecology. *Oikos* **130**, 821–830 (2021).
46. Suppes, P. *Representation and Invariance of Scientific Structures*. (Center for the Study of Language and Information, Chicago, IL, 2001).
47. Van Fraassen, B. C. Aim and Structure of Scientific Theories. in 307–318 (1986).
48. Felix, G. M., Pinheiro, R. B. P., Poulin, R., Krasnov, B. R. & Mello, M. A. R. The compound topology of host–parasite networks is explained by the integrative hypothesis of specialization. *Oikos* **2022**, (2022).
49. Laurindo, R. de S. *et al.* Drivers of bat roles in Neotropical seed dispersal networks: abundance is more important than functional traits. *Oecologia* **193**, 189–198 (2020).
50. Devriese, A., Peeters, G., Brys, R. & Jacquemyn, H. Interactions outside local patches contribute to the compound topology of plant–pollinator networks in fragmented dune slacks. *Oikos* (2025) doi:10.1002/oik.11176.
51. Nakamura, D. Y. M., Koffler, S., Mello, M. A. R. & Franco, T. M. Resin foraging interactions in stingless bees: an ecological synthesis using multilayer networks. *Apidologie (Celle)* **55**, 34–34 (2024).
52. Martins Santos, F. *et al.* Armadillo burrows: a meeting point for biodiversity in the Pantanal Wetland. *Mamm. Biol.* (10 2024) doi:10.1007/s42991-024-00466-8.
53. Angarita-Báez, J., Delfino, H. & Carlos, C. American shorebird-prey network patterns: Geographic variation in non-breeding habitats. **43**, (05 2025).
54. Santos, M., Rodrigues, G., Fransozo, M. L. & Fransozo, V. Resource partitioning and adequacy among ontogenetic groups in a hermit crab and gastropod shell network. *Aquat. Ecol.* **55**, (03 2021).

- 669 55. Domínguez-García, V. & Kéfi, S. The structure and robustness of ecological
670 networks with two interaction types. *PLoS Comput. Biol.* **20**, e1011770 (2024).
- 671 56. Gonçalves, M. *et al.* Traditional Agriculture and Food Sovereignty: Quilombola
672 Knowledge and Management of Food Crops. *J. Ethnobiol.* **42**, (07 2022).
- 673 57. Gaston, K. J. *Rarity*. (Springer, Dordrecht, Netherlands, 2012). doi:10.1007/978-
674 94-011-0701-3.
- 675 58. Keddy, P. A. & Shipley, B. Competitive hierarchies in herbaceous plant
676 communities. *Oikos* **54**, 234 (1989).
- 677 59. Frank, S. A. The common patterns of nature. *J. Evol. Biol.* **22**, 1563–1585 (2009).
- 678 60. Lopezaraiza-Mikel, M. E., Hayes, R. B., Whalley, M. R. & Memmott, J. The
679 impact of an alien plant on a native plant-pollinator network: an experimental
680 approach. *Ecol. Lett.* **10**, 539–550 (2007).
- 681 61. Cantor, G. Ueber unendliche, lineare Punktmannichfaltigkeiten. *Math. Ann.* **21**,
682 545–591 (1883).
- 683 62. Olesen, J. M. *et al.* Missing and forbidden links in mutualistic networks. *Proc.*
684 *Biol. Sci.* **278**, 725–732 (2011).
- 685 63. Hubbell, S. P. *The Unified Neutral Theory of Biodiversity and Biogeography*.
686 (Princeton University Press, Princeton, NJ, 2001).
- 687 64. Barabasi, A.-L. & Albert, R. Emergence of scaling in random networks. *Science*
688 **286**, 509–512 (1999).
- 689 65. van Valen, L. A new evolutionary law. *Evolutionary Theory* **1**, 1–30 (1973).
- 690 66. Landim, A., Guimarães, P., Fernandez, F. & Dias, A. A framework for the
691 restoration of seed dispersal and pollination. *Restor. Ecol.* (2024)
692 doi:10.1111/rec.14151.
- 693 67. Ulrich, J. & Peters, V. Integrating network theory and biodiversity conservation:

- do different species selection approaches result in different recommendations?
Biodivers. Conserv. **32**, (2023).
68. Plowright, R. *et al.* Ecological countermeasures to prevent pathogen spillover and subsequent pandemics. *Nat. Commun.* **15**, (2024).
69. United Nations. *Transforming Our World: The 2030 Agenda for Sustainable Development*. <https://sdgs.un.org> (2015).
70. Adisasmito, W. B. *et al.* One Health: A new definition for a sustainable and healthy future. *PLoS Pathog.* **18**, e1010537–e1010537 (2022).
71. Rafferty, N. & Cosma, C. Sustainable nature-based solutions require establishment and maintenance of keystone plant-pollinator interactions. *J. Ecol.* (2024) doi:10.1111/1365-2745.14353.
72. Gibb, R. *et al.* The anthropogenic fingerprint on emerging infectious diseases. *medRxiv* (2024) doi:10.1101/2024.05.22.24307684.
73. Tao, Y., Hite, J. L., Lafferty, K. D., Earn, D. J. D. & Bharti, N. Transient disease dynamics across ecological scales. *Theor. Ecol.* **14**, 625–640 (2021).
74. Carlson, C. J. *et al.* The Global Virome in One Network (VIRION): an Atlas of Vertebrate-Virus Associations. *MBio* (2022) doi:10.1128/mbio.02985-21.
75. Muylaert, R. L. *et al.* Connections in the Dark: Social-Ecological Networks as a Promising Tool for Bat Conservation and One Health. *bioRxiv* (2025) doi:10.1101/2025.07.22.666214.
76. Kluger, L. C., Gorris, P., Kochalski, S., Mueller, M. S. & Romagnoni, G. Studying human–nature relationships through a network lens: A systematic review. *People Nat. (Hoboken)* **2**, 1100–1116 (2020).
77. Bodin, Ö. Social-Ecological Networks: What Are They, Why Are They Useful And How Can I Use Them? in *The Sage Handbook of Social Network Analysis*

- 719 (eds. McLevey, J., Scott, J. & Carrington, P. J.) 75–75 (SAGE Publications Ltd,
720 Thousand Oaks, 2023).
- 721 78. Cirtwill, A. R. *et al.* A review of species role concepts in food webs. *Food Webs*
722 **16**, e00093–e00093 (2018).
- 723 79. Allasino, M. L., Haedo, J. P., Lázaro, A., Torretta, J. P. & Marrero, H. J. Positive
724 relationship between crop centrality and pollination service. *Agric. Ecosyst.*
725 *Environ.* **343**, 108279–108279 (2023).
- 726 80. Madotto, A. & Liu, J. Super-spreader identification using meta-centrality. *Sci. Rep.*
727 **6**, (2016).
- 728 81. Pilosof, S., Porter, M. A., Pascual, M. & Kéfi, S. The multilayer nature of
729 ecological networks. *Nat. Ecol. Evol.* **1**, 0101–0101 (2017).
- 730 82. Lotfi, N., Requejo, H. S., Rodrigues, F. A. & Mello, M. A. R. A new centrality
731 index designed for multilayer networks. *Methods Ecol. Evol.* **15**, 204–213 (2024).
- 732 83. Stone, L. & Roberts, A. The checkerboard score and species distributions.
733 *Oecologia* **85**, 74–79 (1990).
- 734 84. Jordano, P. Patterns of mutualistic interactions in pollination and seed dispersal -
735 connectance, dependence asymmetries, and coevolution. *Am. Nat.* **129**, 657–677
736 (1987).
- 737 85. Blüthgen, N., Menzel, F., Hovestadt, T., Fiala, B. & Blüthgen, N. Specialization,
738 Constraints, and Conflicting Interests in Mutualistic Networks. *Curr. Biol.* **17**,
739 341–346 (2007).
- 740 86. Ulrich, W., Almeida-Neto, M. & Gotelli, N. J. A consumer's guide to nestedness
741 analysis. *Oikos* **118**, 3–17 (2009).
- 742 87. Beckett, S. J. Improved community detection in weighted bipartite networks. *R.*
743 *Soc. Open Sci.* **3**, 140536–140536 (2016).

- 744 88. Platt, J. R. Strong Inference: Certain systematic methods of scientific thinking may
745 produce much more rapid progress than others. *Science* **146**, 347–353 (1964).
- 746 89. Pinheiro, R. B. P., Dormann, C. F., Felix, G. M. & Mello, M. A. R. A novel
747 perspective on the meaning of nestedness with conceptual and methodological
748 solutions. *bioRxiv* 2021.04.05.438470–2021.04.05.438470 (2021).
- 749