

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/233819496>

# Umbria–Marche Basin, Central Italy: A Reference Section for the Aptian–Albian Interval at Low Latitudes

Article in *Scientific Drilling* · April 2012

DOI: 10.5194/sd-13-42-2012

CITATIONS

14

11 authors, including:



**Rodolfo Coccioni**

Università degli Studi di Urbino "Carlo Bo"

234 PUBLICATIONS 4,618 CITATIONS

[SEE PROFILE](#)



**Giuseppe Bancala**

Schlumberger Limited

14 PUBLICATIONS 150 CITATIONS

[SEE PROFILE](#)

READS

154



**Luigi Jovane**

University of São Paulo

126 PUBLICATIONS 1,321 CITATIONS

[SEE PROFILE](#)



**Gerson Fauth**

Universidade do Vale do Rio dos Sinos

117 PUBLICATIONS 373 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Magnetostratigraphic analysis of ferromanganese deposits [View project](#)



Foraminifera and thecamoebians from tropical estuaries [View project](#)

# The Poggio Le Guaine Core (Umbria-Marche Basin, Central Italy): A Reference Section for the Aptian-Albian Interval at Low Latitudes

by Rodolfo Coccioni, Luigi Jovane, Giuseppe Bancalà, Carla Bucci, Gerson Fauth, Fabrizio Frontalini, Liliane Janikian, Jairo Savian, Renato Paes de Almeida, Grasiane Luz Mathias, and Ricardo Ivan Ferreira da Trindade

doi:10.2204/iodp.sd.13.07.2011

## Abstract

The Poggio le Guaine core from the Umbria–Marche Basin in central Italy penetrated 98.72 m of pelagic sediments consisting mainly of limestones, calcareous marls, marls, clays, and black shales. The cored interval extends from the Albian–Cenomanian boundary down to the uppermost Barremian (99.6–126 Ma). The core was completely logged and continuously drilled with essentially 100% recovery of excellent quality material. The cored sequence recorded from OAE1a to OAE1d and regional equivalents of widely distributed Aptian–Albian oceanic anoxic events. The Poggio le Guaine core can be considered a 'reference section' for the Aptian–Albian interval at low latitudes and is designated to provide high-ranking informative records for that critical time interval through high-resolution multiproxy studies.

## Introduction

Within the Cretaceous Period, the Aptian–Albian interval (125–99.6 Ma, Ogg et al., 2008) was a critical time on a global scale. This is evident from 1) changes in the nature of the ocean–climate system brought about by increased ocean crust production coupled with active midplate and plate margin volcanism in a shifting paleogeography (Skelton et al., 2003); 2) cyclic deposition and preservation of common 'black shales', some of them termed Oceanic Anoxic Events

(OAE1a to OAE1d) (Schlanger and Jenkyns, 1976; Arthur et al., 1990); 3) periodic changes in redox conditions at the ocean bottom (Oceanic Red Beds, ORBs) (Wang et al., 2009); and 4) rapid biotic radiations and turnovers (Leckie et al., 2002). The Aptian–Albian time is also of interest for one of the most noteworthy geomagnetic events, namely the post-M0r 'Cretaceous Quiet Zone'. This long and constant normal polarity superchron without any convincing true reversal to date (Satolli et al., 2008) precludes usage of reversals magnetostratigraphy from the Aptian through the Santonian.

The Poggio le Guaine core was designed to provide the following: 1) a high-resolution relative magnetic paleointensity reference curve for the Aptian–Albian interval of the long normal Cretaceous superchron that would allow correlation of the relative paleointensity curves from other places in the world; 2) a high resolution age model constrained by astronomical tuning and absolute dating, into which the sequence of the geological, biogeochemical, oceanographic and climatic events throughout the Aptian–Albian critical time can be placed; and 3) understanding of the causal linkages among these events as well as their consequences. The core was drilled at Poggio le Guaine, where the most continuous, complete, and best preserved Aptian–Albian succession is exposed throughout the Umbria–Marche Basin (UMB) of the northern Apennines of central Italy (Fig. 1). It represents a continuous record of fossiliferous pelagic rocks extending from the Albian–Cenomanian boundary down to the uppermost Barremian (99.6–126 Ma). In this progress report we present the first preliminary findings of this ongoing project.

In this progress report we present the first preliminary findings of this ongoing project.

## Geological and stratigraphic setting

After three decades of dedicated research, the Aptian–Albian pelagic succession of the UMB has become a classical reference section for studies at regional to global scale. It deposited well above the calcite compensation depth at middle to lower bathyal depths (1000–1500 m) and at ~20°N paleolatitude over the southern margin of the western Tethys Ocean (Arthur and Premoli Silva, 1982; Coccioni et al., 1987, 1989, 1990, 1992; Erba, 1988, 1992; Cresta et al., 1989; Tornaghi et al., 1989; Coccioni,



**Figure 1.** Location of the Poggio le Guaine drill site in the Umbria–Marche Basin (northern Apennines, central Italy). The location of the Piobbico drill site—where an 84-m-thick core extending from the upper Albian down to the uppermost Barremian was drilled in 1982—is also shown.


1996; Coccioni and Galeotti, 1993; Satolli et al., 2008; Tiraboschi et al., 2009; Turchyn et al., 2009).

This succession extends from the uppermost part of the Maiolica Formation (Tithonian to lower Aptian) to the lower part of the Scaglia Bianca Formation (uppermost Albian to lowermost Turonian) and includes the entire Marne a Fucoidi Formation (Fig. 2). The uppermost part of the Maiolica Formation is represented by thin-bedded white to gray limestones interbedded with black shales. The lower part of the Scaglia Bianca Formation is characterized by thin-bedded yellowish-gray limestones with subordinate reddish limestones and some discrete thin black shales, which are the regional sedimentary expression of the latest Albian OAE1d (Fig. 2). Within the Cretaceous succession of the UMB, the Aptian-Albian Marne a Fucoidi Formation represents a distinctive varicolored interlude with more shale. This formation consists of thinly interbedded pale reddish to dark reddish, pale olive to dark reddish brown and pale olive to grayish olive marlstones and calcareous marlstones together with dark gray to black organic carbon-rich shales, usually with a low carbonate content, and yellowish-gray to light gray marly limestones and limestones (Fig. 2). Several distinctive organic-rich black shale and marl marker beds occur within the Aptian-Albian interval, some of which have been identified as the regional sedimentary expression of OAE1a to OAE1d (Coccioni et al., 1987, 1989, 1990; Coccioni, 1996, 2001).

Following the timescale of Ogg et al. (2008), we estimate the sediment accumulation rate to be  $\sim 0.24$  cm kyr<sup>-1</sup> and  $\sim 0.5$  cm kyr<sup>-1</sup> in the Aptian and Albian, respectively.

## Location of the drilling site and drilling operations

The Poggio le Guaine succession was selected as the reference section for the Aptian-Albian interval following pre-




**Figure 2.** Stratigraphic framework of the Poggio le Guaine core with real stratigraphic depths according to bed dip measurements. Also shown are the occurrence and distribution of the organic-rich black shales including the marker beds resulting from Oceanic Anoxic Events (OAEs) and the discrete intervals where reddish colored beds become dominant (Cretaceous Oceanic Red Beds, CORBs). Integrated plankton biostratigraphy is adapted after Coccioni et al. (1987, 1989, 1990, 1992, 2007), Erba (1988, 1992), Tornaghi et al. (1989), Tiraboschi et al. (2009) and unpublished data. Abbreviations: Lithos. = Lithostratigraphic, Barr. = Barremian, Cen. = Cenomanian, H. = Hedbergella, L. = Leupoldina, G. = Globigerinelloides, P. = Paraticinella, Ps. = Pseudoplanomalina, T. = Ticinella, B. = Biticinella, Psth. = Pseudothalmanninella, Pth. = Parathalmanninella, Th. = Thalmanninella, C. = Chiastozygus, Pa. = Parahbdolithus, N. = Nannoconus, Pr. = Predicosphaera, A. = Axopodorhabdus, R. = Rhagodiscus, E. = Eiffellithus. Timescale is after Ogg et al. (2008). Following Huber and Leckie (2011), the Aptian-Albian boundary is placed at the highest occurrence of Paraticinella eubejaouensis.

vious detailed stratigraphic investigation (Coccioni et al., 1987, 1989, 1990; Coccioni, 1996).

The Poggio le Guaine drill site (lat. 43°32'42.72"N; long. 12°32'40.92"E) is located on the Monte Nerone ridge at 888 m above sea level, 6 km west of the town of Cagli (Regione Marche, Italy) (Fig. 1). The outcropping beds show a strike of 310° and a dip of 30° to the northeast.

The drilling campaign took place on 1–14 September 2010. Drilling operations were performed with the Atlas Copco Mustang 5-F4 surface core drilling rig (Fig. 3). Coring





**Figure 4.** Photographs of selected cores recovered at Poggio le Guaine. [A] Cores 20 (20.90–22.45 m) to 22 (23.83–25.39 m) penetrated the marly limestones, marls, and black shales of the upper Albian Amadeus Segment (OAE1c) (22.77–24.74 m); [B] Cores 46 (58.42–60.01 m) to 49 (62.68–64.33 m) recovered the marly clays, clays, marls, and black shales of the lower Albian Monte Nerone (57.07–60.01 m) and Kilian (62.64–63.02 m) Levels (OAE1b); [C] Cores 49 (62.68–64.33 m) to 52 (67.18–68.77 m) penetrated latest Aptian marls and calcareous marls together with the black shales of the uppermost Aptian Jacob Level (OAE1b) (67.44–67.52); [D] Cores 54 (70.32–71.89 m) to 57 (74.94–76.45 m) recovered late Aptian marls, marly clays, and calcareous marls; [E] Cores 66 (88.00–89.60 m) to 68 (90.85–91.99 m) penetrated the lower Aptian Selli Level (OAE1a) (89.24–91.19).

was accomplished with a T2 double corer, using narrow-kerf, sawtoothed drill bits that cut a 101-cm-diameter hole and approximately 8-cm-diameter cores. The target interval was successfully encountered below the weathering zone, which is only a couple of decimeters thick. Lithologies were logged and digital color photographs taken, as each core section was recovered (Fig. 4). Later, the seventy-one cores drilled were packed, labeled, and put in PVC

plastic boxes to prevent contamination and loss of moisture. During core packing, care was taken in collecting the fragments from the few fractured portions in plastic bags. The entire core is stored at the core repository of the Department of Earth, Life and Environmental Sciences, University of Urbino.



**Figure 3.** Drilling operations at Poggio le Guaine.

The Poggio le Guaine cored interval extends from the Albian-Cenomanian boundary down to the uppermost Barremian including the upper transition to the Scaglia Bianca Formation and the lower transition to the Maiolica Formation. The total length of the core is 98.72 m with essentially 100% recovery of excellent quality material throughout the drilling. For each core a set of dip measurements was taken directly at the drill site with four measurements on average for each meter of core. Marked changes of dip angle were observed in the uppermost and lowermost portions of the core, the latter particularly affected by folding and minor faulting. Taking into account the dip measurements, the corrected thickness of the drilled section is 96.02 m, with 82.53 m corresponding to the Marne a Fucoidi Formation, 3.51 m corresponding to the underlying Maiolica Formation, and 9.98 m corresponding to the overlying Scaglia Bianca Formation (Fig. 2).

### Litho-, Bio-, and Chronostratigraphy of the Poggio le Guaine Core

On the basis of dominant colors determined with the Munsell Rock Color Chart, the pattern of alternating colors, the calcium carbonate content, the occurrence of black shales, chert and radiolarian-rich beds, and by direct comparison with previous results from an adjacent outcrop (Coccioni et al., 1987, 1989, 1990), the Poggio le Guaine core was subdivided into twenty-three lithological units, which are recognizable on a regional scale (Coccioni, 1996; Fig. 2). Unit 1 to part of Unit 4 corresponds to the Scaglia Bianca Formation, and part of Unit 4 to the upper part of Unit 23 corresponds to the Marne a Fucoidi Formation. The Maiolica Formation is equated with most of Unit 23.

A total of two hundred forty organic-rich black shales and marls, millimeter to decimeter thick, is recorded in the cored sequence. These layers are not evenly distributed throughout the core. They are frequent to common in Units 2, 5 to 7, 9 to 15, 22 and 23 that are dominated by lithotypes that are pale olive to grayish olive and yellowish gray to light gray, and occasionally by pale olive and pale reddish brown to dark reddish brown (Fig. 2). Some of these organic-rich beds record OAE1a to OAE1d, namely the lower Aptian Selli Level (OAE1a, ~124 Ma), the uppermost Aptian Jacob Level, the lowermost Albian Kilian Level, the lower Albian Monte Nerone Level and the Urbino Level (all added to OAE1b, ~110–113 Ma), the upper Albian Amadeus Segment (equivalent to part of OAE1c, ~105 Ma) and the uppermost Albian Piali Level (OAE1d, ~100.5 Ma). Some others are interpreted as regional equivalents of widely distributed OAEs, namely the lower Aptian Noir Level and the middle to upper Aptian Fallot Level (Herrle et al., 2004; Figs. 2, 4).

Seven discrete intervals dominantly reddish are also recognized in the cored sequence, which correlate well with those identified as ORBs in the Aptian-Albian succession of the UMB (Hu et al., 2005; Fig. 2).

In agreement with previous detailed stratigraphic investigations from Aptian-Albian land-based sections throughout the UMB (Coccioni et al., 1987, 1989, 1990, 1992; Cresta et al., 1989; Coccioni and Galeotti, 1993; Coccioni 1996, 2001, 2002) and from a 84-m-thick core drilled at Piobbico (Fig. 1) in 1982 and extending from the upper Albian down to the uppermost Barremian (Erba, 1988, 1992; Tornaghi et al., 1989), the cored succession biostratigraphically represents the interval from the *Hedbergella excelsa* to the base of the *Thalmaninella globotruncanoides* planktonic foraminiferal zones and from the *Chiastozygus litterarius* (NC6) to the *Eiffelithus turriseiffelii* (NC10) calcareous nannofossil zones. Accordingly, from a chronostratigraphic point of view the Poggio le Guaine core encompasses the latest Barremian to earliest Cenomanian interval (Fig. 2).

## Objectives

The Poggio le Guaine core, which can be considered as a reference section for the Aptian-Albian interval at low latitudes, is designated to provide high-ranking informative records for that critical time interval through high-resolution multiproxy studies carried by a consortium of Italian and Brazilian researchers. The studies already in progress include integrated stratigraphic and sedimentological analyses, multidisciplinary magneto-, bio-, and chemostratigraphy, as well as clay and organic geochemistry investigations.

The data set collected will supply the framework for the following three activities. 1) Computing a high-resolution magnetostratigraphy, in particular a relative paleointensity reference curve for the Aptian–Albian interval of the long normal Cretaceous superchron, will allow correlation of rela-

tive paleointensity curves from other places in the world. 2) A unique and original age model can be built up, at high resolution and using absolute dating and astronomical tuning. Into this the sequence of the geological, biogeochemical, oceanographic and climatic events that are recorded throughout the Aptian–Albian critical time can be placed. 3) The causal linkages, as well as their consequences, among these events can be defined.

## Acknowledgments

The Poggio le Guaine core is an integral part of the Magnetic Anomaly of Relative Intensity for the Aptian (MARIA) project. Coring was funded by Petróleo Brasileiro S.A. - Petrobras. The 'Università Agraria di Secchiano' (Cagli, Regione Marche, Italy) generously granted access and facilitated drilling.

## References

- Arthur, M.A., and Premoli Silva, I., 1982. Development of widespread organic carbon-rich strata in the Mediterranean Tethys. In Schlanger, S.O., and Cita, M.B. (Eds.), *Nature of Cretaceous Carbon-Rich Facies*: San Diego (Academic Press), 7–54.
- Arthur, M.A., Jenkyns, H.C., Brumsack, H.-J., and Schlanger, S.O., 1990. Stratigraphy, geochemistry, and paleoceanography of organic-carbon-rich Cretaceous sequences. In Ginsburg, R.N., and Beaudoin, B. (Eds.), *Cretaceous Resources, Events and Rhythms. NATO ASI Ser. 304*: Dordrecht, Netherlands (Kluwer Acad.), 75–119.
- Coccioni, R., 1996. The Cretaceous of the Umbria-Marche Apennines (Central Italy). Jost Wiedmann Symposium on Cretaceous Stratigraphy, Paleobiology and Paleobiogeography, Tübingen, 7–10 March 1996, 129–136.
- Coccioni, R., 2001. The "Piali Level" from the latest Albian of the Umbria-Marche Apennines (Italy). *Federazione Italiana di Scienze della Terra, Geoitalia 2001*, 192–193.
- Coccioni, R., and Galeotti, S., 1993. Orbitally induced cycles in benthonic foraminiferal morphogroups and trophic structures distribution patterns from the Late Albian "Amadeus Segment" (Central Italy). *J. Micropaleontol.*, 12:227–239, doi:10.1144/jm.12.2.227.
- Coccioni, R., Nesci, O., Tramontana, M., Wezel, F.C., and Moretti, E., 1987. Descrizione di un livello-guida "radiolaritico-bituminoso-ittiolitico" alla base delle Marne a Fucoidi nell'Appennino umbro-marchigiano. *Boll. Soc. Geol. Italiana*, 106:183–192.
- Coccioni, R., Franchi, R., Nesci, O., Wezel, F.C., Battistini, F., and Pallecchi, P., 1989. Stratigraphy and mineralogy of the Selli Level (Early Aptian) at the base of the Marne a Fucoidi in the Umbro-Marchean Apennines, Italy. In Wiedmann, J. (Ed.), *Cretaceous of the Western Tethys. Proceedings 3rd International Cretaceous Symposium*, Stuttgart (E. Schweizerbart'sche Verlagsbuchhandlung), 563–584.
- Coccioni, R., Franchi, R., Nesci, O., Perilli, N., Wezel, F.C., and Battistini, F., 1990. Stratigrafia, micropaleontologia e mineralogia delle Marne a Fucoidi delle sezioni di Poggio le Guaine e del Fiume Bosso (Appennino umbro-marchigiano). Atti 2° Convegno Internazionale "Fossili,



- Evoluzione, Ambiente", Pergola, 25–30 ottobre 1987, Tecnostampa, 163–201.
- Coccioni, R., Erba, E., and Premoli Silva, I., 1992. Barremian-Aptian calcareous plankton biostratigraphy from the Gorgo a Cerbara section (Marche, Central Italy) and implications for plankton evolution. *Cret. Res.*, 13:517–537, doi:10.1016/0195-6671(92)90015-I.
- Coccioni, R., Premoli Silva, I., Marsili, A., and Verga, D., 2007. First radiation of Cretaceous planktonic foraminifera with radially elongate chambers at Angles (Southeastern France) and biostratigraphic implications. *Rev. micropaléontologie*, 50:215–224.
- Cresta, S., Monechi, S., and Parisi, G., 1989. Stratigrafia del Mesozoico al Cenozoico nell'area Umbro-Marchigiana. *Mem. Descr. Carta Geol. Italia*, 34:185.
- Erba, E., 1988. Aptian-Albian calcareous nannofossil biostratigraphy of the Scisti a Fucoidi cored at Piobbico (central Italy). *Riv. Ital. Paleont. Strat.*, 94:249–284.
- Erba, E., 1992. Calcareous nannofossil distribution in pelagic rhythmic sediments (Aptian-Albian Piobbico core, central Italy). *Riv. Ital. Paleont. Strat.*, 97:455–484.
- Herrle, J.O., Kössler, P., Friedrich, O., Erlenkeuser, H., and Hemleben, C., 2004. High-resolution carbon isotope records of the Aptian to Lower Albian from SE France and the Mazagan Plateau (DSDP Site 545): A stratigraphic tool for paleoceanographic and paleobiologic reconstruction. *Earth Planet. Sci. Lett.*, 218:149–161, doi:10.1016/S0012-821X(03)00646-0.
- Hu, X., Jansa, L., and Sarti, M., 2005. Mid-Cretaceous oceanic red beds in the Umbria-Marche Basin, central Italy: Constraints on paleoceanography and paleoclimate. *Paleogeogr. Paleoclimatol. Paleoecol.* 233:163–186, doi:10.1016/j.palaeo.2005.10.003.
- Huber, B.T., and Leckie, R.M., 2011. Planktic foraminiferal species turnover across deep-sea Aptian/Albian boundary sections. *J. Foram. Res.*, 41:53–95, doi:10.2113/gsjfr.41.1.53.
- Leckie, R.M., Bralower, T.J., and Cashman, R., 2002. Oceanic anoxic events and plankton evolution: Biotic response to tectonic forcing during the mid-Cretaceous. *Paleoceanogr.*, 17:13–13-29, doi: 10.1029/2001PA000623.
- Ogg, J.G., Ogg, G., and Gradstein, F.M., 2008. *The Concise Geologic Time Scale*: Cambridge (Cambridge University Press).
- Satolli, S., Besse, J., and Calamita, F., 2008. Paleomagnetism of Aptian-Albian sections from the Northern Apennines (Italy): Implications for the 150–100 Ma apparent polar wander of Adria and Africa. *Earth Planet. Sci. Lett.*, 276:115–128, doi:10.1016/j.epsl.2008.09.013.
- Schlanger, S.O., and Jenkyns, H.C., 1976. Cretaceous oceanic anoxic events: Causes and consequences. *Geol. Mijnbouw*, 55:179–184.
- Skelton, P.W., Spicer, R.A., Kelley, S.P., and Gilmour, I., 2003. *The Cretaceous World*: Cambridge (Cambridge University Press).
- Tiraboschi, D., Erba, E., and Jenkyns, H.C., 2009. Origin of rhythmic Albian black shales (Piobbico core, central Italy): Calcareous nannofossil quantitative and statistical analyses and paleoceanographic reconstructions. *Paleoceanogr.*, 24:PA2222, doi: 10.1029/2008PA001670.
- Tornaghi, M.E., Premoli Silva, I., and Ripepe, M., 1989. Lithostratigraphy and planktonic foraminiferal biostratigraphy of the Aptian-Albian “Scisti a Fucoidi” in the Piobbico core, Marche, Italy: Background for cyclostratigraphy. *Riv. Ital. Paleont. Strat.*, 95:223–264.
- Wang, C., Hu, X., Huang, Y., Scott, R.W., and Wagreich, M., 2009. Overview of Cretaceous Oceanic Red Beds (CORBs): A window on global oceanic and climate change. In Hu, X., Wang, C., Scott, W., Wagreich, M., and Jansa, L. (Eds.), *Oceanic Red Beds: Stratigraphy, Composition, Origins, and Paleoclimatographic and Paleoclimatic Significance*. Society for Sedimentary Geology, Special Publication, 91:13–33.

## Authors

**Rodolfo Coccioni**, Dipartimento di Scienze della Terra, della Vita e dell'Ambiente dell'Università degli Studi “Carlo Bo,” Campus Scientifico “E. Mattei,” Località Crocicchia, 61029 Urbino, Italy, Email: rodolfo.coccioni@uniurb.it.

**Luigi Jovane**, Departamento de Oceanografia Física, Instituto Oceanográfico, Universidade de São Paulo, 05508-900 São Paulo, Brazil.

**Giuseppe Bancalà**, Dipartimento di Scienze della Terra, della Vita e dell'Ambiente dell'Università degli Studi “Carlo Bo,” Campus Scientifico “E. Mattei,” Località Crocicchia, 61029 Urbino, Italy.

**Carla Bucci**, Dipartimento di Scienze della Terra, della Vita e dell'Ambiente dell'Università degli Studi “Carlo Bo,” Campus Scientifico “E. Mattei,” Località Crocicchia, 61029 Urbino, Italy.

**Gerson Fauth**, Centro de Ciências Exatas e Tecnológicas, Área de Conhecimento e Aplicação de Geociências, Universidade do Vale do Rio dos Sinos, São Leopoldo, Brazil.

**Fabrizio Frontalini**, Dipartimento di Scienze della Terra, della Vita e dell'Ambiente dell'Università degli Studi “Carlo Bo,” Campus Scientifico “E. Mattei,” Località Crocicchia, 61029 Urbino, Italy.

**Liliane Janikian**, Departamento de Geofísica, Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, 05508-090 São Paulo, Brazil.

**Jairo Savian**, Departamento de Geofísica, Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, 05508-090 São Paulo, Brazil.

**Renato Paes de Almeida**, Departamento de Geologia Sedimentar e Ambiental, Instituto de Geociências, Universidade de São Paulo, 05508-090 São Paulo, Brazil.

**Grasiane Luz Mathias**, Departamento de Geofísica, Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, 05508-090 São Paulo, Brazil.

**Ricardo Ivan Ferreira da Trindade**, Departamento de Geofísica, Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, 05508-090 São Paulo, Brazil.

## Photo Credits

Fig. 3:

Fig. 4: