

Nº 23

Group rings with units of bounded
exponent over the center

Sônia P. Coelho

Junho 1981

GROUP RINGS WITH UNITS OF BOUNDED EXPONENT OVER THE CENTER

Sônia P. Coelho

Instituto de Matemática e Estatística
Universidade de São Paulo - Brasil

Let KG be the group ring of a group G over a field K , and U its group of units. Given a group H , we shall denote by $\xi(H)$ the center of H and by $T(H)$ the set of all its torsion elements.

The following question appears in [5, p. 231]: When is $U^n \subset \xi(U)$, for some n ? It was considered by G. Cliff and S.K. Sehgal in [2], where G is assumed to be a solvable group. A complete answer at characteristic zero is given there. Also they obtain partial results at characteristic $p \neq 0$, with certain restrictions on the exponent n .

In this note, we shall answer the question at characteristic p assuming that G is either a solvable or an FC-group. In fact, we shall need specially the following property which is common to both these families of groups: if H is a finitely generated subgroup of G such that $H/\xi(H)$ is torsion, then both $T(H)$ and H' , the derived group of H , are finite groups [1, Lemma 1.5, p. 116 and 2, Lemma 2.1, p. 147].

In § 1, we answer the question for torsion groups assuming only that G is locally finite (Theorem A), and in § 3 we give the answer for non torsion groups that are either solvable or FC (Theorem C).

First, we introduce some notation. We will denote $T(G)$ simply by T , and the integer $p \neq 0$ will always denote the characteristic of K . For an element t in a group, we shall say that t is a p -element if $o(t)$, the order of t , is a power of p , and that t is a p' -element if $o(t)$ is not divisible by p . Similarly, a group H will be called a p' -group if every element of H is a p' -element.

§ 1. The torsion subgroup of G

LEMMA 1.1: If $U^n \subseteq \xi(U)$ for some n and G has a non central p' -element, then K is finite and the orders of the p' -elements of G are bounded.

PROOF: We shall show first that the orders of the p' -elements of G are bounded.

It is enough to show that if u is a central p' -element of G , then $o(u) \leq n$. If not, take such an u , with $o(u) > n$. Then $K\langle u \rangle = \bigoplus_i K_i$, a direct sum of fields. For every i , denote by $\pi_i: K\langle u \rangle \rightarrow K_i$ the natural projection (that is, if e_i is the unity element of K_i , then $\pi_i(u) = (u e_i)$). Clearly, at least one of the $\pi_i(u)$, say $\pi_1(u)$, has multiplicative order equal to $o(u)$. As a consequence, K_1 has more than n elements.

Now, if t is a non central p' -element of G , we consider $K\langle u, t \rangle = \bigoplus_i K_i[t]$, where $K_i[t]$ denotes the smallest subalgebra of KG that contains K_i and t .

We claim that $K_1[t]$ is not contained in the center of KG . In fact, suppose that $K_1[t]$ is central and

let $e = e_1$ be the unity element of K_1 . Then et is central. Now take $x \in G$ such that $xtx^{-1} \neq t$. Then, $xetx^{-1} = extx^{-1} = et$.

By considering supports in the last equality we get: $u^i xtx^{-1} = u^j t$, $0 \leq i, j < o(u)$, $i \neq j$. Hence $xtx^{-1} = u^{j-i}t$.

Again, in the last equality, we obtain: $eu^{j-i}t = et$, or $eu^{j-i} = e$. Then, $(eu)^{j-i} = e$. But $eu = \pi_1(u)$, and hence the multiplicative order of $\pi_1(u)$ divides $|j-i| < o(u)$, a contradiction.

Now, since t is a p' -element, we have: $K_1[t] = \bigoplus_i L_i$, a direct sum of fields which are Galois extensions of K_1 . But $K_1[t]$ is not central, hence some of the L_i , say L_1 , is not contained in the center of KG . Let \bar{L}_1 be the subfield of L_1 consisting of its central elements, and let $\phi \neq 1$ be an \bar{L}_1 -automorphism of L_1 .

Since $L_1 = K(\zeta)$, with $\zeta^{o(t)} = 1$, we have that $\phi(\zeta) = \zeta^i$, for some i .

Now, take an arbitrary element $k \in \bar{L}_1$. Since $U^n \subset \xi(U)$, we get that $(\zeta+k)^n \in \bar{L}_1$. Then, $(\phi(\zeta+k))^n = \phi((\zeta+k)^n) = (\zeta+k)^n$, and $\phi(\zeta+k)$ is a root of $X^n - (\zeta+k)^n$; from this we see that $\phi(\zeta+k) = \alpha(\zeta+k)$, $\alpha^n = 1$, $\alpha \neq 1$.

On the other hand, $\phi(\zeta+k) = \zeta^i + k$, and thus $\zeta^i + k = \alpha(\zeta+k)$.

Solving this equation for k , we have that $k = \frac{\zeta^i - \alpha k}{\alpha - 1}$. Here, only α depends on k . Since α can take at most $n-1$ values, we see that $|L_1|$, the number of elements

of L_1 , is at most $n-1$. But $\bar{L}_1 \supset K_1$, and $|K_1| > n$, a contradiction.

It still remains to prove that K is finite. If not, replace K_1 by K in the proof above. Again, we have that $K_1[t] = K[t]$ is not central, and we can repeat the argument to obtain a contradiction. \blacktriangle

LEMMA 1.2: Assume that $U^n \subset \xi(U)$ for some n . Then, there exists a positive integer m , which is a power of p , such that x^m is central in KG , for every nilpotent element x in KG .

PROOF: Let $x \in KG$ be a nilpotent element, and let r be such that $x^{p^r} = 0$. Then, $1+x$ is a p -element of U and by hypothesis $(1+x)^n \in \xi(U)$. Writing $n = p^t n'$, with $(n', p) = 1$, it is easy to see that x^{p^t} is central in KG . \blacktriangle

LEMMA 1.3: Assume that $U^n \subset \xi(U)$ for some n , and let $n = p^t n'$, with $(n', p) = 1$. If G has a p -element of order greater than $2p^{3t}$, then $G^{p^t} \subset \xi(G)$.

PROOF: From the proof of Lemma 1.2, we see that x^{p^t} is central for every nilpotent element $x \in KG$. So, set $m = p^t$, and take a p -element $h \in G$ such that $o(h) > 2m^3$. Since $h-1$ is nilpotent, we have that $(h-1)^m = h^m - 1$ is central, hence h^m is central.

Set $h' = h^m$, take $x, y \in G$ and consider the nilpotent element $y(h'-1)$. Again, by Lemma 1.2, we have that $(y(h'-1))^m = y^m(h'^m - 1)$ is central. Hence:

$$xy^m(h'^m - 1) = y^m(h'^m - 1)x,$$

$$xy^m h'^m - xy^m = y^m h'^m x - y^m x.$$

Since $o(h') > 2m^2$, we know that $h'^{m-1} \neq 0$ and hence we have two elements of G in the support of the above element. If $p \neq 2$, we see immediately that $xy^m = y^m x$, thus $y^m \in \xi(G)$. If $p = 2$, we may have:

$$xy^m = y^m h'^m x,$$

$$xy^m h'^m = y^m x.$$

Using the fact that h'^m is central and replacing $y^m x$ in the first equation by its value in the second one, we get that $xy^m = xy^m h'^m h'^m$, or $(h')^{2m} = 1$, which contradicts the fact that $o(h') > 2m^2$. \blacktriangle

LEMMA 1.4: Let m be a power of p . If $G^m \subset \xi(G)$ and G contains a normal p -abelian subgroup ϕ such that G/ϕ is a finite p -group, then G is nilpotent.

PROOF: It follows as in [5, 6.6, pp. 157-158]. \blacktriangle

LEMMA 1.5: Let S be a commutative ring with identity, I a nil ideal of S , of bounded exponent, and Q a finite group. Let $S(Q, \rho, \sigma)$ be a crossed product of Q over S , with an arbitrary factor system ρ and σ such that $\sigma_t(I) \subset I$, for every $t \in Q$. Then, IQ is a nil ideal of $S(Q, \rho, \sigma)$, of bounded exponent.

PROOF: It is immediate to verify that IQ is an ideal of $S(Q, \rho, \sigma)$.

Let $Q = \{t_1, t_2, \dots, t_n\}$ and choose m such that $s^m = 0$, for every $s \in I$.

Take $r > m(n+1)^2$ and $x = s_1 t_1 + \dots + s_n t_n$ an arbitrary element of IQ , $s_i \in I$, $1 \leq i \leq n$.

We want to prove that $x^r = 0$. It is enough to show that any product of r elements from the set $\{s_1 t_1, \dots, s_n t_n\}$ is zero. Then, let $y = s_{i_1} t_{i_1} \dots s_{i_r} t_{i_r}$ be such a product.

It is easy to see that there exists an index j such that $s_{i_j} t_{i_j}$ occurs in y k times, with $k > m(n+1)$, and we may suppose without loss of generality that $s_{i_j} t_{i_j} = s_1 t_1$.

As the products $s_i t_i s_j t_j$ still have the form st , $s \in I$, $t \in Q$, and $ts_1 = \sigma_t(s_1)t$, for every $\sigma \in T$, we can write y in the form $y = \left(\prod_{i=1}^k z_i \right) \gamma$, with $z_i \in \{\sigma_t(s_1) | t \in Q\} \cup \{s_1\}$, $\gamma \in IQ$. Since the above set has at most $n+1$ elements and $k > m(n+1)$, there must exist an index j such that z_j occurs in y more than m times. Now, $z_j \in I$, therefore $z_j^m = 0$, and hence $y = 0$. \blacktriangle

LEMMA 1.6: Let $G = T$, a locally finite group. If $U^n \subseteq \xi(U)$ for some n , then KT satisfies a polynomial identity.

PROOF: Let m be as in Lemma 1.2, and consider $2m$ arbitrary elements of KT , say x_1, x_2, \dots, x_{2m} . By considering the subgroup generated by the supports of these elements, we may suppose that T is finite.

Denote by $J(KT)$ the Jacobson radical of KT . Then $KT/J(KT) = \bigoplus_i M_{n_i}(D_i)$, a direct sum of full matrix rings over division rings D_i .

Set x' for the image of an element $x \in KT$ through the natural epimorphism $KT \rightarrow KT/J(KT)$. For a given index i , take x_i an arbitrary nilpotent element in $M_{n_i}(D_i)$,

and choose any element $y_i \in KT$ such that $(y_i)' = x_i$. Then y_i is nilpotent, provided $J(KT)$ is nilpotent because T is finite. By Lemma 1.2, y_i^m is central in KT . Hence $x_i^m = (y_i^m)'$ is a central nilpotent element of $KT/J(KT)$, so it must be zero.

Now it is easy to see that the size of the matrices is bounded by m , that is, $n_i \leq m$, for every i .

On the other hand, given i and $d_i \neq 0$ in D_i , we can choose $u \in U$ such that $u' = d_i$ (see [5, Lemma 3.3, p. 179]). As $u^n \in \xi(U)$, d_i^n is central in D_i , and hence D_i is a field, by [3, Theorem 3.22, p.79].

Therefore, $KT/J(KT)$ satisfies $S_{2m}(X_1, X_2, \dots, X_{2m})$, the standard polynomial of degree $2m$ in the non commuting variables X_1, X_2, \dots, X_{2m} . Again, since $J(KT)$ is nilpotent, we can use Lemma 1.2 to obtain, for every $z \in KT$:

$$(S_{2m}(x_1, \dots, x_{2m}))^m z = z (S_{2m}(x_1, \dots, x_{2m}))^m. \quad \blacktriangle$$

We may now obtain a characterization of the fact of U being of bounded exponent over the center when G is a locally finite group.

Theorem A: Let $G = T$, a locally finite group. Then, $U^n \subset \xi(U)$ for some n if and only if the following conditions hold:

- (i) $T^\ell \subset \xi(U)$ for some ℓ .
- (ii) T contains a normal p -abelian subgroup of finite index.
- (iii) either every p' -element of T is central or T is of bounded exponent and K is finite.

PROOF: Suppose $U^n \subset \xi(U)$ for some n . Then (i) is trivial and (ii) follows from Lemma 1.6 and a Theorem by Passman [1, Corollary 3.10, p. 197].

To prove (iii), assume that not every p' -element of T is central. By Lemma 1.1, K is finite and $t^r = 1$, for a suitable r and for every p' -element $t \in T$. Now, if $n = p^t \cdot n'$, with $(n', p) = 1$, and T has a p -element of order greater than $2p^{3t}$, then $T^{p^t} \subset \xi(T)$ by Lemma 1.3, and hence every p' -element is central, a contradiction.

So, $t^s = 1$ for a suitable s and for every p -element $t \in T$.

Now take $x \in T$ and let T_0 be a normal p -abelian subgroup of index u , as in (ii). Then, $x^u \in T_0$, and we may write: $x^u = yz$, where $y, z \in T_0$, y is a p -element and z is a p' -element. Since T_0/T_0' is abelian, taking $(rs)^{\text{th}}$ -powers, we get: $x^{urs} \equiv y^{rs} z^{rs} \pmod{T_0'} \equiv 1 \pmod{T_0'}$. But T_0' is finite, so we have that $x^{urs|T_0'|} = 1$, and (iii) is proved.

Assume now that conditions (i), (ii) and (iii) hold, and let T_0 be a normal p -abelian subgroup of finite index in T , as in (ii), and A the set: $\{t \in T \mid t \text{ is a } p'\text{-element}\}$.

Suppose first that A is a central subgroup of T . Then, as T is locally finite, it is easy to see that $T = PxA$, where $P = \{t \in T \mid t \text{ is a } p\text{-element}\}$ is a subgroup of T .

Considering the subgroup $\phi = T_0 \cdot A$, it is easy to see that T satisfies the conditions of Lemma 1.4 and

hence it is nilpotent. Furthermore, T contains a normal p -abelian subgroup ϕ such that T/ϕ is a finite p -group, and we conclude from [5, Theorem 6.1, p. 155] that KT is Lie m -Engel for a suitable m . Hence, $U^n \subset \xi(U)$ for some n by [5, Lemma 4.3, p. 151].

Suppose now that A is a non central subset. By (iii), $T^S = 1$, for some s , and K is finite. Because T'_0 is a finite p -group, it is easy to see that $P_0 = \{t \in T_0 \mid t \text{ is a } p\text{-element}\}$ is a normal subgroup of T .

We claim that $\Delta(T, P_0)$, the kernel of the natural epimorphism $KT \rightarrow KT/P_0$, is nil of bounded exponent. Indeed, $\Delta(T_0/T'_0, P_0/T'_0)$ is nil of bounded exponent because T_0/T'_0 is abelian and P_0/T'_0 is of bounded exponent. Setting $S = KT_0/T'_0$, $Q = (T/T'_0)/(T_0/T'_0) \cong T/T_0$, we see that KT/T'_0 is the crossed product $S(Q, \rho, \sigma)$, with ρ and σ as usual. If $I = \Delta(T_0/T'_0, P_0/T'_0)$, by Lemma 1.5 we conclude that $IQ = \Delta(T/T'_0, P_0/T'_0)$ is nil of bounded exponent. Since T'_0 is a finite p -group, we see that $\Delta(T_0, T'_0)$ is nilpotent and hence it is easy to see that $\Delta(T, P_0)$ is nil of bounded exponent by considering the natural epimorphism $KT \rightarrow KT/T'_0$.

Now, T_0/P_0 is a normal subgroup of T/P_0 , of finite index, say, r . By [1, Lemma 1.10, p. 176], we get that $KT/P_0 \in M_r(KT_0/P_0)$.

Pick now $u \in U$. Considering the subgroup generated by the support of u , we may suppose that T is finite. Hence T_0/P_0 is a finite abelian p' -group, such that $(T_0/P_0)^S = 1$

Therefore, $KT_0/P_0 = \bigoplus_i K_i$, a direct sum of fields, all of them contained in $K(\zeta)$, with $\zeta^s=1$. Hence, $M_r(KT_0/P_0) = M_r(\bigoplus_i K_i) = \bigoplus_i M_r(K_i)$, and we have that $KT/P_0 \subset \bigoplus_i M_r(F_i)$, with $F_i = K(\zeta)$, for every i . Set S for $\bigoplus_i M_r(F_i)$ and u' for the image of u by the composition map of the natural epimorphism $f: KT \rightarrow KT/P_0$ followed by the inclusion $g: KT/P_0 \rightarrow S$. As K is finite, the group of non-singular matrices of $M_r(K(\zeta))$ is finite, say of order q , depending on r and s only. So, $u'^q = 1$ and we get: $u^q = 1 + \delta$, $\delta \in \Delta(T, P_0)$.

As $\Delta(T, P_0)$ is nil of bounded exponent, we can take m a power of p such that $x^m = 0$ for all $x \in \Delta(T, P_0)$. Now we can conclude that $u^{qm} = 1 + \delta^m = 1 \in \xi(U)$. \blacktriangle

COROLLARY; Let $G = T$, a locally finite group, and assume that the set of all p -elements of T is not of bounded exponent. Then the following conditions are equivalent:

- (i) $U^n \subset \xi(U)$ for some n .
- (ii) KT is Lie m -Engel for some m .

PROOF: First suppose that $U^n \subset \xi(U)$ for some n . By the preceding theorem, we get that (i), (ii) and (iii) hold.

Furthermore, every p' -element of T is central by Lemma 1.3. Follow now the "only if" - part of the proof of the theorem to conclude that KT is Lie m -Engel for some m .

By [5, Lemma 4.3, p. 151], the converse is obvious. \blacktriangle

§ 2. A certain nil ideal of KG

In this section, G will be either a solvable or an FC-group. As we mentioned in the introduction, if $G/\xi(G)$ is torsion, then we can conclude that T is a locally finite subgroup of G and G' is contained in T .

We shall denote by A the set of all p' -elements of G and by P the set of all p -elements of G .

LEMMA 2.1: Suppose that $U^n \leq \xi(U)$ for some n and G has an element of infinite order. Then, every idempotent of KG is central. ▲

PROOF: See [2, Lemma 2.4, p. 148].

LEMMA 2.2: Suppose that $U^n \leq \xi(U)$ for some n and G has an element of infinite order. Then:

- (i) A is an abelian subgroup of G .
- (ii) If A is non central, then K is finite and for every $x \in G$ and every $t \in A$ there exists an integer r such that $xtx^{-1} = t^{p^r}$, and $(K:\mathbb{F}_p) \mid r$.
- (iii) P is a subgroup of G .
- (IV) $T = PxA$.

PROOF: For the proof of (i) see [2, Corollary 2.5, p. 148].

To prove (ii) we notice that if A is non central, then K is finite by Lemma 1.1.

Now, take $x \in G$ and $t \in A$ such that $xtx^{-1} \neq t$. We have that $K \langle t \rangle = \bigoplus K_i$, a direct sum of fields such that at least one of them, say K_1 , is of the form $K_1 = K(\zeta)$, where

ζ is a root of unity whose order is equal to the order of t , and the natural projection $K\langle t \rangle \rightarrow K_1$ maps t on ζ .

Since, by Lemma 2.1, every idempotent is central, we must have $xtx^{-1} = t^i$ for some i (this can be seen by considering the idempotent $e = (o(t))^{-1} (1+t+\dots+t^{o(t)-1})$). Hence, conjugation by x defines an automorphism ϕ of K_1 . By the above, $\phi(\zeta) = \zeta^i$.

On the other hand, since K_1 is finite, ϕ is a power of the Frobenius automorphism of K_1 , F , given by: $F(k) = k^p$, for all $k \in K_1$. If $\phi = F^r$, we have that $\phi(\zeta) = \zeta^{p^r} = \zeta^i$, from which we conclude that $o(t) = o(\zeta)$ divides $p^r - i$. Then, $p^r \equiv i \pmod{o(t)}$, and $xtx^{-1} = t^i = t^{p^r}$.

Furthermore, as every element of K is fixed by ϕ , we have that $k^{p^r} = k$ for every $k \in K$, and hence K is contained in a field with p^r elements, that is: $(K:\mathbb{F}_p) \mid r$.

For (iii), we observe first that every p -element commutes with every p' -element. If not, then by Lemma 1.1 K is finite. Now take $\pi \in P$ and $t \in A$ such that $\pi t \pi^{-1} \neq t$ and proceed like in [2, 3.2, p. 152] to conclude that this implies the existence of a non central idempotent, which contradicts Lemma 2.1.

As T is locally finite, the proof of (iv) is now trivial. ▲

LEMMA 2.3: Let A_1 be an abelian p' -subgroup of G , and K a finite field. If, for every $t \in A_1$ and every $x \in G$, there exists an integer r such that $xtx^{-1} = t^{p^r}$, and $(K:\mathbb{F}_p) \mid r$, then every idempotent of KA_1 is central in KG .

PROOF: Let $e \in KA_1$ be an idempotent, and let $x \in G$. By considering the subgroup generated by the support of e , we may suppose that A_1 is finite.

Let $A_1 = \langle t_1 \rangle \times \dots \times \langle t_s \rangle$, a direct product of cyclic groups. It is easy to see that we may choose an integer r such that $xt_i x^{-1} = t_i^{p^r}$, for every i .

We have that $e = f(t_1, \dots, t_s)$, where $f(x_1, \dots, x_s)$ is a polynomial in the commuting variables x_1, \dots, x_s , with coefficients in K . Conjugating by x , we have: $x e x^{-1} = f(t_1^{p^r}, \dots, t_s^{p^r})$. But by hypothesis every element $k \in K$ satisfies $k^{p^r} = k$, therefore this is true for the coefficients of f . Hence $f(t_1^{p^r}, \dots, t_s^{p^r}) = (f(t_1, \dots, t_s))^{p^r}$ and $x e x^{-1} = e^{p^r} = e$, as we wished to prove. \blacktriangle

We can now give a partial characterization of the fact of U being of bounded exponent over the center, when G is non torsion.

Theorem B: Suppose that G has an element of infinite order. Then, $U^n \subset \xi(U)$ for some n if and only if the following conditions hold:

- (i) $G^l \subset \xi(G)$ for some l .
- (ii) A is an abelian subgroup of G and, if A is non central, then K is finite, A is of bounded exponent and for every $t \in A$ and every $x \in G$ there exists an integer r such that $xtx^{-1} = t^{p^r}$, where $(K: \mathbb{F}_p) \mid r$.
- (iii) P is a subgroup of G contained in the centralizer of A .
- (iv) There exists an integer m , which is a

power of p , such that x^m is central in KG , for every $x \in \Delta(G, P)$.

PROOF: Suppose first that $U^n \subset \xi(U)$ for some n . (i) is trivial, (ii) follows from Lemma 1.1 and Lemma 2.2, and (iii) follows from Lemma 2.2.

To prove (iv), let $x \in \Delta(G, P)$. We may suppose that G is finitely generated and hence P is a finite normal subgroup of G . Therefore, x is nilpotent and we can apply Lemma 1.2 to obtain the thesis.

Suppose now conditions (i) to (iv) hold, and pick $u \in U$. Again we may suppose that G is finitely generated and hence T is finite.

We observe that if $KA = \bigoplus_i K_i$, a direct sum of fields, then $K(AxP) = \bigoplus_i K_i P$.

Consider now the natural epimorphism $KG \rightarrow KG/P$, with kernel $\Delta(G, P)$.

Setting S' for the image of a subset S of KG through this epimorphism, we have: $(K(AxP))' = \bigoplus_i K_i'$ where each K_i' is a field. Furthermore, $T(G/P) = (AxP)/P$, and hence $KT(G/P) = \bigoplus_i K_i'$.

Since, by condition (ii) and Lemma 2.3, every idempotent of KA is central in KG , and $G' \subset T$, we may apply [5, Lemma 3.22, p. 194], and u can be written in the form:

$$(*) \quad u = \sum_i f_i g_i + \delta, \quad f_i \in K_i, \quad g_i \in G, \quad \delta \in \Delta(G, P).$$

Now we consider both cases separately.

Suppose first that A is central. Taking the

$(\ell)^{\text{th}}$ - power in (*), we have: $u^\ell = \sum_i f_i^\ell g_i^\ell + \delta'$, $\delta' \in \Delta(G, P)$.
 Now, $\sum_i f_i^\ell g_i^\ell$ is central, and it is sufficient to apply condition (iv).

Suppose now that K is finite and A is of bounded exponent s . Then, $K_i \subset K(\zeta)$, with $\zeta^s = 1$, for all i .

Computing $(\ell)^{\text{th}}$ - powers in (*), we obtain:
 $u^\ell = \sum_i f_i' g_i^\ell + \delta'$, $f_i' \in K_i$, $g_i^\ell \in \xi(G)$, $\delta' \in \Delta(G, P)$.

Since $K_i \subset K(\zeta)$, for all i , we have that $f_i'^r = 1$ for every $f_i' \neq 0$ and a suitable r which depends on K and s only.

Taking $(r)^{\text{th}}$ - powers above, we have: $u^{\ell r} = \sum_i g_i^{\ell r} + \delta''$, $\delta'' \in \Delta(G, P)$.

Now, $\sum_i g_i^{\ell r}$ is central and it suffices to use condition (iv). ▲

COROLLARY: Suppose that the set of all p -elements of G is not of bounded exponent. Then the following conditions are equivalent:

- (i) $U^n \subset \xi(U)$ for some n .
- (ii) KG is Lie ℓ -Engel for some ℓ .

PROOF: Suppose first $U^n \subset \xi(U)$ for some n . By the corollary to theorem A, KT is Lie m -Engel for some m and hence T is nilpotent.

As $G \subset T$, we can conclude that G is solvable (even if it were an FC-Group).

By Lemma 1.3, $G^{p^t} \subset \xi(G)$ for some t and by

[2, Lemma 2.2, p. 148] G' is a p -group. Hence, as we noted before, $\Delta(G, G')$ is a nil ideal.

Given $x, y \in KG$, $xy - yx \in \Delta(G, G')$, By (iv) of Theorem B, for every $z \in KG$ we have that $(xy - yx)^m z = z(xy - yx)^m$.

Hence, KG satisfies a polynomial identity. By Passman's theorem [4, Theorem 1.1, p. 142], setting ϕ for the FC-subgroup of G , we have: $|G/\phi| < \infty$, $|\phi'| < \infty$.

Furthermore, it is easy to see that both are p -groups. By Lemma 1.4, G is nilpotent. We conclude from [5, Theorem 6.1, p. 155] that KG is Lie ℓ -Engel for some ℓ .

As we noted before, the converse is immediate by [5, Lemma 4.3, p. 151]. ▲

§ 3. Augmentation ideals of bounded exponent

We shall now discuss condition (iv) of Theorem B.

In all this section, except in Theorem C, G will be either a solvable or an FC-group, such that T is a locally finite subgroup of G , and $G' \subset T$.

Furthermore, we shall assume that T has the form: $T = PxA$, where $A = \{t \in G \mid t \text{ is a } p'\text{-element}\}$ is an abelian subgroup of G and $P = \{t \in G \mid t \text{ is a } p\text{-element}\}$ is a subgroup of G , of bounded exponent.

We introduce some notation. Given a group H , $\phi(H)$ will denote the FC-subgroup of H . The group $\phi(G)$ will

be denoted simply by ϕ .

LEMMA 3.1: Suppose that there exists an integer m , which is a power of p , such that x^m is central in KG for all $x \in \Delta(G, P)$. Then G contains a normal subgroup H , of finite index, containing ϕ , such that $H \cap P$ is finite.

PROOF: By Passman [4, Theorem 1.1, p. 142], if KG satisfies a polynomial identity, then $|G/\phi| < \infty$, $|\phi'| < \infty$. Since $G' \subset PxA$, and $(\phi.A)/A \subset \phi(G/A)$, it is enough to prove that $K(G/A)$ satisfies a polynomial identity.

In the group ring $K(G/A)$, the ideal $\Delta(G/A, (PxA)/A)$ is the image of $\Delta(G, P)$ by the natural epimorphism $KG \rightarrow K(G/A)$ and hence it also satisfies the hypothesis.

As $(G/A)' \subset (PxA)/A$, we have that $\Delta(G/A, (G/A)') \subset \Delta(G/A, (PxA)/A)$ and it follows easily, as in the proof of the corollary to Theorem B, that $K(G/A)$ satisfies the polynomial $(XY-YZ)^m Z - Z(XY-YX)^m$. \blacktriangle

In the following lemmas, H will be a normal subgroup of G , of finite index, containing ϕ , as in the previous lemma, P_0 will be $H \cap P$ (hence a finite group) and P_1 will be $H \cap P$.

LEMMA 3.2: Suppose that there exists a positive integer m , which is a power of p , such that x^m is central in KG , for all x in $\Delta(G, P)$: Then, $\Delta(G, P)$ is nil of bounded exponent.

PROOF: We observe first that it is enough to prove that $\Delta(H, P_1)$ is nil of bounded exponent. In fact, suppose that this were proved. Given $x \in \Delta(G, P)$, then x^m is a central element. But every central element of $\Delta(G, P)$ is contained in $\Delta(G, P) \cap K\phi$, which is contained in $\Delta(H, P_1)$. Then $\Delta(G, P)$

is nil of bounded exponent.

Let us prove now that $\Delta(H, P_1)$ is nil of bounded exponent. As P_0 is finite, $\Delta(H, P_0)$ is nilpotent and hence it suffices to prove that $\Delta(H/P_0, P_1/P_0)$ is nil of bounded exponent.

We have that $(H/P_1)'$ is a p' -group, P_1/P_0 is a central subgroup of H/P_0 , and for all $x \in \Delta(H/P_0, P_1/P_0)$, x^m is central in KH/P_0 . Furthermore, all hypothesis on G carry on to H/P_0 . Therefore, in order to prove that $\Delta(H/P_0, P_1/P_0)$ is nil of bounded exponent, we may replace H/P_0 by H and P_1/P_0 by P_1 and assume in addition that H' is a p' -group and P_1 is central.

Now, we shall see which is the form of a central element of KH that belongs to $\Delta(H, P_1)$.

Let S' be a transversal of $T(H)$ in H . Observing that $T(H) = P_1 x A_1$, where $A_1 = \{t \in H \mid t \text{ is a } p'\text{-element}\}$, it is easy to see that $S = \{ab \mid a \in A_1, b \in S'\}$ is a transversal of P_1 in H .

Now, let $x \in \phi(H)$, $y \in H$, $x = hab$, $h \in P_1$, $ab \in S$. Since $y(ab)y^{-1}(ab)^{-1} \in H' \subset A_1$ and P_1 is central, xyx^{-1} has the form: $xyx^{-1} = ha'b$, $a' \in A_1$. So denoting by $\gamma(x)$ the sum of the elements of the conjugacy class of x , we have that $\gamma(hab) = h(a_1 + \dots + a_r)b$, $a_j \in A_1$, $1 \leq j \leq r$. Therefore, we may write every central element $z \in KH$ in the form:

$$(1) \quad z = \sum_i \alpha_i b_i,$$

where $\alpha_i \in KT(H)$, $\alpha_i b_i$ is central in KH for all i , and the b_i are distinct elements of S' .

We claim that $\alpha_i b_i = b_i \alpha_i$, for all i . In fact, as $\alpha_i b_i$ is central, we have that $\alpha_i b_i = b_i (\alpha_i b_i) b_i^{-1} = b_i \alpha_i$.

If in addition $z \in \Delta(H, P_1)$, then $\alpha_i \in \Delta(T(H), P_1)$, for all i . In fact, denote by x' the image of an element $x \in KH$ by the natural epimorphism $KH \rightarrow KH/P_1$. Since $z \in \Delta(H, P_1)$, $z' = \sum_i \alpha_i' b_i' = 0$. But the b_i' are distinct elements of a basis of the $K(T(H)/P_1)$ -module KH/P_1 . Hence $\alpha_i' = 0$, for all i , which means that $\alpha_i \in \Delta(H, P_1) \cap KT(H) = \Delta(T(H), P_1)$.

Therefore every central element z of $\Delta(H, P_1)$ has the form (1), and furthermore $\alpha_i \in \Delta(T(H), P_1)$, for all i .

Call s the exponent of P_1 . Since $T(H)$ is abelian, computing the $(s)^{\text{th}}$ -power of z in (1) we have that:

$$z^s = \sum_i (\alpha_i b_i)^s = \sum_i \alpha_i^s b_i^s = 0.$$

We have proved that $z^s = 0$, for a fixed s and every central element z of $\Delta(H, P_1)$. As every $x \in \Delta(H, P_1)$ is such that x^m is central, we can conclude that $\Delta(H, P_1)$ is nil of bounded exponent. \blacktriangle

LEMMA 3.3: If $\Delta(H, P_1)$ is nil of bounded exponent, then either P_1 is finite or H contains a characteristic p -abelian subgroup of finite index.

PROOF: Suppose that P_1 is infinite. By Passman [1, Corollary 3.10, p. 197], it is enough to prove that KH satisfies a polynomial identity. Since $\Delta(H, P_0)$ is nilpotent it

suffices to prove that KH/P_0 satisfies a polynomial identity.

We observe that all the hypothesis of the lemma carry on to KH/P_0 . In fact, $\Delta(H/P_0, P_1/P_0)$ is nil of bounded exponent and P_1/P_0 is infinite (because P_0 is finite). Furthermore, $(H/P_0)'$ is a p' -group and P_1/P_0 is central in H/P_0 .

Therefore, to prove that KH/P_0 satisfies a polynomial identity we may replace H/P_0 by H and assume in addition that H' is a p' -group and P_1 is a central subgroup of H .

Now, pick $n > 0$ such that $x^n = 0$ for all $x \in \Delta(H, P_1)$. We want to show that there exist elements x_1, x_2, \dots, x_n in $\Delta(P_1) = \Delta(P_1, P_1)$ such that $x_i^2 = 0$, for all i , but $x_1 x_2 \dots x_n \neq 0$. In fact, as P_1 is abelian of bounded exponent, it is a direct product of cyclic groups [4, Theorem 11.2, p. 44]. P_1 is infinite and hence the number of cyclic groups is infinite. Choose h_1, \dots, h_n generators in different cyclic subgroups and set $m_i = o(h_i) - 1$. The elements $x_i = (h_i - 1)^{m_i}$ are easily verified to satisfy the required conditions.

Take now $S = \{g_i\}_{i \in I}$ a transversal of P_1 in H , and elements $g_1, \dots, g_n \in S$. Then, the element $\alpha = g_1 x_1 + \dots + g_n x_n$ belongs to $\Delta(H, P)$, hence $\alpha^n = 0$.

Since the x_i are central and $x_i^2 = 0$, computing α^n we get: $\alpha^n = F(g_1, \dots, g_n) x_1 x_2 \dots x_n$, where $F(X_1, \dots, X_n)$ is a polynomial in the non-commuting variables X_1, X_2, \dots, X_n , namely: $F(X_1, \dots, X_n) = \sum X_{\sigma(1)} \dots X_{\sigma(n)}$, the sum running over

all $\sigma \in S_n$, the symmetric group on n elements.

We want to show that $F(g_1, \dots, g_n) = 0$.

First we make some observations. If

$g_{\sigma(1)} g_{\sigma(2)} \cdots g_{\sigma(n)} \equiv g_{\tau(1)} g_{\tau(2)} \cdots g_{\tau(n)} \pmod{P_1}$ for $\sigma, \tau \in S_n$, then $g_{\sigma(1)} g_{\sigma(2)} \cdots g_{\sigma(n)} = a g_{\tau(1)} g_{\tau(2)} \cdots g_{\tau(n)}$, for some $a \in P_1$.

On the other other hand, since $g_i g_j = \alpha(i, j) g_j g_i$, where $\alpha(i, j) \in H'$, we can write the product $g_{\sigma(1)} g_{\sigma(2)} \cdots g_{\sigma(n)}$ in the form $b g_{\tau(1)} g_{\tau(2)} \cdots g_{\tau(n)}$ for some $b \in H'$.

Therefore, we obtain: $a = b$, with $a \in P_1$ and $b \in H'$, which is a p' -group. Thus $a = b = 1$.

We have shown that $g_{\sigma(1)} g_{\sigma(2)} \cdots g_{\sigma(n)} \equiv g_{\tau(1)} g_{\tau(2)} \cdots g_{\tau(n)} \pmod{P_1}$ if and only if $g_{\sigma(1)} g_{\sigma(2)} \cdots g_{\sigma(n)} = g_{\tau(1)} g_{\tau(2)} \cdots g_{\tau(n)}$.

We now define an equivalent relation \sim in S_n setting: for $\sigma, \tau \in S_n$, $\sigma \sim \tau$ if and only if $g_{\sigma(1)} \cdots g_{\sigma(n)} = g_{\tau(1)} \cdots g_{\tau(n)}$. We denote by S_1, S_2, \dots, S_t the equivalence classes of this relation. Choosing, for each i , an element $\sigma_i \in S_i$, it follows from the above that we may write:

$$F(g_1, \dots, g_n) = g_{\sigma_1(1)} \cdots g_{\sigma_1(n)} |S_1| + \cdots + g_{\sigma_t(1)} \cdots g_{\sigma_t(n)} |S_t|.$$

Now, since $g_{\sigma_i(1)} \cdots g_{\sigma_i(n)} \not\equiv g_{\sigma_j(1)} \cdots g_{\sigma_j(n)} \pmod{P_1}$ if $i \neq j$, from $F(g_1, \dots, g_n) x_1 x_2 \cdots x_n = 0$, we get: $g_{\sigma_i(1)} \cdots g_{\sigma_i(n)} |S_i| x_1 x_2 \cdots x_n = 0$, for all i .

But $g_{\sigma_i(1)} \cdots g_{\sigma_i(n)}$ is invertible and $x_1 x_2 \cdots x_n \neq 0$; hence this can happen only if $|S_i| = 0$, for all i (that is, $p \mid |S_i|$, for all i).

Therefore, $F(g_1, \dots, g_n) = 0$, for arbitrary elements $g_1, \dots, g_n \in S$.

Now, KH is a left module over the central subalgebra KP having S as a basis, and $F(X_1, \dots, X_n)$ is a multilinear polynomial. By [1, Lemma 1.2, p. 171], F is a polynomial identity for KH . \blacktriangle

COROLLARY: If $\Delta(G, P)$ is nil of bounded exponent, then either P is finite or G contains a normal p -abelian subgroup of finite index.

PROOF: Suppose that P is infinite, and take H as in Lemma 3.1. Since $|G/H| < \infty$, we have that $|PH/H| < \infty$ and hence $|P/P_1| < \infty$. Therefore, P_1 must be infinite. By Lemma 3.3, H contains a characteristic p -abelian subgroup of finite index, and thus the thesis follows. \blacktriangle

LEMMA 3.4: If G contains a normal p -abelian subgroup of finite index, then $\Delta(G, P)$ is nil of bounded exponent.

PROOF: Let L be such a subgroup of G . We have that L/L' is abelian; hence $\Delta(L/L', PL/L')$ is nil of bounded exponent.

Setting $S = KL/L'$, $Q = (G/L')/(L/L') \cong G/L$, we see that $K(G/L')$ is the crossed product $S(Q, \rho, \sigma)$, with ρ and σ as usual.

If $I = \Delta(L/L', PL/L')$, applying Lemma 1.5 we conclude that $IQ = \Delta(G/L', (P \cap L)/L')$ is nil of bounded exponent. Since L' is a finite p -group, $\Delta(G, L')$ is nilpotent and hence $\Delta(G, P \cap L)$ is nil of bounded exponent.

Now, let us consider the natural epimorphism $\phi: KG \rightarrow K(G/(P \cap L))$. We have that $\phi(\Delta(G, P)) = \Delta(G/(P \cap L), P/(P \cap L))$. But $P/(P \cap L) \cong PL/L$, and PL/L is a finite p -group since it is contained in G/L . Therefore, there exists an integer n such that $\Delta(G, P)^n \subset \Delta(G, P \cap L)$. Since we have shown that this ideal is nil of bounded exponent, the Lemma is proved. \blacktriangle

We can now give a complete answer to the initial question.

Theorem C: Suppose that G is non-torsion and either solvable or FC. Then, $U^n \subset \xi(U)$ for some n if and only if either KG is Lie m -Engel for some m or the following conditions hold:

(i) $G^l \subset \xi(G)$ for some l .

(ii) A is an abelian subgroup of G and, if A is non central, then K is finite, A is of bounded exponent and for every $x \in G$ and every $t \in A$ there exists an integer r such that $xtx^{-1} = t^{p^r}$, where $(K: \mathbb{F}_p) \mid r$.

(iii) P is a subgroup of G of bounded exponent, contained in the centralizer of A and, if P is not finite, then G contains a normal p -abelian subgroup of finite index.

PROOF: Suppose that $U^n \subset \xi(U)$ for some n and that KG is not Lie m -Engel.

By theorem B, the conditions (i) and (ii) hold, and by (iii) of theorem B, P is a subgroup of G contained in the centralizer of A . If P is not of bounded ex-

ponent, by the corollary to Theorem B, KG is Lie m -Engel, for some m , a contradiction. Hence, P is of bounded exponent.

Also, since condition (iv) of theorem B holds, we have that $\Delta(G, P)$ is nil of bounded exponent by lemma 3.2. By the corollary to lemma 3.3, the remainder of condition (iii) holds.

Suppose now KG is Lie m -Engel for some m . Then, as we have noted before, $U^n \subset \xi(U)$ for a suitable n [see 5, Lemma 4.3, p. 151].

Finally, suppose that conditions (i), (ii) and (iii) hold. If P is finite, then $\Delta(G, P)$ is nilpotent and condition (iv) of theorem B holds. If G contains a normal p -abelian subgroup of finite index, then using lemma 3.4, again condition (iv) of theorem B holds. Since conditions (i), (ii) and (iii) of this theorem are verified, then there exists an n such that $U^n \subset \xi(U)$. ▲

The author is grateful to Prof. S.K. Sehgal for suggesting this problem.

References

- [1] D.S. Passman, The Algebraic Structure of Group Rings, Wiley-Interscience, New York, 1977.
- [2] G.H. Cliff and S.K. Sehgal, Group rings with torsion units over the center, Manuscripta Math. 33, 145-158 (1980).
- [3] I.N. Herstein, Non Commutative Rings, Mathematical Association of America, United States of America, 1973.
- [4] L. Fuchs, Abelian Groups, Hungary, 1960.
- [5] S.K. Sehgal, Topics in Group Rings, Pure and applied mathematics, v.50, Dekker, New York, 1978.

TRABALHOS

DO

DEPARTAMENTO DE MATEMATICA

TITULOS PUBLICADOS

- 8001 - PLETCH, A. Local freeness of profinite groups. São Paulo, IME-USP, 1980. 10p.
- 8002 - PLETCH, A. Strong completeness in profinite groups. São Paulo, IME-USP, 1980. 8p.
- 8003 - CARNIELLI, W.A. & ALCANTARA, L.P. de. Transfinite induction on ordinal configurations. São Paulo, IME-USP, 1980. 22p.
- 8004 - JONES RODRIGUES, A.R. Integral representations of cyclic p-groups. São Paulo, IME-USP, 1980. 13p.
- 8005 - CORRADA, M. & ALCANTARA, L.P. de. Notes on many-sorted systems. São Paulo, IME-USP, 1980. /25/p.
- 8006 - POLCINO MILIES, F.C. & SEHGAL, S.K. FC-elements in a group ring. São Paulo, IME-USP, 1980. /10/p.
- 8007 - CHEN, C.C. On the Ricci condition and minimal surfaces with constantly curved Gauss map. São Paulo, IME-USP, 1980. 10p.
- 8008 - CHEN, C.C. Total curvature and topological structure of complete minimal surfaces. São Paulo, IME-USP, 1980. 21p.
- 8009 - CHEN, C.C. On the image of the generalized Gauss map of a complete minimal surface in R^4 . São Paulo, IME-USP, 1980. 8p.
- 8110 - JONES RODRIGUES, A.R. Units of ZCp^n . São Paulo, IME-USP, 1981. 7p.
- 8111 - KOTAS, J. & COSTA, N.C.A. da. Problems of modal and discussive logics. São Paulo, IME-USP, 1981. 35p.

- 8112 - BRITO, F.B. & GONÇALVES, D.L. Algebras não associativas, sistemas diferenciais polinomiais homogêneos e classes características. São Paulo, IME-USP, 1981. 7p.
- 8113 - POLCINO MILIES, F.C. Group rings whose torsion units form a subgroup II. São Paulo, IME-USP, 1981. 1v. (não paginado)
- 8114 - CHEN, C.C. An elementary proof of Calabi's theorems on holomorphic curves. São Paulo, IME-USP, 1981. 5p.
- 8115 - COSTA, N.C.A. da & ALVES, E.H. Relations between paraconsistent logic and many-valued logic. São Paulo, IME-USP, 1981. 8p.
- 8116 - CASTILHA, M.S.A.C. On Przymusiński's theorem. São Paulo, IME-USP, 1981. 6p.
- 8117 - CHEN, C.C. & GOFS, C.C. Degenerate minimal surfaces in R^4 . São Paulo, IME-USP, 1981. 2p.
- 8118 - CASTILHA, M.S.A.C. Imagens inversas de algumas aplicações fechadas. São Paulo, IME-USP, 1981. 11p.
- 8119 - ARAGONA VALEJO, A.J. & EXEL FILHO, P. An infinite dimensional version of Hartogs' extension theorem. São Paulo, IME-USP, 1981. 9p.
- 8120 - GONÇALVES, J.Z. Groups rings with solvable unit groups. São Paulo, IME-USP, 1981. 15p.
- 8121 - CARNIELLI, W.A. & ALCANTARA, L.P. de. Paraconsistent algebras. São Paulo, IME-USP, 1981. 16p.
- 8122 - GONÇALVES, D.L. Nilpotent actions. São Paulo, IME-USP, 1981. 10p.