

Superselective transcatheter arterial embolization of iatrogenic inferior epigastric artery after paracentesis: Unusual manifestation of hemoperitoneum

Tulio Fabiano de Oliveira Leite

Medical School of Ribeirão Preto, University of São Paulo, Rua Rafael Rinaldi, 365, APT 601, Uberlândia, Minas Gerais, 38400.384, Brazil

ARTICLE INFO

Article history:

Received 16 June 2020

Received in revised form 7 July 2020

Accepted 8 July 2020

Available online 2 August 2020

Keywords:

Acute hemorrhage

Inferior epigastric artery

Iatrogenic injury

Transcatheter arterial embolization

Paracentesis

ABSTRACT

INTRODUCTION: Paracentesis is a safe procedure and can be performed as a therapy or diagnosis in cancer patients, liver cirrhosis, heart or liver failure.

PRESENTATION OF CASE: 59-year-old man with alcoholic liver cirrhosis with ascites and coagulation abnormalities. After diagnostic paracentesis he presented hemodynamic instability with signs of hypovolemic shock and hemoperitoneum. Computed angiography with signs of active bleeding and pseudoaneurysm at the site of paracentesis.

DISCUSSION: The interventional radiology unit was referred and submitted to arteriography, which demonstrated active bleeding from the left lower epigastric artery. It was successfully treated by transcatheter embolization with 100–300 μ m PVA particles.

CONCLUSION: Transcatheter embolization with PVA particles is a fast, safe, minimally invasive, reliable method with a high technical success rate for the treatment of active bleeding resulting from injury to the lower epigastric artery.

© 2020 Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The lower left quadrant of the abdomen is the most commonly used site for paracentesis [1]. Therapeutic paracentesis is considered safe procedure and is performed commonly for the control of massive ascites in patients with liver cirrhosis with a 1% overall complication rate [2]. Hepatopathies may present collateral circulation and coagulopathy, that increase the chance of bleeding during paracentesis. Acute hematoma in the anterior abdominal wall due to hemorrhage of the inferior epigastric artery (IEA) is an uncommon cause of abdominal pain and frequently self-limiting. The fail in the diagnostic can lead to life-threatening complications in patients with comorbidities [3]. The objective of the case report is to describe an unusual event of hemoperitoneum and the efficacy in transcatheter embolization for bleeding from the IEA. Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

2. Case report

Patient 59 years old man, cirrhotic by alcohol CHILD B9 and MELD 17 with regular general condition, ascites, pale, tachycardic, tachydyspneic, jaundice, afebrile, with worsening renal function. Laboratory findings showed white blood cells 4730, Hemoglobin 10 g / dL, platelets 83,000 / mm³, INR (international normalized ratio) 1.86; Na + 135 mEq / L; K + 4.4mEq / L; aspartate aminotransferase 69 IU / L; total bilirubin 9.6 mg / dL; direct bilirubin 3.6 mg / dL; albumin 2.5 g / dL; urea 30 mg / dL. A physician performed a left lower quadrant therapeutic paracentesis with an 18 gauge needle without image guidance. Manual compression was applied to the puncture site. After paracentesis, the patient complains of worsening abdominal pain and left wall hematoma associated with hypotension and worsening tachycardia. Contrast-enhanced abdominopelvic computed tomography scan was carried and revealed extravasation of the contrast from the IEA. (Fig. 1: A and B) Patient referred to the interventional radiology unit and angiography was performed by puncturing the common right femoral artery. Pelvic arteriography was performed and bleeding was identified from the left lower epigastric artery. For microcatheterization of the lower epigastric artery was used catheter Progret 2.8 (Terumo) and embolization was performed with PVA 100–300 μ m. (Fig. 2: A and B) The endpoints of the embolization was considered complete when no further extravasation of contrast after embolization of the IEA. (Fig. 3) A successful procedural outcome was defined by the return

E-mail address: tuliofabiano@hotmail.com




Fig. 1. A: Contrast-enhanced abdominal CT scan obtained after paracentesis reveals ascites and focus of extravasation of the contrast medium in the left lower. (yellow arrow) B: Reconstruction of abdominal computed tomography angiography with a focus on extravasation of the contrast medium in the lower left. (yellow arrow).




Fig. 2. A: Angiography carried out after selective catheterization with a microcatheter of the left inferior epigastric artery shows active hemorrhage with extravasation of contrast. (yellow arrow) B: Angiography carried out after super selective catheterization with a microcatheter of the left inferior epigastric artery with active hemorrhage. (yellow arrow).

of hemodynamic stability and the stabilization of hematocrit. The work was reported according to the Consensus Surgical Case REport (SCARE) Guidelines, International Journal of Surgery 2018 [4].

3. Discussion

The treatment of ascites in liver disease is clinical with diuretics and salt restriction. Paracentesis is a treatment option in advanced stage of cirrhosis, in general is a safe, minimally invasive, especially when performed using sonographic guidance. The lower left quadrant of the abdomen, 4–5 cm medial to the upper anterior iliac spine, is the most common site for the procedure⁵. As a result of its anatomy, the IEA may potentially be traumatized during percutaneous procedures in until 40 %.

Besides, there are others etiologies of injury to the IEA, including inadvertent puncture during common femoral artery catheterization, placement of peritoneal dialysis catheters, surgical trauma including the use of sutures in the abdominal wall (15 %), blunt trauma (10 %), subcutaneous injection (5%), stabbing (5%) [6,7].

Patients with a normal coagulogram rarely experience complications such as bleeding and when the endovascular technique occurs it can be a treatment option. The use of ultrasound is able to reduce adverse events during paracentesis, for example, (1.4 % vs 4.7 %, $p < 0.0004$), post-paracentesis infection (0.41 % vs 2.44 %, $p < 0.001$), hematoma (0.0 % vs 0.87 %, $p < 0.01$), and seroma (0.14 % vs 1.05 %, $p < 0.03$) [8]. Lesion of the lower epigastric artery usually appears as a hematoma in the abdominal wall, abdominal pain and a drop in hematimetry.

Diagnosis can be based on the combination of patient history, physical examination, ultrasonography, and contrast-enhanced computed tomography Computed tomography (CT) with contrast has been reported to be an accurate, noninvasive technique for identifying arterial hemorrhage in patients with blunt abdominal or pelvic trauma with sensitivity and specificity of the 70 % and 100 %, respectively [9]. Angiotomography may be useful to lateralize a suspected IEA before angiography. The presence of the extravasation of contrast material on contrast-enhanced CT can suggest the presence of arterial injury that will require treatment. For patients




Fig. 3. Angiogram shows complete occlusion of the left inferior epigastric artery after embolization with PVA 100–300 μm without extravasation of contrast. (red arrow).

with chronic renal failure with high levels of creatinine or who are allergic to contrast, Doppler ultrasound may be an alternative in the diagnosis of active bleeding.

There are various options of treatment of injury IEA such as surgery, transcatheter embolization, ultrasound-guided thrombin injection and ultrasound-guided compression [5].

Lesions of the lower epigastric artery, endovascular techniques are preferable, especially in patients with portal hypertension, liver failure and coagulopathy. Our option was to perform embolization with PVA, as this technique has less morbidity and mortality and it would be possible to perform it with local anesthesia.

There are various endovascular therapeutic options including embolization with polyvinyl alcohol (PVA), gelfoam, coils, autologous coagulum, and n-butyl cyanoacrylate, Onix or associated combinations [10]. Our interventional radiology department has no standard protocol for embolization of active hemorrhages of IEA. The use of embolization materials at our department is still solely based on the interventional radiologist personal experience and individual case decision.

In conclusion, hemoperitoneum is a rare complication in paracentesis, which can be severe in patients with advanced liver cirrhosis. The use of ultrasound has been shown to reduce complications inherent to paracentesis. The endovascular approach with superselective embolization is a safe, effective and definitive procedure in lesions of the lower epigastric artery, especially in cirrhotic patients.

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request

Declaration of Competing Interest

The author declare no conflict of interest in preparing this article.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Ethical approval

Case reports are not required to be approved by the institution's ethics committee.

Consent

The document is attached.

Author contribution

TÚLIO FABIANO DE OLIVEIRA LEITE is a Vascular Surgeon from Faculdade Medicina de Botucatu (UNESP), Interventional Radiologist and Endovascular Surgeon from Universidade de São Paulo (USP). Research Member of the Research Group on Medical Sciences and Vascular Pathophysiology at Universidade Federal Fluminense (UFF / CAPES / CNPQ).

The study was carried out at the University of São Paulo.

- 1) Conception and design of the study: TFOL
- 2) Analysis and interpretation of data: TFOL
- 3) Data collection: TFOL
- 4) Writing of the article: TFOL
- 5) Critical revision of the text: TFOL
- 6) Final approval of the article *: TFOL
- 7) Statistical analysis: TFOL
- 8) General responsibility for the study: TFOL
- 9) Financing information: There was no financing

Registration of research studies

Not applicable.

Guarantor

Túlio Fabiano de Oliveira Leite.

Provenance and peer review

Not commissioned, externally peer-reviewed.

References

- [1] Y.J. Park, S.Y. Lee, S.H. Kim, I.H. Kim, S.W. Kim, S.O. Lee, Transcatheter coil embolization of the inferior epigastric artery in a huge abdominal wall hematoma caused by paracentesis in a patient with liver cirrhosis, *Korean J. Hepatol.* 17 (September(3)) (2011) 233.
- [2] P.R. Sobkin, A.I. Bloom, M.W. Wilson, J.M. LaBerge, G.S. Hastings, R.L. Gordon, et al., Massive abdominal wall hemorrhage from injury to the inferior epigastric artery: a retrospective review, *J. Vasc. Interv. Radiol.* 19 (March (3)) (2008) 327–332.
- [3] M.C. Wick, J. Klocker, C. Grundtman, W. Jaschke, A.P. Chemelli, Transcatheter embolization for the management of acute active inferior epigastric artery hemorrhages, *J. Endovasc. Ther.* 20 (August(4)) (2013) 561–567.
- [4] R.A. Agha, M.R. Borrelli, R. Farwana, K. Koshy, A.J. Fowler, D.P. Orgill, H. Zhu, A. Alsawadi, A. Noureldin, A. Rao, A. Enam, The SCARE 2018 statement: updating consensus Surgical Case REport (SCARE) guidelines, *Int. J. Surg.* 1 (December (60)) (2018) 132–136.
- [5] T.F. Leite, L.A. Pires, K. Goke, J.G. Silva, C.A. Chagas, Corona Mortis: anatomical and surgical description on 60 cadaveric hemipelvises, *Revista do Colégio Brasileiro de Cirurgiões* 44 (December(6)) (2017) 553–559.
- [6] T.F. de Oliveira Leite, E. Bortolini, B. Linard, B.A. Boueri, F.C. Carnevale, C.H. Nomura, et al., Evaluation of morphological and clinical factors related to failure of percutaneous treatment with thrombin injection of femoral pseudoaneurysms from cardiac catheterization, *Ann. Vasc. Surg.* 1 (August (59)) (2019) 173–183.
- [7] P.R. Sobkin, A.I. Bloom, M.W. Wilson, J.M. LaBerge, G.S. Hastings, R.L. Gordon, et al., Massive abdominal wall hemorrhage from injury to the inferior epigastric artery: a retrospective review, *J. Vasc. Interv. Radiol.* 19 (March (3)) (2008) 327–332.
- [8] P.A. Patel, F.R. Ernst, C.L. Gunnarsson, Evaluation of hospital complications and costs associated with using ultrasound guidance during abdominal paracentesis procedures, *J. Med. Econ.* 15 (January (1)) (2012) 1–7.
- [9] S. Yalamanchili, S.M. Harvey, A. Friedman, J.N. Shams, J.E. Silberzweig, Transarterial embolization for inferior epigastric artery injury, *Vasc. Endovascular Surg.* 42 (October(5)) (2008) 489–493.
- [10] T.F. Leite, O.I. Pereira, Superselective transcatheter arterial embolization in the treatment of Angiodysplasia, *Clin. Med. Insights Case Rep.* 12 (May (2019)), 1179547619842581.

Open Access

This article is published Open Access at [sciencedirect.com](https://www.sciencedirect.com). It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.