

ON THE EXISTENCE OF PERIODIC SOLUTIONS FOR THE EQUATION

$$\ddot{x} + f(x)\dot{x} + g(x) = 0$$

Hamilton Luiz Guidorizzi

Instituto de Matemática e Estatística

Universidade de São Paulo - Cx Postal 20570

(Ag. Iguatemi) 01498 - São Paulo - Brasil

Abstract

We establish in this work sufficient conditions for existence (and for non existence) of periodic solutions for the Liénard equation

$$\ddot{x} + f(x)\dot{x} + g(x) = 0.$$

1. The Definite Positive Function V_α . Auxiliary Lemmas.

Throughout this work we assume $f, g : \mathbb{R} \rightarrow \mathbb{R}$ are functions satisfying the following conditions:

- a) f is continuous and g is of class C^1 ;
- b) $xg(x) > 0$ for $x \neq 0$;
- c) $\int_0^{+\infty} g(x)dx = +\infty = \int_0^{-\infty} g(x)dx$.

Let α be a given real. We indicate by Ω_α the following open set:

$$\Omega_\alpha = \{(x, y) \in \mathbb{R}^2 \mid y > -\frac{1}{\alpha}\} \quad \text{for } \alpha > 0;$$

$$\Omega_\alpha = \{(x, y) \in \mathbb{R}^2 \mid y < -\frac{1}{\alpha}\} \quad \text{for } \alpha < 0;$$

$$\Omega_\alpha = \mathbb{R}^2 \quad \text{for } \alpha = 0.$$


We indicate by V_α the definite positive function given by

$$V_\alpha(x, y) = \int_0^x g(u)du + \int_0^y \frac{s}{\alpha s + 1} ds, \quad (x, y) \in \Omega_\alpha.$$

It can be immediately verified that, for $\alpha \neq 0$,

$$\int_0^{+\infty} \frac{s}{\alpha s + 1} ds = +\infty = \int_0^{-\frac{1}{\alpha}} \frac{s}{\alpha s + 1} ds.$$

It can also be immediately verified that the level curves of V_α are all closed curves and that $V_\alpha(x, 0)$ is strictly increasing in $[0, +\infty[$. Such curves show the following aspect: (see [1])



The equation

$$\ddot{x} + f(x)\dot{x} + g(x) = 0 \tag{1}$$

is equivalent to the system:

$$\begin{cases} \dot{x} = y \\ \dot{y} = -f(x)y - g(x) \end{cases} \tag{2}$$

The condition a) ensures existence and uniqueness of solution of (2). The condition b) ensures that $(0, 0)$ is the only point of equilibrium for system (2). It can be immediately verified that the derivative of V_α relative to system (2) is:

$$\dot{V}_\alpha(x, y) = -\frac{[f(x) - \alpha g(x)]}{\alpha y + 1} y^2, \quad (x, y) \in \Omega_\alpha. \tag{3}$$

Because $\alpha y + 1 > 0$ is true for all $(x, y) \in \Omega_\alpha$, it follows that the sign of \dot{V}_α depends only of $f(x) - \alpha g(x)$.

Lemma 1. Assume there are $\alpha > 0$ and $b > 0$ such that for all $x \geq b$,

$$f(x) \geq \alpha g(x).$$

Let $y_0 > 0$, $L = V_\alpha(b, y_0)$ and

$$K = \{(x, y) \in \Omega_\alpha \mid x \geq b \text{ and } V_\alpha(x, y) \leq L\}.$$

Let $\gamma(t) = (x(t), y(t))$ be the solution of (2) so that $\gamma(t_0) = (b, y_1)$, with $0 < y_1 < y_0$. Then, there is $t_1 > t_0$ such that

$$\gamma(t) \in K, \quad t_0 \leq t \leq t_1$$

and

$$\gamma(t_1) = (b, y_2),$$

with $-\frac{1}{\alpha} < y_2 < 0$.

Demonstration.

From $\dot{x}(t_0) = y_1 > 0$, it follows there is $t_2 > t_0$ so that

$$\gamma(t) \in K, \quad t_0 \leq t \leq t_2.$$

On the other hand, being $\dot{x}(t) > 0$ on the half plane $y > 0$, $\dot{x}(t) < 0$ on the half plane $y < 0$, $\dot{y}(t) < 0$ on the positive half-axis $0x$ and $(0, 0)$ the only point of equilibrium, there must exist $t_3 > t_2$ such that $\gamma(t_3) \notin K$.

Let

$$t_1 = \max\{u > t_0 \mid \gamma(t) \in K, \quad t_0 \leq t \leq u\}.$$

From the hypothesis

$$f(x) \geq \alpha g(x), \quad x \geq b,$$

and from (3) it follows that


$$\dot{V}_\alpha(\gamma(t)) \leq 0, \quad t_0 \leq t \leq t_1.$$

Because $V_\alpha(\gamma(t_0)) = V_\alpha(b, y_1) < L$, it follows that $V(\gamma(t_1)) < L$. So, $\gamma(t_1)$ does not belong to the arc given by

$$x \geq b \text{ and } V_\alpha(x, y) = L.$$

Because $\dot{x}(t) > 0$ on the $y > 0$ half-plane, it follows that

$$\gamma(t_1) = (b, y_2), \text{ with } -\frac{1}{\alpha} < y_2 < 0.$$



In a similar way, we can demonstrate the following lemmas:

Lemma 2. Assume the exist $\alpha < 0$ and $a < 0$ such that, for all $x \leq a$,

$$f(x) \geq \alpha g(x).$$

Let $y_0 < 0$, $L = V_\alpha(a, y_0)$ and

$$K = \{(x, y) \in \Omega_\alpha \mid x \leq a \text{ and } V_\alpha(x, y) \leq L\}.$$

Let $\gamma(t) = (x(t), y(t))$ the solution of (2) such that $\gamma(t_0) = (a, y_1)$, with $y_0 < y_1 < 0$.

Then there is $t_1 > t_0$ so that

$$\gamma_1(t) \in K, \quad t_0 \leq t \leq t_1$$

and

$$\gamma(t_1) = (a, y_2)$$

with $0 < y_2 < -\frac{1}{\alpha}$.

Lemma 3. Assume there exists $a < 0$ such that for all $x \leq a$,

$$f(x) \geq 0.$$

Let $y_0 < 0$, $L = V_0(a, y_0)$ and

$$K = \{(x, y) \in \mathbb{R}^2 \mid x \leq a \text{ and } V_0(x, y) \leq L\}.$$

Let $\gamma(t) = (x(t), y(t))$ the solution of (2) such that $\gamma(t_0) = (a, y_1)$, with $y_0 < y_1 < 0$. Then, there is $t_1 > t_0$ such that $\gamma(t) \in K$, $t_0 \leq t \leq t_1$ and

$$\gamma(t_1) = (a, y_2)$$

with

$$0 < y_2 < |y_0|.$$

Lemma 4. Assume there is $b > 0$ such that

$$f(x) \geq 0, \quad x \geq b.$$

Let $y_0 > 0$, $L = V_0(b, y_0)$ and

$$K = \{(x, y) \in \mathbb{R}^2 \mid x \geq b \text{ and } V_0(x, y) \leq L\}.$$

Let $\gamma(t) = (x(t), y(t))$ be the solution of (2) such that $\gamma(t_0) = (b, y_1)$, with $0 < y_1 < y_0$. Then there is $t_1 > t_0$ such that

$$\gamma(t) \in K, \quad t_0 \leq t \leq t_1$$

and

$$\gamma(t_1) = (b, y_2)$$

with $-y_0 < y_2 < 0$.

To close this section, we prove that the solutions of (2) do not admit vertical asymptotes. It is enough, to this end, to show that all solutions of the equation

$$\frac{dy}{dx} = -f(x) - \frac{g(y)}{y}, \quad y \neq 0 \tag{4}$$

do not admit vertical asymptotes.

Let us assume that (4) has a solution

$$y = y(x), \quad a \leq x < b$$

such that

$$\lim_{x \rightarrow b^-} y(x) = +\infty. \tag{5}$$

We can assume with no loss of generality, that $0 < y(a) \leq y(x)$ for $a \leq x < b$.

Let

$$A = \max_{a \leq x \leq b} |f(x)| \quad \text{and} \quad B = \max_{a \leq x \leq b} |g(x)|.$$

It follows from the mean value theorem that, for $a < x < b$,

$$y(x) - y(a) \leq \left[A + \frac{B}{y(a)} \right] (b - a)$$

which is in clear contradiction with (5). The other situations can be analysed in a similar way.

2. Sufficient conditions for the existence of periodic solutions.

Theorem 1. Consider the equation

$$\ddot{x} + f(x)\dot{x} + g(x) = 0 \tag{1}$$

where f, g satisfy the conditions a), b) and c) of the previous section. Assume also, that the following hypotheses are satisfied:

- 1) There are $\alpha > 0$ and $b > 0$ such that for all $x \geq b$,

$$f(x) \geq \alpha g(x);$$

- 2) The origin is repulsive;

- 3) There is $a < 0$ such that for all $x \in [c, a]$,

$$f(x) \geq 0$$

where $V_0(c, 0) = V_0(a, r)$, $r = \frac{1}{\alpha} + (A + \alpha B)(b - a)$,

$$A = \max_{a \leq x \leq b} |f(x)| \quad \text{and} \quad B = \max_{a \leq x \leq b} |g(x)|.$$

Under these conditions, the equation (1) will admit at least one non trivial periodic solution.

Demonstration.

The equation (1) is equivalent to the system

$$\begin{cases} \dot{x} = y \\ \dot{y} = -f(x)y - g(x) \end{cases} \quad (2)$$

Let $\gamma(t) = (x(t), y(t))$ the solution of (2) that at time $t = 0$ is at the position $\gamma(0) = (b, -\frac{1}{\alpha})$. Because γ does not admit vertical asymptotes and the origin is repulsive, there is a smaller time $t_1 > 0$ such that

$$\gamma(t_1) = (a, y_1), \quad y_1 < 0,$$

or


$$\gamma(t_1) = (x_1, 0), \quad a < x_1 < 0.$$

It can be immediately shown that

$$-\frac{1}{\alpha} - [A + \alpha B](b - a) < y_1 < 0.$$

(Indeed: Assuming $y_1 < -\frac{1}{\alpha}$, let $y = y(x)$ the solution of $\frac{dy}{dx} = -f(x) - \frac{g(x)}{y}$ such that $y(a) = y_1$ and $y(b) = -\frac{1}{\alpha}$. There is $x_0 \in]a, b[$ such that $y(x_0) = -\frac{1}{\alpha}$ and $y(x) < -\frac{1}{\alpha}$, $a \leq x \leq x_0$; by the mean value theorem, $y(x_0) - y(a) < [A + \alpha b](x_0 - a)$.)

Let $t_2 > 0$ the smallest value of t when γ crosses the y negative half-axis: $\gamma(t_2) = (0, y_2)$, $y_2 < 0$. The hypotheses 1), 2) and 3) together with lemmas 1 and 3 ensure that $\gamma(t)$ will again cross the y negative half-axis at a point $(0, y_3)$ with $y_2 < y_3 < 0$.



($L_1 = V_0(c, 0) = V_0(a, r)$ and $L = V_a(b, y_3)$). By the Theorem of Poincaré-Bendixon, the equation will admit at least one periodic solution. ■

Remark 1. One possible value for y_3 is

$$y_3 = r + \left[A + \frac{B}{r}\right](b - a).$$

Let $m > b$ such that $V_a(m, 0) = V_a(b, y_3)$. The hypothesis 1) can be weakened: it is enough to assume

$$f(x) \geq \beta g(x), \quad b \leq x \leq m.$$

Remark 2. The hypotheses 1) and 3) can be replaced for:

1') There are $\alpha < 0$ and $a < 0$ such that for all $x \leq a$,

$$f(x) \geq \alpha g(x);$$

3') There are $b > 0$ such that, for all $x \in [b, c]$,

$$f(x) \geq 0$$

where $V_0(c, 0) = V_0(b, r)$, $r = -\frac{1}{\alpha} + [A - \alpha B](b - a)$,

$$A = \max_{a \leq x \leq b} |f(x)| \quad \text{and} \quad B = \max_{a \leq x \leq b} |g(x)|.$$

Remark 3. A sufficient condition for the origin to be repulsive is that there exist β and s reals, $s > 0$, such that

$$f(x) < \beta g(x), \quad 0 < |x| < s,$$

for, in this case, we will have

$$V_\beta(x, y) < 0 \quad \text{for} \quad 0 < |x| < s$$

what implies the origin is repulsive [see (1)].

Theorem 2. Consider the equation

$$\ddot{x} + f(x)\dot{x} + g(x) = 0$$

where f and g satisfy the conditions a), b) and c) of the previous section. Let us assume, also, that the following hypotheses are satisfied:

1) There are $\alpha > 0$ and $b > 0$ such that, for all $x \geq b$,

$$f(x) \geq \alpha g(x);$$

- 2) The origin is repulsive;
 3) There is $a < 0$ such that, for all $x \leq a$, $f(x) \geq \beta g(x)$ where $\frac{1}{\beta} = \frac{1}{\alpha} + (A + \alpha B)(b - a)$, $A = \max_{a \leq x \leq b} |f(x)|$ and $B = \max_{a \leq x \leq b} |g(x)|$.

Under these conditions, the equation will admit at least one non trivial periodic solution.

Demonstration.

Let $\gamma(t) = (x(t), y(t))$ be the solution of (2) that at time $t = 0$ is at the position

$$\gamma(0) = (b, -\frac{1}{\alpha}).$$

By the same reasoning as in Theorem 1, there will be a smaller value $t_1 > 0$ such that

$$\gamma(t_1) = (x_1, 0), \quad a < x_1 < 0$$

or


$$\gamma(t_1) = (a, y_1)$$

where $-\frac{1}{\beta} < y_1 < 0$. Let $-\frac{1}{\beta} < y_4 < y_1$. Suppose $\gamma(t_1) = (a, y_1)$. The hypothesis 3) ensures that $\gamma(t)$ cannot leave the compact set

$$K = \{(x, y) \in \Omega_\beta \mid x \leq a, V_\beta(x, y) \leq V_\beta(a, y_4)\}$$

by crossing the arc

$$x \leq a \quad \text{and} \quad V_\beta(x, y) = V_\beta(a, y_4) = L_2$$



The demonstration is completed following the same reasoning as in Theorem 1. ■

Remark 4. The hypothesis 3) of Theorem 2 can be replaced by:

3') There is $a < 0$ such that, for all $x \in [c, a]$,

$$f(x) \geq \beta g(x)$$

where $\frac{1}{\beta} > r = \frac{1}{\alpha} + (A + \alpha B)(b - a)$, $c < a$ is such that $V_\beta(c, 0) = V_\beta(a, -r)$, $A = \max_{a \leq x < b} |f(x)|$ and $B = \max_{a \leq x \leq b} |g(x)|$.

When the hypothesis 3') is satisfied, we can make y_3 equal to

$$y_3 = y_5 + \left(A + \frac{B}{y_5}\right)(b - a)$$

where $y_5 > 0$ is such that $V_\beta(a, -r) = V_\beta(a, y_5)$.

In this case, it is enough to assume in hypothesis 1) that

$$f(x) \geq \alpha g(x), \quad b \leq x \leq m$$

where $m > b$ is such that $V_a(b, y_3) = V_a(m, 0)$.

Remark 5. The hypotheses 1) and 3) of Theorem 2 can be replaced by:

1") There are $\alpha < 0$ and $a < 0$ such that, for all $x \leq a$,

$$f(x) \geq \alpha g(x);$$

3") There is $b > 0$ such that, for all $x \geq b$,

$$f(x) \geq \beta g(x)$$

where $-\frac{1}{\beta} = -\frac{1}{\alpha} + (A - \alpha B)(b - a)$, $A = \max_{a \leq x \leq b} |f(x)|$ and $B = \max_{a \leq x \leq b} |g(x)|$.

3. Sufficient Condition for the Non Existence of Periodic Solution.

Theorem 3. Consider the equation

$$\ddot{x} + f(x)\dot{x} + g(x) = 0$$

where f and g satisfy the conditions a), b) and c) of section 1. Let us assume, also, there exists α real such that, for all $x \neq 0$,

$$f(x) > \alpha g(x).$$

Then the origin will be asymptotically stable in the Liapunov sense. Besides, if $\gamma(t)$ is a solution of system (2) such that $\gamma(t_0) \in \Omega_\alpha$ for some t_0 , then

$$\gamma(t) \in \Omega_\alpha, \quad t \geq t_0$$

and $\gamma(t)$ will approach the origin when $t \rightarrow +\infty$.

Demonstration.

It is sufficient to observe that the origin is the largest invariant set included in the set

$$E = \{(x, y) \in \Omega_\alpha \mid \dot{V}_\alpha(x, y) = 0\}$$

and apply the La Salle theorem [4]. ■

Remark 6. The hypothesis

$$f(x) > \alpha g(x), \quad x \neq 0,$$

can be weakened. It is sufficient to assume, for all x , $f(x) \geq \alpha g(x)$, with the condition that the origin must be the largest invariant set included in

$$E = \{(x, y) \in \Omega_\alpha \mid \dot{V}_\alpha(x, y) = 0\}.$$

Corollary. Satisfied the hypothesis of Theorem 3, all non trivial solution of equation

$$\ddot{x} + f(x)\dot{x} + g(x) = 0$$

will be non periodic.

References

- [1] H.L. Guidorizzi. *Contribuições ao estudo das equações diferenciais ordinárias de 2ª ordem*. Tese de Doutorado, IME-USP, (1988).
- [2] H.L. Guidorizzi. *Sobre existência de solução periódica para a equação de Liénard $\ddot{x} + f(x)\dot{x} + g(x) = 0$* . 28º Seminário Brasileiro de Análise, 144-153, (1988).
- [3] G. Villari. *On the qualitative behavior of solutions of Liénard equation*. J. of Differential Equations, 67, 269-277, (1987).
- [4] J. La Salle and S. Lefschetz. *Stability by Liapunov's Direct Method, with applications*. Academic Press, (1964).