

RT-MAE 2002-17

**AN IMPROVED TEST FOR  
HETEROSKEDASTICITY USING ADJUSTED  
MODIFIED PROFILE LIKELIHOOD INFERENCE**

*by*

**Silvia L.P. Ferrari, Audrey H.M.A. Cysneiros  
and  
Francisco Cribari-Neto**

**Palavras-Chave:** Bartlett correction; heteroskedasticity; likelihood ratio test; nuisance parameters; profile likelihood.  
**Classificação AMS:** 62J05, 62F04.  
(AMS Classification)

# AN IMPROVED TEST FOR HETEROSKEDASTICITY USING ADJUSTED MODIFIED PROFILE LIKELIHOOD INFERENCE

SILVIA L.P. FERRARI

Departamento de Estatística/IME  
Universidade de São Paulo  
Caixa Postal 66281, São Paulo/SP, 05311-970, Brazil  
email: sferrari@ime.usp.br

AUDREY H.M.A. CYSNEIROS

Departamento de Estatística, CCEN  
Universidade Federal de Pernambuco  
Cidade Universitária, Recife/PE, 50740-540, Brazil  
email: audrey@de.ufpe.br

FRANCISCO CRIBARI-NETO

Departamento de Estatística, CCEN  
Universidade Federal de Pernambuco  
Cidade Universitária, Recife/PE, 50740-540, Brazil  
email: cribari@ufpe.br

**ABSTRACT.** This paper develops an inference strategy for detecting heteroskedasticity in linear regression models. We develop a Bartlett adjustment to the modified profile likelihood ratio test (Cox and Reid, 1987) for heteroskedasticity in the linear regression model. Our results generalize those in Ferrari and Cribari-Neto (2002), since they allow for a vector-valued structure for the parameter that defines the skedastic function. Monte Carlo evidence shows that the proposed test displays reliable finite-sample behavior, outperforming the original likelihood ratio test, the Bartlett-corrected likelihood ratio test, and the modified profile likelihood ratio test.

**KEYWORDS AND PHRASES:** Bartlett correction; heteroskedasticity; likelihood ratio test; nuisance parameters; profile likelihood.

## 1. INTRODUCTION

Linear regression models are oftentimes used to model the behavior of a variable of interest conditional on a set of explanatory variables. In practical applications, it is common for these models to include a heteroskedastic structure, thus indicating that the conditional variances are not constant across observations. Since the modeling strategies are different when such variances are not constant, it is important to first test whether heteroskedasticity is present in the data. The most commonly used tests for heteroskedasticity are based on first order asymptotics, since they rely on a large sample approximation. The asymptotic approximation used to perform the tests, however, may not deliver accurate inference when the sample size is not large. In particular, likelihood ratio tests for constancy of variances in linear regression models tend to be severely oversized in samples of typical size. It is thus important to develop inference strategies that are more reliable and display superior finite-sample behavior. Cordeiro (1993) proposed a Bartlett adjustment to the likelihood ratio test for heteroskedasticity, where the test statistic is modified using an adjustment factor that typically reduces finite-sample size discrepancies. The original likelihood ratio statistic is distributed as  $\chi^2$  up to an error of order  $n^{-1}$ , whereas the transformed statistic follows the same limiting distribution up to an error of order  $n^{-2}$ . Simonoff and Tsai (1994) used the modified profile likelihood inference approach proposed by Cox and Reid (1987) to develop a test for heteroskedasticity aiming at reducing the effect of nuisance parameters on the resulting inference. The proposed test typically delivers more reliable inference in finite samples, but the test statistic is still  $\chi^2$  distributed, under the null hypothesis, up to an error of order  $n^{-1}$ . Ferrari and Cribari-Neto (2002) have Bartlett-corrected this test to obtain an adjusted modified profile likelihood test that is second order accurate. Their results, however, have an important limitation: they are only valid for situations where the parameter that defines the heteroskedastic behavior of the data is scalar. Since in most applications practitioners use variance specifications that are based on more than just one covariate, it is important to develop second order accurate adjusted modified profile likelihood inference in wider generality. This is the goal of the present paper.

The paper unfolds as follows. Section 2 presents the model of interest and the aforementioned tests for heteroskedasticity. Section 3 develops the main result in the article, namely the second order accurate inference based on adjusted modified profile likelihood methods. Numerical evidence on the finite-sample behavior of the different tests is presented in Section 4. The numerical evidence favors the test proposed in this paper. Section 5 discusses an application to real data. Concluding remarks are given in Section 6.

## 2. SOME TESTS FOR HETEROSKEDASTICITY

The model of interest is

$$y = X\beta + u, \quad (1)$$

where  $y$  is an  $n$ -vector of observations on the dependent variable,  $X$  is an  $n \times k$  nonrandom matrix of covariates with full column rank,  $\beta$  is a  $k$ -vector of unknown regression parameters, and  $u$  is an  $n$ -vector of random errors. It is assumed that  $u \sim \mathcal{N}(0, \sigma^2 W)$ , where  $W$  is an  $n \times n$  diagonal matrix with  $l$ th entry given by  $w_l = w(z_l, \delta) > 0$ ,  $z_l'$  being the  $l$ th row of the  $n \times p$  matrix  $Z$  of variance covariates,  $\sigma^2$  is a strictly positive and finite unknown constant, and  $\delta$  is a  $p \times 1$  unknown parameter vector. The desired inference is obtained by testing the null hypothesis  $\mathcal{H}_0 : \delta = \delta_0$  (homoskedasticity), where  $\delta_0$  is a given  $p$ -vector of constants such that  $w(z_l, \delta_0) = 1$  for  $l = 1, \dots, n$ , against a two-sided alternative hypothesis. The number of parameters of interest is therefore  $p$ , and the number of nuisance parameters is  $k + 1$ . The resulting log-likelihood function is

$$L = L(y; \delta, \beta, \sigma^2) = -\frac{n}{2} \log(\sigma^2) - \frac{1}{2} \sum_{l=1}^n \log w(z_l, \delta) - \frac{1}{2\sigma^2} (y - X\beta)' W^{-1} (y - X\beta) + \text{constant}.$$

The maximum likelihood estimator (MLE) of  $\delta$  can be obtained by maximizing the profile log-likelihood function

$$L_p(y; \delta) = L(y; \delta, \hat{\beta}_\delta, \hat{\sigma}_\delta^2), \quad (2)$$

where

$$\hat{\beta}_\delta = (X'W^{-1}X)^{-1}X'W^{-1}y \quad \text{and} \quad \hat{\sigma}_\delta^2 = \frac{1}{n}(y - X\hat{\beta}_\delta)'W^{-1}(y - X\hat{\beta}_\delta).$$

That is,

$$L_p(y; \delta) = -\frac{n}{2} \log(\hat{\sigma}_\delta^2) - \frac{1}{2} \sum_{l=1}^n \log w(z_l, \delta) + \text{constant}.$$

The likelihood ratio statistic for the test at hand is

$$LR = -2\{L_p(y; \delta_0) - L_p(y; \hat{\delta})\}, \quad (3)$$

where  $\hat{\delta}$  is the MLE of  $\delta$ . The null distribution of  $LR$  in large samples may be approximated by a  $\chi_p^2$  distribution. However, it is well known that this approximation may be very poor if the sample size is not large enough. This shortcoming can be circumvented by modifying the test statistic using a Bartlett correction in order to improve on the approximation. The Bartlett-corrected likelihood ratio statistic,

$$LR^* = \frac{LR}{1 + c},$$

where  $c$  is a suitably chosen constant of order  $n^{-1}$  such that  $E(LR) = p\{(1+c) + O(n^{-3/2})\}$ , has a  $\chi_p^2$  null distribution with an error of order  $n^{-2}$  (see Barndorff-Nielsen and Hall, 1988). The gain in accuracy stems from the fact that  $\text{pr}(LR^* > x_\alpha) = \alpha + O(n^{-2})$  whereas for the unmodified statistic  $\text{pr}(LR > x_\alpha) = \alpha + O(n^{-1})$ , where  $x_\alpha$  is the  $1 - \alpha$  upper point from a  $\chi_p^2$  distribution. A general expression for  $c$  can be found in Lawley (1956); see Cribari-Neto and Cordeiro (1996). The Bartlett adjustment factor needs to be tailored for each application of interest, and Cordeiro (1993) has obtained a closed form expression for such a constant for the test for heteroskedasticity in linear regression. His main result is reproduced by Ferrari and Cribari-Neto (2002, p. 355).

As indicated earlier, Simonoff and Tsai (1994) have used the modified profile likelihood approach proposed by Cox and Reid (1987) to obtain an alternative homoskedasticity test statistic. To that end, it is required that  $\delta$  be orthogonal to the remaining parameters. For the model of interest here, the transformation  $(\delta', \beta', \sigma^2)' \rightarrow (\delta', \beta', \gamma)'$ , with

$$\sigma^2 = \frac{\gamma}{(\prod_{l=1}^n w_l)^{1/n}},$$

delivers the desired orthogonality. The log-likelihood function for the reparameterized model is given by

$$L^* = L^*(y; \delta, \beta, \gamma) = -\frac{n}{2} \log \gamma - \frac{1}{2\gamma} (y - X\beta)' V (y - X\beta) + \text{constant},$$

where  $V = \text{diag}\{v_1, \dots, v_n\}$ , with

$$v_l = v_l(\delta) = \left( \prod_{m=1}^n w_m \right)^{1/n} / w_l. \quad (4)$$

The corresponding profile log-likelihood function for  $\delta$  is

$$L_p^* = L_p^*(y; \delta) = -\frac{n}{2} \log \hat{\gamma}_\delta + \text{constant},$$

where  $\hat{\gamma}_\delta = \hat{\sigma}_\delta^2 (\prod_{l=1}^n w_l)^{1/n}$ . The modified test statistic is given by equation (3) but with  $L_p(y; \delta)$  replace by

$$L_p^*(y; \delta) - \frac{1}{2} \log [\det \{j^*(\delta; \hat{\beta}_\delta, \hat{\gamma}_\delta)\}],$$

where  $j^*$  is the observed information matrix in the reparameterized model for the nuisance parameters  $(\beta', \gamma)'$  evaluated at  $(\hat{\beta}_\delta, \hat{\gamma}_\delta)'$ . The resulting test statistic is

$$LR_m = \frac{n-k-2}{n} LR + \log \left\{ \frac{\det(X'X)}{\det(\hat{X}_m' \hat{X}_m)} \right\},$$

where  $\hat{X}_m = \hat{G}^{-1/2} X$ . Here,  $\hat{G}$  is an  $n \times n$  diagonal matrix with  $l$ th entry given by  $w(z_l, \hat{\delta}) / \{\prod_{m=1}^n w(z_m, \hat{\delta})\}^{1/n}$ .

### 3. SECOND ORDER ACCURATE PROFILE LIKELIHOOD INFERENCE

The modified profile likelihood ratio test presented in Section 2 typically delivers improved inference relative to the original likelihood ratio test, but both tests are based on first order asymptotics. It is possible to develop a Bartlett adjustment to such a test in order to obtain a new test that is second order accurate. This will be done in this section using the results in DiCiccio and Stern (1994). We find a correction factor  $c_m$  that defines the transformed statistic

$$LR_m^* = \frac{LR_m}{1 + c_m}$$

such that  $\text{pr}(LR_m^* > x_\alpha) = \alpha + \mathcal{O}(n^{-2})$ .

The notation used can be summarized as follows, where indices  $a, b, \dots$  range over  $1, \dots, p$ , and indices  $r, s, t, \dots$  range over  $1, \dots, p+k+1$ . Let  $\theta = (\delta', \beta', \gamma)'$ ,  $\theta^r$  denoting the  $r$ th element of  $\theta$ ,  $\lambda_{rs} = E(\partial^2 L / \partial \theta^r \partial \theta^s)$ ,  $\lambda_{rst} = E(\partial^3 L / \partial \theta^r \partial \theta^s \partial \theta^t)$ , etc. Define  $(\lambda_{rs})_t = \partial \lambda_{rs} / \partial \theta^t$ ,  $(\lambda_{rs})_{tu} = \partial^2 \lambda_{rs} / \partial \theta^t \partial \theta^u$ , etc. The  $(r, s)$  element of Fisher's information matrix is thus given by  $-\lambda_{rs}$ , and the corresponding element of its inverse is  $-\lambda^{rs}$ . Additionally,  $\tau^{rs} = \lambda^{ra} \lambda^{sb} \sigma_{ab}$ , where  $(\sigma_{ab})$  is the  $p \times p$  matrix inverse of  $(\lambda^{ab})$ , and  $\nu^{rs} = \lambda^{rs} - \tau^{rs}$ . Note that the entries of the  $(p+k+1) \times (p+k+1)$  matrix  $(\nu^{rs})$  are all zero except for its lower right-hand  $(k+1) \times (k+1)$  submatrix which is the inverse of Fisher's information matrix for the nuisance parameters  $(\beta', \gamma)$  keeping  $\delta$  fixed. The implicit summation convention is used throughout, i.e., indices repeated as subscripts and superscripts indicate summation over the appropriate range.

DiCiccio and Stern (1994, p. 404, equation (25)) obtained a general expression for  $c_m$ , that can be written as

$$\begin{aligned} c_m = & \frac{1}{4} \tau^{ru} \tau^{st} \lambda_{rstu} - \lambda^{ru} \tau^{st} (\lambda_{rst})_u + (\lambda^{ru} \lambda^{st} - \nu^{ru} \nu^{st}) (\lambda_{rs})_{tu} \\ & - \left( \frac{1}{4} \lambda^{ru} \tau^{st} \tau^{vw} + \frac{1}{2} \lambda^{ru} \tau^{sw} \tau^{tv} - \frac{1}{3} \tau^{ru} \tau^{sw} \tau^{tv} \right) \lambda_{rst} \lambda_{uvw} \\ & + (\lambda^{ru} \tau^{st} \lambda^{vw} + \lambda^{ru} \lambda^{sw} \lambda^{tv} - \nu^{ru} \lambda^{sw} \nu^{tv}) \lambda_{rst} (\lambda_{uv})_w \\ & - (\lambda^{ru} \lambda^{st} \lambda^{vw} - \nu^{ru} \nu^{st} \nu^{vw} + \lambda^{ru} \lambda^{sw} \lambda^{tv} - \nu^{ru} \nu^{sw} \nu^{tv}) (\lambda_{rs})_t (\lambda_{uv})_w. \end{aligned}$$

In our setup, i.e., where the parameter vector is written in partitioned form as  $(\delta', \beta', \gamma)'$  and  $\gamma$  is scalar, we have that  $\nu^{ab} = \nu^{a\gamma} = \nu^{a i} = 0$ ,  $\tau^{ab} = \lambda^{ab}$ ,  $\tau^{a\gamma} = \lambda^{a\gamma}$ ,  $\tau^{a i} = \lambda^{a i}$ , and  $\tau^{i\gamma} = \lambda^{i\gamma}$ , where  $\gamma$  is used to represent the index corresponding to the last component

of the parameter vector  $(\delta', \beta', \gamma)'$ , and  $i, j, \dots$  range over  $p+1, \dots, p+k$ . Using the orthogonality between  $\delta, \beta$  and  $\gamma$ , it can be shown that  $\tau^{ij} = \tau\gamma\gamma = \lambda_{ai} = \lambda_{a\gamma} = \lambda_{i\gamma} = \lambda^{ai} = \lambda^{a\gamma} = \lambda^{i\gamma} = 0$ ,  $\nu^{ij} = \lambda^{ij}$  and  $\lambda\gamma\gamma = \nu\gamma\gamma$ . After long algebra, we obtain the following expression for  $c_m$ :

$$\begin{aligned} c_m = & \frac{1}{4}\lambda^{ab}\lambda^{cd}\lambda_{abcd} - \lambda^{ab}\lambda^{cd}(\lambda_{acd})_b + \lambda^{ab}\lambda^{cd}(\lambda_{ac})_{db} - \lambda^{ij}\lambda^{ab}(\lambda_{iab})_j \\ & - \lambda\gamma\gamma\lambda^{ab}(\lambda_{ab\gamma})_\gamma - \left(\frac{1}{4}\lambda^{ab}\lambda^{cd}\lambda^{ef} + \frac{1}{2}\lambda^{ab}\lambda^{cf}\lambda^{de} - \frac{1}{3}\lambda^{ab}\lambda^{cf}\lambda^{de}\right)\lambda_{acd}\lambda_{bef} \\ & + (\lambda^{ab}\lambda^{cd}\lambda^{ef} + \lambda^{ab}\lambda^{cf}\lambda^{de})\lambda_{acd}(\lambda_{be})_f - (\lambda^{ab}\lambda^{cd}\lambda^{ef} + \lambda^{ab}\lambda^{cf}\lambda^{de})(\lambda_{ac})_d(\lambda_{be})_f \\ & - \left(\frac{1}{4}\lambda^{ij}\lambda^{ab}\lambda^{cd} + \frac{1}{2}\lambda^{ij}\lambda^{ad}\lambda^{bc}\right)\lambda_{iab}\lambda_{jcd} + \lambda^{ij}\lambda^{ab}\lambda^{kl}\lambda_{iab}(\lambda_{jkl})_l \\ & - \left(\frac{1}{4}\lambda\gamma\gamma\lambda^{ab}\lambda^{cd} + \frac{1}{2}\lambda\gamma\gamma\lambda^{ad}\lambda^{bc}\right)\lambda_{ab\gamma}\lambda_{cd\gamma} + (\lambda\gamma\gamma\lambda^{ab}\lambda^{i\gamma})\lambda_{ab\gamma}(\lambda_{i\gamma\gamma})_\gamma. \end{aligned} \quad (5)$$

It is noteworthy that the expression we derived, expression (5), can be used to obtain closed-form expressions for the Bartlett adjustment factor in any class of models that uses the partition of the parameter vector as here and where orthogonality holds. In that sense, equation (5) is quite general and can be used to Bartlett-adjust modified profile likelihood ratio tests in classes of models other than the one we focus in this paper.

In what follows we consider the framework described in the previous section. It can be easily shown that

$$\frac{\partial L^*}{\partial \delta^a} = -\frac{1}{2\gamma}(y - X\beta)'V_{R_1}(y - X\beta).$$

The expression for  $V_{R_1}$  follows from

$$V_{R_m} = \text{diag} \left\{ \frac{\partial^m v_1}{\partial \delta^{a_1} \dots \partial \delta^{a_m}}, \dots, \frac{\partial^m v_n}{\partial \delta^{a_1} \dots \partial \delta^{a_m}} \right\}$$

with  $m = 1$ , where  $v_1, \dots, v_n$  are given in (4). Using the fact that if  $E(z) = \mu$  and  $\text{cov}(z) = \Sigma$  for an  $n \times 1$  random vector  $z$ , then  $E(z'Az) = \text{tr}(A\Sigma) + \mu'A\mu$ ,  $A$  being an  $n \times n$  nonrandom matrix, we obtain

$$E \left( \frac{\partial L^*}{\partial \delta^a} \right) = -\frac{1}{2} \text{tr}(V_{R_1} V^{-1}) = -\frac{1}{2} \sum_{l=1}^n \frac{v_{la}}{v_l},$$

where  $v_{la} = \partial v_l / \partial \delta^a$ . It follows from regularity conditions that the above expected value equals zero, so the  $w$ 's must satisfy  $\sum_{r=1}^n v_{ra} / v_r = 0$ . Note that

$$E \left( \frac{\partial^2 L^*}{\partial \delta^a \partial \gamma} \right) = \frac{1}{2\gamma^2} E \{ (y - X\beta)' V_{R_1} (y - X\beta) \} = \frac{1}{2\gamma} \text{tr}(V_{R_1} V^{-1}) = \frac{1}{2\gamma} \sum_{l=1}^n \frac{v_{la}}{v_l} = 0,$$

so that orthogonality between  $\delta$  and  $\gamma$  is preserved, as desired. The second derivative of  $L^*$  with respect to the elements of  $\delta$  is

$$\frac{\partial^2 L^*}{\partial \delta^a \partial \delta^b} = -\frac{1}{2\gamma}(y - X\beta)' V_{R_2}(y - X\beta).$$

Therefore,

$$\lambda_{ab} = -\frac{1}{2} \text{tr}(V_{R_2} V^{-1}) = -\frac{1}{2} \sum_{l=1}^n \frac{v_{lab}}{v_l},$$

where  $v_{lab} = \partial^2 v_l / (\partial \delta^a \partial \delta^b)$ ; in what follows,  $v_{lab} = \partial^3 v_l / (\partial \delta^a \partial \delta^b \partial \delta^c)$  and  $v_{labcd} = \partial^4 v_l / (\partial \delta^a \partial \delta^b \partial \delta^c \partial \delta^d)$ . After some algebra, we also obtain

$$\begin{aligned} \lambda_{abc} &= -\frac{1}{2} \sum_{l=1}^n \frac{v_{lab}}{v_l}, \quad \lambda_{abcd} = -\frac{1}{2} \sum_{l=1}^n \frac{v_{labcd}}{v_l}, \quad \lambda_{ab\gamma} = \frac{1}{2\gamma} \sum_{l=1}^n \frac{v_{lab}}{v_l}, \\ (\lambda_{ab\gamma})_\gamma &= -\frac{1}{2\gamma^2} \sum_{l=1}^n \frac{v_{lab}}{v_l}, \quad \lambda_{\gamma\gamma} = -\frac{n}{2\gamma^2}, \quad \lambda^{\gamma\gamma} = -\frac{2\gamma^2}{n}, \quad (\lambda_{\gamma\gamma})_\gamma = \frac{n}{\gamma^3}, \\ (\lambda_{abc})_d &= -\frac{1}{2} \sum_{l=1}^n \frac{\partial}{\partial \delta^d} \left( \frac{v_{lab}}{v_l} \right), \quad (\lambda_{ab})_{cd} = -\frac{1}{2} \sum_{l=1}^n \frac{\partial^2}{\partial \delta^c \partial \delta^d} \left( \frac{v_{lab}}{v_l} \right), \\ (\lambda_{ab})_c &= -\frac{1}{2} \sum_{l=1}^n \frac{\partial}{\partial \delta^c} \left( \frac{v_{lab}}{v_l} \right), \quad \lambda_{iab} = 0. \end{aligned}$$

It is now possible to further simplify the expression given for  $c_m$ :

$$\begin{aligned} c_m &= \lambda^{ab} \lambda^{cd} \left\{ \frac{1}{4} \lambda_{abcd} - (\lambda_{acd})_b + (\lambda_{ac})_{db} - \frac{1}{4} \lambda^{\gamma\gamma} \lambda_{ab\gamma} \lambda_{cd\gamma} \right\} \\ &\quad - \lambda^{ab} \lambda^{cd} \lambda^{ef} \left\{ \frac{1}{4} \lambda_{acd} \lambda_{bef} + \lambda_{acd} (\lambda_{be})_f - (\lambda_{ac})_d (\lambda_{be})_f \right\} \\ &\quad - \lambda^{ab} \lambda^{cf} \lambda^{de} \left\{ \frac{1}{6} \lambda_{acd} \lambda_{bef} + \lambda_{acd} (\lambda_{be})_f - (\lambda_{ac})_d (\lambda_{be})_f \right\} \\ &\quad - \lambda^{ab} \left\{ \lambda^{\gamma\gamma} (\lambda_{ab\gamma})_\gamma - (\lambda^{\gamma\gamma})^2 \lambda_{ab\gamma} (\lambda_{\gamma\gamma})_\gamma \right\} \\ &\quad - \frac{1}{2} \lambda^{ad} \lambda^{bc} \lambda^{\gamma\gamma} \lambda_{ab\gamma} \lambda_{cd\gamma}. \end{aligned} \tag{6}$$

In what follows, we shall consider the case of multiplicative heteroskedasticity, i.e., the special case where  $w_l = \exp\{z_l' \delta\}$ . Here,

$$v_l = \exp\{-(z_l - \bar{z})' \delta\}, \quad v_{la} = -(z_l - \bar{z})_a \exp\{-(z_l - \bar{z})' \delta\} = -(z_l - \bar{z})_a v_l,$$



$v_{lab} = (z_l - \bar{z})_{ab} v_l$ ,  $v_{labc} = -(z_l - \bar{z})_{abc} v_l$  and  $v_{labcd} = (z_l - \bar{z})_{abcd} v_l$ , with  $\bar{z} = (\bar{z}_1, \dots, \bar{z}_p)'$  and  $\bar{z}_a = n^{-1} \sum_{l=1}^n z_{la}$  for  $a = 1, \dots, p$ . Here,  $(z_l - \bar{z})_a = z_{la} - \bar{z}_a$ ,  $(z_l - \bar{z})_{ab} = (z_{la} - \bar{z}_a)(z_{lb} - \bar{z}_b)$ , and so on. We have that

$$\lambda_{abcd} = -\frac{1}{2} \sum_{l=1}^n (z_l - \bar{z})_{abcd}, \quad \lambda_{abc} = \frac{1}{2} \sum_{l=1}^n (z_l - \bar{z})_{abc}, \quad \lambda_{ab} = -\frac{1}{2} \sum_{l=1}^n (z_l - \bar{z})_{ab},$$

$$\lambda_{ab\gamma} = \frac{1}{2\gamma} \sum_{l=1}^n (z_l - \bar{z})_{ab}, \quad (\lambda_{ab\gamma})_\gamma = -\frac{1}{2\gamma^2} \sum_{l=1}^n (z_l - \bar{z})_{ab},$$

$$\lambda_{\gamma\gamma} = -\frac{2\gamma^2}{n}, \quad (\lambda_{acd})_b = (\lambda_{ac})_{db} = (\lambda_{bc})_f = 0.$$

Plugging these quantities into equation (6) it is possible to write  $c_m$  as

$$\begin{aligned} c_m = & -\frac{1}{8} \sum_{l=1}^n (z_l - \bar{z})_a \lambda^{ab} (z_l - \bar{z})_b (z_l - \bar{z})_c \lambda^{cd} (z_l - \bar{z})_d \\ & + \frac{1}{8n} \sum_{l=1}^n (z_l - \bar{z})_a \lambda^{ab} (z_l - \bar{z})_b \sum_{m=1}^n (z_m - \bar{z})_c \lambda^{cd} (z_m - \bar{z})_d \\ & - \frac{1}{16} \sum_{l=1}^n \sum_{m=1}^n (z_l - \bar{z})_a \lambda^{ab} (z_l - \bar{z})_b (z_l - \bar{z})_c \lambda^{cd} (z_m - \bar{z})_d (z_m - \bar{z})_e \lambda^{ef} (z_m - \bar{z})_f \\ & - \frac{1}{24} \sum_{l=1}^n \sum_{m=1}^n (z_l - \bar{z})_a \lambda^{ab} (z_m - \bar{z})_b (z_l - \bar{z})_c \lambda^{cd} (z_m - \bar{z})_d (z_l - \bar{z})_e \lambda^{ef} (z_m - \bar{z})_f \\ & + \frac{1}{n} \sum_{l=1}^n (z_l - \bar{z})_a \lambda^{ab} (z_l - \bar{z})_b \\ & + \frac{1}{4n} \sum_{l=1}^n \sum_{m=1}^n (z_l - \bar{z})_a \lambda^{ab} (z_m - \bar{z})_b (z_l - \bar{z})_c \lambda^{cd} (z_m - \bar{z})_d. \end{aligned}$$

Let  $H = \{h_{lm}\} = (Z - \bar{Z})[(Z - \bar{Z})'(Z - \bar{Z})]^{-1}(Z - \bar{Z})'$ , with  $(Z - \bar{Z}) = (z_1 - \bar{z}, \dots, z_n - \bar{z})'$ . Note that  $(z_l - \bar{z})_a \lambda^{ab} (z_m - \bar{z})_b = -2 h_{lm}$ . It is then possible to reduce the above expression for  $c_m$  to

$$\begin{aligned} c_m = & -\frac{1}{2} \sum_{l=1}^n h_{ll}^2 + \frac{1}{2n} \left( \sum_{l=1}^n h_{ll} \right)^2 + \frac{1}{2} \sum_{l=1}^n \sum_{m=1}^n h_{ll} h_{lm} h_{mm} + \frac{1}{3} \sum_{l=1}^n \sum_{m=1}^n h_{lm}^3 \\ & - \frac{2}{n} \sum_{l=1}^n h_{ll} + \frac{1}{n} \sum_{l=1}^n \sum_{m=1}^n h_{lm}^2. \end{aligned}$$

It is noteworthy that  $\sum_{l=1}^n h_{ll} = \text{tr}(H) = p$ . Therefore,  $c_m$  can be written in matrix form as

$$c_m = -\frac{1}{2}\text{tr}(H_d H_d) + \frac{1}{2n}p^2 + \frac{1}{2}\iota' H_d H H_d \iota + \frac{1}{3}\iota' H^{(3)} \iota - \frac{2}{n}p + \frac{1}{n}\iota' H^{(2)} \iota, \quad (7)$$

where  $\iota$  is an  $n$ -vector of ones,  $H_d = \text{diag}\{h_{11}, \dots, h_{nn}\}$ ,  $H^{(2)} = (h_{lm}^2)$  and  $H^{(3)} = (h_{lm}^3)$ .

Equation (7) gives a simple matrix expression for the Bartlett adjustment factor. The expression only involves simple matrix operations, and can be easily implemented into any computer algebra system or statistical software that can carry out simple linear algebra operations. It can thus be used with minimal effort by practitioners. We also note that the  $c_m$  in (7) depends only on the matrix  $Z$  of covariates used to model the fluctuations in the conditional variance of the response, on the number of unknown parameters in the skedastic function and on the number of observations. In particular, it is important to note that the Bartlett adjustment factor does not depend on the number of nuisance parameters (nor does it depend on any unknown parameters). This can be interpreted in light of the fact that the Cox and Reid adjustment to the profile likelihood aims at reducing the impact of nuisance parameters on the resulting inference, and this adjustment was used prior to the Bartlett correction of the test statistic. Finally, we note that expression (7) generalizes the result in Ferrari and Cribari-Neto (2002, p. 358), which only holds for  $p = 1$ , as indicated earlier.

#### 4. NUMERICAL EVIDENCE

The simulation results are based on the linear regression model

$$y_l = \beta_1 + \beta_2 x_{l2} + \dots + \beta_k x_{lk} + u_l, \quad l = 1, \dots, n,$$

where  $u_l \sim \mathcal{N}(0, \sigma^2 \exp\{\delta_1 z_{l1} + \dots + \delta_p z_{lp}\})$ , and  $\text{cov}(u_l, u_m) = 0$  for all  $l \neq m$ . When  $\delta_1 = \dots = \delta_p = 0$  the model is homoskedastic, with heteroskedasticity arising when  $\min\{|\delta_1|, \dots, |\delta_p|\} \neq 0$ . The simulations were carried out for different values of  $p$  and  $k$ . The covariates  $x_2, \dots, x_k$  were generated as independent draws from a standard uniform distribution  $\mathcal{U}(0, 1)$ . When  $p < k$ , the matrix  $Z$  was formed using columns 2,  $\dots$ ,  $p+1$  of  $X$ ; when  $p \geq k$ , the extra columns of  $Z$  were created using independent draws from the  $\mathcal{U}(0, 1)$  distribution.

We shall report the null rejection rates of the original likelihood ratio test ( $LR$ ), its Bartlett-corrected version ( $LR^*$ ), the modified profile likelihood ratio test ( $LR_m$ ), and its Bartlett-corrected version ( $LR_m^*$ ) for the test of the null hypothesis  $\mathcal{H}_0 : \delta_1 = \dots = \delta_p = 0$ . The number of replications was set at 10,000 and the following levels of significance were considered:  $\alpha = 0.100, 0.050, 0.010, 0.005$ . The simulations were performed using the

Ox matrix programming language (Doornik, 2001). The nonlinear maximizations of the relevant log-likelihoods were carried out using the BFGS quasi-Newton algorithm, which is generally perceived as the best performing method. All entries in the tables that follow are percentages.

Table 1 presents results for  $n = 35$ ,  $p = 2$  and different values for  $k$ . We vary  $k$  to analyze the effect of the number of nuisance parameters on the different tests. At the outset, we note that the original likelihood ratio test is considerably oversized, especially as the number of nuisance parameters increases. For instance, when  $k = 7$  and  $\alpha = 5\%$ , the null rejection rate exceeds 16%. The tendency of the test to overreject is attenuated by the Bartlett correction, the resulting Bartlett-adjusted test displaying smaller size distortions. For example, its null rejection rate for the same situation was 9.0%. The modified profile likelihood ratio test tends to overcorrect the liberal tendency of the original test, displaying null rejection rates that are smaller than the nominal level of the test. Again, when  $k = 7$  and  $\alpha = 5\%$ , its null rejection rate was 3.2%. The Bartlett correction applied to this test brings its empirical type I error probability closer to the nominal size of the test. For the situation singled out above, its null rejection rate was 3.5%. The above conclusions also hold for other nominal levels, even for very small ones.

Table 2 contains results for the situation where  $n = 35$ ,  $k = 5$  and  $p = 1, \dots, 8$ . Again, the results show that the original likelihood ratio test is considerably oversized, the more so the larger the number of covariates used in the specification of the skedastic function; e.g., when  $p = 6$  and  $\alpha = 5\%$ , the rejection rate of the test under the null hypothesis was nearly 26%, i.e., more than five times the nominal level selected for the test. The other conclusions drawn from Table 1 also hold here. For the same situation ( $p = 6$  and  $\alpha = 5\%$ ), the null rejection rates of the Bartlett-corrected likelihood ratio test, the modified profile likelihood ratio test, and of its Bartlett-adjusted version were 13.0%, 2.8% and 3.4%, respectively.

In Table 3, we fix the value of  $p$  at 2 and vary the sample size:  $n = 35, 50, 100$  for  $k = 5, 8$ . As the sample increases, the null rejection rates of all four tests become closer to the nominal level of the test, as expected. We also note that even for  $n = 100$  the original likelihood ratio test is still oversized. As an illustration, when  $k = 8$  and  $\alpha = 5\%$ , the null rejection rates of the likelihood ratio test, its Bartlett-corrected version, the modified profile likelihood ratio test, and its Bartlett-adjusted version were, respectively, 8.1%, 5.7%, 4.9% and 5.0%.

The test of equality of variances for the one factor and three levels model is considered in Table 4. Here, there are three normal populations with means  $\mu_1, \mu_2, \mu_3$  and variances  $\sigma_1^2, \sigma_2^2, \sigma_3^2$ , and the null hypothesis of interest is  $\mathcal{H}_0 : \sigma_1^2 = \sigma_2^2 = \sigma_3^2$ . Note that here  $p = 2$

Table 1. Null rejection rates,  $n = 35$  and  $p = 2$ .

$k$	$\alpha = 10\%$				$\alpha = 5\%$				$\alpha = 1\%$				$\alpha = 0.5\%$			
	$LR$	$LR^*$	$LR_m$	$LR_m^*$	$LR$	$LR^*$	$LR_m$	$LR_m^*$	$LR$	$LR^*$	$LR_m$	$LR_m^*$	$LR$	$LR^*$	$LR_m$	$LR_m^*$
2	13.4	10.6	8.7	9.7	7.5	5.4	4.2	4.8	1.6	1.1	0.7	0.9	0.9	0.5	0.4	0.5
3	14.5	10.9	8.8	9.6	8.2	5.9	4.4	4.9	2.3	1.2	0.8	0.9	1.2	0.7	0.4	0.5
4	16.3	11.2	7.9	8.8	9.5	5.7	3.9	4.4	2.7	1.3	0.8	0.9	1.7	0.8	0.4	0.4
5	19.2	12.8	8.0	8.9	11.9	7.2	3.7	4.3	3.8	1.6	0.6	0.8	2.4	0.9	0.3	0.4
6	22.3	14.4	6.4	7.4	14.6	7.8	3.0	3.5	5.2	2.1	0.5	0.7	3.3	1.3	0.3	0.3
7	24.9	15.6	7.3	7.8	16.6	9.0	3.2	3.5	6.3	2.5	0.5	0.7	4.1	1.5	0.2	0.3
8	30.4	18.9	5.3	5.8	21.8	11.7	2.3	2.6	9.7	3.7	0.4	0.4	6.9	2.3	0.2	0.2

Table 2. Null rejection rates,  $n = 35$  and  $k = 5$ .

$p$	$\alpha = 10\%$				$\alpha = 5\%$				$\alpha = 1\%$				$\alpha = 0.5\%$			
	$LR$	$LR^*$	$LR_m$	$LR_m^*$	$LR$	$LR^*$	$LR_m$	$LR_m^*$	$LR$	$LR^*$	$LR_m$	$LR_m^*$	$LR$	$LR^*$	$LR_m$	$LR_m^*$
1	16.3	11.6	8.5	9.2	9.6	6.1	4.1	4.4	3.0	1.6	0.7	0.9	1.8	0.9	0.4	0.4
2	19.2	12.8	8.0	8.9	11.9	7.2	3.7	4.3	3.8	1.6	0.6	0.8	2.4	0.9	0.3	0.4
3	21.8	13.9	8.0	8.7	14.0	7.5	3.6	4.2	4.7	1.9	0.7	0.8	2.9	1.0	0.3	0.4
4	26.6	15.8	7.4	8.4	17.2	9.1	3.2	3.8	6.3	2.4	0.7	0.8	4.1	1.4	0.4	0.5
5	29.6	17.3	7.0	7.8	20.0	9.9	3.4	3.8	8.0	2.8	0.6	0.7	5.3	1.7	0.3	0.4
6	36.0	21.2	6.1	6.9	25.9	13.0	2.8	3.4	11.2	4.4	0.7	0.8	8.0	2.8	0.3	0.4
7	46.1	27.6	4.5	4.9	34.9	18.0	2.0	2.2	17.3	6.9	0.4	0.5	12.9	4.2	0.1	0.2
8	49.4	28.8	4.0	4.4	37.4	19.2	1.6	1.9	18.9	7.2	0.4	0.4	14.1	4.7	0.1	0.2

Table 3. Null rejection rates,  $p = 2$  and several sample sizes.

$n$	$\alpha$	$k = 5$				$k = 8$			
		$LR$	$LR^*$	$LR_m$	$LR_m^*$	$LR$	$LR^*$	$LR_m$	$LR_m^*$
30	10.0	25.0	16.4	6.7	7.8	36.6	23.4	4.0	4.5
	5.0	16.7	9.9	3.1	3.6	27.5	14.9	1.6	1.8
	1.0	6.6	2.8	0.6	0.7	13.4	4.8	0.3	0.3
	0.5	4.2	1.5	0.3	0.4	9.5	3.1	0.1	0.1
50	10.0	16.2	11.2	8.9	9.5	21.6	13.8	8.3	8.8
	5.0	9.2	5.8	4.1	4.5	13.6	7.7	3.7	4.1
	1.0	2.5	1.3	0.8	0.9	4.7	2.1	0.8	0.9
	0.5	1.5	0.7	0.4	0.4	3.0	1.1	0.4	0.4
100	10.0	12.0	9.8	9.0	9.9	14.5	10.8	9.6	9.8
	5.0	6.2	5.0	4.6	5.0	8.1	5.7	4.9	5.0
	1.0	1.6	1.0	0.9	1.0	2.2	1.3	1.0	1.0
	0.5	0.8	0.5	0.4	0.5	1.3	0.6	0.5	0.5

Table 4. Null rejection rates, model with one factor and three levels,  $p = 2$  and  $k = 3$ .

$n$	$\alpha$	$LR$	$LR^*$	$LR_m$	$LR_m^*$
9	10.0	28.2	14.4	5.4	6.8
	5.0	18.6	8.1	2.4	3.2
	1.0	7.4	2.2	0.3	0.5
	0.5	5.0	1.2	0.1	0.2
18	10.0	16.6	10.6	8.5	9.2
	5.0	9.6	5.8	4.2	4.7
	1.0	3.1	1.3	0.8	0.9
	0.5	1.9	0.7	0.4	0.5
	10.0	14.6	10.6	9.3	9.9
27	5.0	8.2	5.5	4.7	5.0
	1.0	2.1	1.1	0.9	1.0
	0.5	1.2	0.6	0.4	0.5
36	10.0	12.8	10.1	9.4	9.8
	5.0	7.2	5.3	4.7	5.0
	1.0	1.8	1.1	0.9	1.1
	0.5	1.0	0.5	0.4	0.5
45	10.0	12.3	10.1	9.3	9.7
	5.0	6.5	5.2	4.7	5.0
	1.0	1.5	1.1	1.0	1.1
	0.5	0.9	0.5	0.5	0.5

and  $k = 3$ . The number of observations from each of the three populations was  $n_1, n_2, n_3$  ( $n_1 + n_2 + n_3 = n$ ). The simulation was performed using  $n_1 = n_2 = n_3 = n/3$ . The figures in Table 4 yield conclusions that are in agreement with the ones drawn from Tables 1, 2 and 3.

Table 5 collects results obtained from simulations carried out under the alternative hypothesis (heteroskedasticity) for  $n = 30$ ,  $p = 2$  and different values of  $\delta_1 = \delta_2 = \delta$ . It is noteworthy that these power simulations correspond to the setting in Table 3 for  $n = 30$ . We only compare the power of the modified profile likelihood ratio test and its Bartlett-corrected variant, since the remaining two tests are considerably oversized and cannot be recommended. The results indicate that there is no loss in power derived from using the Bartlett adjustment derived in this paper. The power of the two tests are similar, with the slight advantage of the corrected test stemming from its smaller size distortion.

Table 5. Nonnull rejection rates,  $n = 30$ ,  $p = 2$ .

$\delta$	$k = 5$		$k = 8$	
	$LR$	$LR^*$	$LR_m$	$LR_m^*$
0.5	9.6	10.5	6.9	7.7
1.0	26.3	28.2	15.0	16.6
1.5	47.2	49.2	28.2	29.0
2.0	76.7	78.2	30.6	31.8
2.5	83.0	84.0	39.0	40.9
3.0	97.4	97.6	46.9	48.5
3.5	98.8	98.9	91.3	91.8

Overall, the likelihood ratio test can be severely liberal, overrejecting the null hypothesis more often than expected based on the selected nominal level for the test. The Bartlett correction to the likelihood ratio test does bring the empirical null rejection rate closer to the nominal level of the test, but it does not fully correct its liberal tendency. The modified profile likelihood ratio test introduces, on the other hand, an overcorrection, thus delivering an undersized test. The Bartlett adjustment to this test brings the size distortions closer to zero; it yields the most accurate of the four tests.

## 5. AN APPLICATION

We shall consider the data analyzed by Montgomery, Peck and Vining (2001, p. 76). The application involves a soft drink bottler who wishes to predict the amount of time required to service the vending machines in an outlet. The service activity includes stocking

the machines with cans and bottles of beverages and performing minor maintenance or housekeeping. The response ( $y$ ) is the time (in minutes) spent on servicing soft drinks machines, and the covariates are the number of beverage cases stocked ( $x_2$ ) and the distance (in feet) travelled ( $x_3$ ). The model used is  $y_l = \beta_1 + \beta_2 x_{l2} + \beta_3 x_{l3} + u_l$ ,  $l = 1, \dots, 25$ . We assume that  $u_l \sim \mathcal{N}(0, \sigma^2 \exp\{\delta_1 x_{l2} + \delta_2 x_{l3}\})$  and  $\text{cov}(u_l, u_m) = 0$  for all  $l \neq m$ . Standard diagnostic analyses suggest that observations 9 and 22 are leverage points, the former also classifying as an influential observation; see Montgomery, Peck and Vining (2001, pp. 210, 213, 215, 216, 217). In what follows we shall work with the incomplete dataset, i.e., we shall exclude observations 9 and 22 from the data. The maximum likelihood parameter estimates are  $\hat{\beta}_1 = 4.643$ ,  $\hat{\beta}_2 = 1.456$  and  $\hat{\beta}_3 = 0.011$ . The estimate of  $\sigma^2$  is  $\hat{\sigma}^2 = 6.163$  and the coefficient of determination,  $R^2$ , equals 0.9072. These estimates were obtained imposing  $\delta_1 = \delta_2 = 0$ . Diagnostic analyses for the incomplete data set do not show evidence of heteroskedasticity.

Our main interest lies in testing  $\mathcal{H}_0 : \delta_1 = \delta_2 = 0$  (homoskedasticity) against a two-sided alternative (heteroskedasticity). Rejection of the null hypothesis would suggest that the non-constant response variance should be modelled as well. The test statistics are  $LR = 4.825$ ,  $LR^* = 3.717$ ,  $LR_m = 4.126$  and  $LM_m^* = 4.352$ ; the respective  $p$ -values are 0.090, 0.156, 0.127 and 0.114. It then follows that the standard likelihood ratio test yields rejection of the null hypothesis (thus suggesting the presence of heteroskedasticity) at the 10% nominal level whereas the null of homoskedasticity is not rejected, at the same nominal level, when the remaining tests (i.e., those based on  $LR^*$ ,  $LR_m$  and  $LM_m^*$ ) are used. That is, the Bartlett-corrected, the adjusted profile likelihood ratio and the corrected adjusted profile likelihood ratio tests lead to inference different from that reached by the standard likelihood ratio test.

## 6. CONCLUDING REMARKS

Practitioners commonly test for the presence of heteroskedasticity when estimating linear regression structures in order to decide what is the appropriate modeling strategy. It is thus important to devise reliable tests for detecting nonconstant conditional variances. In this paper we have derived a Bartlett correction that can be applied to likelihood ratio statistics obtained using modified profile likelihood. The results allow for skedastic functions that involve more than one parameter, and thus generalize results available in the literature. The numerical results presented compared the finite-sample performance of four tests, namely: the original likelihood ratio test, its Bartlett-corrected version, the modified profile likelihood ratio test, and the Bartlett-adjusted modified profile likelihood

ratio test we proposed. Overall, the numerical results favor the latter, i.e., they favor the test obtained from applying a Bartlett correction to the statistic of the modified profile likelihood ratio test. In short, the modification of the profile likelihood attenuates the effects of nuisance parameters on the inference, and the Bartlett correction yields a faster convergence rate of the test statistic to its first order asymptotic distribution. We, therefore, encourage practitioners to use the test proposed in the present paper in applications.

## ACKNOWLEDGEMENT

We gratefully acknowledge partial financial support from CAPES, CNPq, FAPESP and FINEP.

## REFERENCES

- Barndorff-Nielsen, O.E. & Hall, P. (1988). On the level-error after Bartlett adjustment of the likelihood ratio statistic. *Biometrika*, 75, 374–378.
- Cordeiro, G.M. (1993). Bartlett corrections and bias correction for two heteroscedastic regression models. *Communications in Statistics, Theory and Methods*, 22, 169–188.
- Cox, D.R. & Reid, N. (1987). Parameter orthogonality and approximate conditional inference. *Journal of the Royal Statistical Society B*, 49, 1–39.
- Cribari-Neto, F. & Cordeiro, G.M. (1996). On Bartlett and Bartlett-type corrections. *Econometric Reviews*, 15, 339–367.
- DiCiccio, T.J. & Stern, S.E. (1994). Frequentist and Bayesian Bartlett correction of test statistics based on adjusted profile likelihoods. *Journal of the Royal Statistical Society B*, 56, 397–408.
- Doornik, J.A. (2001). *Ox: an Object-oriented Matrix Programming Language*, 4th ed. London: Timberlake Consultants and Oxford: <http://www.nuff.ox.ac.uk/Users/Doornik/>.
- Ferrari, S.L.P. & Cribari-Neto, F. (2002). Corrected modified profile likelihood heteroskedasticity tests. *Statistics and Probability Letters*, 57, 353–361.
- Lawley, D.N. (1956). A general method for approximating to the distribution of the likelihood ratio criteria. *Biometrika*, 71, 233–244.
- Montgomery, D.C., Peck, E. A. & Vining, G.G. (2001). *Introduction to Linear Regression Analysis*, 3rd ed. New York: Wiley.
- Simonoff, J.S. & Tsai, C.-H. (1994). Use of modified profile likelihood for improved tests of constancy of variance in regression. *Applied Statistics*, 43, 357–370.



## ÚLTIMOS RELATÓRIOS TÉCNICOS PUBLICADOS

- 2002-01 - BUENO, V.C. Asymptotic reliability for a coherent system with an infinite number of components. 2002. 10p. (RT-MAE-2002-01)
- 2002-02 - TAVARES, H.R., ANDRADE, D.F. Item Response Theory for Longitudinal Data: Item and Population Ability Parameters Estimation. 2002. 20p. (RT-MAE-2002-02)
- 2002-03 - BOLFARINE, H., VALENÇA, D.M. Score tests for Weibull-regression models with random effect. 2002. 16P. (RT-MAE-2002-03)
- 2002-04 - BUENO, V.C. Using a second order stochastic dominance for active redundancy allocation in a K-out-of-n system. 2002. 7p. (RT-MAE-2002-4)
- 2002-05 - FERRARI, S.L.P., LUCAMBIO, F., CRIBARI-NETO, F. Adjusted and Bartlett-Adjusted profile likelihood ratio tests. 2002. 19p. (RT-MAE-2002-05).
- 2002-06 - PEREIRA, C.A.B., STERN, J.M. Full Bayesian Significance Test: Invariant Formulation. 2002. 10p. (RT-MAE-2002-06).
- 2002-07 - BUENO, V.C. Optimal arrangement of components for systems of dependent components. 2002. 13p. (RT-MAE-2002-07)
- 2002-08 - KOLEV, N., PAIVA, D. Multinomial Latent Model for Random Sums. 2002. 9p. (RT-MAE-2002-08)
- 2002-09 - BUENO, V.C. A Note on a Hazard Processes Ordering: Application on a K-out-of-N System. 2002. 9P. (RT-MAE-2002-09)
- 2002-10 - FERRARI, S.L.P., DAVID, J.S.E., ANDRÉ, A.P., PEREIRA, L.A.A. Use of overdispersed regression models in analysing the association between air pollution and human health. 2002. 15p. (RT-MAE-2002-10)

2002-11 - PERES, A.A.Q., BOTTER, D.A.B. Association between air pollution and human health: filling in the gaps of the pollutant series. 2002. 13p. (RT-MAE-2002-11)

2002-12 - CYSNEIROS, F.J.A., PAULA, A.G. One-Sided Tests in Linear Models with Multivariate t-Distribution. 2002. 33p. (RT-MAE-2002-12)

2002-13 - CRIBARI-NETO, F., FERRARI, S.L.P., OLIVEIRA, W.A.S.C. A numerical evaluation of tests based on different heteroskedasticity-consistent covariance matrix estimators. 2002. 19p. (RT-MAE-2002-13)

2002-14 - BARROSO, L.P., CORDEIRO, G.M. Bartlett Corrections in Heteroskedastic t Regression Models. 2002. 12p. (RT-MAE-2002-14)

2002-15 - KOLEV, N. Volodya, I Miss You (Two Correlated Collective Risk Models) 2002. 20p. (RT-MAE-2002-15)

2002-16 - BUENO, V.C. Burn-in a coherent system in a random environment. 2002. 7p. (RT-MAE-2002-16)

The complete list of "Relatórios do Departamento de Estatística", IME-USP, will be sent upon request.

Departamento de Estatística  
IME-USP

Caixa Postal 66.281

05311-970 - São Paulo, Brasil