



## Correction: Exudate droplets incorporated on eggs by *Raoiella indica* Hirst female during oviposition may avoid the predation of *Amblyseius largoensis* (Muma)

Érica C. Calvet<sup>1</sup> · Nataly de La Pava<sup>2</sup> · Ramony K. B. Oliveira<sup>1</sup> · Arodí P. Favaris<sup>3</sup> · José M. S. Bento<sup>3</sup> · Manoel G. C. Gondim Jr<sup>4</sup>

Published online: 16 October 2024

© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2024

### Correction: Experimental and Applied Acarology

<https://doi.org/10.1007/s10493-024-00964-x>

In this article there was a typo in the name of the last author Manoel G. C. Gondim Jr.  
The original article has been corrected.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s10493-024-00964-x>.

✉ Érica C. Calvet  
[ericacalvet@gmail.com](mailto:ericacalvet@gmail.com)

<sup>1</sup> Department of Phytotechnics, Federal University of Ceará, Fortaleza, Ceará, Brazil

<sup>2</sup> Faculty of Engineering, Agronomic Engineering program, University of Magdalena, Magdalena, Santa Marta, Colombia

<sup>3</sup> Department of Entomology and Acarology, Luiz de Queiroz College of Agriculture, University of São Paulo, Piracicaba, São Paulo, Brazil

<sup>4</sup> Department of Agronomy– Entomology, Federal Rural University of Pernambuco, Recife, Pernambuco, Brazil