

TECHNICAL REPORT • OPEN ACCESS

Tools for estimating fake/non-prompt lepton backgrounds with the ATLAS detector at the LHC

To cite this article: G. Aad *et al* 2023 *JINST* **18** T11004

View the [article online](#) for updates and enhancements.

You may also like

- [Sensor response and radiation damage effects for 3D pixels in the ATLAS IBL Detector](#)
G. Aad, E. Aakvaag, B. Abbott et al.
- [Fast *b*-tagging at the high-level trigger of the ATLAS experiment in LHC Run 3](#)
G. Aad, B. Abbott, K. Abeling et al.
- [Development of the CMS detector for the CERN LHC Run 3](#)
A. Hayrapetyan, A. Tumasyan, W. Adam et al.



ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

247th ECS Meeting
Montréal, Canada
May 18-22, 2025
Palais des Congrès de Montréal

Showcase your science!

Abstract submission deadline extended: December 20

ECS UNITED

RECEIVED: November 30, 2022

REVISED: February 10, 2023

ACCEPTED: February 27, 2023

PUBLISHED: November 30, 2023

TECHNICAL REPORT

Tools for estimating fake/non-prompt lepton backgrounds with the ATLAS detector at the LHC



The ATLAS collaboration

E-mail: atlas.publications@cern.ch

ABSTRACT: Measurements and searches performed with the ATLAS detector at the CERN LHC often involve signatures with one or more prompt leptons. Such analyses are subject to ‘fake/non-prompt’ lepton backgrounds, where either a hadron or a lepton from a hadron decay or an electron from a photon conversion satisfies the prompt-lepton selection criteria. These backgrounds often arise within a hadronic jet because of particle decays in the showering process, particle misidentification or particle interactions with the detector material. As it is challenging to model these processes with high accuracy in simulation, their estimation typically uses data-driven methods. Three methods for carrying out this estimation are described, along with their implementation in ATLAS and their performance.

KEYWORDS: Analysis and statistical methods; Particle identification methods

ARXIV EPRINT: [2211.16178](https://arxiv.org/abs/2211.16178)



© 2023 CERN for the benefit of the ATLAS collaboration. Published by IOP Publishing Ltd on behalf of Sissa Medialab. Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

<https://doi.org/10.1088/1748-0221/18/11/T11004>

Contents

1	Introduction	1
2	Methods	1
2.1	Matrix method	2
2.1.1	Asymptotic matrix method	3
2.1.2	Poisson likelihood matrix method	3
2.2	Fake-factor method	4
2.3	Generalisation for multi-lepton final states	5
2.4	Use with weighted events	6
2.5	Performance studies	7
3	The ATLAS detector	10
4	Lepton selection criteria	11
4.1	Electron reconstruction and identification	11
4.2	Muon reconstruction and identification	12
4.3	Lepton isolation	12
4.4	Removing overlaps between jets and leptons	14
5	Monte Carlo simulation samples	14
6	Sources of fake/non-prompt leptons	15
7	Measurement of real and fake/non-prompt lepton efficiencies	18
7.1	Real-lepton efficiencies	19
7.2	Fake/non-prompt lepton efficiencies	19
8	Systematic uncertainties	21
8.1	Statistical uncertainties in the measured efficiencies	22
8.2	Systematic uncertainties in the measured efficiencies	22
8.3	Uncertainties in the modelling of real-lepton processes	23
8.4	Uncertainties due to biases in the Poisson likelihood matrix method	23
9	Examples of application in ATLAS analyses	23
9.1	Measurement of the $t\bar{t}Z$ cross-section in final states with three or four leptons	23
9.1.1	Real-lepton efficiencies	24
9.1.2	Fake/non-prompt lepton efficiencies	24
9.1.3	Results in the fake/non-prompt lepton validation regions	27
9.2	Model-independent search for new phenomena in multi-lepton final states	29
9.2.1	Fake/non-prompt lepton selection	30
9.2.2	Fake factors	31
9.2.3	Validation regions	33

1 Introduction

Many measurements and searches for new physics performed with the ATLAS detector [1] at the Large Hadron Collider (LHC) at CERN require the presence of one or more leptons (electrons or muons) to indicate that a high-energy electroweak process occurred in the collision. Lepton candidates are reconstructed from signals in the inner tracker, calorimeters, and muon spectrometer [2, 3]. Identification criteria are then applied to suppress candidates originating from physical objects other than leptons from the hard-scattering process of the event. These background candidates fall into two categories: *i*) “non-prompt leptons” from the semileptonic decay of hadrons, or from photon (γ) conversions in detector material, and *ii*) “fake leptons” where the reconstructed object is not, in fact, due to a lepton. In contrast to the aforementioned categories, “real leptons” are defined as electrons or muons produced either directly in the hard-scattering process or directly in the decay of a short-lived non-hadronic resonance (such as a W/Z boson).

The rates at which fake or non-prompt leptons are selected are difficult to model accurately from simulation. They can depend strongly on details of the physics simulation, including in non-perturbative regions where the simulation would not be expected to be reliable. They also depend on the modelling of the material composition and response of the detector. In addition, fake leptons sometimes occur with low probability as a result of misidentifying a hadronic jet in multi-jet events. The computing resources required to simulate these processes with a sufficient sample size would be prohibitive. Therefore, “data-driven” approaches are commonly used to estimate these backgrounds.

To simplify the adoption of such methods, and to ensure that they are applied uniformly, a set of standard tools and prescriptions has been developed for use in ATLAS physics analyses that are subject to fake/non-prompt lepton backgrounds. The principles and performance of these tools are described in this paper. The motivation and mathematical basis of the methods are explained in section 2; a description of the relevant features of the ATLAS detector is given in section 3; the criteria used to select leptons are given in section 4; the simulated signal and background processes, as well as the different processes that can lead to fake/non-prompt leptons, are discussed in sections 5 and 6; the procedures used to measure the efficiencies for real and fake/non-prompt leptons are described in section 7; the systematic uncertainties associated with the methods are described in section 8; and section 9 provides examples of the application of the fake/non-prompt lepton estimation methods in two published ATLAS physics analyses, with details that are not included in the existing publications.

2 Methods

The fake/non-prompt lepton estimation methods considered in this paper depend on defining two tiers of lepton selection criteria, called the “baseline” and “tight” criteria. The tight criteria are

used to select the signal leptons in a physics analysis, while the baseline criteria accept all of the tight lepton candidates as well as an additional set of candidates with a higher rate of fake/non-prompt contributions. Candidates that satisfy the baseline criteria but not the tight criteria are called “loose” leptons. If the two sets of criteria are chosen well, the fraction of real leptons in the baseline sample that satisfy the tight criteria will be substantially higher than the corresponding fraction for fake/non-prompt leptons. These fractions are called the “real efficiency” (ε_r) and “fake efficiency” (ε_f), respectively.

The ε_r values are generally taken from Monte Carlo (MC) simulated events that are corrected to account for differences between data and the simulation, while the ε_f values are typically measured in a data sample consisting of different events from the ones that are used for the data analysis, as detailed in section 7.

2.1 Matrix method

With the efficiencies known, a simple counting of the numbers of lepton candidates that satisfy the tight and loose criteria provides an estimate for the number of fake/non-prompt leptons. In the simplest case, where an analysis selects signal events containing exactly one tight lepton candidate and no loose lepton candidates, the relationship between the numbers of tight and loose leptons observed in data and the composition of the sample in terms of real and fake/non-prompt leptons is:

$$\begin{pmatrix} N^t \\ N^l \end{pmatrix} = \begin{pmatrix} \varepsilon_r & \varepsilon_f \\ 1 - \varepsilon_r & 1 - \varepsilon_f \end{pmatrix} \begin{pmatrix} N_r^b \\ N_f^b \end{pmatrix}, \quad (2.1)$$

where N^t and N^l are the numbers of events with tight and loose lepton candidates, and N_r^b and N_f^b are the unknown numbers of real and fake/non-prompt leptons in the baseline sample. In matrix notation, the relationship is given by:

$$\mathbf{N}_{\text{tl}} = \mathbf{M}_{\varepsilon} \mathbf{N}_{\text{rf}}^b. \quad (2.2)$$

The fact that the unknown values (N_r^b and N_f^b) and the observed yields (N^t and N^l) are related via the matrix $\mathbf{M}_{\varepsilon}$ gives rise to the name of this method: the “matrix method” [4]. Inversion of the matrix allows N_f^b to be determined:

$$N_f^b = \frac{1}{\varepsilon_r - \varepsilon_f} [(\varepsilon_r - 1)N^t + \varepsilon_r N^l]. \quad (2.3)$$

In the typical use case, the quantity of interest is the number of events in the tight sample where the lepton is fake/non-prompt, N_f^t . This is related to the number of such events in the baseline sample, N_f^b , by:

$$N_f^t = \varepsilon_f N_f^b. \quad (2.4)$$

Similarly, the number of real leptons in the tight sample is

$$N_r^t = \varepsilon_r N_r^b, \quad (2.5)$$

and these can be treated as elements of a column matrix $\mathbf{N}_{\text{rf}}^t$.

The fact that N^t appears in eq. (2.1) means that information about the content of the analysis signal region is used in the estimate, and therefore an analysis is not completely blinded when using this approach.

Generally, the values of ε_r and ε_f depend on factors such as the lepton candidate's momentum or its proximity to other objects. Details of how these variations are accounted for in the estimation are given below.

2.1.1 Asymptotic matrix method

In this method, events in the baseline sample are considered one at a time, and a “fake weight” w_i is defined for each event, corresponding to the two terms in eq. (2.3) via eq. (2.4):

$$w_i = \begin{cases} \frac{\varepsilon_{fi}}{\varepsilon_{ri} - \varepsilon_{fi}}(\varepsilon_{ri} - 1) & \text{if lepton candidate } i \text{ is tight (so } N^t = 1 \text{ and } N^l = 0) \\ \frac{\varepsilon_{fi}}{\varepsilon_{ri} - \varepsilon_{fi}}\varepsilon_{ri} & \text{if lepton candidate } i \text{ is loose (so } N^t = 0 \text{ and } N^l = 1) \end{cases},$$

where ε_{ri} and ε_{fi} are the values of ε_r and ε_f that are appropriate for lepton i . Since ε_{ri} is always less than one, the weight for an event with a tight lepton candidate is negative. By extension, the total fake/non-prompt lepton background in the tight sample is then estimated by

$$N_f^t = \sum_{\text{events}} w_i.$$

This approach is convenient since the w_i need only be calculated once and then can be stored with the event, allowing the distribution of the fake-lepton yield to be binned in any variable of interest, as well as a simple re-computation of N_f^t if the event selection is modified. One drawback is that since the value of w_i may be negative, there is no guarantee that N_f^t will be positive. The value of N_f^t is also sensitive to fluctuations in the input ε_{ri} and ε_{fi} values.

The statistical uncertainty of N_f^t is given by:

$$\sigma_{N_f^t} = \sqrt{\sum_{\text{events}} w_i^2}. \quad (2.6)$$

The method that makes use of the w_i is known as the “asymptotic matrix method”, since eq. (2.6) is only valid in the asymptotic limit with a large number of events.

2.1.2 Poisson likelihood matrix method

In this method [5]¹ the elements of $\mathbf{N}_{\text{rf}}^t$ are treated as free parameters, which are varied to maximise the likelihood of the observed $\mathbf{N}_{\text{tl}}$ values. By doing so, the Poisson-distributed nature of the $\mathbf{N}_{\text{tl}}$ values is taken into account, so there is no need to use the asymptotic approximation. In the fit, the $\mathbf{N}_{\text{rf}}^t$ values are converted to $\mathbf{N}_{\text{rf}}^b$ using eqs. (2.4) and (2.5), where the entries in the matrix $\mathbf{M}_{\varepsilon}$ are calculated using the averages of the prompt and fake/non-prompt lepton efficiencies in the baseline sample:

$$\mathbf{M}_{\varepsilon} = \begin{pmatrix} \langle \varepsilon_r \rangle & \langle \varepsilon_f \rangle \\ 1 - \langle \varepsilon_r \rangle & 1 - \langle \varepsilon_f \rangle \end{pmatrix}. \quad (2.7)$$

¹An earlier variant of the likelihood matrix method is described in ref. [6].

The resulting N_{rf}^{b} values are used to obtain the expectation values for N_{tl} using eq. (2.2). These expectation values are denoted by $N_{\text{tl,exp}}$. The N_{rf}^{t} parameters are adjusted (subject to the constraint that they must be non-negative) to maximise the joint Poisson likelihood

$$L(N_{\text{rf}}^{\text{t}}) \equiv \prod_i P[N_{\text{tl}_i} | N_{\text{tl,exp}_i}(N_{\text{rf}}^{\text{t}})], \quad (2.8)$$

where the product is over the elements of N_{tl} ; N_{tl_i} and $N_{\text{tl,exp}_i}$ are the i th elements of N_{tl} and $N_{\text{tl,exp}}$, respectively; and $P[n|\mu]$ is the Poisson probability for observing n events when μ are expected. The output of the fit consists of an estimate of the number of fake/non-prompt leptons and the uncertainty in this quantity, which is obtained by noting the value for which $-\ln L$ exceeds its minimum by 0.5.

The primary advantages of the Poisson likelihood approach are that the result is constrained to be non-negative, and the uncertainty is a better approximation to the range that gives 68% coverage, particularly in samples with few events. In addition, in some scenarios it can provide smaller statistical uncertainties than the asymptotic matrix method or the fake-factor method (described in section 2.2). The main drawback is that the estimated yield must be calculated for the sample as a whole rather than from a sum of individual event weights, which complicates the process of producing a distribution of the fake-lepton yield, as required for any differential measurement (to do so, the likelihood must be applied in every bin of the distribution).

2.2 Fake-factor method

The fact that real-lepton kinematic distributions and efficiencies are generally modelled well in simulation, and that scale factors can be applied to account for any differences observed between the values in simulation and in data control samples, leads to an alternative method that uses simulation, rather than the data, to measure the real-lepton contribution to the loose lepton sample.

The number of fake-lepton events in the loose sample is

$$N_{\text{f}}^{\text{l}} = N^{\text{l}} - N_{\text{r}}^{\text{l}} = (1 - \varepsilon_{\text{f}})N_{\text{f}}^{\text{b}}$$

and the number of fake-lepton events in the tight sample is

$$N_{\text{f}}^{\text{t}} = \varepsilon_{\text{f}}N_{\text{f}}^{\text{b}} = F(N^{\text{l}} - N_{\text{r}}^{\text{l}}),$$

where the “fake factor” F is defined as $F \equiv \varepsilon_{\text{f}}/(1 - \varepsilon_{\text{f}})$. Thus, in the fake-factor method, the number of tight fake/non-prompt leptons for a given analysis can be computed using the fake factor F , the total number of loose lepton candidates N^{l} , and the number of real leptons in the loose lepton sample N_{r}^{l} , where the latter quantity can be estimated using MC simulated samples, and its contribution subsequently subtracted from the quantity FN^{l} observed in the data. In practice, the calculation is performed on an event-by-event basis to account for potential variations in F due to properties of the lepton:

$$N_{\text{f}}^{\text{t}} = \sum_{\text{data}, i=1}^{N^{\text{l}}} F_i - \sum_{\text{MC}, j=1}^{N_{\text{MC}}^{\text{l}}} w_{\text{MC}j} F_j,$$

where F_i is the fake factor appropriate for lepton i , all sources of prompt leptons are considered in the sum over MC simulated events, N_{MC}^{l} is the number of MC events in the loose sample, and $w_{\text{MC}j}$

is the weight assigned to simulated event j , based on the cross-section of the simulated process and any corrections to the selection efficiency that may be needed to reflect the performance on data events.

The main advantage of the fake-factor method is that this result does not depend on N^t , i.e. the yield in the analysis signal region. This means that unlike the matrix method, the fake-factor method can be applied while remaining “blind” to the contents of the signal region. However, the method does have some of the same drawbacks as the asymptotic matrix method, namely the possibility of N_f^t being negative, and sensitivity to fluctuations in the ε_f values.

2.3 Generalisation for multi-lepton final states

The above methods can be generalised to cases where multiple baseline lepton candidates are considered in each event. For the matrix methods, this is done by increasing the dimensionality of $\mathbf{M}_\varepsilon$, $\mathbf{N}_{\text{tl}}$, and $\mathbf{N}_{\text{rf}}^{(t,b)}$ to 2^{n_b} , where n_b is the number of baseline lepton candidates in each event. The estimated fake/non-prompt yield depends on the requirements of a particular analysis in three ways: first, from the requirement placed on the desired number n_t of tight lepton candidates per event; second, whether or not events with additional loose lepton candidates are vetoed; and third, on the minimum number of fake/non-prompt leptons m_f defining the background to be evaluated with one of the data-driven methods.² The consideration of m_f is reflected in the transition from the number of fake/non-prompt lepton events in the baseline sample to the number in the tight sample, in a generalisation of eq. (2.4):

$$N_f^t = \sum_i g_i(\varepsilon_r, \varepsilon_f) N_{\text{rf}i}^b. \quad (2.9)$$

Here, the sum is over all combinations of real and fake/non-prompt leptons that include at least m_f fake/non-prompt leptons, g_i is a function of the real and fake/non-prompt efficiencies that will result in the required number of tight lepton candidates for a given set of real and fake/non-prompt leptons, and $N_{\text{rf}i}^b$ is the i th element of $\mathbf{N}_{\text{rf}}^b$.

To address the requirements on n_t and the possible presence of additional loose lepton candidates, the analysis must consider all events in the baseline sample with lepton multiplicities up to the sum of the allowed numbers of tight and loose lepton candidates in the signal region.

As an example, consider the case where $n_b = 2$. If an analysis selects signal events containing exactly two tight lepton candidates and no loose lepton candidates, and the background with $m_f \geq 1$ is being evaluated, then

$$N_f^t = \varepsilon_{r1}\varepsilon_{f2}N_{r_1f_2}^b + \varepsilon_{f1}\varepsilon_{r2}N_{f_1r_2}^b + \varepsilon_{f1}\varepsilon_{f2}N_{f_1f_2}^b,$$

where $N_{r_1f_2}^b$ is the number of events in the baseline sample where the first lepton candidate is real and the second is fake/non-prompt, and $N_{f_1r_2}^b$ and $N_{f_1f_2}^b$ are defined correspondingly. The ordering of the lepton candidates is typically according to p_T , but the method does not depend on the ordering used.

²An example of a case where m_f is greater than one would be a dilepton analysis where there are backgrounds from both $W(\rightarrow \ell\nu) + j$ events where the jet forms a fake/non-prompt lepton, and dijet events where both jets form fake/non-prompt leptons. The analysers may choose to estimate the first contribution from MC simulation, and use data-driven methods for the second contribution, and therefore setting $m_f = 2$ for the data-driven approach is required to avoid double-counting.

Another example may be an analysis accepting dilepton events ($n_b = 2$) with one tight lepton and another baseline (not necessarily tight) lepton. If, again, the background being evaluated consists of events with at least one fake lepton ($m_f \geq 1$), the coefficients g_i in eq. (2.9) change such that the expression becomes:

$$N_f^t = (\varepsilon_{r1} + \varepsilon_{f2} - \varepsilon_{r1}\varepsilon_{f2})N_{r1f2}^b + (\varepsilon_{f1} + \varepsilon_{r2} - \varepsilon_{f1}\varepsilon_{r2})N_{f1r2}^b + (\varepsilon_{f1} + \varepsilon_{f2} - \varepsilon_{f1}\varepsilon_{f2})N_{f1f2}^b,$$

where the additional terms are needed to account for the additional ways that fake/non-prompt leptons might satisfy the signal selection (and the terms involving products of the efficiencies are subtracted to avoid double-counting). For an analysis that imposes the same lepton candidate requirements but only the background with $m_f = 2$ is being evaluated, the expression is:

$$N_f^t = (\varepsilon_{f1} + \varepsilon_{f2} - \varepsilon_{f1}\varepsilon_{f2})N_{f1f2}^b.$$

The methods can also be extended to cases where there are more than two levels of lepton selection criteria, or distinct categories of fake/non-prompt leptons, as in ref. [7].

As with the matrix method, the fake-factor method can be generalised to higher lepton candidate multiplicities. In the dilepton final state, the number of events with two tight lepton candidates, of which at least one is fake/non-prompt, is:

$$N_f^t = N^{t1t2} - N_{r1r2}^{t1t2} = F_1(N^{l1t2} - N_{r1r1}^{l1t2}) + F_2(N^{t1l2} - N_{r1r2}^{t1l2}) - F_1F_2(N^{l1l2} - N_{r1r2}^{l1l2}), \quad (2.10)$$

where $N^{t(1)t(1)2}$ is the number of events where the first lepton candidate is tight (loose) and the second lepton candidate is tight (loose), $N_{r1r2}^{t(1)t(1)2}$ is the contribution to $N^{t(1)t(1)2}$ from events where both lepton candidates are real leptons, and F_1 and F_2 correspond to the fake factors associated with the first and second lepton candidate, respectively.

However, the algebraic simplification that leads to eq. (2.10), where the result depends simply on products of the fake factors and the observed tight and loose lepton candidate yields, with a correction term that depends only on events where all the leptons are real, does not hold for all possible event selections nor all values of m_f ; such a simplification is restricted to cases where the baseline and tight candidate lepton multiplicities are the same in all events, and where $m_f = 1$.

2.4 Use with weighted events

In some cases, such as for self-consistency tests using simulated events, it may be advantageous to weight the events that are input to the fake/non-prompt background estimate. This is straightforward for the fake-factor and asymptotic matrix methods, since the weight returned by the method for each event can be multiplied by the event weight w_{evt_i} . For the likelihood matrix method, this is handled by using the scaled Poisson distribution [8], in which the values of $N_{\text{tl},\text{exp}_i}$ and $N_{\text{tl},\text{evt}_i}$ from eq. (2.8) are scaled according to the event weights w_{evt_i} :

$$s_{\text{tl}} \equiv \frac{\sum_{\text{tl,obs}} w_{\text{evt}_i}}{\sum_{\text{tl,obs}} w_{\text{evt}_i}^2},$$

so that the likelihood becomes

$$L(\mathbf{N}_{\text{rf}}^t) = \prod_i P[N_{\text{tl}_i}/s_{\text{tl}} | N_{\text{tl},\text{exp}_i}(\mathbf{N}_{\text{rf}}^t)/s_{\text{tl}}]$$

2.5 Performance studies

MC simulations of experiments are utilised to assess the statistical performance of the methods. These simulations consist of pseudoexperiments that mimic scenarios that might occur in an actual analysis. Pseudoexperiments with sample sizes of 10 or 1000 dilepton events in the baseline sample are considered. The fraction of fake/non-prompt leptons in the baseline lepton sample is varied for each pseudoexperiment, with a uniform distribution between 0 and 100%. The values of ε_r and ε_f for each lepton are drawn from Gaussian distributions with specified means and widths (limits are imposed such that the values are always between zero and one, and ε_f is always at least 10% less than ε_r). Each lepton is randomly assigned as fake/non-prompt or real, in accord with the fraction of fake/non-prompt leptons assumed for the pseudoexperiment. Then each lepton is judged to either meet or fail to meet the tight selection criteria, based on whether or not it is a real lepton and the values of ε_r and ε_f assigned to it. The set of simulated leptons is input to the data-driven algorithms, and the estimated fake yield and its statistical uncertainty are determined for each pseudoexperiment. These values are then compared with the expectation value for the number of fake/non-prompt lepton events in each pseudoexperiment, which is determined by the numbers of real and fake/non-prompt baseline leptons, the values of ε_r and ε_f for each lepton, and the value of m_f for the simulated analysis. For example, in a dilepton sample with $m_f \geq 1$, the expectation value is:

$$\langle N_f^t \rangle = \sum_{\text{rf events}} \varepsilon_{r1,i} \varepsilon_{f2,i} + \sum_{\text{fr events}} \varepsilon_{f1,i} \varepsilon_{r2,i} + \sum_{\text{ff events}} \varepsilon_{f1,i} \varepsilon_{f2,i}.$$

For the Poisson likelihood matrix method, the likelihood maximisation is implemented using the MINUIT function minimisation package [9] to minimise the negative log likelihood. The interface to MINUIT is provided by the TMinuit class in ROOT [10].

The fake-factor method requires a two-step process, where the contribution from real-lepton events is subtracted in the second step. In an actual physics analysis, this subtraction is done using MC simulation of the prompt lepton contribution. For the simple MC simulation used here, the second step is modelled by running each pseudoexperiment a second time with the same parameters but a statistically independent sample of events, and running the fake-factor method only on the events that do not have fake/non-prompt leptons. The second sample has ten times more events than the primary sample, to be consistent with the usual case where the MC simulation sample has a multiple of the number of events in the data sample. The result of this second run is then scaled down by a factor of ten and subtracted from the result when using the initial set of simulated events.

As an initial example, one can investigate the performance for dilepton events under conditions that are favourable for estimating the fake/non-prompt lepton background. This means that the samples are large (1000 events per pseudoexperiment), and the values of ε_r and ε_f are on average well-separated (here, $\langle \varepsilon_r \rangle = 0.90$ and $\langle \varepsilon_f \rangle = 0.10$). The ratios of the estimated to true fake yields are shown versus $\langle N_f^t \rangle$ in figure 1(a). The average statistical uncertainties in the estimates for each method are shown in figure 1(b), and the fraction of pseudoexperiments in which the true fake yield lies within the uncertainty reported for each method is shown in figure 1(c). The performance of the methods for dilepton analyses with low statistical precision (10 events per pseudoexperiment) are shown in figure 2. Finally, to represent a more challenging situation, the case where there is less separation between the values of ε_r and ε_f (due, for example, to the application of stricter online




Figure 1. Performance of the three methods in pseudoexperiments with dilepton events where the leptons had an average $\varepsilon_r = 0.9$ and $\varepsilon_f = 0.1$, both values were varied according to a Gaussian distribution of width 0.1 when simulating each lepton, and there were 1000 events per pseudoexperiment. The quantity $\langle N_f^t \rangle$ is the expectation value for the number of fake/non-prompt lepton events in each pseudoexperiment. Plot (a) shows the ratio of the estimated to expected fake-lepton yields, (b) shows the absolute uncertainty estimate for each method (for the Poisson likelihood method the average of the upward and downward uncertainties is taken), and (c) shows the fraction of pseudoexperiments where the true fake yield lies within the reported one standard deviation (1σ) range.




Figure 2. Performance of the three methods in pseudoexperiments with dilepton events where the leptons had an average $\varepsilon_r = 0.9$ and $\varepsilon_f = 0.1$, both values were varied according to a Gaussian distribution of width 0.1 when simulating each lepton, and there were 10 events per pseudoexperiment. The quantity $\langle N_f^t \rangle$ is the expectation value for the number of fake/non-prompt lepton events in each pseudoexperiment. Plot (a) shows the ratio of the estimated to expected fake-lepton yields, (b) shows the absolute uncertainty estimate for each method (for the Poisson likelihood method the average of the upward and downward uncertainties is taken), and (c) shows the fraction of pseudoexperiments where the true fake yield lies within the reported one standard deviation (1σ) range.

lepton selection criteria that might be required when the LHC runs at higher luminosities) is also explored. Figure 3 shows the results when $\langle \varepsilon_r \rangle = 0.70$ and $\langle \varepsilon_f \rangle = 0.30$, and there are 10 events per pseudoexperiment.

These studies show that all three methods give accurate estimates, with nearly equivalent performance, in high-statistics samples with a large separation between ε_r and ε_f (as shown in




Figure 3. Performance of the three methods in pseudoexperiments with dilepton events where the leptons had an average $\varepsilon_r = 0.7$ and $\varepsilon_f = 0.3$, both values were varied according to a Gaussian distribution of width 0.1 when simulating each lepton, and there were 10 events per pseudoexperiment. The quantity $\langle N_f^t \rangle$ is the expectation value for the number of fake/non-prompt lepton events in each pseudoexperiment. Plot (a) shows the ratio of the estimated to expected fake-lepton yields, (b) shows the absolute uncertainty estimate for each method (for the Poisson likelihood method the average of the upward and downward uncertainties is taken), and (c) shows the fraction of pseudoexperiments where the true fake yield lies within the reported one standard deviation (1σ) range.

figure 1). One notable feature is the dip in the uncertainty values for all three methods near $\langle N_f^t \rangle = 10$ in figure 1(b). This occurs because there are two ways for the model to produce an expectation around 10: either a very low fake fraction in the baseline to start with, or a very large fake fraction so that there are few real leptons and most of the background is from events with two fake/non-prompt leptons, which gives a minimum value of $\varepsilon_f^2 \cdot 1000 = 10$ when $\varepsilon_f = 0.10$. These two processes will have very different uncertainties. There is also a bias in the Poisson likelihood matrix method toward low values when the true number of fakes is large. This bias arises due to the averaging of efficiencies over the entire baseline sample (see eq. (2.7)) when in fact the efficiencies are on average different for real and fake leptons. Such differences in the averages occur randomly in the ‘toy’ MC tests, but can occur systematically in a physics analysis if, for example, the real and fake efficiencies have different kinematic distributions. The biases can be mitigated by binning the baseline sample in the variables for which the real-lepton and fake-lepton distributions may differ, and performing the likelihood fit separately in each bin. As an example of the effect of such a binning, the pseudoexperiments can be run with the results binned according to the values of ε_f . The effect of using two such bins in the value of ε_f for each lepton is shown in figure 4.

When the situation becomes more challenging, such as in figures 2 and 3, the characteristics of each method become more distinct. For low-statistics samples, the Poisson likelihood approach tends to exhibit a bias toward high values when the true number of fakes is small, a natural consequence of the fact that it cannot return negative values. The coverage of the true value by the estimated uncertainty is, however, still reasonable. A clear distinction between the precision of the methods also appears in figures 2 and 3, where the Poisson likelihood approach has the smallest statistical uncertainty, followed by the asymptotic matrix method and then the fake-factor method. This is because the Poisson likelihood approach considers lepton efficiencies averaged




Figure 4. Ratio of the estimated to expected fake-lepton yields for the Poisson likelihood matrix method in pseudoexperiments with dilepton events where the leptons had an average $\varepsilon_r = 0.9$ and $\varepsilon_f = 0.1$, both values were varied according to a Gaussian distribution of width 0.1 when simulating each lepton, and there were 1000 events per pseudoexperiment. The triangles show the result with a single bin, while the circles show the result where the input events are binned according to the values of ε_f for each lepton. Two bins are used for each of the two leptons, and the results from all four bins are summed to give the estimated N_{fake} value.

over the entire sample and is therefore less susceptible to event-by-event fluctuations in the values of ε_r and ε_f .

Despite the differences between them, all the approaches are valid, as shown by the statistical coverage plots: except in extreme cases, the confidence intervals built from the estimates and their statistical uncertainty do contain the true fake yield in at least 68% of pseudoexperiments, and the visible overcoverage is due mostly to the fact that the uncertainties are computed under the assumption that all of the $\mathbf{N}_{\text{fl}}$ values are independent, while the pseudoexperiments were generated with a fixed number of baseline events for each, which means there was some anticorrelation among the $\mathbf{N}_{\text{fl}}$.

When selecting which method to use, the analyser needs to consider the relative benefits and complexities of implementing the methods, along with the size of the uncertainty in the fake/non-prompt lepton background yield relative to other uncertainties in the analysis.

3 The ATLAS detector

While the above description of the matrix and fake-factor methods is general, the remainder of this paper discusses the application of these methods to ATLAS physics analyses, and therefore a brief description of the experimental apparatus follows.

The ATLAS detector [1] at the LHC covers nearly the entire solid angle around the collision point.³ It consists of an inner tracking detector surrounded by a thin superconducting solenoid,

³ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis

electromagnetic (EM) and hadronic calorimeters, and a muon spectrometer (MS) incorporating three large superconducting toroidal magnets. The inner-detector system is immersed in a 2 T axial magnetic field and provides charged-particle tracking in the range $|\eta| < 2.5$.

The high-granularity silicon pixel detector covers the vertex region and typically provides four measurements per track, the first hit normally being in the insertable B-layer (IBL) installed before Run 2 [11, 12]. It is followed by the silicon microstrip tracker, which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker (TRT), which enables radially extended track reconstruction up to $|\eta| = 2.0$. The TRT also provides electron identification information based on the fraction of hits (typically 30 in total) above a higher energy-deposit threshold corresponding to transition radiation.

The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. Within the region $|\eta| < 3.2$, EM calorimetry is provided by barrel and endcap high-granularity lead/liquid-argon (LAr) calorimeters, with an additional thin LAr presampler covering $|\eta| < 1.8$ to correct for energy loss in material upstream of the calorimeters. Hadronic calorimetry is provided by the steel/scintillator-tile calorimeter, segmented into three barrel structures within $|\eta| < 1.7$, and two copper/LAr hadronic endcap calorimeters. The solid angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimised for EM and hadronic measurements respectively.

The MS comprises separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by the superconducting air-core toroids. The field integral of the toroids ranges between 2.0 and 6.0 Tm across most of the detector. A set of precision chambers covers the region $|\eta| < 2.7$ with three layers of monitored drift tubes, complemented by cathode-strip chambers in the forward region, where the background is highest. The muon trigger system covers the range $|\eta| < 2.4$ with resistive-plate chambers in the barrel, and thin-gap chambers in the endcap regions. Interesting events are selected to be recorded by the first-level trigger system implemented in custom hardware, followed by selections made by algorithms implemented in software in the high-level trigger [13]. The first-level trigger accepts events from the 40 MHz bunch crossings at a rate below 100 kHz, which the high-level trigger reduces in order to record events to disk at about 1 kHz.

An extensive software suite [14] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

4 Lepton selection criteria

Full descriptions of the electron and muon reconstruction algorithms and available selection criteria used in ATLAS are provided in refs. [2] and [3], respectively. Here the features most relevant to the fake/non-prompt lepton background estimation are summarised briefly.

4.1 Electron reconstruction and identification

Electron candidates are reconstructed within $|\eta| < 2.47$ as tracks in the inner detector matched to energy clusters in the EM calorimeter.

points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$.

In order to separate true electrons from other unwanted reconstructed candidates, electron identification (ID) algorithms are used. These rely upon a set of variables that quantify the distribution of energy in the calorimeter, the quality of the spatial match between the calorimeter deposit and the associated track, and the transition radiation signal in the TRT (see table 1 of ref. [2] for a complete list). Rather than place individual requirements on these variables, they are combined into a likelihood discriminant based upon the probability density functions of the variables measured for prompt electrons in $Z \rightarrow e^+e^-$ events and for background candidates reconstructed in inclusive collision events.

Since different analyses have different requirements for electron selection efficiency and background rejection, several “working points” (WPs) are defined by different values of the likelihood. The likelihood threshold values are varied according to the p_T and $|\eta|$ of the electron candidate, so that the selection efficiency varies smoothly with the electron p_T . The most commonly used ID WPs (and their average efficiencies⁴ measured for typical electroweak processes) are “Loose” (93%), “Medium” (88%) and “Tight” (80%). In addition to the listed WPs, there is another (“LooseAnd-BLayer”) WP that imposes the same requirement on the likelihood as the “Loose” WP, but also requires that the electron track have a hit in the IBL to suppress candidates originating from photon conversions.

Often, additional requirements are imposed on the impact parameter of the electron’s track: the impact parameter in the transverse plane, d_0 , with respect to the centre of the beamspot must satisfy $|d_0| < 5\sigma(d_0)$, where $\sigma(d_0)$ is its estimated uncertainty, while the longitudinal separation z_0 between the point where d_0 is measured and the chosen primary vertex of the event, multiplied by a moderating factor $\sin(\theta)$ which accounts for reduced z_0 accuracy for more forward tracks, cannot exceed 0.5 mm in absolute value.

4.2 Muon reconstruction and identification

Muon candidates are reconstructed in the region $|\eta| < 2.5$ by combining MS tracks with matching inner-detector tracks. The muon reconstruction efficiency is approximately 98% per muon in simulated $Z \rightarrow \mu^+\mu^-$ events. After reconstruction, high-quality muon candidates used for physics analyses are selected by a set of requirements on the number of hits in the different inner subdetectors and different MS stations, on the track fit properties, and on variables that test the compatibility of the individual measurements in the two detector systems, as detailed in ref. [3]. These criteria reduce the background from in-flight decays of light-flavour hadrons, which often result in kinked tracks. The most commonly used muon ID WPs (and their efficiencies measured in $t\bar{t}$ MC events) are “Medium” (98%) and “HighPt” (80%), the latter optimised to offer the best momentum resolution for $p_T > 100$ GeV. The same impact parameter requirements as defined for electrons are also often applied to muon candidates, with a tighter condition in the transverse plane: $|d_0| < 3\sigma(d_0)$.

4.3 Lepton isolation

In addition to the ID criteria mentioned above, most analyses place requirements on the isolation of the lepton from other detector activity. This is especially helpful in reducing the contribution from

⁴Those efficiencies, as well as those quoted for muons in the next section, do not include requirements for the leptons to be successfully identified in the context of the trigger decision, which may include small additional inefficiencies [15, 16].

leptons produced in heavy-flavour decays, or muons from $\pi^\pm$ or $K^\pm$ decays within jets⁵ as there are often other components of the jet that are near the lepton in these cases. In all of the methods described in section 2, the baseline lepton selection usually does not require isolation, while the tight lepton selection usually does. However, many of the single-lepton triggers [15, 16] used in ATLAS require isolation, so analyses that depend on such triggers cannot avoid applying isolation requirements for baseline leptons.

The calorimeter isolation [2, 3] is calculated from the sum of transverse energies of calorimeter energy clusters within $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = \text{XX}/100$ of the lepton candidate, not including the contribution expected from the candidate itself. Typical values of XX are 20, 30 or 40. Expected residual contributions to the isolation from the lepton candidate, as well as expected contributions from particles produced by additional proton–proton (pp) interactions, are subtracted [20], resulting in the variable E_T^{coneXX} .

The track-based isolation [2, 3], denoted by p_T^{coneXX} , is based on tracks near the lepton candidate with either $p_T > 0.5$ or 1 GeV that satisfy basic track-quality requirements and are spatially consistent with the primary vertex of the event. The scalar sum of the transverse momenta of such tracks, excluding tracks associated with the lepton candidate, is compared with the p_T of the candidate to assess the isolation. For muon candidates, only the single track associated with the candidate is excluded; for electron candidates, additional tracks consistent with pair-production from a bremsstrahlung photon are also excluded. The track isolation can also be defined with a variable cone size ($p_T^{\text{varconeXX}}$). In this case, the size of the cone around the lepton candidate within which tracks are considered is varied as a function of the p_T of the candidate:

$$\Delta R = \min\left(\frac{10 \text{ GeV}}{p_T [\text{GeV}]}, R_{\text{max}}\right),$$

where R_{max} is the maximum cone size (typically 0.2 or 0.3).

Combining selections on track-based and calorimeter-based isolation provides even better fake/non-prompt lepton background rejection, as the two isolation variables use complementary information. Track-based isolation was found to be less sensitive to detector noise and pile-up effects than calorimeter-based isolation, and the inner detector provides a better p_T measurement than the calorimeters for individual soft hadrons. On the other hand, calorimeter-based isolation includes neutral particles as well as some particles below the inner detector’s track- p_T threshold, which are ignored when computing track isolation. However, track and calorimeter isolation variables measure hadronic activity in a redundant manner, since charged particles are measured by both the calorimeters and the inner detector, and simple selection cuts applied independently to those two variables may not achieve optimal rejection power. To avoid this, an analysis can use a “particle-flow” algorithm, which allows removal of overlapping contributions from the track-based and calorimeter-based isolation, decreasing the correlation between the two variables. For the time being, particle-flow-based isolation variables are defined only for muons, and discussed in ref. [3].

For analyses where the fake/non-prompt lepton background may be dominated by non-prompt electrons and muons from the decays of b - and c -hadrons [21, 22], isolation WPs using a boosted

⁵Jets are reconstructed from clusters of topologically connected calorimeter cells (topo-clusters), as described in ref. [17]. The anti- k_t algorithm [18, 19] is used to form jets from the topo-clusters, with the radius parameter R usually set to 0.4; when reconstructing jets formed from the merged decay products of boosted resonances, a larger R value, typically 1.0, is used. Typically, jets with $p_T > 20$ GeV and $|\eta| < 4.5$ are considered in physics analyses.

decision tree (BDT) discriminant based on isolation and secondary vertex information, referred to as the non-prompt lepton BDT, are also used.

Several isolation WPs based on tracking, a combination of calorimetry and tracking, particle-flow, or a non-prompt lepton BDT are defined to allow for consistency across analyses that require different levels of lepton isolation. They are described in refs. [2] and [3].

4.4 Removing overlaps between jets and leptons

In some cases the same object can result in multiple signatures in the detector. For example, an electron will deposit energy in the calorimeter, and that energy will generally also be clustered into a jet. In addition, sometimes objects will be spatially correlated due to the underlying physics, as when a muon is produced by heavy-flavour decay within a jet. To avoid double-counting, and to select only the isolated leptons that are of interest in physics analyses, an overlap removal procedure is applied to resolve these ambiguities. The procedure used for most physics analyses is as follows, where the lepton candidates are those that satisfy the baseline ID criteria for the analysis:

- All electron candidates that are within $\Delta R = 0.01$ of a muon candidate (or share a track with a muon candidate) are removed.
- All jets that are within $\Delta R = 0.2$ of any remaining electron candidates are removed.
- All electron candidates that are within $\Delta R = 0.4$ of any remaining jet are removed.
- Cases where a remaining jet is within $\Delta R = 0.4$ of a muon candidate are examined to determine the number of tracks associated with the jet. If there are more than two such tracks the muon candidate is removed; otherwise the jet is removed.

Variations of this procedure are also supported, primarily for analyses that focus on heavy-flavour jets or that select boosted massive particles using large- R jets.

5 Monte Carlo simulation samples

The studies of fake/non-prompt leptons that are presented in sections 6 and 7 make use of large MC samples of simulated events. The production of $t\bar{t}$ events at next-to-leading order (NLO) in the quantum chromodynamics (QCD) coupling constant α_s is described in ref. [23] and relies on the POWHEG BOX v2 event generator [24] interfaced with PYTHIA 8.230 [25] for parton showering and subsequent steps, with the A14 set of tuned parameters [26]. The parton distribution functions used for matrix element calculation and parton showering are NNPDF3.0_{NLO} [27] and NNPDF2.3_{LO} [28] respectively. Generated events were filtered such that at least one of the top quarks decays semileptonically. The EVTGEN 1.2.0 program [29] was used to model heavy-flavour hadron decays.

The production of Drell–Yan $Z/\gamma^* \rightarrow \ell^+\ell^-$ events ($\ell = e, \mu, \tau$) at NLO in α_s is described in ref. [30] and relies on the SHERPA 2.2 event generator [31] with the dedicated set of tuned parameters and the NNPDF3.0_{NNLO} [27] parton distribution function. Events were generated according to a partition of the phase space described in ref. [30], resulting in a set of orthogonal samples which

were combined with weights corresponding to the NLO cross-section calculated by the generator. The $\ell^+\ell^-$ invariant mass was required to be at least 40 GeV.

A full ATLAS detector simulation [32] based on GEANT4 [33] was then used to faithfully reproduce particle interactions with the detector and its response. Additional pp interactions in the same or neighbouring bunch crossings were also simulated in order to reproduce the conditions of data-taking during the LHC Run 2 operation. Simulated events are weighted to reproduce the measured performances of various selection criteria, such as electron identification [2]. In addition, reconstructed lepton and jet momenta are transformed [2, 34, 35] to match the real detector resolution.

The source of a lepton candidate in simulation, discussed in section 6, is identified by using preserved generator-level information to match the track of each reconstructed lepton candidate to a suitable stable particle produced by the generator or the GEANT4 simulation, among those preserved in the event record (sections 3.6 and 5.3 in ref. [32]). The matching is performed by evaluating the contribution of each stable particle to the simulated ID and (for muons) MS hits used to reconstruct the track, and requiring that the matched particle accounts for at least half of those hits (section 6.3 in ref. [36]). While the latter is a general procedure used for all tracks, for leptons specifically the result is discarded if the reconstructed track has less than three combined hits in the pixel and SCT detectors, or if the track and the stable particle do not satisfy $\Delta R < 0.2$. If the stable particle is a charged lepton (or, in the case of electron candidates, may also be a photon), its origin is checked so that the different sources of non-prompt leptons can be distinguished.

In section 9, other simulated processes are used. The corresponding MC samples are obtained with a similar workflow. Complete information about the MC generators and their configurations, as well as the cross-section calculations used to normalise the simulated samples, is available in the references provided in that section.

6 Sources of fake/non-prompt leptons

The relative contributions of fake/non-prompt leptons from different sources to the sample of selected lepton candidates depend on the energy and spatial location of the candidate in the detector, the identification, impact parameter and isolation criteria applied, the overlap removal procedure, and the nature of the selected final state (e.g. the presence or absence of heavy-flavour hadrons).

Figures 5 and 6 present for illustration the relative contributions of the different sources of fake/non-prompt muons and electrons respectively, as a function of the transverse momentum of the candidate, as measured in MC simulated events. They are shown for two different processes: $t\bar{t}$ production, leading to final states rich in heavy-flavour hadrons, and Drell–Yan production of e^+e^- or $\mu^+\mu^-$ pairs; for $t\bar{t}$ events at least one of the two W bosons produced in the $t\bar{t}$ decay is required to decay leptonically ($W \rightarrow \ell\nu$). For these particular figures, the reconstruction/ID of electron and muons and the general event selection follow those described in ref. [37] for baseline lepton candidates, which correspond to rather loose criteria (in particular, no isolation nor transverse impact parameter requirements are applied, and no overlap removal is done). For both processes, events are considered only if they contain a pair of reconstructed baseline leptons with identical charges, a signature for which fake/non-prompt leptons usually represent a non-negligible source of background.




Figure 5. Relative contributions of fake or non-prompt muons from different sources as a function of p_T in simulated (a) $t\bar{t}$ and (b) Drell–Yan processes. Events are required to contain at least two leptons with identical charges. Punch-through hadrons are charged hadrons reaching the MS and leading to the reconstruction of a fake muon candidate. Some other minor sources are not displayed; the sum of their contributions is less than 2% in every bin of (a), and less than 4% in bins of (b) up to $p_T = 100$ GeV. The muons are required to satisfy minimal track-quality criteria specified in the text. Error bars represent statistical uncertainties of the simulation.




Figure 6. Relative contributions of fake or non-prompt electrons from different sources as a function of p_T in simulated (a) $t\bar{t}$ and (b) Drell–Yan processes. Events are required to contain at least two leptons with identical charges. Some other minor sources are not displayed; the sum of their contributions is less than 0.5% in every bin. These do not include electron candidates corresponding to high-energy muon ionisation signal in the calorimeters, or from conversions of final-state radiation (FSR) photons with small angular separation from a genuine electron or muon. The electrons are required to satisfy loose ID criteria specified in the text and ref. [37]. Error bars represent statistical uncertainties of the simulation.

The “prompt γ -conversion” category only includes electron candidates where the photon is separated by $\Delta R > 0.1$ from any generator-level high- p_T electron from the hard-scatter interaction (photons emitted at smaller separation are generally reconstructed as part of the electron candidates).

Non-prompt leptons are those arising from electroweak decays of hadrons. Heavy-flavour b - and c -hadrons decay close to the interaction point and the resulting leptons are distinguishable from real leptons mostly by isolation and impact parameter.

Light-flavour hadrons can also be a major source of fake leptons via decay-in-flight in the tracker volume. This happens mainly in final states for which QCD multi-jet production is a significant contributor. A charged hadron stopping early in the calorimeter, and generating a narrower-than-average shower, can mimic the experimental signature of an electron. Electron ID criteria are particularly powerful in rejecting these candidates rather than those from other sources, based notably on three-dimensional profiles of the shower, but since many orders of magnitude more hadrons than leptons are produced in collisions at the LHC, a substantial number of such fake electrons may be selected in a physics analysis. With regard to muons, the depth of the ATLAS calorimeter is sufficient to stop most pions and kaons before they can reach the MS. Muon candidates arising from kaons decaying semileptonically in flight before reaching the MS are a more important contribution to the fake/non-prompt lepton background.

Among hadrons faking electrons, one significant class is neutral pion decays into photons ($\pi^0 \rightarrow \gamma\gamma$); the collimated photons create a single energy deposit in the EM calorimeter, while the associated track might be provided by the conversion of one of the photons in the upstream detector material. Due to the importance of this phenomenon, the electron ID criteria specifically discriminate against these by attempting to identify a two-peak structure in the distribution of the cell energies matched to the electron cluster [2], which is not present for real electrons. Other contributions such as Dalitz decay of pions [38] also create comparable experimental signatures.

The conversion of photons into electron–positron pairs represents the last important class of non-prompt electrons. These γ -conversions must typically occur early on (e.g. in the beam pipe) and be largely asymmetric in the splitting of the momentum between the two electrons; otherwise, the conversion vertex can be reconstructed, or the candidate electron’s track lacks hits in the first layers of the inner detector, both leading to the proper classification of the reconstructed object as a photon instead of an electron [2]. The origin of the photon itself influences the characterisation of the candidate as non-prompt or real: photons emitted close to a real electron, either due to bremsstrahlung or as higher-order quantum electrodynamic (QED) corrections to the production process, are typically considered part of the electron candidate (the calorimeter energy deposits tend to overlap to the extent that a single cluster is reconstructed); furthermore, from the perspective of quantum field theory, well-defined electrons must include extra radiation (“dressed leptons” [39]). The electron reconstruction procedure [2] accounts for bremsstrahlung, in particular by allowing kinks in the track consistent with bremsstrahlung emission in dense material regions. In contrast, photons from other origins, such as initial-state radiation, QED processes not involving leptons or where photons are sufficiently separated from leptons, or hadronic jet fragmentation, may be considered as sources of fake electrons.

It can be seen in figure 5 that non-prompt muons constitute the only substantial contribution to the fake/non-prompt muon background, while for electrons figure 6 shows more variety: in general, non-prompt electrons are particularly represented in the lower p_T range, especially in processes involving the production of heavy-flavour hadrons, while hadron fakes and converted photons populate the higher p_T range.⁶

The different sources of fake/non-prompt leptons have distinct probabilities to satisfy the tight lepton selection criteria described in section 4. Figure 7 illustrates those differences for the particular

⁶However, for the range $p_T < 10$ GeV, not shown in the figure, hadron fakes are also the dominant contribution.




Figure 7. Rejection (defined as $1/\varepsilon_f$) as a function of p_T for different sources of fake/non-prompt (a) muons and (b) electrons in simulated $t\bar{t}$ events containing two leptons with identical charges. The efficiency ε_f is the fraction of lepton candidates selected using the same baseline criteria as in figures 5 and 6 that also pass the tight selection criteria of ref. [37]. Error bars represent statistical uncertainties of the simulation, and for purposes of clarity, values with relative uncertainties greater than 30% are not shown.

example of signal leptons definitions used in ref. [37] (including lepton–jet overlap removal) and simulated $t\bar{t}$ events containing two leptons with identical charges. Such variability between sources is unwelcome, as the fake efficiencies required for the application of the methods described in section 2 then depend upon the relative contributions of each source to the regions of interest, which may not be easy to assess. It is therefore desirable to measure the efficiencies in regions similar in composition to the regions where the background estimate is needed; otherwise large extrapolation uncertainties may apply.

Precise simulation of these various sources of background, including their relative contributions, is indeed very challenging, as it relies heavily on the modelling of the soft-QCD regime by event generators, including modelling of fragmentation and hadronisation processes, hadron decay modelling, and soft emissions and detector modelling. Another issue is that only a small fraction of fake or non-prompt lepton candidates survive the ID and isolation requirements, so the simulation of a very large number of events is needed to obtain a statistically accurate prediction. For inclusive processes with large cross-sections (e.g. multi-jet production), this is often impractical.

For these reasons, many of the fake/non-prompt lepton background predictions used in ATLAS publications are based on methods using the data, such as the ones described in this paper. They rely on common properties shared to some extent by the different sources of fake/non-prompt leptons that differentiate them from real leptons, such as a high likelihood to not meet the combination of ID and isolation criteria.

7 Measurement of real and fake/non-prompt lepton efficiencies

The methods described in section 2 both rely on knowledge of the efficiency for leptons that pass the baseline selection to also pass the tight selection. For the fake-factor method, only the efficiency for fake/non-prompt leptons is used explicitly in the calculation, while for the other methods the efficiency for real leptons must also be measured. In many cases these efficiencies depend on the

properties of the lepton (such as its p_T or angular distance from a jet) or on the event in which it is found (such as the overall activity of the event, as measured for example by the number of reconstructed primary vertices).

7.1 Real-lepton efficiencies

The efficiencies of specific working points of lepton selections are calibrated precisely in ATLAS for general purposes, primarily with “tag-and-probe methods” based e.g. on $Z \rightarrow \ell\ell$ events. By performing these measurements on both data and MC-simulated events, “scale factors” (SFs) that account for differences between the efficiencies observed in data and simulation are derived. These SFs can then be applied to simulated events using the selection criteria that are relevant to a given analysis to determine the appropriate real-lepton efficiencies. This MC-based approach is valid as long as both the baseline and tight lepton selection criteria are taken from the set for which SFs have been measured; only in extraordinary cases would an analysis utilise different selection criteria. Since the efficiencies depend more strongly on the environment than the SFs do, the main advantage of this approach over purely data-based measurements is to allow efficiencies to be obtained directly in the desired environment (i.e. the region in which the fake/non-prompt background estimate is needed), rather than being extrapolated from a more distant region which would be needed for reliable measurements in data.

Details of the real-electron and real-muon efficiency measurements can be found in refs. [2] and [3], respectively. The real efficiencies are often parameterised with respect to the p_T and $|\eta|$ of the leptons, and measured separately for electrons and muons.

7.2 Fake/non-prompt lepton efficiencies

The fake/non-prompt lepton efficiencies are specific to each analysis, primarily since there are several sources for such leptons (see section 6), which will contribute with different weights depending on the chosen selection criteria. In general, though, the first step in the efficiency measurement is to identify a region that has a large contribution from fake leptons. Two approaches are commonly used. In the first, events with a pair of leptons with the same electric charge are selected. Since such lepton pairs are only rarely produced at the LHC (via processes such as WZ + jets, ZZ + jets, and $t\bar{t} + X$ ($X = W/Z$) production) it is likely that one of the two leptons is fake/non-prompt. By placing stringent quality criteria on one lepton in these events, the probability that the remaining lepton is fake/non-prompt is enhanced. The second approach is to use single-lepton events, where criteria are imposed to suppress the contribution from real leptons. Examples of such criteria are requiring the missing transverse momentum⁷ E_T^{miss} , or transverse mass⁸ m_T , to be below specific thresholds, thereby reducing the contribution from W +jets or $t\bar{t}$ events, or requiring via the track impact parameters that the lepton originate from a position inconsistent with the primary event vertex, thereby enhancing the contribution of leptons from heavy-flavour decay.

⁷ E_T^{miss} is defined as the negative vector sum of the p_T of the reconstructed and calibrated objects in the event, with a correction applied for inner detector tracks that originate from the primary collision vertex and are not associated with any other objects, and E_T^{miss} is defined as the magnitude of E_T^{miss} [40].

⁸Here $m_T \equiv \sqrt{2 p_T^\ell E_T^{\text{miss}} (1 - \cos \Delta\phi_{\ell, E_T^{\text{miss}}})}$, where $\Delta\phi_{\ell, E_T^{\text{miss}}}$ is the azimuthal angle between the lepton and E_T^{miss} directions.




Figure 8. Probability for a muon produced in the decay of a b -hadron to satisfy a track-based isolation requirement, in simulated $t\bar{t}$ events, shown as a function of (a) the muon p_T ; and (b) the sum $p_T + E_T^{\text{cone40}}$ of the muon p_T and nearby calorimeter energy deposits. In each case, the universality of these probabilities is gauged by changing the kinematic properties of the parent hadron (markers of various colours), achieved indirectly by selecting events in different ranges of global transverse energy H_T . Precise definitions of the isolation criteria, the $p_T + E_T^{\text{cone40}}$ and H_T variables, as well as descriptions of the muon and jets selections, are given in section 7.2.

In either approach, there will be a residual contribution from events with only real leptons in the selected sample. This contribution is typically estimated using MC simulation and subtracted separately from both the tight and baseline samples before the ratio of these samples is taken to measure the efficiency.

As with the real-lepton efficiencies, the fake/non-prompt lepton efficiencies depend on properties of the lepton candidates or of the event in which they are found. Therefore, it is generally helpful to bin the efficiencies in the lepton p_T and $|\eta|$, and possibly in terms of other quantities as well. The optimal binning to be used is chosen in the context of each physics analysis, considering the given numbers of events and potential changes in the relative contributions from the different sources of fake/non-prompt leptons; illustrative examples are provided in section 9.

Adopting a parameterisation of the efficiency with respect to variables other than p_T and η can sometimes be beneficial. For example, the probability that a fake or non-prompt lepton satisfies the isolation criteria is correlated with the fraction of the jet’s visible momentum carried by the lepton. Therefore parametrising the fake efficiency as a function of the lepton p_T amounts to assuming that, for a particular lepton p_T , the distribution of the parent jet momentum is similar in the regions where the efficiencies are measured and the regions where the background estimates are needed. If this assumption does not hold, it can be useful to adopt instead a parameterisation as a function of the parent jet momentum. Since this quantity is not easily accessed experimentally (unlike the lepton p_T), proxy observables are used in practice. An example of successful application is the analysis in ref. [41], which employed the sum of the lepton’s p_T and the transverse energy in a cone around the lepton as a proxy.

The preceding discussion is also illustrated in figure 8 for non-prompt muons produced in the decay of b -hadrons in simulated $t\bar{t}$ events. The probabilities for such muons to satisfy a track-based isolation requirement, as defined in section 4, are shown for two alternative parameterisations: one

based on the muon p_T , and the other on the scalar sum of the muon p_T and the transverse energy deposited in calorimeter-cell clusters within a cone of size $\Delta R = 0.4$ around the muon (referred to as E_T^{cone40} in section 4). This scalar sum serves as a proxy for the parent jet’s transverse momentum.

In the second parameterisation, the fake efficiency is the fraction of jets with a non-prompt muon and visible momentum $p_T + E_T^{\text{cone40}}$ in which the muon is mostly isolated, i.e. $p_T \gg E_T^{\text{cone40}}$. While one might consider jets with arbitrarily soft muons in the denominator of this fraction, for practical reasons figure 8(b) only includes events where muons satisfy $p_T > 10 \text{ GeV}$. To study the dependence of these efficiencies on the momentum distribution of the underlying jet, different regions of jet momentum are emphasised by imposing different requirements on the global transverse energy H_T in the event, defined for this purpose as the scalar p_T sum of all jets with $p_T > 25 \text{ GeV}$ and $|\eta| < 2.8$ that are a distance $\Delta R > 0.6$ from the muon. This quantity is indeed partially correlated with the kinematics of the muon’s parent jet, via the momentum of the top quarks producing all these jets. The reconstruction, calibration and selection of jets and muons for this figure are otherwise those detailed in ref. [37].

It can be observed that for the case of a p_T -dependent parameterisation, the efficiencies vary strongly with H_T , although large differences occur mostly for $p_T > 40 \text{ GeV}$. Since most non-prompt muons are produced at low p_T , the overall impact of this non-captured dependency might be small, unless regions of interest in the analysis specifically select high- p_T leptons. In contrast, the parameterisation as a function of $p_T + E_T^{\text{cone40}}$ is much less influenced by H_T , making the measured efficiencies less dependent on the event topology. In practice, a compromise has to be found between this observation and other elements evoked above justifying a p_T -dependent parameterisation, especially for electrons because ID criteria are usually employed in addition to isolation. The direct dependency of efficiencies on other variables that are also correlated with the event topology (e.g. E_T^{miss}) may also be reduced by such a parameterisation.

As detailed in the next section, suitable uncertainties must be assigned to the use of the fake efficiencies in different regions than those in which they are measured, in particular to account for potential differences in relative contributions of the different sources of fake/non-prompt leptons. To minimize these uncertainties, it has sometimes been found beneficial to use an approach closer to that of section 7.1, in which efficiencies are evaluated in the simulation and supplemented by data-driven correction factors that depend on the source of the fake/non-prompt lepton. These correction factors are derived using dedicated control regions that are enriched in a particular source of fake/non-prompt leptons. The main assumptions are then the universality of the correction factors across different processes, and the ability of the simulation to adequately predict the relative contributions of each source in the regions of interest. Such an approach has for example been used in ref. [42].

8 Systematic uncertainties

Systematic uncertainties in the fake/non-prompt lepton background estimates from the matrix method and the fake-factor method arise from uncertainties in the values of ε_r and ε_f . These uncertainties can be traced to statistical uncertainties from the samples used to measure the efficiencies, to potential biases that may cause the efficiencies in the signal region for a particular analysis to differ from the values obtained from control samples (such as differences in the origin of fake/non-prompt leptons between these regions), and to uncertainties in the modelling of contamination from real-

lepton processes in the samples used to measure the fake efficiency. Details of these uncertainties and their estimation are provided below. The overall impact of variations in ε_r and ε_f depends on the characteristics of the analysis. For example, in the matrix method Equations 2.3–2.5 imply that for analyses with

$$N_r^t \times \frac{\Delta\varepsilon_r}{\varepsilon_r^2} > N_f^t \times \frac{\Delta\varepsilon_f}{\varepsilon_f^2}$$

the uncertainty in ε_r will dominate, and vice versa.

8.1 Statistical uncertainties in the measured efficiencies

Statistical uncertainties in the real-lepton and fake-lepton efficiencies can be accounted for either by analytically propagating the uncertainties through to the estimated event yields or by varying the efficiencies input to the nominal yield calculation by their statistical uncertainties and using the resulting difference in the estimated fake/non-prompt background yield as the resulting uncertainty. The real- and fake-efficiency uncertainties are generally uncorrelated since the efficiencies are measured in statistically independent samples. In the usual case where the efficiencies are measured in bins of one or more quantities, the bins are uncorrelated, so the variations in each bin are applied separately. The total statistical uncertainty in the fake/non-prompt lepton background estimate is given by the sum in quadrature of the uncertainties from all variations. This source of systematic uncertainty is usually not dominant.

8.2 Systematic uncertainties in the measured efficiencies

Systematic uncertainties in ε_f are generally larger and more challenging to assess than for ε_r :

1. Real-lepton efficiencies have only slight variations due to event environment effects since real leptons have small contributions to their measurements from underlying event and jet activity. This also means there are a wide variety of samples with which they can be calibrated in great detail.
2. There are several sources of fake/non-prompt leptons, and the efficiencies may differ between these sources. Therefore, any differences in the fake/non-prompt lepton composition between the sample used to measure the efficiencies and the signal region for an analysis may lead to a bias in the efficiencies.

Several methods are used to estimate systematic uncertainties in the fake-lepton efficiencies. One is simply to vary the selection criteria for events in the control region used to measure the fake-lepton efficiencies, since the composition of fake/non-prompt leptons in the standard and alternative control regions may differ. A more sophisticated approach is to use MC simulation to estimate the fake/non-prompt lepton compositions in both the control and analysis regions. This information, combined with the MC-estimated selection efficiencies for each source of fake/non-prompt leptons, can be used to provide an estimate of the uncertainty.

8.3 Uncertainties in the modelling of real-lepton processes

When measuring the fake-lepton efficiencies, a correction must be applied to account for contamination from processes with real leptons in the control sample used for the measurement:

$$\varepsilon_f = \frac{N^t - N_r^t}{(N^t + N^l) - (N_r^t + N_r^l)}$$

where N_r^t and N_r^l are the numbers of real leptons in the selected tight and loose samples, respectively. MC simulation of real-lepton processes is generally used to estimate N_r^t and N_r^l , with corrections applied to account for known differences in object selection efficiencies between simulation and data. Nonetheless, several sources of systematic uncertainty in the real-lepton contamination remain: uncertainties in the cross-sections of the real-lepton processes, uncertainties in the correction factors, and uncertainties in the parameters, e.g. parton distribution functions (PDFs) and factorisation/renormalisation scales (μ_f , μ_r), used in the simulation.

8.4 Uncertainties due to biases in the Poisson likelihood matrix method

The biases that occur in some situations for the Poisson likelihood matrix method (see section 2.5) may also be considered as a source of systematic uncertainty, especially when the signal region contains many bins with few events in each. The magnitude of the bias can be estimated either by repeating the analysis with coarser bins, or by constraining the total fake/non-prompt lepton yield estimate to be the value returned by applying the Poisson likelihood matrix method to the entire unbinned sample. Any resulting differences in the binned estimates can be taken as a systematic uncertainty.

9 Examples of application in ATLAS analyses

In this section the application of the fake/non-prompt lepton background estimation methods is described using two example ATLAS analyses. The first is a measurement of the $t\bar{t}Z$ differential cross-section using events that contain three or four lepton candidates, and the second is a model-independent search for “beyond the Standard Model” (BSM) phenomena in events with three or more lepton candidates. In both cases, the high lepton multiplicity suppresses the Standard Model (SM) backgrounds, which makes the relative contribution of the fake/non-prompt lepton background larger.

Data from $\sqrt{s} = 13$ TeV pp collisions recorded by the ATLAS detector between 2015 and 2018 are used to perform these analyses. In this period, the LHC delivered colliding beams with a peak instantaneous luminosity of $L = 2.1 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$, achieved in 2018, and an average number of pp interactions per bunch crossing of 33.7. After applying beam, detector and data-quality criteria, the total integrated luminosity of the dataset is 139 fb^{-1} [43]. The uncertainty in the combined 2015–2018 integrated luminosity is 1.7% [44], obtained using the LUCID-2 detector [45] for the primary luminosity measurements.

9.1 Measurement of the $t\bar{t}Z$ cross-section in final states with three or four leptons

This analysis measured the differential production cross-section of the $t\bar{t}Z$ process in final states with a total of three or four electron and muon candidates. The tools described above are used for

the lepton efficiency measurements and the application of the Poisson likelihood matrix method (described in section 2.1.2). In addition to the fake/non-prompt lepton estimation in the signal regions of the analysis, the results are checked in several validation regions enriched in fake/non-prompt leptons. The Poisson likelihood matrix method was chosen for this analysis since the number of fake/non-prompt leptons in the signal regions with three or four leptons is expected to be very low, and the Poisson likelihood matrix method provides more stable results for binned estimations, which are necessary for differential background predictions in these low-statistics signal regions. The fraction of events with more than one fake/non-prompt lepton in the signal or validation regions has been checked and found to be negligible. More details about the analysis and the fake/non-prompt lepton background estimation can be found in ref. [46].

9.1.1 Real-lepton efficiencies

The first step in the application of the matrix method is to measure the efficiencies for real and fake/non-prompt leptons that satisfy the baseline criteria to also satisfy the tight criteria. In the $t\bar{t}Z$ cross-section measurement, baseline electrons are required to satisfy the “LooseAndBLayer” ID WP, whereas tight electrons are required to satisfy the stricter “Medium” ID criteria and to be isolated from nearby tracks and calorimeter energy deposits. Both the baseline and tight muons are required to satisfy the “Medium” WP, and tight muons are in addition required to be isolated from nearby tracks.

As discussed in section 7.1, the real-lepton efficiencies are obtained using MC simulation, corrected to match the performance seen in data control samples. Those efficiencies are shown in figure 9, binned in lepton p_T and $|\eta|$. To check for potential dependencies on the number of additional jets in the events used for the measurements, the efficiencies are derived for different jet multiplicities. No significant differences between the real-lepton efficiencies are observed.


9.1.2 Fake/non-prompt lepton efficiencies

The fake/non-prompt lepton efficiencies are measured with same-charge electron–muon ($e\mu$) or muon–muon ($\mu\mu$) data events using a tag-and-probe method, where one “tag lepton” with very stringent requirements on momentum and isolation ($p_T > 40$ GeV, $\max(p_T^{\text{cone20}}, E_T^{\text{cone20}})/p_T < 0.01$) is selected and the remaining “probe lepton” is used for the efficiency measurement. Events with more than two leptons are not considered. Only the aforementioned baseline electron or muon requirements are used to select the probe lepton. The samples are dominated by events containing at least one fake/non-prompt lepton, and are orthogonal to the signal regions, which require a minimum of three tight lepton candidates. In addition, $t\bar{t}Z$ production is negligible in these samples.


The definition of the regions used for the fake-efficiency measurements is summarised in table 1. For the electron fake efficiencies an $e\mu$ signature is used, with the muon being used as the tag lepton. For the muon fake efficiencies, the $\mu\mu$ region is used,⁹ since the $e\mu$ region also contains unwanted events where an electron with an incorrectly measured charge is selected as the tag lepton and paired with a real muon.

Real-lepton background processes (primarily diboson and $t\bar{t}W$ production) leading to the same-charge dilepton signature are estimated using MC-simulated events and are subtracted from data

⁹In $\mu\mu$ events with both muons satisfying the tag-selection, the one with the higher p_T is chosen as the tag lepton.




(a)




(b)

Figure 9. The two-dimensional real-lepton efficiencies obtained for (a) electrons and (b) muons, in bins of p_T and $|\eta|$ of the leptons. The last p_T bin is inclusive. The real-lepton efficiencies are obtained using MC simulation, but corrected to match the performance seen in data control samples.



(a)



(b)

Figure 10. The two-dimensional fake/non-prompt lepton efficiencies measured for (a) electrons and (b) muons, in bins of p_T and $|\eta|$ of the leptons. The last p_T bin is inclusive. The indicated uncertainties show only the statistical errors in the given bins. Compared to the real-lepton efficiencies shown in figure 9, the fake/non-prompt lepton efficiencies depend much more on the specifications of the analysis.

Table 1. Definition of the fake-lepton control regions used for the electron (e -fakes-CR) and muon (μ -fakes-CR) fake-efficiency measurements. N_ℓ is the number of leptons (ℓ), while N_{jets} ($N_{b\text{-jets}}$) is the number of jets (b -tagged jets, see text), respectively. For additional details of the definitions of physics objects, see ref. [46].

Region	N_ℓ ($\ell = e, \mu$)	p_T (ℓ)	p_T (jets)	N_{jets}	$N_{b\text{-jets}}$
e -fakes-CR	$= 2 (e^\pm \mu^\pm)$	$> 7 \text{ GeV}$	$> 25 \text{ GeV}$	≥ 1	≥ 1
μ -fakes-CR	$= 2 (\mu^\pm \mu^\pm)$	$> 7 \text{ GeV}$	$> 25 \text{ GeV}$	≥ 1	≥ 1

to obtain an unbiased efficiency measurement. The contribution from electrons with misassigned charge in the same-charge $e\mu$ region is also subtracted using estimates from MC-simulated events.

After this subtraction, the dominant source of fake/non-prompt leptons is found to be heavy-flavour hadron decays. The fake efficiencies, binned in lepton p_T and $|\eta|$, are shown in figure 10.

It is assumed that since the loose and tight lepton selection criteria depend on quantities related to the lepton itself or to its immediate surroundings, the chosen parameterisation captures the main variations in the fake efficiencies, and residual dependencies on the event environment can be covered by systematic uncertainties (see section 8). Indeed, in the simulation, the dependence of the fake efficiencies on the number of light-flavour jets or b -tagged¹⁰ jets is mild. The uncertainties are evaluated by comparing, in the simulation, the ε_f values in the control regions described in table 1 to the values in the signal regions. These differences, evaluated as a function of p_T and $|\eta|$, are applied as a systematic uncertainty of the fake-efficiency measurement (as discussed in section 8), and are of the order of 10%–20%, except for muons with $p_T > 50 \text{ GeV}$, for which they reach 40%. Furthermore, normalisation uncertainties are considered for the real-lepton background processes, which are subtracted in the fake-efficiency measurement. They are evaluated by scaling the real-lepton background processes upwards and downwards within their cross-section uncertainties before the subtraction and using the differences between the modified and nominal efficiencies as uncertainties, which are added in quadrature to the aforementioned uncertainties.

9.1.3 Results in the fake/non-prompt lepton validation regions

To validate the performance of the method, predictions are obtained and compared with data in two dedicated validation regions called “VR-3 ℓ -1 b 3 j ” and “VR-3 ℓ -1 b 3 j -noZ”, which have a larger proportion of fake/non-prompt leptons than is expected in the signal regions. The definitions of these two validation regions are summarised in table 2. No charge requirements are placed on the reconstructed lepton candidates in these regions.

The variable $m_{\ell\ell}^{\text{SF}}$ refers to the invariant mass of the same-flavour opposite-charge (SFOC) lepton pair with the invariant mass closest to the Z boson mass. VR-3 ℓ -1 b 3 j is a region similar to the actual signal regions defined in ref. [46], but without a requirement on the Z mass for the (SFOC) lepton candidate pair. Therefore, it contains a higher fraction of fake/non-prompt leptons after the selection. To further enhance the fake/non-prompt lepton contribution, the third-highest- p_T

¹⁰Jets containing b -hadrons are identified (tagged) by the MV2c10 b -tagging algorithm [47]. The algorithm uses a multivariate discriminant with quantities such as the impact parameters of associated tracks, and well-reconstructed secondary vertices.

Table 2. Definition of the fake/non-prompt lepton enriched validation regions, VR-3 ℓ -1 b 3 j and VR-3 ℓ -1 b 3 j -noZ. N_ℓ stands for the number of leptons (ℓ), while N_{jets} ($N_{b\text{-jets}}$) is the number of jets (b -tagged jets), respectively. The leptons are ordered by decreasing p_T ($\ell_{1,2,3}$). The mass (m) variables are discussed in the text. For additional details of the object definitions, see ref. [46].

Region	N_ℓ ($\ell = e, \mu$)	p_T ($\ell_{1,2,3}$)	p_T (jets)	$ m_{\ell\ell}^{\text{SF}} - m_Z $	N_{jets}	$N_{b\text{-jets}}$
VR-3 ℓ -1 b 3 j	= 3 (2 tight, 1 loose)	> 27, 20, 20 GeV	> 25 GeV	–	= 3	= 1
VR-3 ℓ -1 b 3 j -noZ	= 3 (all tight)	> 27, 20, 20 GeV	> 25 GeV	> 10 GeV	= 3	= 1

lepton that satisfies the baseline selection criteria must not satisfy the tight criteria. An additional validation region, VR-3 ℓ -1 b 3 j -noZ, is defined by requiring all three leptons to satisfy the tight selection criteria, but placing a veto on SFOC lepton pairs that have an invariant mass consistent with the Z boson, thereby enhancing the fake/non-prompt lepton fraction in this region. Both regions are orthogonal to the analysis signal regions and intended to validate the predictions of the matrix method for different levels of fake/non-prompt lepton contamination.

Some example distributions are shown in figure 11 for VR-3 ℓ -1 b 3 j and figure 12 for VR-3 ℓ -1 b 3 j -noZ. The processes with three real leptons (modelled with MC simulations) plus the prediction from the matrix method can be compared with data in these regions.

The hatched bands in figures 11 and 12 show only the statistical uncertainties of the MC prediction and the uncertainties associated with the fake/non-prompt lepton estimates (i.e. no theoretical or detector-related systematic uncertainties are included). The total uncertainty associated with the fake/non-prompt lepton estimate itself contains a systematic component, which is evaluated from variations of the input fake/real efficiencies (described in the previous section), and the statistical uncertainty of the data sample to which the Poisson likelihood matrix method is applied.¹¹

There is generally good agreement between the data and the total background estimate, except that the background is overestimated at low $\Delta R(\ell_1, \ell_2)$. One contribution to that difference is that the two leading (two highest- p_T) lepton candidates are likely to be real, yet when they are near each other they have a lower efficiency for satisfying the isolation criteria, and thus are misinterpreted as fake/non-prompt leptons by the matrix method. Analyses that are sensitive to such issues may benefit from imposing a minimum ΔR requirement between leptons. A discrepancy is also observed in the higher p_T bins of figure 11(d). Events in these bins have three leptons with p_T above 50 GeV. As shown in figures 9 and 10, only a single p_T bin above 50 GeV is available for measuring ε_r and ε_f , due to the limited number of events in the control regions, so variations above 50 GeV may be missed. This point was not investigated thoroughly since only a small fraction of events in the analysis were impacted.

¹¹The total uncertainties of the fake/non-prompt lepton estimates may be different from the uncertainties reported in ref. [46], as the number of loose leptons in these validation regions is larger than in the signal region of the $t\bar{t}Z$ analysis and, therefore, the statistical uncertainties are smaller.




Figure 11. (a) Angular separation between the leading and second leading (in p_T) lepton candidates, $\Delta R(\ell_1, \ell_2)$, (b) angular separation between the second- and third-leading lepton candidates, $\Delta R(\ell_2, \ell_3)$, (c) missing transverse momentum in the event, E_T^{miss} , and (d) the p_T of the third-leading lepton in VR-3 ℓ -1 b 3 j . The processes with three real leptons are modelled with MC simulation, while the contribution from fake/non-prompt leptons (dark red) comes from the Poisson likelihood matrix method as described above. The hatched band shows the uncertainty from the MC statistics and the fake/non-prompt background estimate. The rightmost bins are inclusive and contain all events above the x -axis ranges. The lower panel shows the ratio of data to the total SM prediction (sum of the real-lepton background contributions estimated with MC samples and the fake/non-prompt lepton contribution estimated with the matrix method). Further details are available in ref. [46].

9.2 Model-independent search for new phenomena in multi-lepton final states

Many interesting new models for BSM physics predict final states with three or more leptons. The general multi-lepton search for new phenomena [48] agnostically considers such final states. Its aim is to be sensitive to BSM phenomena in often-overlooked corners of phase space. The background estimation for the multi-lepton search uses MC predictions to account for events that contain only real leptons, and the fake-factor method for events containing at least one fake/non-prompt lepton. The dominant sources of fake/non-prompt leptons are semileptonic heavy-flavour decays (primarily of b -hadrons), light-hadron decays, and misidentification of light hadrons as leptons. These mainly arise in Z + jets and $t\bar{t}$ events.




Figure 12. Comparisons of the predicted and observed yields in VR-3 ℓ -1 $b3j$ -noZ, with respect to the p_T of the (a) leading and (b) subleading lepton candidates, (c) the scalar sum of the lepton and jet transverse momenta, H_T , and (d) the missing transverse momentum in the event, E_T^{miss} . The processes with three real leptons are modelled with MC simulation, while the contribution from fake/non-prompt leptons (dark red) comes from the Poisson likelihood matrix method as described above. The hatched band shows the uncertainty from MC statistics and the fake/non-prompt background estimate. The rightmost (leftmost) bins are inclusive and contain all events above (below) the x -axis ranges. The lower panel shows the ratio of data to the total SM prediction (sum of the real-lepton background contributions estimated with MC samples and the fake/non-prompt lepton contribution estimated with the matrix method). Further details are available in ref. [46].

9.2.1 Fake/non-prompt lepton selection

The fake factors are measured using events with a single lepton candidate, where the selection targets QCD dijet events. In order to prevent a bias in the fake factor due to trigger selection criteria [15, 16], the selected events are required to have fired a loose single-lepton trigger where isolation requirements are not imposed. However, due to the high rate of events that pass such triggers, a prescale factor is applied, which reduces the number of events available for measuring the fake factors (the luminosity of the prescaled samples is $\sim 0.5 \text{ fb}^{-1}$ for electrons and $\sim 3.7 \text{ fb}^{-1}$ for muons). Baseline lepton candidates must pass a common object selection, as detailed in ref. [48]. Electron candidates are required to pass either the “Loose” ID WP with calorimeter- and

track-based isolation requirements, or the “Tight” ID WP with no isolation requirement. Baseline muon candidates are required to pass the “Medium” ID WP (“HighPt” for $p_T > 300$ GeV). Only leptons with $p_T > 25$ GeV and satisfying the longitudinal and transverse track impact parameter requirements (see sections 4.1 and 4.2) are considered.

More stringent selection criteria are imposed for tight lepton candidates. For muons, where fake/non-prompt candidates are mainly muons from semileptonic heavy-flavour decays, only the isolation criteria is modified: the tight selection requires that muons satisfy track-based isolation criteria. For electrons, where both light- and heavy-flavour hadrons are a non-negligible source of fake/non-prompt candidates, both the ID and isolation criteria are modified. Tight electrons must satisfy both the “Tight” ID WP and calorimeter- and track-based isolation requirements.

Additional selection requirements are imposed on the single-lepton sample to ensure a high purity of fake/non-prompt leptons, which reduces the statistical uncertainty of the computed fake factor: the E_T^{miss} is required to be < 25 GeV (< 40 GeV) for events with electron (muon) candidates, and the number of jets in the event is required to be ≥ 1 (2) for events with electron (muon) candidates. For muon-candidate events, there must also be at least one jet with $p_T > 35$ GeV (the “tag jet”), and the azimuthal angle between the muon candidate and this jet is required to be > 2.7 radians.

The fake factor is binned in both the p_T and $|\eta|$ of the lepton candidate. The bins are defined to have tolerable statistical uncertainties while preventing sizeable differences between the fake-factor values in adjacent bins. The highest muon- p_T bin includes all muon candidates with $p_T > 80$ GeV, as there are very few fake/non-prompt muons at higher transverse momenta.

9.2.2 Fake factors

The fake factors calculated for the multi-lepton analysis are shown in figures 13 and 14 for electrons and muons. The uncertainties in the fake factors are categorised and evaluated as described below.

There are statistical uncertainties due to the data and MC sample sizes in each fake-factor bin, which due to their small size are summed in quadrature into a single uncertainty.

In the regions enriched in fake/non-prompt leptons, MC predictions are used to subtract the real-lepton contribution from the data. The uncertainties in the MC contributions are propagated to the fake factors. The main contributions from real leptons in the single-lepton regions are from Z + jets, W + jets and $t\bar{t}$ events. For the Z/W + jets processes, an uncertainty of 5% is applied to the cross-section. For the $t\bar{t}$ process, uncertainties on the cross section due to renormalization and factorization scales ($\sim 3\%$) and PDF ($\sim 4\%$) are applied.

Extrapolating the fake factors from the single-lepton sample to the multi-lepton samples used in the analysis introduces an uncertainty because these samples differ in the kinematic distributions of fake/non-prompt leptons, and possibly also in the fake/non-prompt lepton composition. Two uncertainties are included to address the bias caused by imposing a E_T^{miss} upper bound in the fake/non-prompt lepton estimation sample, and by imposing a p_T requirement on the tag jet in the fake/non-prompt muon estimation sample. These uncertainties are estimated by varying the requirements on these variables upwards and downwards by 10 GeV.¹² Plots showing the impact of these systematic effects on the fake factors are given in figures 13 and 14.

¹²Only the E_T^{miss} variation is displayed separately in the plots, as the variation of the jet p_T results in an uncertainty that is too small to be visible.




Figure 13. Measured fake factors for electrons as a function of p_T for different $|\eta|$ ranges, and their dependence on individual variations of parameters of the measurement, used to determine systematic uncertainties. The “MC syst” uncertainty covers variations in the model used to subtract the real-lepton contribution in the control regions used to measure ε_f . The combined impact of statistical and systematic uncertainties added in quadrature is indicated by the shaded yellow area (the grey area represents only the statistical uncertainty).

Finally, a direct assessment of the uncertainty in the composition of the fake/non-prompt lepton background in the multi-lepton sample is made. Since fake/non-prompt leptons can come from both light- and heavy-flavour sources, it is possible that the relative abundances from these sources can vary between samples. This possibility is addressed through an additional uncertainty which leverages the different event signatures produced by light- and heavy-flavour hadrons, the latter consisting primarily of b -hadrons. To evaluate this uncertainty, an alternative set of fake factors is computed which, in addition to the binning in p_T and $|\eta|$, are binned according to the presence or absence of b -tagged jets in the event. This alternative set of fake factors is shown in figure 15.

The composition uncertainty in the total event yields is then derived from the difference between estimates calculated with either the nominal fake factors oblivious to the presence of b -tagged jets in the events, or the alternative set that requires such a jet. This uncertainty is found to have a negligible impact on the analysis, since most of the jets in the signal region and in the sample used to measure the fake factors are not b -tagged.




Figure 14. Measured fake factors for muons as a function of p_T for different $|\eta|$ ranges, and their dependence on individual variations of parameters of the measurement, used to determine systematic uncertainties. The combined impact of statistical and systematic uncertainties added in quadrature is indicated by the shaded yellow area (the grey area represents only the statistical uncertainty).

9.2.3 Validation regions

Validation regions are defined using appropriate sub-selections of $ee\mu$ and $e\mu\mu$ events. These are used to check that the computed fake factors extrapolate correctly from the regions where they are calculated to the regions in which they are applied. The “on- Z ” validation region requires an SFOC lepton pair with a dilepton mass within 10 GeV of the Z boson mass. The “off- Z ” validation region also requires a SFOC pair of leptons, but requires the dilepton mass to fall outside of the Z -mass window. Only mixed-flavour final states are selected for these validation regions so that which lepton to choose as the third one, assumed to be the fake/non-prompt lepton, is unambiguous. The sources of fake/non-prompt leptons contribute in different ratios to the on- Z and off- Z validation regions: the on- Z validation region is more sensitive to Z + jets events than the off- Z region, while the inverse is true for $t\bar{t}$ events, although in absolute terms, Z + jets events are more numerous than $t\bar{t}$ events in both cases. Both validation regions target, through a m_T requirement of $m_T(\ell, E_T^{\text{miss}}) < 40$ GeV, a third lepton that is likely to be fake/non-prompt. The union of the on- Z and off- Z validation regions is called the “fakes validation region”.

The variables of primary importance for this analysis are the invariant mass of all lepton candidates in the event (m_{inv}) and the E_T^{miss} . The signal regions are separated according to the




Figure 15. Fake factors measured in bins of the lepton candidate’s p_T and $|\eta|$ for events with (a,c) electron candidates and (b,d) muon candidates in events (a,b) without and (c,d) with b -tagged jets.

values of these quantities, as discussed in ref. [48]. The m_{inv} distributions in the two validation regions are shown in figure 16, while the E_T^{miss} distributions are shown in figure 17. Lastly, the electron and muon candidate p_T distributions are shown in the fakes validation region in figure 18. The comparisons in figure 16 were also presented in ref. [48] (albeit in logarithmic scale) and thus include fitted normalisation factors for the WZ + jets and ZZ + jets backgrounds from their respective control regions (“post-fit”). On the other hand, the complementary distributions shown in figures 17 and 18 were obtained independently of this statistical analysis, and thus employ the unconstrained SM background normalisations and uncertainties (“pre-fit” distributions). The background estimate is consistent with the data within the statistical and systematic uncertainties.




Figure 16. Comparison between data and prediction for the m_{inv} distribution in the (a) on-Z and (b) off-Z validation regions, after fitting the normalisation factors for the WZ + jets and ZZ + jets backgrounds and systematic uncertainties [48]. All uncertainties, systematic and statistical, are included. The leftmost (rightmost) bin is inclusive and contains all events with $m_{\text{inv}} < 200 \text{ GeV}$ ($> 500 \text{ GeV}$). The hatched grey area in these figures shows the total uncertainty. Further details are available in ref. [48].




Figure 17. Comparison between data and prediction for the E_T^{miss} distribution in the (a) on-Z and (b) off-Z validation regions. All uncertainties, systematic and statistical, are included. The rightmost bin is inclusive and contains all events with $E_T^{\text{miss}} > 475 \text{ GeV}$. The hatched grey area in these figures shows the total uncertainty. Further details are available in ref. [48].




Figure 18. Comparison between data and prediction for the p_T distribution in the fakes validation region (which is the union of the on-Z and off-Z validation regions). Shown are (a) the electron p_T in $e\mu\mu$ and (b) the muon p_T in $ee\mu$ events. All uncertainties, systematic and statistical, are included. The rightmost bin is inclusive and contains all events with $p_T > 275$ GeV. The hatched grey area in these figures shows the total uncertainty. Further details are available in ref. [48].

10 Conclusions

For physics analyses exploring signatures with one or more prompt leptons, background contributions due to fake/non-prompt leptons are often difficult to estimate in simulations. Therefore, data-driven methods are commonly used. Three related methods have been adopted by the ATLAS Collaboration as recommended tools: the asymptotic matrix method, the Poisson likelihood matrix method, and the fake-factor method. All three approaches depend on defining two categories of leptons, one of which (“tight”) is subject to the same identification and selection criteria as are used in the analysis. The other category (“loose”) adds additional lepton candidates with less stringent selection requirements. The union of the two sets is called the “baseline” sample. The criteria are typically defined such that the probability for a real baseline lepton to satisfy the tight criteria is substantially higher than the corresponding probability for a fake/non-prompt lepton. Then, the relative numbers of loose and tight leptons in the analysis sample can be used to estimate the contribution of fake/non-prompt leptons, either inclusively or differentially in any variables of interest.

Despite their similarities, the methods each have their own strengths and drawbacks. The asymptotic matrix method and fake-factor method provide a fake/non-prompt lepton weight for each event, which is convenient for analyses. However, these methods are subject to large uncertainties if the efficiency for baseline fake/non-prompt leptons to satisfy the tight criteria is large in parts of the analysis phase space. The Poisson likelihood matrix method returns a smaller uncertainty in such cases, and avoids any possibility of producing a negative estimate for the event yield, but does not provide a per-event weight, introducing difficulties e.g. for differential estimations. The fake-factor method uses simulation rather than data to incorporate the contribution from events

where all leptons are tight. This method can therefore be employed while the signal region for an analysis is fully blinded, although it may induce additional simulation-related uncertainties in the background estimate.

The systematic uncertainties for all three methods arise from similar sources, with the largest contributions related to the extrapolation of the efficiencies measured in the control samples to events in the analysis sample. Differences in the fake/non-prompt lepton composition in the samples must be accounted for and appropriate uncertainties must be assigned to this extrapolation.

The performance of the Poisson likelihood matrix method and the fake-factor method has been demonstrated in a differential $t\bar{t}Z$ cross section measurement and in a model-independent search for BSM phenomena in multi-lepton final states, respectively. In both cases, the chosen method was shown to provide a reliable estimate of the fake/non-prompt lepton background, as measured in validation regions.

Acknowledgments

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, Compute Canada and CRC, Canada; PRIMUS 21/SCI/017 and UNCE SCI/013, Czech Republic; COST, ERC, ERDF, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; Norwegian Financial Mechanism 2014-2021, Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in ref. [49].

References

- [1] ATLAS collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, 2008 *JINST* **3** S08003 [[INSPIRE](#)].
- [2] ATLAS collaboration, *Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton-proton collision data*, 2019 *JINST* **14** P12006 [[arXiv:1908.00005](#)] [[INSPIRE](#)].
- [3] ATLAS collaboration, *Muon reconstruction and identification efficiency in ATLAS using the full Run 2 pp collision data set at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **81** (2021) 578 [[arXiv:2012.00578](#)] [[INSPIRE](#)].
- [4] D0 collaboration, *Extraction of the width of the W boson from measurements of $\sigma(p\bar{p} \rightarrow W + X) \times B(W \rightarrow e\nu)$ and $\sigma(p\bar{p} \rightarrow Z + X) \times B(Z \rightarrow ee)$ and their ratio*, *Phys. Rev. D* **61** (2000) 072001 [[hep-ex/9906025](#)] [[INSPIRE](#)].
- [5] J. Erdmann et al., *Reformulation of a likelihood approach to fake-lepton estimation in the framework of Bayesian inference*, *Nucl. Instrum. Meth. A* **1021** (2022) 165939 [[arXiv:2106.13628](#)] [[INSPIRE](#)].
- [6] T.P.S. Gillam and C.G. Lester, *Improving estimates of the number of ‘fake’ leptons and other mis-reconstructed objects in hadron collider events: BoB’s your UNCLE*, *JHEP* **11** (2014) 031 [[arXiv:1407.5624](#)] [[INSPIRE](#)].
- [7] ATLAS collaboration, *Observation of inclusive electrons in the ATLAS experiment at $\sqrt{s} = 7$ TeV*, *ATLAS-CONF-2010-073* (2010).
- [8] G. Bohm and G. Zech, *Statistics of weighted Poisson events and its applications*, *Nucl. Instrum. Meth. A* **748** (2014) 1 [[arXiv:1309.1287](#)] [[INSPIRE](#)].
- [9] F. James and M. Roos, *Minuit: A System for Function Minimization and Analysis of the Parameter Errors and Correlations*, *Comput. Phys. Commun.* **10** (1975) 343 [[INSPIRE](#)].
- [10] R. Brun and F. Rademakers, *ROOT: An object oriented data analysis framework*, *Nucl. Instrum. Meth. A* **389** (1997) 81 [[INSPIRE](#)].
- [11] ATLAS collaboration, *ATLAS Insertable B-Layer Technical Design Report*, *ATLAS-TDR-2010-19* (2010).
- [12] ATLAS IBL collaboration, *Production and integration of the ATLAS Insertable B-Layer*, 2018 *JINST* **13** T05008 [[arXiv:1803.00844](#)] [[INSPIRE](#)].
- [13] ATLAS collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317 [[arXiv:1611.09661](#)] [[INSPIRE](#)].
- [14] ATLAS collaboration, *The ATLAS collaboration software and firmware*, *ATL-SOFT-PUB-2021-001* (2021).
- [15] ATLAS collaboration, *Performance of the ATLAS muon triggers in Run 2*, 2020 *JINST* **15** P09015 [[arXiv:2004.13447](#)] [[INSPIRE](#)].
- [16] ATLAS collaboration, *Performance of electron and photon triggers in ATLAS during LHC Run 2*, *Eur. Phys. J. C* **80** (2020) 47 [[arXiv:1909.00761](#)] [[INSPIRE](#)].
- [17] ATLAS collaboration, *Electron and photon energy calibration with the ATLAS detector using LHC Run 1 data*, *Eur. Phys. J. C* **74** (2014) 3071 [[arXiv:1407.5063](#)] [[INSPIRE](#)].
- [18] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].

- [19] M. Cacciari, G.P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [20] M. Cacciari and G.P. Salam, *Pileup subtraction using jet areas*, *Phys. Lett. B* **659** (2008) 119 [[arXiv:0707.1378](#)] [[INSPIRE](#)].
- [21] ATLAS collaboration, *Evidence for the associated production of the Higgs boson and a top quark pair with the ATLAS detector*, *Phys. Rev. D* **97** (2018) 072003 [[arXiv:1712.08891](#)] [[INSPIRE](#)].
- [22] ATLAS collaboration, *Search for chargino–neutralino pair production in final states with three leptons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 1118 [[arXiv:2106.01676](#)] [[INSPIRE](#)].
- [23] ATLAS collaboration, *Improvements in $t\bar{t}$ modelling using NLO+PS Monte Carlo generators for Run 2*, *ATL-PHYS-PUB-2018-009* (2018).
- [24] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [25] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [26] ATLAS collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, *ATL-PHYS-PUB-2014-021* (2014).
- [27] NNPDF collaboration, *Parton distributions for the LHC Run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [28] R.D. Ball et al., *Parton distributions with LHC data*, *Nucl. Phys. B* **867** (2013) 244 [[arXiv:1207.1303](#)] [[INSPIRE](#)].
- [29] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [30] ATLAS collaboration, *ATLAS simulation of boson plus jets processes in Run 2*, *ATL-PHYS-PUB-2017-006* (2017).
- [31] SHERPA collaboration, *Event Generation with Sherpa 2.2*, *SciPost Phys.* **7** (2019) 034 [[arXiv:1905.09127](#)] [[INSPIRE](#)].
- [32] ATLAS collaboration, *The ATLAS Simulation Infrastructure*, *Eur. Phys. J. C* **70** (2010) 823 [[arXiv:1005.4568](#)] [[INSPIRE](#)].
- [33] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [34] ATLAS collaboration, *Studies of the muon momentum calibration and performance of the ATLAS detector with pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **83** (2023) 686 [[arXiv:2212.07338](#)] [[INSPIRE](#)].
- [35] ATLAS collaboration, *Jet energy scale and resolution measured in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 689 [[arXiv:2007.02645](#)] [[INSPIRE](#)].
- [36] ATLAS collaboration, *Charged-particle distributions in $\sqrt{s} = 13$ TeV pp interactions measured with the ATLAS detector at the LHC*, *Phys. Lett. B* **758** (2016) 67 [[arXiv:1602.01633](#)] [[INSPIRE](#)].
- [37] ATLAS collaboration, *Search for squarks and gluinos in final states with same-sign leptons and jets using 139 fb^{-1} of data collected with the ATLAS detector*, *JHEP* **06** (2020) 046 [[arXiv:1909.08457](#)] [[INSPIRE](#)].

- [38] R.H. Dalitz, *On an alternative decay process for the neutral pi-meson*, Letters to the Editor, *Proc. Phys. Soc. A* **64** (1951) 667 [[INSPIRE](#)].
- [39] ATLAS collaboration, *Proposal for particle-level object and observable definitions for use in physics measurements at the LHC*, [ATL-PHYS-PUB-2015-013](#) (2015).
- [40] ATLAS collaboration, *Performance of missing transverse momentum reconstruction with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **78** (2018) 903 [[arXiv:1802.08168](#)] [[INSPIRE](#)].
- [41] ATLAS collaboration, *Measurement of $W^{\pm}W^{\pm}$ vector-boson scattering and limits on anomalous quartic gauge couplings with the ATLAS detector*, *Phys. Rev. D* **96** (2017) 012007 [[arXiv:1611.02428](#)] [[INSPIRE](#)].
- [42] ATLAS collaboration, *Search for supersymmetry in events with four or more charged leptons in 139 fb^{-1} of $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *JHEP* **07** (2021) 167 [[arXiv:2103.11684](#)] [[INSPIRE](#)].
- [43] ATLAS collaboration, *ATLAS data quality operations and performance for 2015-2018 data-taking*, *2020 JINST* **15** P04003 [[arXiv:1911.04632](#)] [[INSPIRE](#)].
- [44] ATLAS collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, [ATLAS-CONF-2019-021](#) (2019).
- [45] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, *2018 JINST* **13** P07017 [[INSPIRE](#)].
- [46] ATLAS collaboration, *Measurements of the inclusive and differential production cross sections of a top-quark-antiquark pair in association with a Z boson at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 737 [[arXiv:2103.12603](#)] [[INSPIRE](#)].
- [47] ATLAS collaboration, *ATLAS b-jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 970 [[arXiv:1907.05120](#)] [[INSPIRE](#)].
- [48] ATLAS collaboration, *Search for new phenomena in three- or four-lepton events in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **824** (2022) 136832 [[arXiv:2107.00404](#)] [[INSPIRE](#)].
- [49] ATLAS collaboration, *ATLAS computing acknowledgements*, [ATL-SOFT-PUB-2021-003](#), (2021).

The ATLAS collaboration

G. Aad ¹⁰², B. Abbott ¹²⁰, D.C. Abbott ¹⁰³, K. Abeling ⁵⁵, S.H. Abidi ²⁹, A. Aboulhorma ^{35e},
H. Abramowicz ¹⁵¹, H. Abreu ¹⁵⁰, Y. Abulaiti ¹¹⁷, A.C. Abusleme Hoffman ^{137a},
B.S. Acharya ^{69a,69b,q}, C. Adam Bourdarios ⁴, L. Adamczyk ^{85a}, L. Adamek ¹⁵⁵,
S.V. Addepalli ²⁶, J. Adelman ¹¹⁵, A. Adiguzel ^{21c}, S. Adorni ⁵⁶, T. Adye ¹³⁴, A.A. Affolder ¹³⁶,
Y. Afik ³⁶, M.N. Agaras ¹³, J. Agarwala ^{73a,73b}, A. Aggarwal ¹⁰⁰, C. Agheorghiesei ^{27c},
J.A. Aguilar-Saavedra ^{130f}, A. Ahmad ³⁶, F. Ahmadov ^{38,ab}, W.S. Ahmed ¹⁰⁴, S. Ahuja ⁹⁵,
X. Ai ⁴⁸, G. Aielli ^{76a,76b}, M. Ait Tamlihat ^{35e}, B. Aitbenchikh ^{35a}, I. Aizenberg ¹⁶⁹,
M. Akbiyik ¹⁰⁰, T.P.A. Åkesson ⁹⁸, A.V. Akimov ³⁷, K. Al Khoury ⁴¹, G.L. Alberghi ^{23b},
J. Albert ¹⁶⁵, P. Albicocco ⁵³, S. Alderweireldt ⁵², M. Aleksa ³⁶, I.N. Aleksandrov ³⁸,
C. Alexa ^{27b}, T. Alexopoulos ¹⁰, A. Alfonsi ¹¹⁴, F. Alfonsi ^{23b}, M. Alhroob ¹²⁰, B. Ali ¹³²,
S. Ali ¹⁴⁸, M. Aliev ³⁷, G. Alimonti ^{71a}, W. Alkahi ⁵⁵, C. Allaire ⁶⁶, B.M.M. Allbrooke ¹⁴⁶,
C.A. Allendes Flores ^{137f}, P.P. Allport ²⁰, A. Aloisio ^{72a,72b}, F. Alonso ⁹⁰, C. Alpigiani ¹³⁸,
M. Alvarez Estevez ⁹⁹, M.G. Alviggi ^{72a,72b}, M. Aly ¹⁰¹, Y. Amaral Coutinho ^{82b}, A. Ambler ¹⁰⁴,
C. Amelung ³⁶, M. Amerl ¹, C.G. Ames ¹⁰⁹, D. Amidei ¹⁰⁶, S.P. Amor Dos Santos ^{130a},
K.R. Amos ¹⁶³, V. Ananiev ¹²⁵, C. Anastopoulos ¹³⁹, T. Andeen ¹¹, J.K. Anders ³⁶,
S.Y. Andrean ^{47a,47b}, A. Andreazza ^{71a,71b}, S. Angelidakis ⁹, A. Angerami ^{41,ae},
A.V. Anisenkov ³⁷, A. Annovi ^{74a}, C. Antel ⁵⁶, M.T. Anthony ¹³⁹, E. Antipov ¹²¹, M. Antonelli ⁵³,
D.J.A. Antrim ^{17a}, F. Anulli ^{75a}, M. Aoki ⁸³, T. Aoki ¹⁵³, J.A. Aparisi Pozo ¹⁶³, M.A. Aparo ¹⁴⁶,
L. Aperio Bella ⁴⁸, C. Appelt ¹⁸, N. Aranzabal ³⁶, V. Araujo Ferraz ^{82a}, C. Arcangeletti ⁵³,
A.T.H. Arce ⁵¹, E. Arena ⁹², J-F. Arguin ¹⁰⁸, S. Argyropoulos ⁵⁴, J.-H. Arling ⁴⁸,
A.J. Armbruster ³⁶, O. Arnaez ¹⁵⁵, H. Arnold ¹¹⁴, Z.P. Arrubarrena Tame ¹⁰⁹, G. Artoni ^{75a,75b},
H. Asada ¹¹¹, K. Asai ¹¹⁸, S. Asai ¹⁵³, N.A. Asbah ⁶¹, J. Assahsah ^{35d}, K. Assamagan ²⁹,
R. Astalos ^{28a}, R.J. Atkin ^{33a}, M. Atkinson ¹⁶², N.B. Atlay ¹⁸, H. Atmani ^{62b}, P.A. Atmasiddha ¹⁰⁶,
K. Augsten ¹³², S. Auricchio ^{72a,72b}, A.D. Auriol ²⁰, V.A. Austrup ¹⁷¹, G. Avner ¹⁵⁰,
G. Avolio ³⁶, K. Axiotis ⁵⁶, M.K. Ayoub ^{14c}, G. Azuelos ^{108,aj}, D. Babal ^{28a}, H. Bachacou ¹³⁵,
K. Bachas ^{152,t}, A. Bachiu ³⁴, F. Backman ^{47a,47b}, A. Badea ⁶¹, P. Bagnaia ^{75a,75b},
M. Bahmani ¹⁸, A.J. Bailey ¹⁶³, V.R. Bailey ¹⁶², J.T. Baines ¹³⁴, C. Bakalis ¹⁰, O.K. Baker ¹⁷²,
P.J. Bakker ¹¹⁴, E. Bakos ¹⁵, D. Bakshi Gupta ⁸, S. Balaji ¹⁴⁷, R. Balasubramanian ¹¹⁴,
E.M. Baldin ³⁷, P. Balek ¹³³, E. Ballabene ^{71a,71b}, F. Balli ¹³⁵, L.M. Baltes ^{63a}, W.K. Balunas ³²,
J. Balz ¹⁰⁰, E. Banas ⁸⁶, M. Bandieramonte ¹²⁹, A. Bandyopadhyay ²⁴, S. Bansal ²⁴, L. Barak ¹⁵¹,
E.L. Barberio ¹⁰⁵, D. Barberis ^{57b,57a}, M. Barbero ¹⁰², G. Barbour ⁹⁶, K.N. Barends ^{33a},
T. Barillari ¹¹⁰, M-S. Barisits ³⁶, T. Barklow ¹⁴³, R.M. Barnett ^{17a}, P. Baron ¹²²,
D.A. Baron Moreno ¹⁰¹, A. Baroncelli ^{62a}, G. Barone ²⁹, A.J. Barr ¹²⁶,
L. Barranco Navarro ^{47a,47b}, F. Barreiro ⁹⁹, J. Barreiro Guimarães da Costa ^{14a}, U. Barron ¹⁵¹,
M.G. Barros Teixeira ^{130a}, S. Barsov ³⁷, F. Bartels ^{63a}, R. Bartoldus ¹⁴³, A.E. Barton ⁹¹,
P.artos ^{28a}, A. Basalaeu ⁴⁸, A. Basan ¹⁰⁰, M. Baselga ⁴⁹, I. Bashta ^{77a,77b}, A. Bassalat ^{66,b},
M.J. Basso ¹⁵⁵, C.R. Basson ¹⁰¹, R.L. Bates ⁵⁹, S. Batlamous ^{35e}, J.R. Batley ³², B. Batool ¹⁴¹,
M. Battaglia ¹³⁶, D. Battulga ¹⁸, M. Bause ^{75a,75b}, P. Bauer ²⁴, A. Bayirli ^{21a}, J.B. Beacham ⁵¹,
T. Beau ¹²⁷, P.H. Beauchemin ¹⁵⁸, F. Becherer ⁵⁴, P. Bechtel ²⁴, H.P. Beck ^{19,s}, K. Becker ¹⁶⁷,
A.J. Beddall ^{21d}, V.A. Bednyakov ³⁸, C.P. Bee ¹⁴⁵, L.J. Beemster ¹⁵, T.A. Beermann ³⁶,
M. Begalli ^{82d}, M. Begel ²⁹, A. Behera ¹⁴⁵, J.K. Behr ⁴⁸, C. Beirao Da Cruz E Silva ³⁶,

J.F. Beirer ^{55,36}, F. Beisiegel ²⁴, M. Belfkir ¹⁵⁹, G. Bella ¹⁵¹, L. Bellagamba ^{23b}, A. Bellerive ³⁴, P. Bellos ²⁰, K. Beloborodov ³⁷, K. Belotskiy ³⁷, N.L. Belyaev ³⁷, D. Benckekroun ^{35a}, F. Bendebba ^{35a}, Y. Benhammou ¹⁵¹, D.P. Benjamin ²⁹, M. Benoit ²⁹, J.R. Bensingier ²⁶, S. Bentvelsen ¹¹⁴, L. Beresford ³⁶, M. Beretta ⁵³, E. Bergeaas Kuutmann ¹⁶¹, N. Berger ⁴, B. Bergmann ¹³², J. Beringer ^{17a}, S. Berlendis ⁷, G. Bernardi ⁵, C. Bernius ¹⁴³, F.U. Bernlochner ²⁴, T. Berry ⁹⁵, P. Berta ¹³³, A. Berthold ⁵⁰, I.A. Bertram ⁹¹, S. Bethke ¹¹⁰, A. Betti ^{75a,75b}, A.J. Bevan ⁹⁴, M. Bhamjee ^{33c}, S. Bhatta ¹⁴⁵, D.S. Bhattacharya ¹⁶⁶, P. Bhattacharai ²⁶, V.S. Bhopatkar ¹²¹, R. Bi^{29,am}, R.M. Bianchi ¹²⁹, O. Biebel ¹⁰⁹, R. Bielski ¹²³, M. Biglietti ^{77a}, T.R.V. Billoud ¹³², M. Bindi ⁵⁵, A. Bingul ^{21b}, C. Bini ^{75a,75b}, A. Biondini ⁹², C.J. Birch-sykes ¹⁰¹, G.A. Bird ^{20,134}, M. Birman ¹⁶⁹, M. Biros ¹³³, T. Bisanz ³⁶, E. Bisceglie ^{43b,43a}, D. Biswas ^{170,m}, A. Bitadze ¹⁰¹, K. Björke ¹²⁵, I. Bloch ⁴⁸, C. Blocker ²⁶, A. Blue ⁵⁹, U. Blumenschein ⁹⁴, J. Blumenthal ¹⁰⁰, G.J. Bobbink ¹¹⁴, V.S. Bobrovnikov ³⁷, M. Boehler ⁵⁴, D. Bogavac ³⁶, A.G. Bogdanchikov ³⁷, C. Bohm ^{47a}, V. Boisvert ⁹⁵, P. Bokan ⁴⁸, T. Bold ^{85a}, M. Bomben ⁵, M. Bona ⁹⁴, M. Boonekamp ¹³⁵, C.D. Booth ⁹⁵, A.G. Borbély ⁵⁹, H.M. Borecka-Bielska ¹⁰⁸, L.S. Borgna ⁹⁶, G. Borissov ⁹¹, D. Bortoletto ¹²⁶, D. Boscherini ^{23b}, M. Bosman ¹³, J.D. Bossio Sola ³⁶, K. Bouaouda ^{35a}, N. Bouchhar ¹⁶³, J. Boudreau ¹²⁹, E.V. Bouhova-Thacker ⁹¹, D. Boumediene ⁴⁰, R. Bouquet ⁵, A. Boveia ¹¹⁹, J. Boyd ³⁶, D. Boye ²⁹, I.R. Boyko ³⁸, J. Bracinik ²⁰, N. Brahimi ^{62d}, G. Brandt ¹⁷¹, O. Brandt ³², F. Braren ⁴⁸, B. Brau ¹⁰³, J.E. Brau ¹²³, K. Brendlinger ⁴⁸, R. Brenner ¹⁶⁹, L. Brenner ¹¹⁴, R. Brenner ¹⁶¹, S. Bressler ¹⁶⁹, D. Britton ⁵⁹, D. Britzger ¹¹⁰, I. Brock ²⁴, G. Brooijmans ⁴¹, W.K. Brooks ^{137f}, E. Brost ²⁹, L.M. Brown ¹⁶⁵, T.L. Bruckler ¹²⁶, P.A. Bruckman de Renstrom ⁸⁶, B. Brüers ⁴⁸, D. Bruncko ^{28b,*}, A. Bruni ^{23b}, G. Bruni ^{23b}, M. Bruschi ^{23b}, N. Bruscino ^{75a,75b}, T. Buanes ¹⁶, Q. Buat ¹³⁸, P. Buchholz ¹⁴¹, A.G. Buckley ⁵⁹, I.A. Budagov ^{38,*}, M.K. Bugge ¹²⁵, O. Bulekov ³⁷, B.A. Bullard ¹⁴³, S. Burdin ⁹², C.D. Burgard ⁴⁹, A.M. Burger ⁴⁰, B. Burghgrave ⁸, J.T.P. Burr ³², C.D. Burton ¹¹, J.C. Burzynski ¹⁴², E.L. Busch ⁴¹, V. Büscher ¹⁰⁰, P.J. Bussey ⁵⁹, J.M. Butler ²⁵, C.M. Buttar ⁵⁹, J.M. Butterworth ⁹⁶, W. Buttinger ¹³⁴, C.J. Buxo Vazquez ¹⁰⁷, A.R. Buzykaev ³⁷, G. Cabras ^{23b}, S. Cabrera Urbán ¹⁶³, D. Caforio ⁵⁸, H. Cai ¹²⁹, Y. Cai ^{14a,14d}, V.M.M. Cairo ³⁶, O. Cakir ^{3a}, N. Calace ³⁶, P. Calafiura ^{17a}, G. Calderini ¹²⁷, P. Calfayan ⁶⁸, G. Callea ⁵⁹, L.P. Caloba ^{82b}, D. Calvet ⁴⁰, S. Calvet ⁴⁰, T.P. Calvet ¹⁰², M. Calvetti ^{74a,74b}, R. Camacho Toro ¹²⁷, S. Camarda ³⁶, D. Camarero Munoz ²⁶, P. Camarri ^{76a,76b}, M.T. Camerlingo ^{72a,72b}, D. Cameron ¹²⁵, C. Camincher ¹⁶⁵, M. Campanelli ⁹⁶, A. Camplani ⁴², V. Canale ^{72a,72b}, A. Canesse ¹⁰⁴, M. Cano Bret ⁸⁰, J. Cantero ¹⁶³, Y. Cao ¹⁶², F. Capocasa ²⁶, M. Capua ^{43b,43a}, A. Carbone ^{71a,71b}, R. Cardarelli ^{76a}, J.C.J. Cardenas ⁸, F. Cardillo ¹⁶³, T. Carli ³⁶, G. Carlino ^{72a}, J.I. Carlotta ¹³, B.T. Carlson ^{129,u}, E.M. Carlson ^{165,156a}, L. Carminati ^{71a,71b}, M. Carnesale ^{75a,75b}, S. Caron ¹¹³, E. Carquin ^{137f}, S. Carrá ^{71a,71b}, G. Carratta ^{23b,23a}, F. Carrio Argos ^{33g}, J.W.S. Carter ¹⁵⁵, T.M. Carter ⁵², M.P. Casado ^{13,j}, A.F. Casha ¹⁵⁵, E.G. Castiglia ¹⁷², F.L. Castillo ^{63a}, L. Castillo Garcia ¹³, V. Castillo Gimenez ¹⁶³, N.F. Castro ^{130a,130e}, A. Catinaccio ³⁶, J.R. Catmore ¹²⁵, V. Cavaliere ²⁹, N. Cavalli ^{23b,23a}, V. Cavasinni ^{74a,74b}, E. Celebi ^{21a}, F. Celli ¹²⁶, M.S. Centonze ^{70a,70b}, K. Cerny ¹²², A.S. Cerqueira ^{82a}, A. Cerri ¹⁴⁶, L. Cerrito ^{76a,76b}, F. Cerutti ^{17a}, A. Cervelli ^{23b}, S.A. Cetin ^{21d}, Z. Chadi ^{35a}, D. Chakraborty ¹¹⁵, M. Chala ^{130f}, J. Chan ¹⁷⁰, W.Y. Chan ¹⁵³, J.D. Chapman ³², B. Chargeishvili ^{149b}, D.G. Charlton ²⁰, T.P. Charman ⁹⁴, M. Chatterjee ¹⁹, S. Chekanov ⁶, S.V. Chekulaev ^{156a}, G.A. Chelkov ^{38,a}, A. Chen ¹⁰⁶, B. Chen ¹⁵¹, B. Chen ¹⁶⁵, H. Chen ^{14c},

H. Chen ^{ib29}, J. Chen ^{ib62c}, J. Chen ^{ib142}, S. Chen ^{ib153}, S.J. Chen ^{ib14c}, X. Chen ^{ib62c}, X. Chen ^{ib14b,ai},
Y. Chen ^{ib62a}, C.L. Cheng ^{ib170}, H.C. Cheng ^{ib64a}, S. Cheong ^{ib143}, A. Cheplakov ^{ib38},
E. Cheremushkina ^{ib48}, E. Cherepanova ^{ib114}, R. Cherkaoui El Moursli ^{ib35e}, E. Cheu ^{ib7}, K. Cheung ^{ib65},
L. Chevalier ^{ib135}, V. Chiarella ^{ib53}, G. Chiarelli ^{ib74a}, N. Chiedde ^{ib102}, G. Chiodini ^{ib70a},
A.S. Chisholm ^{ib20}, A. Chitan ^{ib27b}, M. Chitishvili ^{ib163}, Y.H. Chiu ^{ib165}, M.V. Chizhov ^{ib38}, K. Choi ^{ib11},
A.R. Chomont ^{ib75a,75b}, Y. Chou ^{ib103}, E.Y.S. Chow ^{ib114}, T. Chowdhury ^{ib33g}, L.D. Christopher ^{ib33g},
K.L. Chu ^{ib64a}, M.C. Chu ^{ib64a}, X. Chu ^{ib14a,14d}, J. Chudoba ^{ib131}, J.J. Chwastowski ^{ib86}, D. Cieri ^{ib110},
K.M. Ciesla ^{ib85a}, V. Cindro ^{ib93}, A. Ciocio ^{ib17a}, F. Ciotto ^{ib72a,72b}, Z.H. Citron ^{ib169,n}, M. Citterio ^{ib71a},
D.A. Ciubotaru ^{ib27b}, B.M. Ciungu ^{ib155}, A. Clark ^{ib56}, P.J. Clark ^{ib52}, J.M. Clavijo Columbie ^{ib48},
S.E. Clawson ^{ib101}, C. Clement ^{ib47a,47b}, J. Clercx ^{ib48}, L. Clissa ^{ib23b,23a}, Y. Coadou ^{ib102},
M. Cobal ^{ib69a,69c}, A. Cocco ^{ib57b}, R.F. Coelho Barrue ^{ib130a}, R. Coelho Lopes De Sa ^{ib103},
S. Coelli ^{ib71a}, H. Cohen ^{ib151}, A.E.C. Coimbra ^{ib71a,71b}, B. Cole ^{ib41}, J. Collot ^{ib60},
P. Conde Muiño ^{ib130a,130g}, M.P. Connell ^{ib33c}, S.H. Connell ^{ib33c}, I.A. Connolly ^{ib59}, E.I. Conroy ^{ib126},
F. Conventi ^{ib72a,ak}, H.G. Cooke ^{ib20}, A.M. Cooper-Sarkar ^{ib126}, F. Cormier ^{ib164}, L.D. Corpe ^{ib36},
M. Corradi ^{ib75a,75b}, E.E. Corrigan ^{ib98}, F. Corriveau ^{ib104,z}, A. Cortes-Gonzalez ^{ib18}, M.J. Costa ^{ib163},
F. Costanza ^{ib4}, D. Costanzo ^{ib139}, B.M. Cote ^{ib119}, G. Cowan ^{ib95}, J.W. Cowley ^{ib32}, K. Cranmer ^{ib117},
S. Crépe-Renaudin ^{ib60}, F. Crescioli ^{ib127}, M. Cristinziani ^{ib141}, M. Cristoforetti ^{ib78a,78b,d}, V. Croft ^{ib158},
G. Crosetti ^{ib43b,43a}, A. Cueto ^{ib36}, T. Cuhadar Donszelmann ^{ib160}, H. Cui ^{ib14a,14d}, Z. Cui ^{ib7},
W.R. Cunningham ^{ib59}, F. Curcio ^{ib43b,43a}, P. Czodrowski ^{ib36}, M.M. Czurylo ^{ib63b},
M.J. Da Cunha Sargedas De Sousa ^{ib62a}, J.V. Da Fonseca Pinto ^{ib82b}, C. Da Via ^{ib101}, W. Dabrowski ^{ib85a},
T. Dado ^{ib49}, S. Dahbi ^{ib33g}, T. Dai ^{ib106}, C. Dallapiccola ^{ib103}, M. Dam ^{ib42}, G. D'amen ^{ib29},
V. D'Amico ^{ib109}, J. Damp ^{ib100}, J.R. Dandoy ^{ib128}, M.F. Daneri ^{ib30}, M. Danninger ^{ib142}, V. Dao ^{ib36},
G. Darbo ^{ib57b}, S. Darmora ^{ib6}, S.J. Das ^{ib29,am}, S. D'Auria ^{ib71a,71b}, C. David ^{ib156b}, T. Davidek ^{ib133},
D.R. Davis ^{ib51}, B. Davis-Purcell ^{ib34}, I. Dawson ^{ib94}, K. De ^{ib8}, R. De Asmundis ^{ib72a}, M. De Beurs ^{ib114},
N. De Biase ^{ib48}, S. De Castro ^{ib23b,23a}, N. De Groot ^{ib113}, P. de Jong ^{ib114}, H. De la Torre ^{ib107},
A. De Maria ^{ib14c}, A. De Salvo ^{ib75a}, U. De Sanctis ^{ib76a,76b}, A. De Santo ^{ib146},
J.B. De Vivie De Regie ^{ib60}, D.V. Dedovich ^{ib38}, J. Degens ^{ib114}, A.M. Deiana ^{ib44}, F. Del Corso ^{ib23b,23a},
J. Del Peso ^{ib99}, F. Del Rio ^{ib63a}, F. Deliot ^{ib135}, C.M. Delitzsch ^{ib49}, M. Della Pietra ^{ib72a,72b},
D. Della Volpe ^{ib56}, A. Dell'Acqua ^{ib36}, L. Dell'Asta ^{ib71a,71b}, M. Delmastro ^{ib4}, P.A. Delsart ^{ib60},
S. Demers ^{ib172}, M. Demichev ^{ib38}, S.P. Denisov ^{ib37}, L. D'Eramo ^{ib115}, D. Derendarz ^{ib86}, F. Derue ^{ib127},
P. Dervan ^{ib92}, K. Desch ^{ib24}, K. Dette ^{ib155}, C. Deutsch ^{ib24}, F.A. Di Bello ^{ib57b,57a},
A. Di Ciaccio ^{ib76a,76b}, L. Di Ciaccio ^{ib4}, A. Di Domenico ^{ib75a,75b}, C. Di Donato ^{ib72a,72b},
A. Di Girolamo ^{ib36}, G. Di Gregorio ^{ib5}, A. Di Luca ^{ib78a,78b}, B. Di Micco ^{ib77a,77b}, R. Di Nardo ^{ib77a,77b},
C. Diaconu ^{ib102}, F.A. Dias ^{ib114}, T. Dias Do Vale ^{ib142}, M.A. Diaz ^{ib137a,137b}, F.G. Diaz Capriles ^{ib24},
M. Didenko ^{ib163}, E.B. Diehl ^{ib106}, L. Diehl ^{ib54}, S. Díez Cornell ^{ib48}, C. Diez Pardo ^{ib141},
C. Dimitriadi ^{ib24,161}, A. Dimitrievska ^{ib17a}, J. Dingfelder ^{ib24}, I-M. Dinu ^{ib27b}, S.J. Dittmeier ^{ib63b},
F. Dittus ^{ib36}, F. Djama ^{ib102}, T. Djobava ^{ib149b}, J.I. Djuvsland ^{ib16}, C. Doglioni ^{ib101,98}, J. Dolejsi ^{ib133},
Z. Dolezal ^{ib133}, M. Donadelli ^{ib82c}, B. Dong ^{ib107}, J. Donini ^{ib40}, A. D'Onofrio ^{ib77a,77b},
M. D'Onofrio ^{ib92}, J. Dopke ^{ib134}, A. Doria ^{ib72a}, M.T. Dova ^{ib90}, A.T. Doyle ^{ib59}, M.A. Draguet ^{ib126},
E. Drechsler ^{ib142}, E. Dreyer ^{ib169}, I. Drivas-koulouris ^{ib10}, A.S. Drobac ^{ib158}, M. Drozdova ^{ib56},
D. Du ^{ib62a}, T.A. du Pree ^{ib114}, F. Dubinin ^{ib37}, M. Dubovsky ^{ib28a}, E. Duchovni ^{ib169}, G. Duckeck ^{ib109},
O.A. Ducu ^{ib27b}, D. Duda ^{ib110}, A. Dudarev ^{ib36}, M. D'uffizi ^{ib101}, L. Duflot ^{ib66}, M. Dührssen ^{ib36},
C. Dülsen ^{ib171}, A.E. Dumitriu ^{ib27b}, M. Dunford ^{ib63a}, S. Dungs ^{ib49}, K. Dunne ^{ib47a,47b},

A. Duperrin ¹⁰², H. Duran Yildiz ^{3a}, M. Düren ⁵⁸, A. Durglishvili ^{149b}, B.L. Dwyer ¹¹⁵,
 G.I. Dyckes ^{17a}, M. Dyndal ^{85a}, S. Dysch ¹⁰¹, B.S. Dziedzic ⁸⁶, Z.O. Earnshaw ¹⁴⁶,
 B. Eckerova ^{28a}, S. Eggebrecht ⁵⁵, M.G. Eggleston ⁵¹, E. Egidio Purcino De Souza ¹²⁷, L.F. Ehrke ⁵⁶,
 G. Eigen ¹⁶, K. Einsweiler ^{17a}, T. Ekelof ¹⁶¹, P.A. Ekman ⁹⁸, Y. El Ghazali ^{35b},
 H. El Jarrari ^{35e,148}, A. El Moussaouy ^{35a}, V. Ellajosyula ¹⁶¹, M. Ellert ¹⁶¹, F. Ellinghaus ¹⁷¹,
 A.A. Elliot ⁹⁴, N. Ellis ³⁶, J. Elmsheuser ²⁹, M. Elsing ³⁶, D. Emelianov ¹³⁴, A. Emerman ⁴¹,
 Y. Enari ¹⁵³, I. Ene ^{17a}, S. Epari ¹³, J. Erdmann ^{49,ag}, A. Ereditato ¹⁹, P.A. Erland ⁸⁶,
 M. Errenst ¹⁷¹, M. Escalier ⁶⁶, C. Escobar ¹⁶³, E. Etzion ¹⁵¹, G. Evans ^{130a}, H. Evans ⁶⁸,
 M.O. Evans ¹⁴⁶, A. Ezhilov ³⁷, S. Ezzarqtouni ^{35a}, F. Fabbri ⁵⁹, L. Fabbri ^{23b,23a}, G. Facini ⁹⁶,
 V. Fadeyev ¹³⁶, R.M. Fakhrutdinov ³⁷, S. Falciano ^{75a}, L.F. Falda Ulhoa Coelho ³⁶, P.J. Falke ²⁴,
 S. Falke ³⁶, J. Faltova ¹³³, Y. Fan ^{14a}, Y. Fang ^{14a,14d}, G. Fanourakis ⁴⁶, M. Fanti ^{71a,71b},
 M. Faraj ^{69a,69b}, Z. Farazpay ⁹⁷, A. Farbin ⁸, A. Farilla ^{77a}, T. Farooque ¹⁰⁷, S.M. Farrington ⁵²,
 F. Fassi ^{35e}, D. Fassouliotis ⁹, M. Fauci Giannelli ^{76a,76b}, W.J. Fawcett ³², L. Fayard ⁶⁶,
 P. Federicova ¹³¹, O.L. Fedin ^{37,a}, G. Fedotov ³⁷, M. Feickert ¹⁷⁰, L. Feligioni ¹⁰², A. Fell ¹³⁹,
 D.E. Fellers ¹²³, C. Feng ^{62b}, M. Feng ^{14b}, Z. Feng ¹¹⁴, M.J. Fenton ¹⁶⁰, A.B. Fenyuk ³⁷,
 L. Ferencz ⁴⁸, R.A.M. Ferguson ⁹¹, S.I. Fernandez Luengo ^{137f}, J. Ferrando ⁴⁸, A. Ferrari ¹⁶¹,
 P. Ferrari ^{114,113}, R. Ferrari ^{73a}, D. Ferrere ⁵⁶, C. Ferretti ¹⁰⁶, F. Fiedler ¹⁰⁰, A. Filipčič ⁹³,
 E.K. Filmer ¹, F. Filthaut ¹¹³, M.C.N. Fiolhais ^{130a,130c,c}, L. Fiorini ¹⁶³, F. Fischer ¹⁴¹,
 W.C. Fisher ¹⁰⁷, T. Fitschen ¹⁰¹, I. Fleck ¹⁴¹, P. Fleischmann ¹⁰⁶, T. Flick ¹⁷¹, L. Flores ¹²⁸,
 M. Flores ^{33d,af}, L.R. Flores Castillo ^{64a}, F.M. Follega ^{78a,78b}, N. Fomin ¹⁶, J.H. Foo ¹⁵⁵,
 B.C. Forland ⁶⁸, A. Formica ¹³⁵, A.C. Forti ¹⁰¹, E. Fortin ¹⁰², A.W. Fortman ⁶¹, M.G. Foti ^{17a},
 L. Fountas ^{9,k}, D. Fournier ⁶⁶, H. Fox ⁹¹, P. Francavilla ^{74a,74b}, S. Francescato ⁶¹,
 S. Franchellucci ⁵⁶, M. Franchini ^{23b,23a}, S. Franchino ^{63a}, D. Francis ³⁶, L. Franco ¹¹³,
 L. Franconi ¹⁹, M. Franklin ⁶¹, G. Frattari ²⁶, A.C. Freegard ⁹⁴, P.M. Freeman ²⁰, W.S. Freund ^{82b},
 N. Fritzsche ⁵⁰, A. Froch ⁵⁴, D. Froidevaux ³⁶, J.A. Frost ¹²⁶, Y. Fu ^{62a}, M. Fujimoto ¹¹⁸,
 E. Fullana Torregrosa ^{163,*}, J. Fuster ¹⁶³, A. Gabrielli ^{23b,23a}, A. Gabrielli ¹⁵⁵, P. Gadow ⁴⁸,
 G. Gagliardi ^{57b,57a}, L.G. Gagnon ^{17a}, G.E. Gallardo ¹²⁶, E.J. Gallas ¹²⁶, B.J. Gallop ¹³⁴,
 R. Gamboa Goni ⁹⁴, K.K. Gan ¹¹⁹, S. Ganguly ¹⁵³, J. Gao ^{62a}, Y. Gao ⁵²,
 F.M. Garay Walls ^{137a,137b}, B. Garcia ^{29,am}, C. García ¹⁶³, J.E. García Navarro ¹⁶³,
 M. Garcia-Sciveres ^{17a}, R.W. Gardner ³⁹, D. Garg ⁸⁰, R.B. Garg ^{143,r}, C.A. Garner ¹⁵⁵,
 V. Garonne ²⁹, S.J. Gasiorowski ¹³⁸, P. Gaspar ^{82b}, G. Gaudio ^{73a}, V. Gautam ¹³, P. Gauzzi ^{75a,75b},
 I.L. Gavrilenko ³⁷, A. Gavriluk ³⁷, C. Gay ¹⁶⁴, G. Gaycken ⁴⁸, E.N. Gazis ¹⁰,
 A.A. Geanta ^{27b,27e}, C.M. Gee ¹³⁶, J. Geisen ⁹⁸, C. Gemme ^{57b}, M.H. Genest ⁶⁰,
 S. Gentile ^{75a,75b}, S. George ⁹⁵, W.F. George ²⁰, T. Gerialis ⁴⁶, L.O. Gerlach ⁵⁵,
 P. Gessinger-Befurt ³⁶, M. Ghasemi Bostanabad ¹⁶⁵, M. Ghneimat ¹⁴¹, K. Ghorbanian ⁹⁴,
 A. Ghosal ¹⁴¹, A. Ghosh ¹⁶⁰, A. Ghosh ⁷, B. Giacobbe ^{23b}, S. Giagu ^{75a,75b}, P. Giannetti ^{74a},
 A. Giannini ^{62a}, S.M. Gibson ⁹⁵, M. Gignac ¹³⁶, D.T. Gil ^{85b}, A.K. Gilbert ^{85a}, B.J. Gilbert ⁴¹,
 D. Gillberg ³⁴, G. Gilles ¹¹⁴, N.E.K. Gillwald ⁴⁸, L. Ginabat ¹²⁷, D.M. Gingrich ^{2,aj},
 M.P. Giordani ^{69a,69c}, P.F. Giraud ¹³⁵, G. Giugliarelli ^{69a,69c}, D. Giugni ^{71a}, F. Giuli ³⁶,
 I. Gkialas ^{9,k}, L.K. Gladilin ³⁷, C. Glasman ⁹⁹, G.R. Gledhill ¹²³, M. Glisic ¹²³, I. Gnesi ^{43b,g},
 Y. Go ^{29,am}, M. Goblirsch-Kolb ²⁶, B. Gocke ⁴⁹, D. Godin ¹⁰⁸, B. Gokturk ^{21a}, S. Goldfarb ¹⁰⁵,
 T. Golling ⁵⁶, M.G.D. Gololo ^{33g}, D. Golubkov ³⁷, J.P. Gombas ¹⁰⁷, A. Gomes ^{130a,130b},
 G. Gomes Da Silva ¹⁴¹, A.J. Gomez Delegido ¹⁶³, R. Goncalves Gama ⁵⁵, R. Gonçalves ^{130a,130c},

G. Gonella ¹²³, L. Gonella ²⁰, A. Gongadze ³⁸, F. Gonnella ²⁰, J.L. Gonski ⁴¹,
R.Y. González Andana ⁵², S. González de la Hoz ¹⁶³, S. Gonzalez Fernandez ¹³,
R. Gonzalez Lopez ⁹², C. Gonzalez Renteria ^{17a}, R. Gonzalez Suarez ¹⁶¹, S. Gonzalez-Sevilla ⁵⁶,
G.R. Gonzalvo Rodriguez ¹⁶³, L. Goossens ³⁶, N.A. Gorasia ²⁰, P.A. Gorbounov ³⁷, B. Gorini ³⁶,
E. Gorini ^{70a,70b}, A. Gorišek ⁹³, A.T. Goshaw ⁵¹, M.I. Gostkin ³⁸, S. Goswami ¹²¹,
C.A. Gottardo ³⁶, M. Gouighri ^{35b}, V. Goumarre ⁴⁸, A.G. Goussiou ¹³⁸, N. Govender ^{33c},
C. Goy ⁴, I. Grabowska-Bold ^{85a}, K. Graham ³⁴, E. Gramstad ¹²⁵, S. Grancagnolo ¹⁸,
M. Grandi ¹⁴⁶, V. Gratchev ^{37,*}, P.M. Gravila ^{27f}, F.G. Gravili ^{70a,70b}, H.M. Gray ^{17a},
M. Greco ^{70a,70b}, C. Grefe ²⁴, I.M. Gregor ⁴⁸, P. Grenier ¹⁴³, C. Grieco ¹³, A.A. Grillo ¹³⁶,
K. Grimm ^{31,o}, S. Grinstein ^{13,w}, J.-F. Grivaz ⁶⁶, E. Gross ¹⁶⁹, J. Grosse-Knetter ⁵⁵, C. Grud ¹⁰⁶,
A. Grummer ¹¹², J.C. Grundy ¹²⁶, L. Guan ¹⁰⁶, W. Guan ¹⁷⁰, C. Gubbels ¹⁶⁴,
J.G.R. Guerrero Rojas ¹⁶³, G. Guerrieri ^{69a,69b}, F. Guescini ¹¹⁰, R. Gugel ¹⁰⁰, J.A.M. Guhit ¹⁰⁶,
A. Guida ⁴⁸, T. Guillemin ⁴, E. Guillon ^{167,134}, S. Guindon ³⁶, F. Guo ^{14a,14d}, J. Guo ^{62c},
L. Guo ⁶⁶, Y. Guo ¹⁰⁶, R. Gupta ⁴⁸, S. Gurbuz ²⁴, S.S. Gurdasani ⁵⁴, G. Gustavino ³⁶,
M. Guth ⁵⁶, P. Gutierrez ¹²⁰, L.F. Gutierrez Zagazeta ¹²⁸, C. Gutsche ⁹⁶, C. Guyot ¹³⁵,
C. Gwenlan ¹²⁶, C.B. Gwilliam ⁹², E.S. Haaland ¹²⁵, A. Haas ¹¹⁷, M. Habedank ⁴⁸, C. Haber ^{17a},
H.K. Hadavand ⁸, A. Hadeef ¹⁰⁰, S. Hadzic ¹¹⁰, E.H. Haines ⁹⁶, M. Haleem ¹⁶⁶, J. Haley ¹²¹,
J.J. Hall ¹³⁹, G.D. Hallowell ¹⁰², L. Halser ¹⁹, K. Hamano ¹⁶⁵, H. Hamdaoui ^{35e}, M. Hamer ²⁴,
G.N. Hamity ⁵², J. Han ^{62b}, K. Han ^{62a}, L. Han ^{14c}, L. Han ^{62a}, S. Han ^{17a}, Y.F. Han ¹⁵⁵,
K. Hanagaki ⁸³, M. Hance ¹³⁶, D.A. Hangal ^{41,ae}, H. Hanif ¹⁴², M.D. Hank ³⁹, R. Hankache ¹⁰¹,
J.B. Hansen ⁴², J.D. Hansen ⁴², P.H. Hansen ⁴², K. Hara ¹⁵⁷, D. Harada ⁵⁶, T. Harenberg ¹⁷¹,
S. Harkusha ³⁷, Y.T. Harris ¹²⁶, N.M. Harrison ¹¹⁹, P.F. Harrison ¹⁶⁷, N.M. Hartman ¹⁴³,
N.M. Hartmann ¹⁰⁹, Y. Hasegawa ¹⁴⁰, A. Hasib ⁵², S. Haug ¹⁹, R. Hauser ¹⁰⁷, M. Havranek ¹³²,
C.M. Hawkes ²⁰, R.J. Hawkings ³⁶, S. Hayashida ¹¹¹, D. Hayden ¹⁰⁷, C. Hayes ¹⁰⁶,
R.L. Hayes ¹⁶⁴, C.P. Hays ¹²⁶, J.M. Hays ⁹⁴, H.S. Hayward ⁹², F. He ^{62a}, Y. He ¹⁵⁴, Y. He ¹²⁷,
M.P. Heath ⁵², V. Hedberg ⁹⁸, A.L. Heggelund ¹²⁵, N.D. Hehir ⁹⁴, C. Heidegger ⁵⁴,
K.K. Heidegger ⁵⁴, W.D. Heidorn ⁸¹, J. Heilman ³⁴, S. Heim ⁴⁸, T. Heim ^{17a}, J.G. Heinlein ¹²⁸,
J.J. Heinrich ¹²³, L. Heinrich ^{110,ah}, J. Hejbal ¹³¹, L. Helary ⁴⁸, A. Held ¹⁷⁰, S. Hellesund ¹²⁵,
C.M. Helling ¹⁶⁴, S. Hellman ^{47a,47b}, C. Helsens ³⁶, R.C.W. Henderson ⁹¹, L. Henkelmann ³²,
A.M. Henriques Correia ³⁶, H. Herde ⁹⁸, Y. Hernández Jiménez ¹⁴⁵, L.M. Herrmann ²⁴,
M.G. Herrmann ¹⁰⁹, T. Herrmann ⁵⁰, G. Herten ⁵⁴, R. Hertenberger ¹⁰⁹, L. Hervas ³⁶,
N.P. Hessey ^{156a}, H. Hibi ⁸⁴, E. Higón-Rodriguez ¹⁶³, S.J. Hillier ²⁰, I. Hinchliffe ^{17a},
F. Hinterkeuser ²⁴, M. Hirose ¹²⁴, S. Hirose ¹⁵⁷, D. Hirschbuehl ¹⁷¹, T.G. Hitchings ¹⁰¹, B. Hiti ⁹³,
J. Hobbs ¹⁴⁵, R. Hobincu ^{27e}, N. Hod ¹⁶⁹, M.C. Hodgkinson ¹³⁹, B.H. Hodgkinson ³²,
A. Hoecker ³⁶, J. Hofer ⁴⁸, D. Hohn ⁵⁴, T. Holm ²⁴, M. Holzbock ¹¹⁰, L.B.A.H. Hommels ³²,
B.P. Honan ¹⁰¹, J. Hong ^{62c}, T.M. Hong ¹²⁹, J.C. Honig ⁵⁴, A. Hönle ¹¹⁰, B.H. Hooberman ¹⁶²,
W.H. Hopkins ⁶, Y. Horii ¹¹¹, S. Hou ¹⁴⁸, A.S. Howard ⁹³, J. Howarth ⁵⁹, J. Hoya ⁶,
M. Hrabovsky ¹²², A. Hrynevich ⁴⁸, T. Hryn'ova ⁴, P.J. Hsu ⁶⁵, S.-C. Hsu ¹³⁸, Q. Hu ⁴¹,
Y.F. Hu ^{14a,14d,al}, D.P. Huang ⁹⁶, S. Huang ^{64b}, X. Huang ^{14c}, Y. Huang ^{62a}, Y. Huang ^{14a},
Z. Huang ¹⁰¹, Z. Hubacek ¹³², M. Huebner ²⁴, F. Huegging ²⁴, T.B. Huffman ¹²⁶, M. Huhtinen ³⁶,
S.K. Huiberts ¹⁶, R. Hulsken ¹⁰⁴, N. Huseynov ^{12,a}, J. Huston ¹⁰⁷, J. Huth ⁶¹, R. Hyneman ¹⁴³,
S. Hyrych ^{28a}, G. Iacobucci ⁵⁶, G. Iakovidis ²⁹, I. Ibragimov ¹⁴¹, L. Iconomidou-Fayard ⁶⁶,
P. Iengo ^{72a,72b}, R. Iguchi ¹⁵³, T. Iizawa ⁵⁶, Y. Ikegami ⁸³, A. Ilg ¹⁹, N. Ilic ¹⁵⁵, H. Imam ^{35a},

T. Ingebretsen Carlson ^{47a,47b}, G. Introzzi ^{73a,73b}, M. Iodice ^{77a}, V. Ippolito ^{75a,75b}, M. Ishino ¹⁵³, W. Islam ¹⁷⁰, C. Issever ^{18,48}, S. Istin ^{21a}, H. Ito ¹⁶⁸, J.M. Iturbe Ponce ^{64a}, R. Iuppa ^{78a,78b}, A. Ivina ¹⁶⁹, J.M. Izen ⁴⁵, V. Izzo ^{72a}, P. Jacka ^{131,132}, P. Jackson ¹, R.M. Jacobs ⁴⁸, B.P. Jaeger ¹⁴², C.S. Jagfeld ¹⁰⁹, P. Jain ⁵⁴, G. Jäkel ¹⁷¹, K. Jakobs ⁵⁴, T. Jakoubek ¹⁶⁹, J. Jamieson ⁵⁹, K.W. Janas ^{85a}, G. Jarlskog ⁹⁸, A.E. Jaspan ⁹², M. Javurkova ¹⁰³, F. Jeanneau ¹³⁵, L. Jeanty ¹²³, J. Jejelava ^{149a,ac}, P. Jenni ^{54,h}, C.E. Jessiman ³⁴, S. Jézéquel ⁴, C. Jia ^{62b}, J. Jia ¹⁴⁵, X. Jia ⁶¹, X. Jia ^{14a,14d}, Z. Jia ^{14c}, Y. Jiang ^{62a}, S. Jiggins ⁵², J. Jimenez Pena ¹¹⁰, S. Jin ^{14c}, A. Jinaru ^{27b}, O. Jinnouchi ¹⁵⁴, P. Johansson ¹³⁹, K.A. Johns ⁷, J.W. Johnson ¹³⁶, D.M. Jones ³², E. Jones ¹⁶⁷, P. Jones ³², R.W.L. Jones ⁹¹, T.J. Jones ⁹², R. Joshi ¹¹⁹, J. Jovicevic ¹⁵, X. Ju ^{17a}, J.J. Junggeburth ³⁶, T. Junkermann ^{63a}, A. Juste Rozas ^{13,w}, S. Kabana ^{137e}, A. Kaczmarzka ⁸⁶, M. Kado ^{75a,75b}, H. Kagan ¹¹⁹, M. Kagan ¹⁴³, A. Kahn ⁴¹, A. Kahn ¹²⁸, C. Kahra ¹⁰⁰, T. Kaji ¹⁶⁸, E. Kajomovitz ¹⁵⁰, N. Kakati ¹⁶⁹, C.W. Kalderon ²⁹, A. Kamenshchikov ¹⁵⁵, S. Kanayama ¹⁵⁴, N.J. Kang ¹³⁶, D. Kar ^{33g}, K. Karava ¹²⁶, M.J. Kareem ^{156b}, E. Karentzos ⁵⁴, I. Karkanas ^{152,f}, S.N. Karpov ³⁸, Z.M. Karpova ³⁸, V. Kartvelishvili ⁹¹, A.N. Karyukhin ³⁷, E. Kasimi ^{152,f}, C. Kato ^{62d}, J. Katzy ⁴⁸, S. Kaur ³⁴, K. Kawade ¹⁴⁰, K. Kawagoe ⁸⁹, T. Kawamoto ¹³⁵, G. Kawamura ⁵⁵, E.F. Kay ¹⁶⁵, F.I. Kaya ¹⁵⁸, S. Kazakos ¹³, V.F. Kazanin ³⁷, Y. Ke ¹⁴⁵, J.M. Keaveney ^{33a}, R. Keeler ¹⁶⁵, G.V. Kehris ⁶¹, J.S. Keller ³⁴, A.S. Kelly ⁹⁶, D. Kelsey ¹⁴⁶, J.J. Kempster ²⁰, K.E. Kennedy ⁴¹, P.D. Kennedy ¹⁰⁰, O. Kepka ¹³¹, B.P. Kerridge ¹⁶⁷, S. Kersten ¹⁷¹, B.P. Kerševan ⁹³, S. Keshri ⁶⁶, L. Keszeghova ^{28a}, S. Ketabchi Haghighat ¹⁵⁵, M. Khandoga ¹²⁷, A. Khanov ¹²¹, A.G. Kharlamov ³⁷, T. Kharlamova ³⁷, E.E. Khoda ¹³⁸, T.J. Khoo ¹⁸, G. Khorauli ¹⁶⁶, J. Khubua ^{149b}, Y.A.R. Khwaira ⁶⁶, M. Kiehn ³⁶, A. Kilgallon ¹²³, D.W. Kim ^{47a,47b}, E. Kim ¹⁵⁴, Y.K. Kim ³⁹, N. Kimura ⁹⁶, A. Kirchhoff ⁵⁵, D. Kirchmeier ⁵⁰, C. Kirfel ²⁴, J. Kirk ¹³⁴, A.E. Kiryunin ¹¹⁰, T. Kishimoto ¹⁵³, D.P. Kisliuk ¹⁵⁵, C. Kitsaki ¹⁰, O. Kivernyk ²⁴, M. Klassen ^{63a}, C. Klein ³⁴, L. Klein ¹⁶⁶, M.H. Klein ¹⁰⁶, M. Klein ⁹², S.B. Klein ⁵⁶, U. Klein ⁹², P. Klimek ³⁶, A. Klimentov ²⁹, F. Klimpel ¹¹⁰, T. Klioutchnikova ³⁶, P. Kluit ¹¹⁴, S. Kluth ¹¹⁰, E. Kneringer ⁷⁹, T.M. Knight ¹⁵⁵, A. Knue ⁵⁴, D. Kobayashi ⁸⁹, R. Kobayashi ⁸⁷, M. Kocian ¹⁴³, P. Kodyš ¹³³, D.M. Koeck ¹⁴⁶, P.T. Koenig ²⁴, T. Koffas ³⁴, M. Kolb ¹³⁵, I. Koletsou ⁴, T. Komarek ¹²², K. Köneke ⁵⁴, A.X.Y. Kong ¹, T. Kono ¹¹⁸, N. Konstantinidis ⁹⁶, B. Konya ⁹⁸, R. Kopeliansky ⁶⁸, S. Koperny ^{85a}, K. Korcyl ⁸⁶, K. Kordas ^{152,f}, G. Koren ¹⁵¹, A. Korn ⁹⁶, S. Korn ⁵⁵, I. Korolkov ¹³, N. Korotkova ³⁷, B. Kortman ¹¹⁴, O. Kortner ¹¹⁰, S. Kortner ¹¹⁰, W.H. Kostecka ¹¹⁵, V.V. Kostyukhin ¹⁴¹, A. Kotskechagia ¹³⁵, A. Kotwal ⁵¹, A. Koulouris ³⁶, A. Kourkouveli-Charalampidi ^{73a,73b}, C. Kourkouvelis ⁹, E. Kourlitis ⁶, O. Kovanda ¹⁴⁶, R. Kowalewski ¹⁶⁵, W. Kozanecki ¹³⁵, A.S. Kozhin ³⁷, V.A. Kramarenko ³⁷, G. Kramberger ⁹³, P. Kramer ¹⁰⁰, M.W. Krasny ¹²⁷, A. Krasznahorkay ³⁶, J.A. Kremer ¹⁰⁰, T. Kresse ⁵⁰, J. Kretschmar ⁹², K. Kreul ¹⁸, P. Krieger ¹⁵⁵, S. Krishnamurthy ¹⁰³, M. Krivos ¹³³, K. Krizka ^{17a}, K. Kroeninger ⁴⁹, H. Kroha ¹¹⁰, J. Kroll ¹³¹, J. Kroll ¹²⁸, K.S. Krowpman ¹⁰⁷, U. Kruchonak ³⁸, H. Krüger ²⁴, N. Krumnack ⁸¹, M.C. Kruse ⁵¹, J.A. Krzysiak ⁸⁶, O. Kuchinskaia ³⁷, S. Kудay ^{3a}, D. Kuechler ⁴⁸, J.T. Kuechler ⁴⁸, S. Kuehn ³⁶, R. Kuesters ⁵⁴, T. Kuhl ⁴⁸, V. Kukhtin ³⁸, Y. Kulchitsky ^{37,a}, S. Kuleshov ^{137d,137b}, M. Kumar ^{33g}, N. Kumari ¹⁰², A. Kupco ¹³¹, T. Kupfer ⁴⁹, A. Kupich ³⁷, O. Kuprash ⁵⁴, H. Kurashige ⁸⁴, L.L. Kurchaninov ^{156a}, Y.A. Kurochkin ³⁷, A. Kurova ³⁷, M. Kuze ¹⁵⁴, A.K. Kvam ¹⁰³, J. Kvita ¹²², T. Kwan ¹⁰⁴, K.W. Kwok ^{64a}, N.G. Kyriacou ¹⁰⁶, L.A.O. Laatu ¹⁰², C. Lacasta ¹⁶³, F. Lacava ^{75a,75b}, H. Lacker ¹⁸, D. Lacour ¹²⁷, N.N. Lad ⁹⁶,

E. Ladygin ³⁸, B. Laforge ¹²⁷, T. Lagouri ^{137e}, S. Lai ⁵⁵, I.K. Lakomic ^{85a}, N. Lalloue ⁶⁰,
 J.E. Lambert ¹²⁰, S. Lammers ⁶⁸, W. Lampl ⁷, C. Lampoudis ^{152,f}, A.N. Lancaster ¹¹⁵,
 E. Lançon ²⁹, U. Landgraf ⁵⁴, M.P.J. Landon ⁹⁴, V.S. Lang ⁵⁴, R.J. Langenberg ¹⁰³,
 A.J. Lankford ¹⁶⁰, F. Lanni ³⁶, K. Lantzsch ²⁴, A. Lanza ^{73a}, A. Lapertosa ^{57b,57a},
 J.F. Laporte ¹³⁵, T. Lari ^{71a}, F. Lasagni Manghi ^{23b}, M. Lassnig ³⁶, V. Latonova ¹³¹, T.S. Lau ^{64a},
 A. Laudrain ¹⁰⁰, A. Laurier ³⁴, S.D. Lawlor ⁹⁵, Z. Lawrence ¹⁰¹, M. Lazzaroni ^{71a,71b}, B. Le ¹⁰¹,
 B. Leban ⁹³, A. Lebedev ⁸¹, M. LeBlanc ³⁶, T. LeCompte ⁶, F. Ledroit-Guillon ⁶⁰, A.C.A. Lee ⁹⁶,
 G.R. Lee ¹⁶, L. Lee ⁶¹, S.C. Lee ¹⁴⁸, S. Lee ^{47a,47b}, T.F. Lee ⁹², L.L. Leeuw ^{33c},
 H.P. Lefebvre ⁹⁵, M. Lefebvre ¹⁶⁵, C. Leggett ^{17a}, K. Lehmann ¹⁴², G. Lehmann Miotto ³⁶,
 M. Leigh ⁵⁶, W.A. Leight ¹⁰³, A. Leisos ^{152,v}, M.A.L. Leite ^{82c}, C.E. Leitgeb ⁴⁸, R. Leitner ¹³³,
 K.J.C. Leney ⁴⁴, T. Lenz ²⁴, S. Leone ^{74a}, C. Leonidopoulos ⁵², A. Leopold ¹⁴⁴, C. Leroy ¹⁰⁸,
 R. Les ¹⁰⁷, C.G. Lester ³², M. Levchenko ³⁷, J. Levêque ⁴, D. Levin ¹⁰⁶, L.J. Levinson ¹⁶⁹,
 M.P. Lewicki ⁸⁶, D.J. Lewis ⁴, A. Li ⁵, B. Li ^{62b}, C. Li ^{62a}, C-Q. Li ^{62c}, H. Li ^{62a}, H. Li ^{62b},
 H. Li ^{14c}, H. Li ^{62b}, J. Li ^{62c}, K. Li ¹³⁸, L. Li ^{62c}, M. Li ^{14a,14d}, Q.Y. Li ^{62a}, S. Li ^{14a,14d},
 S. Li ^{62d,62c,e}, T. Li ^{62b}, X. Li ¹⁰⁴, Z. Li ^{62b}, Z. Li ¹²⁶, Z. Li ¹⁰⁴, Z. Li ⁹², Z. Li ^{14a,14d},
 Z. Liang ^{14a}, M. Liberatore ⁴⁸, B. Liberti ^{76a}, K. Lie ^{64c}, J. Lieber Marin ^{82b}, H. Lien ⁶⁸,
 K. Lin ¹⁰⁷, R.A. Linck ⁶⁸, R.E. Lindley ⁷, J.H. Lindon ², A. Linss ⁴⁸, E. Lipeles ¹²⁸,
 A. Lipniacka ¹⁶, A. Lister ¹⁶⁴, J.D. Little ⁴, B. Liu ^{14a}, B.X. Liu ¹⁴², D. Liu ^{62d,62c}, J.B. Liu ^{62a},
 J.K.K. Liu ³², K. Liu ^{62d,62c}, M. Liu ^{62a}, M.Y. Liu ^{62a}, P. Liu ^{14a}, Q. Liu ^{62d,138,62c}, X. Liu ^{62a},
 Y. Liu ^{14c,14d}, Y.L. Liu ¹⁰⁶, Y.W. Liu ^{62a}, M. Livan ^{73a,73b}, J. Llorente Merino ¹⁴², S.L. Lloyd ⁹⁴,
 E.M. Lobodzinska ⁴⁸, P. Loch ⁷, S. Loffredo ^{76a,76b}, T. Lohse ¹⁸, K. Lohwasser ¹³⁹,
 M. Lokajicek ¹³¹, J.D. Long ¹⁶², I. Longarini ¹⁶⁰, L. Longo ^{70a,70b}, R. Longo ¹⁶², I. Lopez Paz ⁶⁷,
 A. Lopez Solis ⁴⁸, J. Lorenz ¹⁰⁹, N. Lorenzo Martinez ⁴, A.M. Lory ¹⁰⁹, X. Lou ^{47a,47b},
 X. Lou ^{14a,14d}, A. Lounis ⁶⁶, J. Love ⁶, P.A. Love ⁹¹, J.J. Lozano Bahilo ¹⁶³, G. Lu ^{14a,14d},
 M. Lu ⁸⁰, S. Lu ¹²⁸, Y.J. Lu ⁶⁵, H.J. Lubatti ¹³⁸, C. Luci ^{75a,75b}, F.L. Lucio Alves ^{14c},
 A. Lucotte ⁶⁰, F. Luehring ⁶⁸, I. Luise ¹⁴⁵, O. Lukianchuk ⁶⁶, O. Lundberg ¹⁴⁴,
 B. Lund-Jensen ¹⁴⁴, N.A. Luongo ¹²³, M.S. Lutz ¹⁵¹, D. Lynn ²⁹, H. Lyons ⁹², R. Lysak ¹³¹,
 E. Lytken ⁹⁸, F. Lyu ^{14a}, V. Lyubushkin ³⁸, T. Lyubushkina ³⁸, M.M. Lyukova ¹⁴⁵, H. Ma ²⁹,
 L.L. Ma ^{62b}, Y. Ma ⁹⁶, D.M. Mac Donell ¹⁶⁵, G. Maccarrone ⁵³, J.C. MacDonald ¹³⁹,
 R. Madar ⁴⁰, W.F. Mader ⁵⁰, J. Maeda ⁸⁴, T. Maeno ²⁹, M. Maerker ⁵⁰, H. Maguire ¹³⁹,
 D.J. Mahon ⁴¹, A. Maio ^{130a,130b,130d}, K. Maj ^{85a}, O. Majersky ^{28a}, S. Majewski ¹²³,
 N. Makovec ⁶⁶, V. Maksimovic ¹⁵, B. Malaescu ¹²⁷, Pa. Malecki ⁸⁶, V.P. Maleev ³⁷, F. Malek ⁶⁰,
 D. Malito ^{43b,43a}, U. Mallik ⁸⁰, C. Malone ³², S. Maltezos ¹⁰, S. Malyukov ³⁸, J. Mamuzic ¹³,
 G. Mancini ⁵³, G. Manco ^{73a,73b}, J.P. Mandalia ⁹⁴, I. Mandić ⁹³, L. Manhaes de Andrade Filho ^{82a},
 I.M. Maniatis ¹⁶⁹, J. Manjarres Ramos ⁵⁰, D.C. Mankad ¹⁶⁹, A. Mann ¹⁰⁹, B. Mansoulie ¹³⁵,
 S. Manzoni ³⁶, A. Marantis ^{152,v}, G. Marchiori ⁵, M. Marcisovsky ¹³¹, C. Marcon ^{71a,71b},
 M. Marinescu ²⁰, M. Marjanovic ¹²⁰, E.J. Marshall ⁹¹, Z. Marshall ^{17a}, S. Marti-Garcia ¹⁶³,
 T.A. Martin ¹⁶⁷, V.J. Martin ⁵², B. Martin dit Latour ¹⁶, L. Martinelli ^{75a,75b}, M. Martinez ^{13,w},
 P. Martinez Agullo ¹⁶³, V.I. Martinez Outschoorn ¹⁰³, P. Martinez Suarez ¹³, S. Martin-Haugh ¹³⁴,
 V.S. Martoiu ^{27b}, A.C. Martyniuk ⁹⁶, A. Marzin ³⁶, S.R. Maschek ¹¹⁰, D. Mascione ^{78a,78b},
 L. Masetti ¹⁰⁰, T. Mashimo ¹⁵³, J. Masik ¹⁰¹, A.L. Maslennikov ³⁷, L. Massa ^{23b},
 P. Massarotti ^{72a,72b}, P. Mastrandrea ^{74a,74b}, A. Mastroberardino ^{43b,43a}, T. Masubuchi ¹⁵³,
 T. Mathisen ¹⁶¹, N. Matsuzawa ¹⁵³, J. Maurer ^{27b}, B. Maček ⁹³, D.A. Maximov ³⁷, R. Mazini ¹⁴⁸,

I. Maznas ^{152,f}, M. Mazza ¹⁰⁷, S.M. Mazza ¹³⁶, C. Mc Ginn ^{29,am}, J.P. Mc Gowan ¹⁰⁴,
 S.P. Mc Kee ¹⁰⁶, E.F. McDonald ¹⁰⁵, A.E. McDougall ¹¹⁴, J.A. Mcfayden ¹⁴⁶, G. Mchedlidze ^{149b},
 R.P. McKenzie ^{33g}, T.C. McLachlan ⁴⁸, D.J. McLaughlin ⁹⁶, K.D. McLean ¹⁶⁵, S.J. McMahon ¹³⁴,
 P.C. McNamara ¹⁰⁵, C.M. Mcpartland ⁹², R.A. McPherson ^{165,z}, T. Megy ⁴⁰, S. Mehlhase ¹⁰⁹,
 A. Mehta ⁹², B. Meirose ⁴⁵, D. Melini ¹⁵⁰, B.R. Mellado Garcia ^{33g}, A.H. Melo ⁵⁵, F. Meloni ⁴⁸,
 E.D. Mendes Gouveia ^{130a}, A.M. Mendes Jacques Da Costa ²⁰, H.Y. Meng ¹⁵⁵, L. Meng ⁹¹,
 S. Menke ¹¹⁰, M. Mentink ³⁶, E. Meoni ^{43b,43a}, C. Merlassino ¹²⁶, L. Merola ^{72a,72b},
 C. Meroni ^{71a}, G. Merz ¹⁰⁶, O. Meshkov ³⁷, J. Metcalfe ⁶, A.S. Mete ⁶, C. Meyer ⁶⁸,
 J-P. Meyer ¹³⁵, M. Michetti ¹⁸, R.P. Middleton ¹³⁴, L. Mijović ⁵², G. Mikenberg ¹⁶⁹,
 M. Mikestikova ¹³¹, M. Mikuž ⁹³, H. Mildner ¹³⁹, A. Milic ³⁶, C.D. Milke ⁴⁴, D.W. Miller ³⁹,
 L.S. Miller ³⁴, A. Milov ¹⁶⁹, D.A. Milstead ^{47a,47b}, T. Min ^{14c}, A.A. Minaenko ³⁷,
 I.A. Minashvili ^{149b}, L. Mince ⁵⁹, A.I. Mincer ¹¹⁷, B. Mindur ^{85a}, M. Mineev ³⁸, Y. Mino ⁸⁷,
 L.M. Mir ¹³, M. Miralles Lopez ¹⁶³, M. Mironova ¹²⁶, M.C. Missio ¹¹³, T. Mitani ¹⁶⁸,
 A. Mitra ¹⁶⁷, V.A. Mitsou ¹⁶³, O. Miu ¹⁵⁵, P.S. Miyagawa ⁹⁴, Y. Miyazaki ⁸⁹, A. Mizukami ⁸³,
 J.U. Mjörnmark ⁹⁸, T. Mkrтчyan ^{63a}, T. Mlinarevic ⁹⁶, M. Mlynarikova ³⁶, T. Moa ^{47a,47b},
 S. Mobius ⁵⁵, K. Mochizuki ¹⁰⁸, P. Moder ⁴⁸, P. Mogg ¹⁰⁹, A.F. Mohammed ^{14a,14d},
 S. Mohapatra ⁴¹, G. Mokgatitwane ^{33g}, B. Mondal ¹⁴¹, S. Mondal ¹³², K. Mönig ⁴⁸,
 E. Monnier ¹⁰², L. Monsonis Romero ¹⁶³, J. Montejo Berlingen ³⁶, M. Montella ¹¹⁹, F. Monticelli ⁹⁰,
 N. Morange ⁶⁶, A.L. Moreira De Carvalho ^{130a}, M. Moreno Llácer ¹⁶³, C. Moreno Martinez ⁵⁶,
 P. Morettini ^{57b}, S. Morgenstern ¹⁶⁷, M. Morii ⁶¹, M. Morinaga ¹⁵³, A.K. Morley ³⁶,
 F. Morodei ^{75a,75b}, L. Morvaj ³⁶, P. Moschovakos ³⁶, B. Moser ³⁶, M. Mosidze ^{149b},
 T. Moskalets ⁵⁴, P. Moskvitina ¹¹³, J. Moss ^{31,p}, E.J.W. Moyse ¹⁰³, O. Mtintsilana ^{33g},
 S. Muanza ¹⁰², J. Mueller ¹²⁹, D. Muenstermann ⁹¹, R. Müller ¹⁹, G.A. Mullier ¹⁶¹, J.J. Mullin ¹²⁸,
 D.P. Mungo ¹⁵⁵, J.L. Munoz Martinez ¹³, D. Munoz Perez ¹⁶³, F.J. Munoz Sanchez ¹⁰¹,
 M. Murin ¹⁰¹, W.J. Murray ^{167,134}, A. Murrone ^{71a,71b}, J.M. Muse ¹²⁰, M. Muškinja ^{17a},
 C. Mwewa ²⁹, A.G. Myagkov ^{37,a}, A.J. Myers ⁸, A.A. Myers ¹²⁹, G. Myers ⁶⁸, M. Myska ¹³²,
 B.P. Nachman ^{17a}, O. Nackenhorst ⁴⁹, A. Nag ⁵⁰, K. Nagai ¹²⁶, K. Nagano ⁸³, J.L. Nagle ^{29,am},
 E. Nagy ¹⁰², A.M. Nairz ³⁶, Y. Nakahama ⁸³, K. Nakamura ⁸³, H. Nanjo ¹²⁴, R. Narayan ⁴⁴,
 E.A. Narayanan ¹¹², I. Naryshkin ³⁷, M. Naseri ³⁴, C. Nass ²⁴, G. Navarro ^{22a},
 J. Navarro-Gonzalez ¹⁶³, R. Nayak ¹⁵¹, A. Nayaz ¹⁸, P.Y. Nechaeva ³⁷, F. Nechansky ⁴⁸,
 L. Nedic ¹²⁶, T.J. Neep ²⁰, A. Negri ^{73a,73b}, M. Negrini ^{23b}, C. Nellist ¹¹³, C. Nelson ¹⁰⁴,
 K. Nelson ¹⁰⁶, S. Nemecek ¹³¹, M. Nessi ^{36,i}, M.S. Neubauer ¹⁶², F. Neuhaus ¹⁰⁰, J. Neundorf ⁴⁸,
 R. Newhouse ¹⁶⁴, P.R. Newman ²⁰, C.W. Ng ¹²⁹, Y.S. Ng ¹⁸, Y.W.Y. Ng ⁴⁸, B. Ngair ^{35e},
 H.D.N. Nguyen ¹⁰⁸, R.B. Nickerson ¹²⁶, R. Nicolaidou ¹³⁵, J. Nielsen ¹³⁶, M. Niemeyer ⁵⁵,
 N. Nikiforou ³⁶, V. Nikolaenko ^{37,a}, I. Nikolic-Audit ¹²⁷, K. Nikolopoulos ²⁰, P. Nilsson ²⁹,
 I. Ninca ⁴⁸, H.R. Nindhito ⁵⁶, A. Nisati ^{75a}, N. Nishu ², R. Nisius ¹¹⁰, J-E. Nitschke ⁵⁰,
 E.K. Nkadimeng ^{33g}, S.J. Noacco Rosende ⁹⁰, T. Nobe ¹⁵³, D.L. Noel ³², Y. Noguchi ⁸⁷,
 T. Nommensen ¹⁴⁷, M.A. Nomura ²⁹, M.B. Norfolk ¹³⁹, R.R.B. Norisam ⁹⁶, B.J. Norman ³⁴,
 J. Novak ⁹³, T. Novak ⁴⁸, O. Novgorodova ⁵⁰, L. Novotny ¹³², R. Novotny ¹¹², L. Nozka ¹²²,
 K. Ntekas ¹⁶⁰, N.M.J. Nunes De Moura Junior ^{82b}, E. Nurse ⁹⁶, F.G. Oakham ^{34,aj}, J. Ocariz ¹²⁷,
 A. Ochi ⁸⁴, I. Ochoa ^{130a}, S. Oerdek ¹⁶¹, J.T. Offermann ³⁹, A. Ogrodnik ^{85a}, A. Oh ¹⁰¹,
 C.C. Ohm ¹⁴⁴, H. Oide ⁸³, R. Oishi ¹⁵³, M.L. Ojeda ⁴⁸, Y. Okazaki ⁸⁷, M.W. O’Keefe ⁹²,
 Y. Okumura ¹⁵³, A. Olariu ^{27b}, L.F. Oleiro Seabra ^{130a}, S.A. Olivares Pino ^{137e},

D. Oliveira Damazio ²⁹, D. Oliveira Goncalves ^{82a}, J.L. Oliver ¹⁶⁰, M.J.R. Olsson ¹⁶⁰,
A. Olszewski ⁸⁶, J. Olszowska ^{86,*}, Ö.O. Öncel ⁵⁴, D.C. O’Neil ¹⁴², A.P. O’Neill ¹⁹,
A. Onofre ^{130a,130e}, P.U.E. Onyisi ¹¹, M.J. Oreglia ³⁹, G.E. Orellana ⁹⁰, D. Orestano ^{77a,77b},
N. Orlando ¹³, R.S. Orr ¹⁵⁵, V. O’Shea ⁵⁹, R. Ospanov ^{62a}, G. Otero y Garzon ³⁰, H. Otono ⁸⁹,
P.S. Ott ^{63a}, G.J. Ottino ^{17a}, M. Ouchrif ^{35d}, J. Ouellette ^{29,am}, F. Ould-Saada ¹²⁵, M. Owen ⁵⁹,
R.E. Owen ¹³⁴, K.Y. Oyulmaz ^{21a}, V.E. Ozcan ^{21a}, N. Ozturk ⁸, S. Ozturk ^{21d}, J. Pacalt ¹²²,
H.A. Pacey ³², K. Pachal ⁵¹, A. Pacheco Pages ¹³, C. Padilla Aranda ¹³, G. Padovano ^{75a,75b},
S. Pagan Griso ^{17a}, G. Palacino ⁶⁸, A. Palazzo ^{70a,70b}, S. Palestini ³⁶, M. Palka ^{85b}, J. Pan ¹⁷²,
T. Pan ^{64a}, D.K. Panchal ¹¹, C.E. Pandini ¹¹⁴, J.G. Panduro Vazquez ⁹⁵, H. Pang ^{14b}, P. Pani ⁴⁸,
G. Panizzo ^{69a,69c}, L. Paolozzi ⁵⁶, C. Papadatos ¹⁰⁸, S. Parajuli ⁴⁴, A. Paramonov ⁶,
C. Paraskevopoulos ¹⁰, D. Paredes Hernandez ^{64b}, T.H. Park ¹⁵⁵, M.A. Parker ³², F. Parodi ^{57b,57a},
E.W. Parrish ¹¹⁵, V.A. Parrish ⁵², J.A. Parsons ⁴¹, U. Parzefall ⁵⁴, B. Pascual Dias ¹⁰⁸,
L. Pascual Dominguez ¹⁵¹, V.R. Pascuzzi ^{17a}, F. Pasquali ¹¹⁴, E. Pasqualucci ^{75a}, S. Passaggio ^{57b},
F. Pastore ⁹⁵, P. Pasuwan ^{47a,47b}, P. Patel ⁸⁶, J.R. Pater ¹⁰¹, T. Pauly ³⁶, J. Pearkes ¹⁴³,
M. Pedersen ¹²⁵, R. Pedro ^{130a}, S.V. Peleganchuk ³⁷, O. Penc ³⁶, E.A. Pender ⁵², C. Peng ^{64b},
H. Peng ^{62a}, K.E. Penski ¹⁰⁹, M. Penzin ³⁷, B.S. Peralva ^{82d}, A.P. Pereira Peixoto ⁶⁰,
L. Pereira Sanchez ^{47a,47b}, D.V. Perepelitsa ^{29,am}, E. Perez Codina ^{156a}, M. Perganti ¹⁰,
L. Perini ^{71a,71b,*}, H. Pernegger ³⁶, S. Perrella ³⁶, A. Perrevoort ¹¹³, O. Perrin ⁴⁰, K. Peters ⁴⁸,
R.F.Y. Peters ¹⁰¹, B.A. Petersen ³⁶, T.C. Petersen ⁴², E. Petit ¹⁰², V. Petousis ¹³²,
C. Petridou ^{152,f}, A. Petrukhin ¹⁴¹, M. Pettee ^{17a}, N.E. Pettersson ³⁶, A. Petukhov ³⁷,
K. Petukhova ¹³³, A. Peyaud ¹³⁵, R. Pezoa ^{137f}, L. Pezzotti ³⁶, G. Pezzullo ¹⁷², T.M. Pham ¹⁷⁰,
T. Pham ¹⁰⁵, P.W. Phillips ¹³⁴, M.W. Phipps ¹⁶², G. Piacquadio ¹⁴⁵, E. Pianori ^{17a},
F. Piazza ^{71a,71b}, R. Piegaia ³⁰, D. Pietreanu ^{27b}, A.D. Pilkington ¹⁰¹, M. Pinamonti ^{69a,69c},
J.L. Pinfold ², B.C. Pinheiro Pereira ^{130a}, C. Pitman Donaldson ⁹⁶, D.A. Pizzi ³⁴,
L. Pizzimento ^{76a,76b}, A. Pizzini ¹¹⁴, M.-A. Pleier ²⁹, V. Plesanovs ⁵⁴, V. Pleskot ¹³³, E. Plotnikova ³⁸,
G. Poddar ⁴, R. Poettgen ⁹⁸, L. Poggioli ¹²⁷, I. Pogrebnyak ¹⁰⁷, D. Pohl ²⁴, I. Pokharel ⁵⁵,
S. Polacek ¹³³, G. Polesello ^{73a}, A. Poley ^{142,156a}, R. Polifka ¹³², A. Polini ^{23b}, C.S. Pollard ¹⁶⁷,
Z.B. Pollock ¹¹⁹, V. Polychronakos ²⁹, E. Pompa Pacchi ^{75a,75b}, D. Ponomarenko ³⁷,
L. Pontecorvo ³⁶, S. Popa ^{27a}, G.A. Popeneciu ^{27d}, D.M. Portillo Quintero ^{156a}, S. Pospisil ¹³²,
P. Postolache ^{27c}, K. Potamianos ¹²⁶, I.N. Potrap ³⁸, C.J. Potter ³², H. Potti ¹, T. Poulsen ⁴⁸,
J. Poveda ¹⁶³, M.E. Pozo Astigarraga ³⁶, A. Prades Ibanez ¹⁶³, M.M. Prapa ⁴⁶, J. Pretel ⁵⁴,
D. Price ¹⁰¹, M. Primavera ^{70a}, M.A. Principe Martin ⁹⁹, R. Privara ¹²², M.L. Proffitt ¹³⁸,
N. Proklova ¹²⁸, K. Prokofiev ^{64c}, G. Proto ^{76a,76b}, S. Protopopescu ²⁹, J. Proudfoot ⁶,
M. Przybycien ^{85a}, J.E. Puddefoot ¹³⁹, D. Pudzha ³⁷, P. Puzo ⁶⁶, D. Pyatizbyantseva ³⁷, J. Qian ¹⁰⁶,
D. Qichen ¹⁰¹, Y. Qin ¹⁰¹, T. Qiu ⁹⁴, A. Quadt ⁵⁵, M. Queitsch-Maitland ¹⁰¹, G. Quetant ⁵⁶,
G. Rabanal Bolanos ⁶¹, D. Rafanoharana ⁵⁴, F. Ragusa ^{71a,71b}, J.L. Rainbolt ³⁹, J.A. Raine ⁵⁶,
S. Rajagopalan ²⁹, E. Ramakoti ³⁷, K. Ran ^{48,14d}, N.P. Rapheeha ^{33g}, V. Raskina ¹²⁷,
D.F. Rassloff ^{63a}, S. Rave ¹⁰⁰, B. Ravina ⁵⁵, I. Ravinovich ¹⁶⁹, M. Raymond ³⁶, A.L. Read ¹²⁵,
N.P. Readioff ¹³⁹, D.M. Rebuffi ^{73a,73b}, G. Redlinger ²⁹, K. Reeves ⁴⁵, J. Reichert ¹²⁸,
J.A. Reidelsturz ¹⁷¹, D. Reikher ¹⁵¹, A. Rej ¹⁴¹, C. Rembser ³⁶, A. Renardi ⁴⁸, M. Renda ^{27b},
M.B. Rendel ¹¹⁰, F. Renner ⁴⁸, A.G. Rennie ⁵⁹, S. Resconi ^{71a}, M. Ressegotti ^{57b,57a},
E.D. Resseguie ^{17a}, S. Rettie ³⁶, J.G. Reyes Rivera ¹⁰⁷, B. Reynolds ¹¹⁹, E. Reynolds ^{17a},
M. Rezaei Estabragh ¹⁷¹, O.L. Rezanova ³⁷, P. Reznicek ¹³³, N. Ribaric ⁹¹, E. Ricci ^{78a,78b},

R. Richter ¹¹⁰, S. Richter ^{47a,47b}, E. Richter-Was ^{85b}, M. Ridel ¹²⁷, S. Ridouani ^{35d}, P. Rieck ¹¹⁷, P. Riedler ³⁶, M. Rijssenbeek ¹⁴⁵, A. Rimoldi ^{73a,73b}, M. Rimoldi ⁴⁸, L. Rinaldi ^{23b,23a}, T.T. Rinn ²⁹, M.P. Rinnagel ¹⁰⁹, G. Ripellino ¹⁴⁴, I. Riu ¹³, P. Rivadeneira ⁴⁸, J.C. Rivera Vergara ¹⁶⁵, F. Rizatdinova ¹²¹, E. Rizvi ⁹⁴, C. Rizzi ⁵⁶, B.A. Roberts ¹⁶⁷, B.R. Roberts ^{17a}, S.H. Robertson ^{104,z}, M. Robin ⁴⁸, D. Robinson ³², C.M. Robles Gajardo ^{137f}, M. Robles Manzano ¹⁰⁰, A. Robson ⁵⁹, A. Rocchi ^{76a,76b}, C. Roda ^{74a,74b}, S. Rodriguez Bosca ^{63a}, Y. Rodriguez Garcia ^{22a}, A. Rodriguez Rodriguez ⁵⁴, A.M. Rodríguez Vera ^{156b}, S. Roe ³⁶, J.T. Roemer ¹⁶⁰, A.R. Roepe-Gier ¹²⁰, J. Roggel ¹⁷¹, O. Røhne ¹²⁵, R.A. Rojas ¹⁰³, B. Roland ⁵⁴, C.P.A. Roland ⁶⁸, J. Roloff ²⁹, A. Romaniouk ³⁷, E. Romano ^{73a,73b}, M. Romano ^{23b}, A.C. Romero Hernandez ¹⁶², N. Rompotis ⁹², L. Roos ¹²⁷, S. Rosati ^{75a}, B.J. Rosser ³⁹, E. Rossi ⁴, E. Rossi ^{72a,72b}, L.P. Rossi ^{57b}, L. Rossini ⁴⁸, R. Rosten ¹¹⁹, M. Rotaru ^{27b}, B. Rottler ⁵⁴, C. Rougier ^{102,ad}, D. Rousseau ⁶⁶, D. Rousso ³², G. Rovelli ^{73a,73b}, A. Roy ¹⁶², A. Rozanov ¹⁰², Y. Rozen ¹⁵⁰, X. Ruan ^{33g}, A. Rubio Jimenez ¹⁶³, A.J. Ruby ⁹², V.H. Ruelas Rivera ¹⁸, T.A. Ruggeri ¹, F. Rühr ⁵⁴, A. Ruiz-Martinez ¹⁶³, A. Rummler ³⁶, Z. Rurikova ⁵⁴, N.A. Rusakovich ³⁸, H.L. Russell ¹⁶⁵, J.P. Rutherford ⁷, K. Rybacki ⁹¹, M. Rybar ¹³³, E.B. Rye ¹²⁵, A. Ryzhov ³⁷, J.A. Sabater Iglesias ⁵⁶, P. Sabatini ¹⁶³, L. Sabetta ^{75a,75b}, H.F.W. Sadrozinski ¹³⁶, F. Safai Tehrani ^{75a}, B. Safarzadeh Samani ¹⁴⁶, M. Safdari ¹⁴³, S. Saha ¹⁰⁴, M. Sahinsoy ¹¹⁰, M. Saimpert ¹³⁵, M. Saito ¹⁵³, T. Saito ¹⁵³, D. Salamani ³⁶, G. Salamanna ^{77a,77b}, A. Salnikov ¹⁴³, J. Salt ¹⁶³, A. Salvador Salas ¹³, D. Salvatore ^{43b,43a}, F. Salvatore ¹⁴⁶, A. Salzburger ³⁶, D. Sammel ⁵⁴, D. Sampsonidis ^{152,f}, D. Sampsonidou ^{62d,62c}, J. Sánchez ¹⁶³, A. Sanchez Pineda ⁴, V. Sanchez Sebastian ¹⁶³, H. Sandaker ¹²⁵, C.O. Sander ⁴⁸, J.A. Sandesara ¹⁰³, M. Sandhoff ¹⁷¹, C. Sandoval ^{22b}, D.P.C. Sankey ¹³⁴, T. Sano ⁸⁷, A. Sansoni ⁵³, L. Santi ^{75a,75b}, C. Santoni ⁴⁰, H. Santos ^{130a,130b}, S.N. Santpur ^{17a}, A. Santra ¹⁶⁹, K.A. Saoucha ¹³⁹, J.G. Saraiva ^{130a,130d}, J. Sardain ⁷, O. Sasaki ⁸³, K. Sato ¹⁵⁷, C. Sauer ^{63b}, F. Sauerburger ⁵⁴, E. Sauvan ⁴, P. Savard ^{155,aj}, R. Sawada ¹⁵³, C. Sawyer ¹³⁴, L. Sawyer ⁹⁷, I. Sayago Galvan ¹⁶³, C. Sbarra ^{23b}, A. Sbrizzi ^{23b,23a}, T. Scanlon ⁹⁶, J. Schaarschmidt ¹³⁸, P. Schacht ¹¹⁰, D. Schaefer ³⁹, U. Schäfer ¹⁰⁰, A.C. Schaffer ^{66,44}, D. Schaile ¹⁰⁹, R.D. Schamberger ¹⁴⁵, E. Schanet ¹⁰⁹, C. Scharf ¹⁸, M.M. Schefer ¹⁹, V.A. Schegelsky ³⁷, D. Scheirich ¹³³, F. Schenck ¹⁸, M. Schernau ¹⁶⁰, C. Scheulen ⁵⁵, C. Schiavi ^{57b,57a}, Z.M. Schillaci ²⁶, E.J. Schioppa ^{70a,70b}, M. Schioppa ^{43b,43a}, B. Schlag ¹⁰⁰, K.E. Schleicher ⁵⁴, S. Schlenker ³⁶, J. Schmeing ¹⁷¹, M.A. Schmidt ¹⁷¹, K. Schmieden ¹⁰⁰, C. Schmitt ¹⁰⁰, S. Schmitt ⁴⁸, L. Schoeffel ¹³⁵, A. Schoening ^{63b}, P.G. Scholer ⁵⁴, E. Schopf ¹²⁶, M. Schott ¹⁰⁰, J. Schovancova ³⁶, S. Schramm ⁵⁶, F. Schroeder ¹⁷¹, H-C. Schultz-Coulon ^{63a}, M. Schumacher ⁵⁴, B.A. Schumm ¹³⁶, Ph. Schune ¹³⁵, H.R. Schwartz ¹³⁶, A. Schwartzman ¹⁴³, T.A. Schwarz ¹⁰⁶, Ph. Schwemling ¹³⁵, R. Schwienhorst ¹⁰⁷, A. Sciandra ¹³⁶, G. Sciolla ²⁶, F. Scuri ^{74a}, F. Scutti ¹⁰⁵, C.D. Sebastiani ⁹², K. Sedlaczek ⁴⁹, P. Seema ¹⁸, S.C. Seidel ¹¹², A. Seiden ¹³⁶, B.D. Seidlitz ⁴¹, C. Seitz ⁴⁸, J.M. Seixas ^{82b}, G. Sekhniaidze ^{72a}, S.J. Sekula ⁴⁴, L. Selem ⁴, N. Semprini-Cesari ^{23b,23a}, S. Sen ⁵¹, D. Sengupta ⁵⁶, V. Senthilkumar ¹⁶³, L. Serin ⁶⁶, L. Serkin ^{69a,69b}, M. Sessa ^{77a,77b}, H. Severini ¹²⁰, F. Sforza ^{57b,57a}, A. Sfyrila ⁵⁶, E. Shabalina ⁵⁵, R. Shaheen ¹⁴⁴, J.D. Shahinian ¹²⁸, D. Shaked Renous ¹⁶⁹, L.Y. Shan ^{14a}, M. Shapiro ^{17a}, A. Sharma ³⁶, A.S. Sharma ¹⁶⁴, P. Sharma ⁸⁰, S. Sharma ⁴⁸, P.B. Shatalov ³⁷, K. Shaw ¹⁴⁶, S.M. Shaw ¹⁰¹, Q. Shen ^{62c,5}, P. Sherwood ⁹⁶, L. Shi ⁹⁶, C.O. Shimmin ¹⁷², Y. Shimogama ¹⁶⁸, J.D. Shinner ⁹⁵, I.P.J. Shipsey ¹²⁶, S. Shirabe ⁶⁰, M. Shiyakova ³⁸,

J. Shlomi ¹⁶⁹, M.J. Shochet ³⁹, J. Shojaii ¹⁰⁵, D.R. Shope ¹²⁵, S. Shrestha ^{119,an}, E.M. Shrif ^{33g}, M.J. Shroff ¹⁶⁵, P. Sicho ¹³¹, A.M. Sickles ¹⁶², E. Sideras Haddad ^{33g}, A. Sidoti ^{23b}, F. Siegert ⁵⁰, Dj. Sijacki ¹⁵, R. Sikora ^{85a}, F. Sili ⁹⁰, J.M. Silva ²⁰, M.V. Silva Oliveira ³⁶, S.B. Silverstein ^{47a}, S. Simion ⁶⁶, R. Simoniello ³⁶, E.L. Simpson ⁵⁹, L.R. Simpson ¹⁰⁶, N.D. Simpson ⁹⁸, S. Simsek ^{21d}, S. Sindhu ⁵⁵, P. Sinervo ¹⁵⁵, S. Singh ¹⁴², S. Singh ¹⁵⁵, S. Sinha ⁴⁸, S. Sinha ^{33g}, M. Sioli ^{23b,23a}, I. Siral ³⁶, S.Yu. Sivoklov ^{37,*}, J. Sjölin ^{47a,47b}, A. Skaf ⁵⁵, E. Skorda ⁹⁸, P. Skubic ¹²⁰, M. Slawinska ⁸⁶, V. Smakhtin ¹⁶⁹, B.H. Smart ¹³⁴, J. Smiesko ³⁶, S.Yu. Smirnov ³⁷, Y. Smirnov ³⁷, L.N. Smirnova ^{37,a}, O. Smirnova ⁹⁸, A.C. Smith ⁴¹, E.A. Smith ³⁹, H.A. Smith ¹²⁶, J.L. Smith ⁹², R. Smith ¹⁴³, M. Smizanska ⁹¹, K. Smolek ¹³², A. Smykiewicz ⁸⁶, A.A. Snesarev ³⁷, H.L. Snoek ¹¹⁴, S. Snyder ²⁹, R. Sobie ^{165,z}, A. Soffer ¹⁵¹, C.A. Solans Sanchez ³⁶, E.Yu. Soldatov ³⁷, U. Soldevila ¹⁶³, A.A. Solodkov ³⁷, S. Solomon ⁵⁴, A. Soloshenko ³⁸, K. Solovieva ⁵⁴, O.V. Solovyanov ⁴⁰, V. Solovyev ³⁷, P. Sommer ³⁶, A. Sonay ¹³, W.Y. Song ^{156b}, J.M. Sonneveld ¹¹⁴, A. Sopczak ¹³², A.L. Sopio ⁹⁶, F. Sopkova ^{28b}, V. Sothilingam ^{63a}, S. Sottocornola ⁶⁸, R. Soualah ^{116b}, Z. Soumami ^{35e}, D. South ⁴⁸, S. Spagnolo ^{70a,70b}, M. Spalla ¹¹⁰, F. Spanò ⁹⁵, D. Sperlich ⁵⁴, G. Spigo ³⁶, M. Spina ¹⁴⁶, S. Spinali ⁹¹, D.P. Spiteri ⁵⁹, M. Spousta ¹³³, E.J. Staats ³⁴, A. Stabile ^{71a,71b}, R. Stamen ^{63a}, M. Stamenkovic ¹¹⁴, A. Stampeki ²⁰, M. Standke ²⁴, E. Stanecka ⁸⁶, M.V. Stange ⁵⁰, B. Stanislaus ^{17a}, M.M. Stanitzki ⁴⁸, M. Stankaityte ¹²⁶, B. Stapf ⁴⁸, E.A. Starchenko ³⁷, G.H. Stark ¹³⁶, J. Stark ^{102,ad}, D.M. Starko ^{156b}, P. Staroba ¹³¹, P. Starovoitov ^{63a}, S. Stärz ¹⁰⁴, R. Staszewski ⁸⁶, G. Stavropoulos ⁴⁶, J. Steentoft ¹⁶¹, P. Steinberg ²⁹, A.L. Steinhebel ¹²³, B. Stelzer ^{142,156a}, H.J. Stelzer ¹²⁹, O. Stelzer-Chilton ^{156a}, H. Stenzel ⁵⁸, T.J. Stevenson ¹⁴⁶, G.A. Stewart ³⁶, M.C. Stockton ³⁶, G. Stoica ^{27b}, M. Stolarski ^{130a}, S. Stonjek ¹¹⁰, A. Straessner ⁵⁰, J. Strandberg ¹⁴⁴, S. Strandberg ^{47a,47b}, M. Strauss ¹²⁰, T. Strebler ¹⁰², P. Strizenec ^{28b}, R. Ströhmer ¹⁶⁶, D.M. Strom ¹²³, L.R. Strom ⁴⁸, R. Stroynowski ⁴⁴, A. Strubig ^{47a,47b}, S.A. Stucci ²⁹, B. Stugu ¹⁶, J. Stupak ¹²⁰, N.A. Styles ⁴⁸, D. Su ¹⁴³, S. Su ^{62a}, W. Su ^{62d,138,62c}, X. Su ^{62a,66}, K. Sugizaki ¹⁵³, V.V. Sulin ³⁷, M.J. Sullivan ⁹², D.M.S. Sultan ^{78a,78b}, L. Sultanaliev ³⁷, S. Sultansoy ^{3b}, T. Sumida ⁸⁷, S. Sun ¹⁰⁶, S. Sun ¹⁷⁰, O. Sunneborn Gudnadottir ¹⁶¹, M.R. Sutton ¹⁴⁶, M. Svatos ¹³¹, M. Swiatlowski ^{156a}, T. Swirski ¹⁶⁶, I. Sykora ^{28a}, M. Sykora ¹³³, T. Sykora ¹³³, D. Ta ¹⁰⁰, K. Tackmann ^{48,x}, A. Taffard ¹⁶⁰, R. Tafirout ^{156a}, J.S. Tafoya Vargas ⁶⁶, R.H.M. Taibah ¹²⁷, R. Takashima ⁸⁸, K. Takeda ⁸⁴, E.P. Takeva ⁵², Y. Takubo ⁸³, M. Talby ¹⁰², A.A. Talyshiev ³⁷, K.C. Tam ^{64b}, N.M. Tamir ¹⁵¹, A. Tanaka ¹⁵³, J. Tanaka ¹⁵³, R. Tanaka ⁶⁶, M. Tanasini ^{57b,57a}, J. Tang ^{62c}, Z. Tao ¹⁶⁴, S. Tapia Araya ^{137f}, S. Tapprogge ¹⁰⁰, A. Tarek Abouelfadl Mohamed ¹⁰⁷, S. Tarem ¹⁵⁰, K. Tariq ^{62b}, G. Tarna ^{102,27b}, G.F. Tartarelli ^{71a}, P. Tas ¹³³, M. Tasevsky ¹³¹, E. Tassi ^{43b,43a}, A.C. Tate ¹⁶², G. Tateno ¹⁵³, Y. Tayalati ^{35e,y}, G.N. Taylor ¹⁰⁵, W. Taylor ^{156b}, H. Teagle ⁹², A.S. Tee ¹⁷⁰, R. Teixeira De Lima ¹⁴³, P. Teixeira-Dias ⁹⁵, J.J. Teoh ¹⁵⁵, K. Terashi ¹⁵³, J. Terron ⁹⁹, S. Terzo ¹³, M. Testa ⁵³, R.J. Teuscher ^{155,z}, A. Thaler ⁷⁹, O. Theiner ⁵⁶, N. Themistokleous ⁵², T. Thevenaux-Pelzer ¹⁸, O. Thielmann ¹⁷¹, D.W. Thomas ⁹⁵, J.P. Thomas ²⁰, E.A. Thompson ⁴⁸, P.D. Thompson ²⁰, E. Thomson ¹²⁸, E.J. Thorpe ⁹⁴, Y. Tian ⁵⁵, V. Tikhomirov ^{37,a}, Yu.A. Tikhonov ³⁷, S. Timoshenko ³⁷, E.X.L. Ting ¹, P. Tipton ¹⁷², S. Tisserant ¹⁰², S.H. Tlou ^{33g}, A. Tnourji ⁴⁰, K. Todome ^{23b,23a}, S. Todorova-Nova ¹³³, S. Todt ⁵⁰, M. Togawa ⁸³, J. Tojo ⁸⁹, S. Tokár ^{28a}, K. Tokushuku ⁸³, O. Toldaiev ⁶⁸, R. Tombs ³², M. Tomoto ^{83,111}, L. Tompkins ^{143,r}, K.W. Topolnicki ^{85b}, P. Tornambe ¹⁰³, E. Torrence ¹²³,

H. Torres ⁵⁰, E. Torró Pastor ¹⁶³, M. Toscani ³⁰, C. Tosciri ³⁹, M. Tost ¹¹, D.R. Tovey ¹³⁹,
A. Traeet ¹⁶, I.S. Trandafir ^{27b}, T. Trefzger ¹⁶⁶, A. Tricoli ²⁹, I.M. Trigger ^{156a},
S. Trincaz-Duvoid ¹²⁷, D.A. Trischuk ²⁶, B. Trocmé ⁶⁰, A. Trofymov ⁶⁶, C. Troncon ^{71a},
L. Truong ^{33c}, M. Trzebinski ⁸⁶, A. Trzupiek ⁸⁶, F. Tsai ¹⁴⁵, M. Tsai ¹⁰⁶, A. Tsiamis ^{152,f},
P.V. Tsiarashka ³⁷, S. Tsigaridas ^{156a}, A. Tsirigotis ^{152,v}, V. Tsiskaridze ¹⁴⁵, E.G. Tskhadadze ^{149a},
M. Tsopoulou ^{152,f}, Y. Tsujikawa ⁸⁷, I.I. Tsukerman ³⁷, V. Tsulaia ^{17a}, S. Tsuno ⁸³, O. Tsur ¹⁵⁰,
D. Tsybychev ¹⁴⁵, Y. Tu ^{64b}, A. Tudorache ^{27b}, V. Tudorache ^{27b}, A.N. Tuna ³⁶, S. Turchikhin ³⁸,
I. Turk Cakir ^{3a}, R. Turra ^{71a}, T. Turtuvshin ^{38,aa}, P.M. Tuts ⁴¹, S. Tzamarias ^{152,f}, P. Tzanis ¹⁰,
E. Tzovara ¹⁰⁰, K. Uchida ¹⁵³, F. Ukegawa ¹⁵⁷, P.A. Ulloa Poblete ^{137c}, E.N. Umaka ²⁹, G. Unal ³⁶,
M. Unal ¹¹, A. Undrus ²⁹, G. Unel ¹⁶⁰, J. Urban ^{28b}, P. Urquijo ¹⁰⁵, G. Usai ⁸, R. Ushioda ¹⁵⁴,
M. Usman ¹⁰⁸, Z. Uysal ^{21b}, L. Vacavant ¹⁰², V. Vacek ¹³², B. Vachon ¹⁰⁴, K.O.H. Vadla ¹²⁵,
T. Vafeiadis ³⁶, A. Vaitkus ⁹⁶, C. Valderanis ¹⁰⁹, E. Valdes Santurio ^{47a,47b}, M. Valente ^{156a},
S. Valentineti ^{23b,23a}, A. Valero ¹⁶³, A. Vallier ^{102,ad}, J.A. Valls Ferrer ¹⁶³, D.R. Van Arneman ¹¹⁴,
T.R. Van Daalen ¹³⁸, P. Van Gemmeren ⁶, M. Van Rijnbach ^{125,36}, S. Van Stroud ⁹⁶,
I. Van Vulpen ¹¹⁴, M. Vanadia ^{76a,76b}, W. Vandelli ³⁶, M. Vandenbroucke ¹³⁵, E.R. Vandewall ¹²¹,
D. Vannicola ¹⁵¹, L. Vannoli ^{57b,57a}, R. Vari ^{75a}, E.W. Varnes ⁷, C. Varni ^{17a}, T. Varol ¹⁴⁸,
D. Varouchas ⁶⁶, L. Varriale ¹⁶³, K.E. Varvell ¹⁴⁷, M.E. Vasile ^{27b}, L. Vaslin ⁴⁰, G.A. Vazquez ¹⁶⁵,
F. Vazeille ⁴⁰, T. Vazquez Schroeder ³⁶, J. Veatch ³¹, V. Vecchio ¹⁰¹, M.J. Veen ¹⁰³,
I. Veliscek ¹²⁶, L.M. Veloce ¹⁵⁵, F. Veloso ^{130a,130c}, S. Veneziano ^{75a}, A. Ventura ^{70a,70b},
A. Verbitskiy ¹¹⁰, M. Verducci ^{74a,74b}, C. Vergis ²⁴, M. Verissimo De Araujo ^{82b}, W. Verkerke ¹¹⁴,
J.C. Vermeulen ¹¹⁴, C. Vernieri ¹⁴³, P.J. Verschuuren ⁹⁵, M. Vessella ¹⁰³, M.C. Vetterli ^{142,aj},
A. Vgenopoulos ^{152,f}, N. Viaux Maira ^{137f}, T. Vickey ¹³⁹, O.E. Vickey Boeriu ¹³⁹,
G.H.A. Viehhauser ¹²⁶, L. Viganì ^{63b}, M. Villa ^{23b,23a}, M. Villaplana Perez ¹⁶³, E.M. Villhauer ⁵²,
E. Vilucchi ⁵³, M.G. Vincker ³⁴, G.S. Virdee ²⁰, A. Vishwakarma ⁵², C. Vittori ^{23b,23a},
I. Vivarelli ¹⁴⁶, V. Vladimirov ¹⁶⁷, E. Voevodina ¹¹⁰, F. Vogel ¹⁰⁹, P. Vokac ¹³², J. Von Ahnen ⁴⁸,
E. Von Toerne ²⁴, B. Vormwald ³⁶, V. Vorobel ¹³³, K. Vorobev ³⁷, M. Vos ¹⁶³, K. Voss ¹⁴¹,
J.H. Vossebeld ⁹², M. Vozak ¹¹⁴, L. Vozdecky ⁹⁴, N. Vranjes ¹⁵, M. Vranjes Milosavljevic ¹⁵,
M. Vreeswijk ¹¹⁴, R. Vuillermet ³⁶, O. Vujanovic ¹⁰⁰, I. Vukotic ³⁹, S. Wada ¹⁵⁷, C. Wagner ¹⁰³,
W. Wagner ¹⁷¹, S. Wahdan ¹⁷¹, H. Wahlberg ⁹⁰, R. Wakasa ¹⁵⁷, M. Wakida ¹¹¹,
V.M. Walbrecht ¹¹⁰, J. Walder ¹³⁴, R. Walker ¹⁰⁹, W. Walkowiak ¹⁴¹, A.M. Wang ⁶¹,
A.Z. Wang ¹⁷⁰, C. Wang ^{62a}, C. Wang ^{62c}, H. Wang ^{17a}, J. Wang ^{64a}, R.-J. Wang ¹⁰⁰,
R. Wang ⁶¹, R. Wang ⁶, S.M. Wang ¹⁴⁸, S. Wang ^{62b}, T. Wang ^{62a}, W.T. Wang ⁸⁰, X. Wang ^{14c},
X. Wang ¹⁶², X. Wang ^{62c}, Y. Wang ^{62d}, Y. Wang ^{14c}, Z. Wang ¹⁰⁶, Z. Wang ^{62d,51,62c},
Z. Wang ¹⁰⁶, A. Warburton ¹⁰⁴, R.J. Ward ²⁰, N. Warrack ⁵⁹, A.T. Watson ²⁰, H. Watson ⁵⁹,
M.F. Watson ²⁰, G. Watts ¹³⁸, B.M. Waugh ⁹⁶, A.F. Webb ¹¹, C. Weber ²⁹, H.A. Weber ¹⁸,
M.S. Weber ¹⁹, S.M. Weber ^{63a}, C. Wei ^{62a}, Y. Wei ¹²⁶, A.R. Weidberg ¹²⁶, J. Weingarten ⁴⁹,
M. Weirich ¹⁰⁰, C. Weiser ⁵⁴, C.J. Wells ⁴⁸, T. Wenaus ²⁹, B. Wendland ⁴⁹, T. Wengler ³⁶,
N.S. Wenke ¹¹⁰, N. Wermes ²⁴, M. Wessels ^{63a}, K. Whalen ¹²³, A.M. Wharton ⁹¹, A.S. White ⁶¹,
A. White ⁸, M.J. White ¹, D. Whiteson ¹⁶⁰, L. Wickremasinghe ¹²⁴, W. Wiedenmann ¹⁷⁰,
C. Wiel ⁵⁰, M. Wielers ¹³⁴, C. Wiglesworth ⁴², L.A.M. Wiik-Fuchs ⁵⁴, D.J. Wilbern ¹²⁰,
H.G. Wilkens ³⁶, D.M. Williams ⁴¹, H.H. Williams ¹²⁸, S. Williams ³², S. Willocq ¹⁰³,
P.J. Windischhofer ¹²⁶, F. Winklmeier ¹²³, B.T. Winter ⁵⁴, J.K. Winter ¹⁰¹, M. Wittgen ¹⁴³,
M. Wobisch ⁹⁷, R. Wölker ¹²⁶, J. Wollrath ¹⁶⁰, M.W. Wolter ⁸⁶, H. Wolters ^{130a,130c},

V.W.S. Wong ¹⁶⁴, A.F. Wongel ⁴⁸, S.D. Worm ⁴⁸, B.K. Wosiek ⁸⁶, K.W. Woźniak ⁸⁶, K. Wraight ⁵⁹, J. Wu ^{14a,14d}, M. Wu ^{64a}, M. Wu ¹¹³, S.L. Wu ¹⁷⁰, X. Wu ⁵⁶, Y. Wu ^{62a}, Z. Wu ^{135,62a}, J. Wuerzinger ¹²⁶, T.R. Wyatt ¹⁰¹, B.M. Wynne ⁵², S. Xella ⁴², L. Xia ^{14c}, M. Xia ^{14b}, J. Xiang ^{64c}, X. Xiao ¹⁰⁶, M. Xie ^{62a}, X. Xie ^{62a}, S. Xin ^{14a,14d}, J. Xiong ^{17a}, I. Xiotidis ¹⁴⁶, D. Xu ^{14a}, H. Xu ^{62a}, H. Xu ^{62a}, L. Xu ^{62a}, R. Xu ¹²⁸, T. Xu ¹⁰⁶, W. Xu ¹⁰⁶, Y. Xu ^{14b}, Z. Xu ^{62b}, Z. Xu ^{14a}, B. Yabsley ¹⁴⁷, S. Yacoob ^{33a}, N. Yamaguchi ⁸⁹, Y. Yamaguchi ¹⁵⁴, H. Yamauchi ¹⁵⁷, T. Yamazaki ^{17a}, Y. Yamazaki ⁸⁴, J. Yan ^{62c}, S. Yan ¹²⁶, Z. Yan ²⁵, H.J. Yang ^{62c,62d}, H.T. Yang ^{62a}, S. Yang ^{62a}, T. Yang ^{64c}, X. Yang ^{62a}, X. Yang ^{14a}, Y. Yang ⁴⁴, Z. Yang ^{62a,106}, W-M. Yao ^{17a}, Y.C. Yap ⁴⁸, H. Ye ^{14c}, H. Ye ⁵⁵, J. Ye ⁴⁴, S. Ye ²⁹, X. Ye ^{62a}, Y. Yeh ⁹⁶, I. Yeletsikh ³⁸, B.K. Yeo ^{17a}, M.R. Yexley ⁹¹, P. Yin ⁴¹, K. Yorita ¹⁶⁸, S. Younas ^{27b}, C.J.S. Young ⁵⁴, C. Young ¹⁴³, Y. Yu ^{62a}, M. Yuan ¹⁰⁶, R. Yuan ^{62b,l}, L. Yue ⁹⁶, X. Yue ^{63a}, M. Zaazoua ^{35e}, B. Zabinski ⁸⁶, E. Zaid ⁵², T. Zakareishvili ^{149b}, N. Zakharchuk ³⁴, S. Zambito ⁵⁶, J.A. Zamora Saa ^{137d,137b}, J. Zang ¹⁵³, D. Zanzi ⁵⁴, O. Zaplatilek ¹³², S.V. Zeiβner ⁴⁹, C. Zeitnitz ¹⁷¹, J.C. Zeng ¹⁶², D.T. Zenger Jr ²⁶, O. Zenin ³⁷, T. Ženiš ^{28a}, S. Zenz ⁹⁴, S. Zerradi ^{35a}, D. Zerwas ⁶⁶, M. Zhai ^{14a,14d}, B. Zhang ^{14c}, D.F. Zhang ¹³⁹, J. Zhang ^{62b}, J. Zhang ⁶, K. Zhang ^{14a,14d}, L. Zhang ^{14c}, P. Zhang ^{14a,14d}, R. Zhang ¹⁷⁰, S. Zhang ¹⁰⁶, T. Zhang ¹⁵³, X. Zhang ^{62c}, X. Zhang ^{62b}, Y. Zhang ^{62c,5}, Z. Zhang ^{17a}, Z. Zhang ⁶⁶, H. Zhao ¹³⁸, P. Zhao ⁵¹, T. Zhao ^{62b}, Y. Zhao ¹³⁶, Z. Zhao ^{62a}, A. Zhemchugov ³⁸, X. Zheng ^{62a}, Z. Zheng ¹⁴³, D. Zhong ¹⁶², B. Zhou ¹⁰⁶, C. Zhou ¹⁷⁰, H. Zhou ⁷, N. Zhou ^{62c}, Y. Zhou ⁷, C.G. Zhu ^{62b}, C. Zhu ^{14a,14d}, H.L. Zhu ^{62a}, H. Zhu ^{14a}, J. Zhu ¹⁰⁶, Y. Zhu ^{62c}, Y. Zhu ^{62a}, X. Zhuang ^{14a}, K. Zhukov ³⁷, V. Zhulanov ³⁷, N.I. Zimine ³⁸, J. Zinsser ^{63b}, M. Ziolkowski ¹⁴¹, L. Živković ¹⁵, A. Zoccoli ^{23b,23a}, K. Zoch ⁵⁶, T.G. Zorbas ¹³⁹, O. Zormpa ⁴⁶, W. Zou ⁴¹, L. Zwalinski ³⁶

¹ Department of Physics, University of Adelaide, Adelaide; Australia

² Department of Physics, University of Alberta, Edmonton AB; Canada

³ (a) Department of Physics, Ankara University, Ankara; (b) Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye

⁴ LAPP, Univ. Savoie Mont Blanc, CNRS/IN2P3, Annecy; France

⁵ APC, Université Paris Cité, CNRS/IN2P3, Paris; France

⁶ High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America

⁷ Department of Physics, University of Arizona, Tucson AZ; United States of America

⁸ Department of Physics, University of Texas at Arlington, Arlington TX; United States of America

⁹ Physics Department, National and Kapodistrian University of Athens, Athens; Greece

¹⁰ Physics Department, National Technical University of Athens, Zografou; Greece

¹¹ Department of Physics, University of Texas at Austin, Austin TX; United States of America

¹² Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan

¹³ Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain

¹⁴ (a) Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (b) Physics Department, Tsinghua University, Beijing; (c) Department of Physics, Nanjing University, Nanjing; (d) University of Chinese Academy of Science (UCAS), Beijing; China

¹⁵ Institute of Physics, University of Belgrade, Belgrade; Serbia

¹⁶ Department for Physics and Technology, University of Bergen, Bergen; Norway

¹⁷ (a) Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; (b) University of California, Berkeley CA; United States of America

¹⁸ Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany

¹⁹ Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland

- ²⁰ School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom
- ²¹ ^(a) Department of Physics, Bogazici University, Istanbul; ^(b) Department of Physics Engineering, Gaziantep University, Gaziantep; ^(c) Department of Physics, Istanbul University, Istanbul; ^(d) Istinye University, Sariyer, Istanbul; Türkiye
- ²² ^(a) Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; ^(b) Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia
- ²³ ^(a) Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; ^(b) INFN Sezione di Bologna; Italy
- ²⁴ Physikalisches Institut, Universität Bonn, Bonn; Germany
- ²⁵ Department of Physics, Boston University, Boston MA; United States of America
- ²⁶ Department of Physics, Brandeis University, Waltham MA; United States of America
- ²⁷ ^(a) Transilvania University of Brasov, Brasov; ^(b) Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; ^(c) Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; ^(d) National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; ^(e) University Politehnica Bucharest, Bucharest; ^(f) West University in Timisoara, Timisoara; ^(g) Faculty of Physics, University of Bucharest, Bucharest; Romania
- ²⁸ ^(a) Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; ^(b) Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic
- ²⁹ Physics Department, Brookhaven National Laboratory, Upton NY; United States of America
- ³⁰ Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina
- ³¹ California State University, CA; United States of America
- ³² Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom
- ³³ ^(a) Department of Physics, University of Cape Town, Cape Town; ^(b) iThemba Labs, Western Cape; ^(c) Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; ^(d) National Institute of Physics, University of the Philippines Diliman (Philippines); ^(e) University of South Africa, Department of Physics, Pretoria; ^(f) University of Zululand, KwaDlangezwa; ^(g) School of Physics, University of the Witwatersrand, Johannesburg; South Africa
- ³⁴ Department of Physics, Carleton University, Ottawa ON; Canada
- ³⁵ ^(a) Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; ^(b) Faculté des Sciences, Université Ibn-Tofail, Kénitra; ^(c) Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; ^(d) LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda; ^(e) Faculté des sciences, Université Mohammed V, Rabat; ^(f) Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco
- ³⁶ CERN, Geneva; Switzerland
- ³⁷ Affiliated with an institute covered by a cooperation agreement with CERN
- ³⁸ Affiliated with an international laboratory covered by a cooperation agreement with CERN
- ³⁹ Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America
- ⁴⁰ LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France
- ⁴¹ Nevis Laboratory, Columbia University, Irvington NY; United States of America
- ⁴² Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark
- ⁴³ ^(a) Dipartimento di Fisica, Università della Calabria, Rende; ^(b) INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy
- ⁴⁴ Physics Department, Southern Methodist University, Dallas TX; United States of America
- ⁴⁵ Physics Department, University of Texas at Dallas, Richardson TX; United States of America
- ⁴⁶ National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece
- ⁴⁷ ^(a) Department of Physics, Stockholm University; ^(b) Oskar Klein Centre, Stockholm; Sweden
- ⁴⁸ Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany
- ⁴⁹ Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany
- ⁵⁰ Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany
- ⁵¹ Department of Physics, Duke University, Durham NC; United States of America
- ⁵² SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom

- ⁵³ INFN e Laboratori Nazionali di Frascati, Frascati; Italy
- ⁵⁴ Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany
- ⁵⁵ II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany
- ⁵⁶ Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland
- ⁵⁷ ^(a) Dipartimento di Fisica, Università di Genova, Genova; ^(b) INFN Sezione di Genova; Italy
- ⁵⁸ II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany
- ⁵⁹ SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom
- ⁶⁰ LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France
- ⁶¹ Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America
- ⁶² ^(a) Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; ^(b) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; ^(c) School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; ^(d) Tsung-Dao Lee Institute, Shanghai; China
- ⁶³ ^(a) Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; ^(b) Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany
- ⁶⁴ ^(a) Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; ^(b) Department of Physics, University of Hong Kong, Hong Kong; ^(c) Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China
- ⁶⁵ Department of Physics, National Tsing Hua University, Hsinchu; Taiwan
- ⁶⁶ IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France
- ⁶⁷ Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona; Spain
- ⁶⁸ Department of Physics, Indiana University, Bloomington IN; United States of America
- ⁶⁹ ^(a) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; ^(b) ICTP, Trieste; ^(c) Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy
- ⁷⁰ ^(a) INFN Sezione di Lecce; ^(b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy
- ⁷¹ ^(a) INFN Sezione di Milano; ^(b) Dipartimento di Fisica, Università di Milano, Milano; Italy
- ⁷² ^(a) INFN Sezione di Napoli; ^(b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy
- ⁷³ ^(a) INFN Sezione di Pavia; ^(b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy
- ⁷⁴ ^(a) INFN Sezione di Pisa; ^(b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy
- ⁷⁵ ^(a) INFN Sezione di Roma; ^(b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy
- ⁷⁶ ^(a) INFN Sezione di Roma Tor Vergata; ^(b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy
- ⁷⁷ ^(a) INFN Sezione di Roma Tre; ^(b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy
- ⁷⁸ ^(a) INFN-TIFPA; ^(b) Università degli Studi di Trento, Trento; Italy
- ⁷⁹ Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria
- ⁸⁰ University of Iowa, Iowa City IA; United States of America
- ⁸¹ Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America
- ⁸² ^(a) Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; ^(b) Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; ^(c) Instituto de Física, Universidade de São Paulo, São Paulo; ^(d) Rio de Janeiro State University, Rio de Janeiro; Brazil
- ⁸³ KEK, High Energy Accelerator Research Organization, Tsukuba; Japan
- ⁸⁴ Graduate School of Science, Kobe University, Kobe; Japan
- ⁸⁵ ^(a) AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow; ^(b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland
- ⁸⁶ Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland
- ⁸⁷ Faculty of Science, Kyoto University, Kyoto; Japan
- ⁸⁸ Kyoto University of Education, Kyoto; Japan
- ⁸⁹ Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka; Japan
- ⁹⁰ Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina
- ⁹¹ Physics Department, Lancaster University, Lancaster; United Kingdom
- ⁹² Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom
- ⁹³ Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia

- ⁹⁴ School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom
- ⁹⁵ Department of Physics, Royal Holloway University of London, Egham; United Kingdom
- ⁹⁶ Department of Physics and Astronomy, University College London, London; United Kingdom
- ⁹⁷ Louisiana Tech University, Ruston LA; United States of America
- ⁹⁸ Fysiska institutionen, Lunds universitet, Lund; Sweden
- ⁹⁹ Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain
- ¹⁰⁰ Institut für Physik, Universität Mainz, Mainz; Germany
- ¹⁰¹ School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom
- ¹⁰² CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France
- ¹⁰³ Department of Physics, University of Massachusetts, Amherst MA; United States of America
- ¹⁰⁴ Department of Physics, McGill University, Montreal QC; Canada
- ¹⁰⁵ School of Physics, University of Melbourne, Victoria; Australia
- ¹⁰⁶ Department of Physics, University of Michigan, Ann Arbor MI; United States of America
- ¹⁰⁷ Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America
- ¹⁰⁸ Group of Particle Physics, University of Montreal, Montreal QC; Canada
- ¹⁰⁹ Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany
- ¹¹⁰ Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany
- ¹¹¹ Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan
- ¹¹² Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America
- ¹¹³ Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands
- ¹¹⁴ Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands
- ¹¹⁵ Department of Physics, Northern Illinois University, DeKalb IL; United States of America
- ¹¹⁶ ^(a) New York University Abu Dhabi, Abu Dhabi; ^(b) University of Sharjah, Sharjah; United Arab Emirates
- ¹¹⁷ Department of Physics, New York University, New York NY; United States of America
- ¹¹⁸ Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan
- ¹¹⁹ Ohio State University, Columbus OH; United States of America
- ¹²⁰ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America
- ¹²¹ Department of Physics, Oklahoma State University, Stillwater OK; United States of America
- ¹²² Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic
- ¹²³ Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America
- ¹²⁴ Graduate School of Science, Osaka University, Osaka; Japan
- ¹²⁵ Department of Physics, University of Oslo, Oslo; Norway
- ¹²⁶ Department of Physics, Oxford University, Oxford; United Kingdom
- ¹²⁷ LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France
- ¹²⁸ Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America
- ¹²⁹ Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America
- ¹³⁰ ^(a) Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa; ^(b) Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; ^(c) Departamento de Física, Universidade de Coimbra, Coimbra; ^(d) Centro de Física Nuclear da Universidade de Lisboa, Lisboa; ^(e) Departamento de Física, Universidade do Minho, Braga; ^(f) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); ^(g) Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal
- ¹³¹ Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic
- ¹³² Czech Technical University in Prague, Prague; Czech Republic
- ¹³³ Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic
- ¹³⁴ Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom
- ¹³⁵ IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France
- ¹³⁶ Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America
- ¹³⁷ ^(a) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; ^(b) Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago; ^(c) Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena; ^(d) Universidad Andres Bello,

- Department of Physics, Santiago;^(e) Instituto de Alta Investigación, Universidad de Tarapacá, Arica;^(f) Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile
- ¹³⁸ Department of Physics, University of Washington, Seattle WA; United States of America
- ¹³⁹ Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom
- ¹⁴⁰ Department of Physics, Shinshu University, Nagano; Japan
- ¹⁴¹ Department Physik, Universität Siegen, Siegen; Germany
- ¹⁴² Department of Physics, Simon Fraser University, Burnaby BC; Canada
- ¹⁴³ SLAC National Accelerator Laboratory, Stanford CA; United States of America
- ¹⁴⁴ Department of Physics, Royal Institute of Technology, Stockholm; Sweden
- ¹⁴⁵ Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America
- ¹⁴⁶ Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom
- ¹⁴⁷ School of Physics, University of Sydney, Sydney; Australia
- ¹⁴⁸ Institute of Physics, Academia Sinica, Taipei; Taiwan
- ¹⁴⁹ ^(a) E. Andronikashvili Institute of Physics, Iv. Javakishvili Tbilisi State University, Tbilisi; ^(b) High Energy Physics Institute, Tbilisi State University, Tbilisi; ^(c) University of Georgia, Tbilisi; Georgia
- ¹⁵⁰ Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel
- ¹⁵¹ Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel
- ¹⁵² Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece
- ¹⁵³ International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan
- ¹⁵⁴ Department of Physics, Tokyo Institute of Technology, Tokyo; Japan
- ¹⁵⁵ Department of Physics, University of Toronto, Toronto ON; Canada
- ¹⁵⁶ ^(a) TRIUMF, Vancouver BC; ^(b) Department of Physics and Astronomy, York University, Toronto ON; Canada
- ¹⁵⁷ Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan
- ¹⁵⁸ Department of Physics and Astronomy, Tufts University, Medford MA; United States of America
- ¹⁵⁹ United Arab Emirates University, Al Ain; United Arab Emirates
- ¹⁶⁰ Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America
- ¹⁶¹ Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden
- ¹⁶² Department of Physics, University of Illinois, Urbana IL; United States of America
- ¹⁶³ Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain
- ¹⁶⁴ Department of Physics, University of British Columbia, Vancouver BC; Canada
- ¹⁶⁵ Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada
- ¹⁶⁶ Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany
- ¹⁶⁷ Department of Physics, University of Warwick, Coventry; United Kingdom
- ¹⁶⁸ Waseda University, Tokyo; Japan
- ¹⁶⁹ Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel
- ¹⁷⁰ Department of Physics, University of Wisconsin, Madison WI; United States of America
- ¹⁷¹ Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany
- ¹⁷² Department of Physics, Yale University, New Haven CT; United States of America

^a Also Affiliated with an institute covered by a cooperation agreement with CERN

^b Also at An-Najah National University, Nablus; Palestine

^c Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America

^d Also at Bruno Kessler Foundation, Trento; Italy

^e Also at Center for High Energy Physics, Peking University; China

^f Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki; Greece

^g Also at Centro Studi e Ricerche Enrico Fermi; Italy

^h Also at CERN, Geneva; Switzerland

ⁱ Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland

^j Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain

- ^k Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece
- ^l Also at Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America
- ^m Also at Department of Physics and Astronomy, University of Louisville, Louisville, KY; United States of America
- ⁿ Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel
- ^o Also at Department of Physics, California State University, East Bay; United States of America
- ^p Also at Department of Physics, California State University, Sacramento; United States of America
- ^q Also at Department of Physics, King's College London, London; United Kingdom
- ^r Also at Department of Physics, Stanford University, Stanford CA; United States of America
- ^s Also at Department of Physics, University of Fribourg, Fribourg; Switzerland
- ^t Also at Department of Physics, University of Thessaly; Greece
- ^u Also at Department of Physics, Westmont College, Santa Barbara; United States of America
- ^v Also at Hellenic Open University, Patras; Greece
- ^w Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain
- ^x Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany
- ^y Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco
- ^z Also at Institute of Particle Physics (IPP); Canada
- ^{aa} Also at Institute of Physics and Technology, Ulaanbaatar; Mongolia
- ^{ab} Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan
- ^{ac} Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia
- ^{ad} Also at L2IT, Université de Toulouse, CNRS/IN2P3, UPS, Toulouse; France
- ^{ae} Also at Lawrence Livermore National Laboratory, Livermore; United States of America
- ^{af} Also at National Institute of Physics, University of the Philippines Diliman (Philippines); Philippines
- ^{ag} Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen; Germany
- ^{ah} Also at Technical University of Munich, Munich; Germany
- ^{ai} Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China
- ^{aj} Also at TRIUMF, Vancouver BC; Canada
- ^{ak} Also at Università di Napoli Parthenope, Napoli; Italy
- ^{al} Also at University of Chinese Academy of Sciences (UCAS), Beijing; China
- ^{am} Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America
- ^{an} Also at Washington College, Maryland; United States of America
- * Deceased