

# 6-Bromopyridine-2-carbaldehyde phenylhydrazone

Rodolfo Moreno-Fuquen,<sup>a\*</sup> Manuel N. Chaur,<sup>a</sup> Elkin L. Romero,<sup>a</sup> Fabio Zuluaga<sup>a</sup> and Javier Ellena<sup>b</sup>

<sup>a</sup>Departamento de Química, Facultad de Ciencias, Universidad del Valle, Apartado 25360, Santiago de Cali, Colombia, and <sup>b</sup>Instituto de Física de São Carlos, IFSC, Universidade de São Paulo, USP, São Carlos, SP, Brazil

Correspondence e-mail: rodimo26@yahoo.es


Received 6 June 2012; accepted 12 June 2012

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.039;  $wR$  factor = 0.110; data-to-parameter ratio = 16.1.

The title compound,  $\text{C}_{12}\text{H}_{10}\text{BrN}_3$ , is essentially planar (r.m.s. deviation of all non-H atoms = 0.0174 Å), with a dihedral angle of 0.5 (2)° between the two aromatic rings. In the crystal, molecules are linked by weak  $\text{N}-\text{H}\cdots\text{N}$  interactions, forming a zigzag chain running parallel to [001].

## Related literature

For bond-length data, see: Allen *et al.* (1987). For related structures, see: Yu *et al.* (2005); Fun *et al.* (2012). For the design of molecular dynamic systems, see: Hirose (2010). For the principles of synthetic molecular structures with dynamic properties, see: Kay *et al.* (2007). For configurational changes by UV light and heat, see: Chaur *et al.* (2011); Lehn (2006); Dugave & Demange (2003). For graph-set notation, see: Etter (1990).



## Experimental

### Crystal data

$\text{C}_{12}\text{H}_{10}\text{BrN}_3$   
 $M_r = 276.13$   
 Orthorhombic,  $Pbca$   
 $a = 14.6418$  (3) Å  
 $b = 7.8407$  (1) Å  
 $c = 20.0645$  (4) Å  
 $V = 2303.44$  (7) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 3.54$  mm<sup>-1</sup>  
 $T = 295$  K  
 $0.33 \times 0.30 \times 0.23$  mm

### Data collection

Nonius KappaCCD diffractometer  
 Absorption correction: multi-scan  
 (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.382$ ,  $T_{\max} = 0.544$   
 28166 measured reflections  
 2339 independent reflections  
 1903 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.048$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.110$   
 $S = 1.03$   
 2339 reflections  
 145 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.44$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.72$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$                          | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|----------------------------------------|-------|-------------|-------------|---------------|
| $\text{N1}-\text{H1}\cdots\text{N3}^i$ | 0.86  | 2.34        | 3.180 (3)   | 166           |

Symmetry code: (i)  $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

RMF is grateful to the Spanish Research Council (CSIC) for the use of a free-of-charge licence to the Cambridge Structural Database. RMF, MNC and FZ also thank the Universidad del Valle, Colombia, for partial financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GG2083).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.  
 Chaur, M. N., Collado, D. & Lehn, J. M. (2011). *Chem. Eur. J.* **17**, 248–258.  
 Dugave, C. & Demange, L. (2003). *Chem. Rev.* **103**, 2475–2532.  
 Etter, M. (1990). *Acc. Chem. Res.* **23**, 120–126.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
 Fun, H.-K., Chantrapromma, S., Nilwanna, B. & Karalai, C. (2012). *Acta Cryst. E* **68**, o704–o705.  
 Hirose, J. (2010). *J. Incl. Phenom. Macrocycl. Chem.* **68**, 1–24.  
 Kay, E. R., Leigh, D. A. & Zerbetto, F. (2007). *Angew. Chem. Int. Ed.* **46**, 72–191.  
 Lehn, J. M. (2006). *Chem. Eur. J.* **12**, 5910–5915.  
 Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.  
 Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.  
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
 Yu, M., Fan, Z., Jing, Z.-L., Chen, X., Diao, C.-H. & Deng, Q.-L. (2005). *Acta Cryst. E* **61**, o3342–o3343.

## supplementary materials

*Acta Cryst.* (2012). E68, o2131 [doi:10.1107/S1600536812026517]

**6-Bromopyridine-2-carbaldehyde phenylhydrazone**

**Rodolfo Moreno-Fuquen, Manuel N. Chaur, Elkin L. Romero, Fabio Zuluaga and Javier Ellena**

**Comment**

The design of molecular dynamic systems that operate by external stimuli without producing chemical waste is one of the questions of great interest in the nanotechnology field (Hirose, 2010), since these molecular systems are molecules that can exhibit structural (configurational) and chemical (constitutional) changes by the modification of external factors (presence of other molecules or metal ions, heat and/or light). Among the compounds that have attracted interest in this regard are the hydrazones which contain the imine group ( $-C=N-$ ) (Kay *et al.*, 2007). This group can undergo reversible configurational changes induced by UV light and heat (Chaur *et al.*, 2011). For hydrazones prepared from pyridine-carboxaldehydes and pyridine or phenyl hydrazines the E/Z photoisomerization is favored by an intermolecular hydrogen bond between the hydrazine and the nitrogen of the pyridinecarboxaldehyde group resulting in a metastable state which can be returned to its original state by heating in solution (Chaur *et al.*, 2011; Lehn, 2006; Dugave & Demange, 2003). Besides, these compounds can exhibit dynamic properties since they can undergo coordination with suitable metals and the reversibility of the imine group gives them other features that can be used in the development of molecular machines or in the storage of information (Chaur *et al.*, 2011). The compound reported in this paper is part of a series of compounds that are currently being prepared in our group and that exhibit dynamic properties such as photoisomerization, constitutional changes and metal coordination in order to build supramolecular systems of multiple dynamics. Herein we report the synthesis and crystal structure of 6-bromo-pyridinecarbaldehyde phenylhydrazone (I), Fig 1. In the molecular structure of (I), the bromopyridyl ring is planar (r.m.s. deviation of all non-hydrogen atoms = 0.0020 Å) like the central bridge (C5/C7/N2/N1/C8) (r.m.s. deviation of all non-hydrogen atoms = 0.005 Å) with a dihedral angle of 0.6 (2)° between these two planes. The central bridge forms a dihedral angle of 0.8 (2)° with the benzene ring and the bromopyridyl and benzene rings form a dihedral angle of 0.5 (2)°. The bond lengths agree with the literature values (Allen *et al.*, 1987) and are comparable with the related structures (Yu *et al.*, 2005; Fun *et al.*, 2012). In the crystal packing (Fig. 2), the molecules are linked by weak  $N-H\cdots N$  interactions (Table 1). Indeed, in this substructure, atom N1 in the molecule at (x,y,z) links to N3 atom in the molecule at (-x+2,+y-1/2,-z+1/2). The propagation of this interaction forms C(6) (Etter, 1990), continuous one-dimensional zigzag chain running parallel to [001].

**Experimental**

The hydrazone under study was prepared by the condensation reaction of 6-bromo-2-pyridinecarboxaldehyde with phenylhydrazine in ethanol according to Fig. 3, obtaining a yellow solid with a yield of 81%.  $^1H$  NMR (400 MHz, DMSO- $d_6$ )  $\delta$ ppm: 10.83 (s, 1H, NH), 7.92 (d,  $J$  = 8.03 Hz, 1H), 7.78 (s, 1H), 7.71 (t,  $J$  = 7.78 Hz, 1H), 7.47 (d,  $J$  = 7.78 Hz, 1H), 7.26 (m, 2H), 7.13 (d,  $J$  = 8.28 Hz, 2H), 6.83 (t,  $J$  = 7.28 Hz, 1H). NMR  $^{13}C$  (100 MHz, DMSO- $d_6$ )  $\delta$ ppm: 156.16, 144.31, 140.77, 139.60, 134.52, 129.27, 126.07, 119.99, 117.80, 112.52. IR (KBr)  $N(cm^{-1})$ : 3228 (N-H), 3038 and 2971 ( $=C-H$ ), 1558 y 1601 ( $C=C$  y  $C=N$ ). Melting point: 462 (1) K. Elemental analysis: Calculated: C 52.20 %, H 3.65 %, N


15.22 %; found: C 51.98 %, H 3.46 %, N 15.07 %.

### Refinement

All H-atoms were positioned geometrically using riding model with [C—H = 0.93 Å with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ ].


### Computing details

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999).




**Figure 1**

Molecular conformation and atom numbering scheme for (I) with displacement ellipsoids drawn at the 50% probability level. H atoms are shown as spheres of arbitrary radius.



**Figure 2**

Part of the crystal structure of (I), showing the formation of one-dimensional zigzag chain running parallel to (001). Symmetry code: (i)  $-x+2, +y-1/2, -z+1/2$ .



**Figure 3**

Synthesis of the 6-bromo-pyridinecarbaldehyde phenylhydrazone

### 6-Bromopyridine-2-carbaldehyde phenylhydrazone

#### Crystal data

$C_{12}H_{10}BrN_3$

$M_r = 276.13$

Orthorhombic, *Pbca*

Hall symbol:  $-P\ 2ac\ 2ab$

$a = 14.6418\ (3)\ \text{\AA}$

$b = 7.8407\ (1)\ \text{\AA}$

$c = 20.0645\ (4)\ \text{\AA}$

$V = 2303.44\ (7)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1104$

$D_x = 1.593\ \text{Mg m}^{-3}$

Melting point:  $497\ (1)\ \text{K}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 9233 reflections

$\theta = 3.6\text{--}26.4^\circ$

$\mu = 3.54\ \text{mm}^{-1}$

$T = 295\ \text{K}$

Block, black

$0.33 \times 0.30 \times 0.23\ \text{mm}$

#### Data collection

Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

CCD rotation images, thick slices scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.382$ ,  $T_{\max} = 0.544$

28166 measured reflections

2339 independent reflections

1903 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.048$ 
 $\theta_{\text{max}} = 26.4^\circ$ ,  $\theta_{\text{min}} = 3.6^\circ$ 
 $h = -18 \rightarrow 17$ 
 $k = -9 \rightarrow 9$ 
 $l = -24 \rightarrow 25$ 

### Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.039$ 
 $wR(F^2) = 0.110$ 
 $S = 1.03$ 

2339 reflections

145 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.052P)^2 + 1.6466P]$ 

where  $P = (F_o^2 + 2F_c^2)/3$ 
 $(\Delta/\sigma)_{\text{max}} < 0.001$ 
 $\Delta\rho_{\text{max}} = 0.44 \text{ e } \text{\AA}^{-3}$ 
 $\Delta\rho_{\text{min}} = -0.72 \text{ e } \text{\AA}^{-3}$ 

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

|     | $x$          | $y$         | $z$           | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|-------------|---------------|----------------------------------|
| Br1 | 0.82948 (2)  | 1.00944 (4) | 0.314727 (18) | 0.07300 (18)                     |
| N1  | 1.06677 (16) | 0.1846 (3)  | 0.32501 (10)  | 0.0535 (5)                       |
| H1  | 1.0780       | 0.2044      | 0.2837        | 0.064*                           |
| N2  | 1.01869 (15) | 0.2984 (3)  | 0.36079 (10)  | 0.0506 (5)                       |
| N3  | 0.91170 (14) | 0.6977 (3)  | 0.33261 (10)  | 0.0468 (5)                       |
| C1  | 0.86323 (18) | 0.8132 (3)  | 0.36454 (13)  | 0.0496 (6)                       |
| C2  | 0.8363 (2)   | 0.8049 (4)  | 0.43023 (14)  | 0.0620 (7)                       |
| H2  | 0.8017       | 0.8907      | 0.4499        | 0.074*                           |
| C3  | 0.8635 (2)   | 0.6620 (4)  | 0.46526 (14)  | 0.0674 (8)                       |
| H3  | 0.8471       | 0.6495      | 0.5098        | 0.081*                           |
| C4  | 0.9144 (2)   | 0.5390 (4)  | 0.43455 (13)  | 0.0577 (7)                       |
| H4  | 0.9332       | 0.4428      | 0.4579        | 0.069*                           |
| C5  | 0.93799 (17) | 0.5597 (3)  | 0.36750 (12)  | 0.0467 (5)                       |
| C7  | 0.99160 (18) | 0.4341 (4)  | 0.33110 (12)  | 0.0506 (6)                       |
| H7  | 1.0061       | 0.4522      | 0.2865        | 0.061*                           |
| C8  | 1.09823 (17) | 0.0360 (3)  | 0.35418 (12)  | 0.0460 (5)                       |
| C9  | 1.0819 (2)   | -0.0034 (3) | 0.42067 (14)  | 0.0568 (7)                       |
| H9  | 1.0468       | 0.0692      | 0.4469        | 0.068*                           |
| C10 | 1.1179 (2)   | -0.1504 (4) | 0.44746 (15)  | 0.0696 (8)                       |
| H10 | 1.1073       | -0.1750     | 0.4921        | 0.084*                           |
| C11 | 1.1688 (2)   | -0.2614 (4) | 0.41011 (18)  | 0.0704 (8)                       |
| H11 | 1.1925       | -0.3603     | 0.4289        | 0.084*                           |
| C12 | 1.1841 (2)   | -0.2232 (4) | 0.34413 (17)  | 0.0670 (8)                       |

|     |              |             |              |            |
|-----|--------------|-------------|--------------|------------|
| H12 | 1.2184       | −0.2976     | 0.3181       | 0.080*     |
| C13 | 1.14946 (19) | −0.0769 (4) | 0.31608 (13) | 0.0553 (7) |
| H13 | 1.1604       | −0.0533     | 0.2714       | 0.066*     |

*Atomic displacement parameters (Å<sup>2</sup>)*

|     | $U^{11}$    | $U^{22}$    | $U^{33}$    | $U^{12}$     | $U^{13}$      | $U^{23}$     |
|-----|-------------|-------------|-------------|--------------|---------------|--------------|
| Br1 | 0.0783 (3)  | 0.0622 (2)  | 0.0785 (3)  | 0.01901 (15) | −0.00590 (15) | 0.00948 (14) |
| N1  | 0.0623 (14) | 0.0528 (13) | 0.0454 (11) | 0.0095 (11)  | 0.0048 (9)    | −0.0002 (10) |
| N2  | 0.0556 (12) | 0.0466 (12) | 0.0495 (11) | 0.0035 (10)  | −0.0006 (10)  | −0.0039 (10) |
| N3  | 0.0477 (11) | 0.0483 (12) | 0.0444 (10) | −0.0002 (9)  | −0.0039 (9)   | −0.0015 (9)  |
| C1  | 0.0487 (13) | 0.0479 (14) | 0.0521 (14) | 0.0009 (11)  | −0.0069 (11)  | −0.0008 (11) |
| C2  | 0.0704 (19) | 0.0600 (17) | 0.0557 (16) | 0.0063 (14)  | 0.0061 (13)   | −0.0107 (13) |
| C3  | 0.088 (2)   | 0.0676 (19) | 0.0460 (14) | 0.0066 (17)  | 0.0106 (14)   | −0.0021 (14) |
| C4  | 0.0731 (18) | 0.0524 (14) | 0.0475 (14) | 0.0013 (14)  | 0.0007 (13)   | 0.0024 (12)  |
| C5  | 0.0491 (13) | 0.0472 (13) | 0.0440 (13) | −0.0023 (11) | −0.0036 (10)  | −0.0007 (11) |
| C7  | 0.0548 (15) | 0.0530 (14) | 0.0439 (12) | 0.0004 (12)  | −0.0016 (11)  | −0.0019 (12) |
| C8  | 0.0439 (13) | 0.0461 (13) | 0.0479 (13) | −0.0012 (10) | −0.0028 (10)  | −0.0045 (11) |
| C9  | 0.0661 (17) | 0.0505 (16) | 0.0539 (15) | 0.0015 (12)  | 0.0063 (13)   | −0.0007 (11) |
| C10 | 0.091 (2)   | 0.0568 (17) | 0.0607 (17) | 0.0001 (16)  | 0.0024 (16)   | 0.0107 (14)  |
| C11 | 0.077 (2)   | 0.0480 (16) | 0.086 (2)   | 0.0088 (14)  | −0.0138 (16)  | 0.0051 (15)  |
| C12 | 0.0624 (17) | 0.0590 (18) | 0.080 (2)   | 0.0144 (14)  | −0.0061 (15)  | −0.0181 (16) |
| C13 | 0.0539 (15) | 0.0590 (17) | 0.0530 (15) | 0.0080 (13)  | −0.0020 (12)  | −0.0096 (12) |

*Geometric parameters (Å, °)*

|           |             |            |           |
|-----------|-------------|------------|-----------|
| Br1—C1    | 1.900 (3)   | C5—C7      | 1.455 (4) |
| N1—C8     | 1.383 (3)   | C7—H7      | 0.9300    |
| N1—H1     | 0.8600      | C8—C13     | 1.389 (4) |
| N2—C7     | 1.282 (3)   | C8—C9      | 1.390 (4) |
| N2—N1     | 1.344 (3)   | C9—C10     | 1.376 (4) |
| N3—C1     | 1.317 (3)   | C9—H9      | 0.9300    |
| N3—C5     | 1.345 (3)   | C10—C11    | 1.369 (4) |
| C1—C2     | 1.377 (4)   | C10—H10    | 0.9300    |
| C2—C3     | 1.381 (4)   | C11—C12    | 1.375 (5) |
| C2—H2     | 0.9300      | C11—H11    | 0.9300    |
| C3—C4     | 1.366 (4)   | C12—C13    | 1.375 (4) |
| C3—H3     | 0.9300      | C12—H12    | 0.9300    |
| C4—C5     | 1.398 (4)   | C13—H13    | 0.9300    |
| C4—H4     | 0.9300      |            |           |
| C7—N2—N1  | 117.7 (2)   | N2—C7—H7   | 120.2     |
| N2—N1—C8  | 120.5 (2)   | C5—C7—H7   | 120.2     |
| N2—N1—H1  | 119.7       | N1—C8—C13  | 118.9 (2) |
| C8—N1—H1  | 119.7       | N1—C8—C9   | 122.4 (2) |
| C1—N3—C5  | 117.0 (2)   | C13—C8—C9  | 118.7 (3) |
| N3—C1—C2  | 125.9 (3)   | C10—C9—C8  | 119.7 (3) |
| N3—C1—Br1 | 116.18 (19) | C10—C9—H9  | 120.2     |
| C2—C1—Br1 | 117.9 (2)   | C8—C9—H9   | 120.2     |
| C1—C2—C3  | 116.3 (3)   | C11—C10—C9 | 121.8 (3) |

|              |             |                 |            |
|--------------|-------------|-----------------|------------|
| C1—C2—H2     | 121.8       | C11—C10—H10     | 119.1      |
| C3—C2—H2     | 121.8       | C9—C10—H10      | 119.1      |
| C4—C3—C2     | 120.0 (3)   | C10—C11—C12     | 118.5 (3)  |
| C4—C3—H3     | 120.0       | C10—C11—H11     | 120.7      |
| C2—C3—H3     | 120.0       | C12—C11—H11     | 120.7      |
| C3—C4—C5     | 119.1 (3)   | C13—C12—C11     | 121.0 (3)  |
| C3—C4—H4     | 120.4       | C13—C12—H12     | 119.5      |
| C5—C4—H4     | 120.4       | C11—C12—H12     | 119.5      |
| N3—C5—C4     | 121.5 (2)   | C12—C13—C8      | 120.4 (3)  |
| N3—C5—C7     | 115.9 (2)   | C12—C13—H13     | 119.8      |
| C4—C5—C7     | 122.5 (2)   | C8—C13—H13      | 119.8      |
| N2—C7—C5     | 119.7 (2)   |                 |            |
| C7—N2—N1—C8  | 179.8 (2)   | N3—C5—C7—N2     | −179.5 (2) |
| C5—N3—C1—C2  | −0.6 (4)    | C4—C5—C7—N2     | 0.2 (4)    |
| C5—N3—C1—Br1 | 178.54 (17) | N2—N1—C8—C13    | −178.5 (2) |
| N3—C1—C2—C3  | 0.3 (4)     | N2—N1—C8—C9     | 0.3 (4)    |
| Br1—C1—C2—C3 | −178.8 (2)  | N1—C8—C9—C10    | −177.6 (3) |
| C1—C2—C3—C4  | 0.2 (5)     | C13—C8—C9—C10   | 1.2 (4)    |
| C2—C3—C4—C5  | −0.3 (5)    | C8—C9—C10—C11   | −0.9 (5)   |
| C1—N3—C5—C4  | 0.4 (4)     | C9—C10—C11—C12  | 0.2 (5)    |
| C1—N3—C5—C7  | −179.9 (2)  | C10—C11—C12—C13 | 0.2 (5)    |
| C3—C4—C5—N3  | 0.0 (4)     | C11—C12—C13—C8  | 0.1 (5)    |
| C3—C4—C5—C7  | −179.6 (3)  | N1—C8—C13—C12   | 178.0 (3)  |
| N1—N2—C7—C5  | 179.0 (2)   | C9—C8—C13—C12   | −0.8 (4)   |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H $\cdots$ <i>A</i>  | <i>D</i> —H | H $\cdots$ <i>A</i> | <i>D</i> $\cdots$ <i>A</i> | <i>D</i> —H $\cdots$ <i>A</i> |
|--------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| N1—H1 $\cdots$ N3 <sup>i</sup> | 0.86        | 2.34                | 3.180 (3)                  | 166                           |

Symmetry code: (i)  $-x+2, y-1/2, -z+1/2$ .