



HADRONs

2025 Porto Alegre

• • • XVI International Workshop on Hadron Physics

10–14 Mar | **Centro Cultural da UFRGS**

Invited speakers

Kenji Fukushima
The University of Tokyo

Chun Shen
Wayne State University

Lisheng Geng
Beihang University

Fernanda Steffens
University of Bonn

National Committee

Kanchan Khemchandani (UNIFESP)
Jun Takahashi (UNICAMP)
Leticia Palhares (UERJ)
Luciano Abreu (UFBA)
Ricardo Sonogo Farias (UFSM)
Victor Gonçalves (UFPeI)
Tiago Nunes (UFSC)

Organizing Committee

Victor Gonçalves (UFPeI) - Chair
Gustavo Gil da Silveira (UFRGS)
Magno Machado (UFRGS)
Ricardo Sonogo Farias (UFSM)

Registrations open from
10 Aug 2024 to 14 Feb 2025

<https://indico.cern.ch/e/hadrons2025>





Começar (Iniciar) 10 de mar. de 2025 09:00
Fim 14 de mar. de 2025 19:00
America/Sao_Paulo



Centro Cultural - UFRGS
Jacarandá Auditorium
Rua Eng. Luiz Englert, 333 - Porto Alegre-RS
[Ir para o mapa](#)



[Victor Goncalves](#)



Distribuído por [Indico](#) v3.3.7-pre

[Ajuda](#) | [Contato](#) | [Termos e condições](#) | [Encurtador de URL](#) | [Privacidade](#)





ID da Contribuição: 97

Tipos: **Poster**

Photon Propagator in Confinement and Deconfinement Phases of Lattice QED

segunda-feira, 10 de março de 2025 17:52 (2 minutos)

Quantum electrodynamics (QED) is the gauge field theory that describes the interactions between photons and charged particles, characterized by gauge invariance under the group $U(1)$ [1]. Understanding how QED works is essential to describe one of the fundamental forces of the universe, eletromagnetism, this can be investigated from various perspectives, including experimental, perturbative [2], and lattice approaches [3]. In this study, we adopt the lattice formulation to simulate and analyze QED, employing Monte Carlo techniques.

The photon propagator, a key object in QED, provides the probability amplitude for a photon to travel from one point in spacetime to another. It also describes how photons transmit their effects through spacetime. Understanding the behavior of the photon propagator is crucial for exploring QED phenomena.

In lattice QED, a unique phenomenon emerges that is not observed in nature: a confinement phase. This divides the lattice QED into two distinct regions—the confinement and deconfinement phases [4]. This phase structure can be attributed to lattice properties and understanding how this phase structure arises and how to deal with it is essential to improve our data and extract physical meaning.

In this study, we compute the photon propagator in both phases on a lattice of size 204. We utilize the Los Alamos algorithm for gauge fixing and the Heat Bath Monte Carlo algorithm for thermalization and configuration generation.

[1] S. Pokorski, Gauge field theories, vol. 136 (Cambridge University Press, 2000).

[2] M. E. Peskin, An introduction to quantum field theory (CRC press, 2018).

[3] J. Smit, Introduction to quantum fields on a lattice (Cambridge University Press, 2003).

[4] L. C. Loveridge, O. Oliveira, and P. J. Silva, Physical Review D 103, 094519 (2021)

Autor: NASCIMENTO MANDELO, Pablo (Universidade de São Paulo - USP)

Co-autor: MENDES, Tereza

Apresentador: NASCIMENTO MANDELO, Pablo (Universidade de São Paulo - USP)

Classificação da Sessão: Poster session