



WE NEED TO TALK ABOUT THE TOPOLOGY OF INTERACTION NETWORKS

Marco Aurelio Ribeiro Mello^{1*}  & Carsten Friedrich Dormann² 

¹ Universidade de São Paulo, Institute of Biosciences, Department of Ecology. Rua do Matão 321/Travessa 14, 05508-090 São Paulo, Brazil.

² Albert-Ludwigs-Universität Freiburg, Department of Biometry and Environmental System Analysis. Tennenbacher Straße 4, 79106 Freiburg, Germany.

E-mails: marmello@usp.br (*corresponding author); carsten.dormann@biom.uni-freiburg.de

Abstract: In the past decades, studies on interaction networks may have zoomed in too much on arbitrary subsets of ecological communities. Consequently, the current “nestedness paradigm” is biased towards restricted taxonomic groups. This bias leads to a myopic understanding of ecological functions and ecosystem services, with serious consequences for basic and applied ecology.

Keywords: assembly rules; ecological synthesis; network science; species interactions; theoretical ecology.

Everything in the world is connected. From public transport lines to smartphones in our pockets, no entity operates alone (Vespignani 2018). Connections are also crucial for reaching most of UN’s Sustainable Development Goals (SDGs, United Nations 2015), considering how interactions between humans, animals, plants, fungi, microbes, and their environments affect crop productivity and disease outbreaks, for example (Stanworth *et al.* 2024). This interconnectedness is scientifically daunting, and zooming into interaction subsets is tempting. However, what if we have zoomed in too much?

Let us zoom out a little. Network science, with roots back in the 18th century, became a pervasive approach to study connections. In ecology, it is widely used to study species interactions. Initially focusing on unweighted unipartite foodwebs in the 1880s, between the 1980s and the 2000s, the hype shifted to weighted bipartite mutualistic networks (Ings & Hawes 2018). Since then, a new paradigm has emerged, as we have assumed that mutualisms were nested like Russian dolls (Bascompte *et al.* 2003). In other words, the interactions of least connected species would represent a subset of the interactions of the most connected species. However, how well

does the nestedness paradigm represent real-world systems? Recent analyses and models show that we need to talk.

To smooth our conversation, let us consider the old Buddhist parable of the elephant and the blind monks. It seems that, like those monks, we all have been looking only at network pieces instead of entire systems, which has led to a myopic perspective. This is worrisome, as the nestedness paradigm predominates not only in basic ecology but is also a fledging standard for conservation and restoration, in the sense that it is assumed that a pristine or fully restored interaction network should have a nested topology, especially in the case of mutualisms (Howe 2016, Windsor *et al.* 2022, Lussier *et al.* 2024). This raises serious concerns, as our studies and many others have shown that nestedness is more like a by-product (Staniczenko *et al.* 2013, Valverde *et al.* 2018) than a true topology observed in natural systems. In addition, nestedness as measured by binary indices is even more context-sensitive than its weighted versions (Corso *et al.* 2015), missing finer details about realized niches (Fründ *et al.* 2016), so network studies at a local scale should prefer using weighted nestedness metrics whenever possible.

Therefore, using nestedness as a proxy for ecosystem health can easily lead to misguided inferences and recommendations.

Gold standards should always be taken with a grain of salt. Divergent discoveries have led different research teams, including ours, to question the nestedness paradigm (Dormann 2023, Blüthgen & Staab 2024). For example, when studying networks formed by oil-collecting bees and oil flowers, we noticed, on the one hand, that they were much more nested and less modular than many pollination networks were (Bezerra *et al.* 2009). On the other hand, pollination networks formed by mixed taxonomic groups were reported to be non-nested, and rather modular, such as the filter bubbles observed on social media (Bezerra *et al.* 2009).

After taking a close look at the divergent evidence discussed in the previous paragraphs, we felt like the blind monks examining the elephant. At that time, in the vast majority of studies, ecologists were analyzing interaction networks at a local scale formed by a single taxonomic group. Therefore, they were looking at unrepresentative pieces but drawing conclusions for the whole (Mello *et al.* 2011). This strong bias caused by spatial scale, taxonomic scope, and sampling methods, which markedly differ between studies conducted by different research groups, was only recently investigated in detail (Brimacombe 2024). Network topology, especially when measured with binary metrics, is strongly dependent on spatial scale, which varies considerably across studies (Corso *et al.* 2015, Poisot *et al.* 2020, Brimacombe *et al.* 2023). Thus, we need to consider that a monolayer interaction network at a local scale should be delimited based on the ecological function, for example pollination, that results from the interaction type under scrutiny. The relationships among spatial scale, taxonomic scope, and ecological function are more intricate than they initially appear, as we discuss in the following paragraphs. The elephant's leg may feel like a temple column, but it is neither an elephant nor a temple.

Sometimes, a leg is just a leg. Over time, we discovered that other mutualistic networks, such as those formed by bats, birds, bees, moths, wasps, ants, and flies, but especially those with mixed taxonomic groups (Donatti *et al.* 2011, Mello *et al.* 2011, Sarmiento *et al.* 2014, Schleuning *et al.* 2014, Costa *et al.* 2016), were incompatible with the

nestedness paradigm. Furthermore, in addition to nestedness and modularity, some networks seem to have a gradient topology, with highly specialized, non-overlapping interactions (Fonseca & Ganade 1996). A fourth structural archetype, a compound topology (later referred to as “in-block nestedness”, Lampo *et al.* 2024), which was proposed years ago as a modular network with internally nested modules (Lewinsohn *et al.* 2006), could be the key to integrate divergent evidence. This realization allowed us to conclude that, although nestedness and modularity are not actually two sides of the same coin (Fortuna *et al.* 2010), there is a balance between those two archetypical topologies, which depends mainly on the diversity contained in the network. As we zoom out from single-taxon systems, embracing whichever groups contribute to the ecological function under scrutiny at a site of interest, nested subnetworks aggregate into the elephant of network topology.

Aiming to look at the whole elephant, the Integrative Hypothesis of Specialization (IHS, first proposed by Pinheiro *et al.* 2016) reconciles divergent evidence and explains the coexistence of seemingly mutually exclusive topologies. For the sake of generality, we refer to the two groups in a bipartite network as consumers and resources. The IHS is based on three principles (formulated in detail by Pinheiro *et al.* 2019). First, the unique characteristics of each resource species affect its use by each consumer species. Second, any consumer mutation that improves the use of one resource also tends to enhance its use of similar resources while diminishing the value of dissimilar resources. Third, a consumer's ability to use a resource at a given time results from its previous adaptations and maladaptations. These three principles articulate five parameters that affect one another: (1) resource species richness, (2) consumer species richness, (3) method to generate the initial matrix, (4) maximum resource dissimilarity, and (5) number of resource clusters. Among those parameters, resource dissimilarity stands out as the most important in determining where one topology ends and another begins.

With this structure, the deductive argument of the IHS leads to the conclusion that the topology of an interaction network is primarily determined by resource dissimilarity from the consumer's perspective. This conclusion leads to secondary deductions, such as that networks with high

resource dissimilarity tend to have a modular topology, whereas networks with low resource dissimilarity become nested. A compound topology would emerge in sufficiently large networks that contain high resource dissimilarity distributed in clusters. After being conceived as a graphical model (Pinheiro *et al.* 2016), which was later turned into an algorithmic model that underwent a proof of concept through computer simulations (Pinheiro *et al.* 2019), predictions deduced from the IHS were submitted to several empirical tests showing that compound topologies are indeed much more common than previously thought (*e.g.*, Pinheiro *et al.* 2022). Interaction networks, therefore, resemble filter bubbles filled with Russian dolls.

A good example of this kind of compound topology was observed in a system formed by small mammals and their ectoparasitic fleas (Felix *et al.* 2022). By considering different orders of mammals together at both the local scale and the continental scale, the interaction constraints were strong for the ectoparasites (*i.e.*, not all flea species can infest all mammals), consequently leading to the formation of modules in most networks (Felix *et al.*, 2022). In addition, those modules had an internally nested structure, as ectoparasites have more freedom of interaction after they overcome those major interaction constraints. A compound topology also becomes evident when the taxa included in a network are even farther phylogenetically apart. This is the case for a nocturnal pollination network formed by bats and hawkmoths (Queiroz *et al.* 2021).

Both its logical structure and the empirical support reported in the studies cited above indicate that rather than being just a hypothesis, the IHS could be seen as a budding scientific theory, offering testable predictions based on operational variables within a coherent framework, in addition to a dedicated toolset (check out a reproducibility tutorial at: <https://github.com/marmello77/Restricted-Null-Model>). Moreover, it could become an axiomatic theory of the semantic type, which is rare in ecology and represents a significant advancement (Travassos-Britto *et al.* 2021). Other models also aim at explaining network assembly but have much broader scope, such as allometric trophic networks (Martinez 2020), or narrower scope, such as a microbial network model (Flores *et al.* 2013). There are even other models focusing on compound topologies, but they differ from the

IHS in that they propose that nestedness limits modularity and not the other way round (Palazzi *et al.* 2019) or assume an intrinsic or necessary relationship between topology and stability (Lampo *et al.* 2024). The IHS, in turn, focuses on explaining the assembly rules of archetypical topologies in the most parsimonious way, without evoking a circular relationship with stability. Its semantic nature and direct mapping to measurable operational variables make it quite helpful for studying real-world systems. Nevertheless, all comprehensive models of the assembly rules of interaction networks, which consider that different topologies may arise from an interplay between key factors, should be studied and thoroughly tested, as they have different scopes and thus serve different purposes.

In summary, together as a community, we have learned a lot in the past decades, coming a bit closer to a Buddhist “middle way”. The elephant offers enough to marvel, even after recognizing its parts as what they are: elements of a bigger picture. It is now time to broaden our perspective, so we can not only improve our theoretical understanding of species interactions but also allow its application to the challenges posed by the SDGs.

ACKNOWLEDGMENTS

We thank all colleagues who have been helping us develop the Integrative Hypothesis of Specialization since 2007, especially Boris Krasnov, Cristina Kita, Elisabeth Kalko, Francisco Rodrigues, Gabriel Felix, Isabel Machado, Marcelo Nogueira, Marco Tschapka, Nastaran Lotfi, Pierre-Michel Forget, Rafael Pinheiro, Renata Muylaert, Robert Poulin, Sharlene Santana, and Tanja Straka. We are also grateful to Jochen Fründ and Nico Blüthgen for the exciting discussions about network topology, assembly rules, and ecological theory. MARM was funded by the Alexander von Humboldt Foundation (AvH, 1134644) and São Paulo Research Foundation (FAPESP, 2018/20695-7, 2023/02881-6, 2023/17728-9, and 2023/03083-6). We are also grateful to FAPESP, National Council for Scientific and Technological Development (CNPq), Coordination for the Improvement of Higher Education Personnel (CAPES), and German Academic Exchange Service (DAAD) for the scholarships and fellowships granted to our students and postdocs.

REFERENCES

- Bascompte, J., Jordano, P., Melián, C. J., & Olesen, J. M. 2003. The nested assembly of plant–animal mutualistic networks. *Proceedings of the National Academy of Sciences*, 100(16), 9383–9387. DOI: 10.1073/pnas.1633576100
- Bezerra, E. L. S., Machado, I. C., & Mello, M. A. R. 2009. Pollination networks of oil-flowers: a tiny world within the smallest of all worlds. *Journal of Animal Ecology*, 78(5), 1096–1101. DOI: 10.1111/j.1365-2656.2009.01567.x
- Blüthgen, N., & Staab, M. 2024. A critical evaluation of network approaches for studying species interactions. *Annual Review of Ecology, Evolution, and Systematics*. DOI: 10.1146/annurev-ecolsys-102722-021904
- Brimacombe, C. 2024. Publication-driven consistency in food web structures: Implications for comparative ecology. *Ecology*. DOI: 10.1002/ecy.4467
- Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. 2023. Shortcomings of reusing species interaction networks created by different sets of researchers. *PLOS Biology*, 21(4), e3002068. DOI: 10.1371/journal.pbio.3002068
- Corso, G., Torres Cruz, C. P., Pinto, M. P., de Almeida, A. M., & Lewinsohn, T. M. 2015. Binary versus weighted interaction networks. *Ecological Complexity*, 23, 68–72. DOI: 10.1016/j.ecocom.2015.04.003
- Costa, F. v., Mello, M. A. R., Bronstein, J. L., Guerra, T. J., Muylaert, R. L., Leite, A. C., & Neves, F. S. 2016. Few ant species play a central role linking different plant resources in a network in rupestrian grasslands. *PLOS ONE*, 11(12), e0167161. DOI: 10.1371/journal.pone.0167161
- Donatti, C. I., Guimarães, P. R., Galetti, M., Pizo, M. A., Marquitti, F. M. D., & Dirzo, R. 2011. Analysis of a hyper-diverse seed dispersal network: modularity and underlying mechanisms. *Ecology Letters*, 14(8), 773–781. DOI: 10.1111/j.1461-0248.2011.01639.x
- Dormann, Carsten F. 2023. The rise, and possible fall, of network ecology. In: C.F. Dormann, P. Batáry, I. Grass, A.-M. Klein, J. Loos, C. Scherber, I. Steffan-Dewenter, & T. C. Wanger (Eds.), *Defining Agroecology: A Festschrift for Teja Tscharntke*. pp. 143–159. 1st ed. Hamburg: Tredition.
- Felix, G. M., Pinheiro, R. B. P., Poulin, R., Krasnov, B. R., & Mello, M. A. R. 2022. The compound topology of host–parasite networks is explained by the integrative hypothesis of specialization. *Oikos*, 2022(1). DOI: 10.1111/oik.08462
- Flores, C. O., Valverde, S., & Weitz, J. S. 2013. Multi-scale structure and geographic drivers of cross-infection within marine bacteria and phages. *The ISME Journal*, 7(3), 520–532. DOI: 10.1038/ismej.2012.135
- Fonseca, C. R., & Ganade, G. 1996. Asymmetries, compartments and null interactions in an amazonian ant-plant community. *The Journal of Animal Ecology*, 65(3), 339. DOI: 10.2307/5880
- Fortuna, M. A., Stouffer, D. B., Olesen, J. M., Jordano, P., Mouillot, D., Krasnov, B. R., Poulin, R., & Bascompte, J. 2010. Nestedness versus modularity in ecological networks: two sides of the same coin? *Journal of Animal Ecology*, 79(4), 811–817. DOI: 10.1111/j.1365-2656.2010.01688.x
- Fründ, J., McCann, K. S., & Williams, N. M. 2016. Sampling bias is a challenge for quantifying specialization and network structure: lessons from a quantitative niche model. *Oikos*, 125(4), 502–513. DOI: 10.1111/oik.02256
- Howe, H. F. 2016. Making dispersal syndromes and networks useful in tropical conservation and restoration. *Global Ecology and Conservation*, 6, 152–178. DOI: 10.1016/j.gecco.2016.03.002
- Ings, T. C., & Hawes, J. E. 2018. The history of ecological networks. In: W. Dáttilo & V. Rico-Gray (Eds.), *Ecological Networks in the Tropics*. pp. 15–28. Cham: Springer International Publishing.
- Lampo, A., Palazzi, M. J., Borge-Holthoefer, J., & Solé-Ribalta, A. 2024. Structural dynamics of plant–pollinator mutualistic networks. *PNAS Nexus*, 3(6). DOI: 10.1093/pnasnexus/pgae209
- Lewinsohn, T. M., Inácio Prado, P., Jordano, P., Bascompte, J., & M. Olesen, J. 2006. Structure in plant–animal interaction assemblages. *Oikos*, 113(1), 174–184. DOI: 10.1111/j.0030-1299.2006.14583.x
- Lussier, N. M., Crafford, R. E., Reid, J. L., & Kwit, C. 2024. Seeding success: Integrating seed dispersal networks in tropical forest restoration. *Biotropica*. DOI: 10.1111/btp.13347
- Martinez, N. D. 2020. Allometric trophic networks from individuals to socio-ecosystems: consumer–resource theory of the ecological elephant in the

- room. *Frontiers in Ecology and Evolution*, 8. DOI: 10.3389/fevo.2020.00092
- Mello, M. A. R., Santos, G. M. de M., Mechi, M. R., & Hermes, M. G. 2011. High generalization in flower-visiting networks of social wasps. *Acta Oecologica*, 37(1), 37–42. DOI: 10.1016/j.actao.2010.11.004
- Mello, M. A. R., Marquitti, F. M. D., Guimarães, P. R., Kalko, E. K. V., Jordano, P., & de Aguiar, M. A. M. 2011. The modularity of seed dispersal: differences in structure and robustness between bat- and bird-fruit networks. *Oecologia*, 167(1), 131. DOI: 10.1007/s00442-011-1984-2
- Palazzi, M. J., Borge-Holthoefer, J., Tessone, C. J., & Solé-Ribalta, A. 2019. Macro- and mesoscale pattern interdependencies in complex networks. *Journal of The Royal Society Interface*, 16(159), 20190553. DOI: 10.1098/rsif.2019.0553
- Pinheiro, R. B. P., Félix, G. M. F., Chaves, A. V., Lacorte, G. A., Santos, F. R., Braga, É. M., & Mello, M. A. R. 2016. Trade-offs and resource breadth processes as drivers of performance and specificity in a host–parasite system: a new integrative hypothesis. *International Journal for Parasitology*, 46(2), 115–121. DOI: 10.1016/j.ijpara.2015.10.002
- Pinheiro, R. B. P., Felix, G. M. F., Dormann, C. F., & Mello, M. A. R. 2019. A new model explaining the origin of different topologies in interaction networks. *Ecology*, 100(9), e02796. DOI: 10.1002/ecy.2796
- Pinheiro, R. B. P., Felix, G. M. F., & Lewinsohn, T. M. 2022. Hierarchical compound topology uncovers complex structure of species interaction networks. *Journal of Animal Ecology*, 91(11), 2248–2260. DOI: 10.1111/1365-2656.13806
- Poisot, T., Bergeron, G., Cazelles, K., Dallas, T., Gravel, D., MacDonald, A., Mercier, B., & Vissault, S. 2020. Environmental biases in the study of ecological networks at the planetary scale. *BioRxiv*, 2020.01.27.921429. DOI: 10.1101/2020.01.27.921429
- Queiroz, J. A., Diniz, U. M., Vázquez, D. P., Quirino, Z. M., Santos, F. A. R., Mello, M. A. R., & Machado, I. C. 2021. Bats and hawkmoths form mixed modules with flowering plants in a nocturnal interaction network. *Biotropica*, 53(2), 596–607. DOI: 10.1111/btp.12902
- Sarmiento, R., Alves-Costa, C. P., Ayub, A., & Mello, M. A. R. 2014. Partitioning of seed dispersal services between birds and bats in a fragment of the Brazilian Atlantic Forest. *Zoologia (Curitiba)*, 31(3), 245–255. DOI: 10.1590/S1984-46702014000300006
- Schleuning, M., Ingmann, L., Strauß, R., Fritz, S. A., Dalsgaard, B., Matthias Dehling, D., Plein, M., Saavedra, F., Sandel, B., Svenning, J.-C., Böhning-Gaese, K., & Dormann, C. F. 2014. Ecological, historical and evolutionary determinants of modularity in weighted seed-dispersal networks. *Ecology Letters*, 17(4), 454–63. DOI: 10.1111/ele.12245
- Staniczenko, P. P. A., Kopp, J. C., & Allesina, S. 2013. The ghost of nestedness in ecological networks. *Nature Communications*, 4(1), 1391. DOI: 10.1038/ncomms2422
- Stanworth, A., Peh, K. S. -H., & Morris, R. J. 2024. Linking network ecology and ecosystem services to benefit people. *People and Nature*. DOI: 10.1002/pan3.10632
- Travassos-Britto, B., Pardini, R., El-Hani, C. N., & Prado, P. I. 2021. Towards a pragmatic view of theories in ecology. *Oikos*, 130(6), 821–830. DOI: 10.1111/oik.07314
- United Nations. 2015. Transforming our world: The 2030 agenda for sustainable development. p. 1–41. New York. (Retrieved on December 5th, 2022, from <https://sdgs.un.org>).
- Valverde, S., Piñero, J., Corominas-Murtra, B., Montoya, J., Joppa, L., & Solé, R. 2018. The architecture of mutualistic networks as an evolutionary spandrel. *Nature Ecology & Evolution*, 2(1), 94–99. DOI: 10.1038/s41559-017-0383-4
- Vespignani, A. 2018. Twenty years of network science. *Nature*, 558(7711), 528–529. DOI: 10.1038/d41586-018-05444-y
- Windsor, F. M., Armenteras, D., Assis, A. P. A., Astegiano, J., Santana, P. C., Cagnolo, L., Carneiro, L. G., Emary, C., Fort, H., Gonzalez, X. I., Kitson, J. J. N., Lacerda, A. C. F., Lois, M., Márquez-Velásquez, V., Miller, K. E., Monasterolo, M., Omacini, M., Maia, K. P., Palacios, T. P., Pocock, M. J. O., Poggio, S. L., Varassin, I. G., Vázquez, D. P., Tavella, J., Rother, D. C., Devoto, M., Guimarães, P. R., & Evans, D. M. 2022. Network science: Applications for sustainable agroecosystems and food security. *Perspectives in Ecology and Conservation*, 20(2), 79–90. DOI: 10.1016/j.pecon.2022.03.001

Submitted: 17 June 2024

Accepted: 26 February 2025

Published online: March 14 2025

Associate Editor: Gudryan Barônio