



Established Invasive Tree Species Offer Opportunities for Forest Resilience to Climate Change

Bart Nyssen^{1,2} · Jan Den Ouden³ · Anja Bindewald⁴ · Pedro Brancalion⁵ · Klaus Kremer⁶ · Katharina Lapin⁷ · Lisa Raats¹ · Elisabeth Schatzdorfer⁸ · John Stanturf⁹ · Kris Verheyen¹⁰ · Bart Muys²

Accepted: 21 August 2024 / Published online: 14 October 2024
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2024

Abstract

Purpose of review A rapidly changing climate is weakening the resilience of forest ecosystems through vitality loss of major native tree species, which reduces the ability of forests to deliver ecosystem services. Established invasive tree species (EITS) may take over the vacant space potentially preventing the regeneration of the preferred native tree species. This paper aims to investigate how expansion of these invasive non-native tree species can be addressed in a context of climate-smart forest management, considering alternatives to costly and often ineffective EITS control measures.

Recent findings We found that forest ecologists increasingly recognize that climate-smart forest management, in particular tree species diversification and close-to-nature forest management, can strengthen the resilience of forests against negative impacts by EITS. In the resulting resilient forest ecosystems, a more closed canopy may deprive EITS of their invasive nature, and EITS may contribute to climate change adaptation.

Summary This review proposes new pathways for forest management transcending the apparent incompatibility between the dominance of EITS and the adaptation capacity of forests and forest management to climate change. Adaptive measures to increase the resilience of forests to climate change may prevent the dominance of EITS. Under such conditions, useful functional traits of these tree species may even contribute to maintenance or enhancement of biodiversity, provisioning of ecosystem services and adaptation to climate change.

Keywords Climate-smart forest management · Non-native species · Invasiveness · Invasibility · Biodiversity · Forest restoration · *Ailanthus altissima* · *Eucalyptus globulus* · *Pinus radiata* · *Prunus serotina* · *Robinia pseudoacacia*

Introduction

Global change, caused by the rapid increase of human activities and the anthropogenic emissions of greenhouse gases, threatens the functioning of forest ecosystems. Climate

warming and biological invasions are among the major drivers of ecosystem change, and they are often connected through positive feedback loops [1, 2]. These stressors may affect forest structure and functioning by changing abiotic conditions, vegetation structure and species composition

✉ Bart Nyssen
b.nyssen@bosgroepen.nl

¹ Bosgroep Zuid Nederland, Huisvenseweg 14, 5591 VD Heeze, The Netherlands

² Department of Earth & Environmental Sciences, KU Leuven, Celestijnenlaan 200E Box 2411, 3001 Leuven, Belgium

³ Forest Ecology and Management Group, Wageningen University, P.O. Box 47, Wageningen, The Netherlands

⁴ Department of Forest Conservation, Forest Research Institute of Baden-Württemberg (FVA), Freiburg, Germany

⁵ Department of Forest Sciences, “Luiz de Queiroz” College of Agriculture, University of São Paulo, Piracicaba, Brazil

⁶ Chair of Silviculture, Institute of Forest Sciences, University of Freiburg, Tennenbacherstr. 4, 79106 Freiburg, Germany

⁷ Austrian Research Centre for Forests BFW, Vienna, Austria

⁸ European Forest Institute, Bonn, EFI, Germany

⁹ Institute of Forestry and Engineering, Estonian University of Life Sciences, Kreutzwaldi 5, 51006 Tartu, Estonia

¹⁰ Forest & Nature Lab, Department of Environment, Ghent University, Geraardsbergsesteenweg 267, 9090 Melle, Gontrode, Belgium

[3–8]. Rapid climate warming triggers more frequent and heavier disturbances—including wildfires, windstorms, droughts, insect, and pathogen outbreaks—threatening the ability of forest ecosystems to provide ecosystem services [1, 9–11]. Also, establishing non-native invasive species cause changes in ecosystem structure and functions [12–25].

Tree species have been moved across continents in the past for various reasons (e.g., landscape aesthetics, ecosystem restoration, enrichment of biodiversity or wood production) and in many cases have become part of regional tree species pools worldwide. Some species stand out because of their competitive behaviour and other impacts on ecosystems, and are referred to as ‘invasive’. Research on the effects of non-native tree species in forests often focuses on the prevention of new invasions. In this review, we focus on invasive tree species that are already well established and widespread, here called; ‘Established Invasive Tree Species’ (EITS).

EITS can be harmful to non-forest native ecosystems, like grasslands, rangelands, and savannas [26–28]. As fast-growing tree species with pioneer traits and colonisation potential, they may turn these vegetation types into forests, resulting in potential local extinctions of light-demanding species [28–30]. Covering the impacts of EITS in this context is out of the scope of the paper. Yet, these species may also have significant impacts on forest-related biodiversity, ecosystem processes and related ecosystem services [25, 28, 31–35]. They may directly or indirectly influence the availability of resources for other species by causing drastic changes in the biotic or abiotic environment (e.g., by shading, competition for belowground resources, effects on biogeochemistry through rich litter, N-fixation) [36–38] hampering the regeneration of native tree species [39–42]. In fire-prone regions, EITS can alter fire regimes by increasing fuel availability and flammability [43, 44]. Additionally, EITS may have a high potential for conflict with nature conservation objectives, since they are usually not part of the target forest composition [45, 46]. Yet, some benefits of EITS have also been reported [47–49]. Such benefits may include contributions to species richness [50–52], enhancement of nutrient availability [53–55], provision of ecosystem services [56–58] and genetic conservation of EITS threatened in their native habitat [59].

As forest managers seek a pragmatic way to deal with EITS, the climatic conditions under which they do so are also changing. Global average temperatures have risen rapidly in recent decades, with ensuing droughts and heat waves [60]. The effects of climate change have caused increased tree mortality on all continents [61–63]. Sometimes forests were unable to recover [64], while in other situations, forests

reacted by shifts in tree species composition [65]. These observations suggest that persistent and progressive changes in global forests may occur in response to climate change so that the provision of ecosystem services, such as climate regulation, biomass production, food, medicine, water supply and purification, pollination, and habitat provision for forest species, may decline [66–68]. Often EITS are better adapted to these changes, possibly due to higher ecological plasticity and thus broader ecological amplitude of tolerated ecological conditions [57, 69, 70].

The disturbances caused by climate change may also increase the susceptibility of forests to biological invasions due to reduced dispersal, establishment and recruitment of late-successional, long-lived tree species. These conditions favour pioneer trees with high turnover rates, which are attributes of most EITS. Moreover, resources released through mortality, such as light, can locally facilitate the expansion of shade-intolerant EITS. [71–77].

In addition to natural forests with drought-intolerant native trees, forests sensitive to climate change often turn out to be highly disturbed ecosystems with altered abiotic conditions and species composition—such as plantations or forest expansion on abandoned industrial and agricultural sites. Such altered forest ecosystems with abiotic and biotic conditions outside the historical range of variability of a given place are examples of novel ecosystems [50, 51, 78–80]. One expected outcome of climate change is that many of these novel systems will emerge worldwide, with increasing occurrence of EITS as a consequence [81–83].

In research on the resilience of forest ecosystems, the presence of these EITS is commonly perceived as a threat [84–86], whereas some non-invasive, non-native tree species are sometimes expected to contribute to biodiversity, ecosystem services and climate change adaptation [87–90]. In management practice, however, this division between non-invasive and invasive established tree species is not that clear-cut [46], and even EITS can contribute to the provision of ecosystem services [67, 91]. Forest managers and researchers notice that under some circumstances EITS can also contribute to the resilience of forests to the changing climate [57, 92, 93].

One of the main contributions to resilience may be filling vacant niches, so that some aspects of structure and functions recover faster after a disturbance. For instance, some EITS may operate as nurse trees—by regulating the forest microclimate and sometimes promoting litter decomposition – and stimulate forest succession [94–97]. In the scientific literature, these benefits of EITS, based on the same processes that were interpreted previously as signs of invasiveness, have received little research attention so far.

Although human-induced climate change and the presence of invasive species are relatively new phenomena, pressure from stressors is part of any forest ecosystem. The set of mechanisms that jointly ensure the survival and renewal of forest ecosystems under environmental stress will determine their resilience. In forest ecology, various resilience definitions are used that do not all agree [98, 99]. In this paper resilience is understood as the capacity of a forest ecosystem to *absorb* (resist) environmental changes (pressures) and natural and anthropogenic disturbances (pulses) without losing its structure and functions while *adapting* to the changing environmental conditions, the so-called ecological resilience [99–101].

Strictly speaking, forest resilience is often considered to be relative to one specific ecosystem service (e.g., wood production) [98, 102]. However, in most cases forests are expected to provide a wide range of ecosystem services [1, 4, 11, 57, 68, 103]. Thus, the concept of resilience commonly refers to the stability in providing a set of ecosystem services under changing conditions.

In this review we investigate how the presence of EITS can be addressed in the context of climate-smart forest management, aiming to strengthen forest resilience to climate change. We consider the presence of EITS to be no longer avoidable in many regions, already present in forests and often in high density. So, in this context, EITS are not considered as potential or actual threats to forests but as elements that are already present, with a potential contribution to forest resilience against climate change. When we refer to changes in external conditions, we specifically mean the effects of ongoing climate change.

Our research question is twofold:

- (1) How can EITS be integrated in forest ecosystems and forest management so that these tree species contribute to forest biodiversity and forest-related ecosystem services?
- (2) Can EITS themselves contribute to the climate change adaptation capacity of these forests?

Materials and methods

We carried out a literature review on the role of EITS in forest resilience enhancement to climate change in three steps:

- (1) Identification of challenges and benefits of EITS and the possibilities of integrating EITS in forest ecosystems (see section ‘EITS challenges and opportunities’);

- (2) Describing the impact of integrated EITS on biodiversity and ecosystem services, with a focus on timber production (see section ‘Contribution of integrated EITS to species diversity and wood production’);
- (3) Exploring the effects of EITS on climate resilience and their potential to contribute to climate change adaptation (see section ‘Contribution of EITS to climate resilience’).

We used an informal iterative workflow in which we interpreted data from the literature using expert knowledge and concepts developed by the authors based on years of experience with EITS in forests. We searched for relevant articles in databases such as PubMed, Scopus, and Web of Science. We also consulted experts in the field and discussed the findings with them. Based on the findings, we created a list of key points and synthesised the results of 466 publications, mainly published in the period after the year 2000.

In this review, we restrict ourselves to the impact of tree species on forest ecosystem functioning and the provision of ecosystem services. Therefore, we do not include societal impacts or origin of a species as is often done in assessments of impacts of biological invasions [104–107] but use an ecological definition of invasiveness in which the term describes the competitive advantages that enable a species to proliferate rapidly and conquer new environments [31, 108–112]. Moreover, since EITS are just as permanently present as native tree species in many parts of the world, we separate ‘invasive’ from ‘non-native’ [113]. The two dual concepts, invasive vs non-invasive and native vs non-native, yield four possible combinations of these terms. ‘Invasive non-native’ and ‘invasive native’ both indicate species that exhibit an invasive character. In this ecological definition of invasiveness, invasive species sustain self-replacing populations over several life cycles, produce reproductive offspring, often in very large numbers, and have the potential to spread over long distances [31, 108–110]. In forests, invasive tree species appear to exhibit typical pioneer behaviour: early seed setting, easy spreading and rapid juvenile growth [35, 92, 114]. These pioneer tree species usually have high light requirements upon establishment [115]. From the extensive lists compiled of invasive tree species [28] we will refer in more detail to five globally widespread EITS, which are each appealing examples in different parts of the world: *Ailanthus altissima* (Mill.) Swingle, *Eucalyptus globulus* Labill., *Pinus radiata* D. Don, *Prunus serotina* Ehrh. and *Robinia pseudoacacia* L. See Table 1 for an overview of their occurrence. A short characterisation of each species, based on this review, is given in separate boxes (see Boxes 1, 2, 3, 4, and 5).

Table 1 EITS with the continent of origin (O), the continent where they are considered invasive (I) [28] and reference to the box with short descriptions of each species

EITS							Box
Continent	Africa	North America	South America	Asia	Europe	Oceania	
<i>Ailanthus altissima</i>	I	I	I	O/I	I	I	Box 1
<i>Eucalyptus globulus</i>		I	I		I	O/I	Box 2
<i>Pinus radiata</i>	I	O	I	I		I	Box 3
<i>Prunus serotina</i>	I	O			I		Box 4
<i>Robinia pseudoacacia</i>	I	O/I	I		I		Box 5

Invasiveness of EITS appears to vary considerably, from the highly invasive *P. serotina* [116–119]—thanks to efficient dispersal by birds and mammals over short and long distances -, over the moderately invasive *A. altissima* [120–123], *P. radiata* [28, 29, 76] and *R. pseudoacacia* [54, 70, 124]—thanks to efficient wind dispersal -, to the limited invasiveness of *E. globulus* [125–131].

To organise all gathered information and gain insight into the possibilities for enhancing forest resilience to EITS and climate change, we first developed a conceptual framework for forest resilience under climate change (see section ‘[Forest resilience framework](#)’). This framework was also valuable to address the abundant information on forest adaptation to climate change, resilience against dominance by EITS, and possible contributions of EITS to forest adaptation to climate change. Moreover, it helped us to bridge the apparent contradiction between reducing EITS by strengthening forest resilience, and the possible positive contribution of these EITS to the resilience of the forest ecosystem against climate change. Finally, in the conclusion section, we formulate scenarios and possible solutions for a way forward for EITS integration based on our findings and synthesis.

Forest resilience framework

Following the previous definition, forest resilience has two major components—absorption and adaptation -, both ecosystem reactions to stress grounded in ecosystem processes [132, 133]. Absorption refers to the plasticity of the ecosystem under relatively stable environmental conditions, both at the level of organisms and of the system as a whole. This concept aligns closely with the definition of *engineering resilience*. Under stable conditions the frequency and intensity of disturbances—such as storms, fires, wood harvesting, diseases, and pests - remain within the absorptive capacity of the ecosystem operating under a dynamic equilibrium. This ensures continuity in the composition, structure and functioning of the ecosystem [10, 134]. But when the resilience of a forest ecosystem is weak—e.g., due to extreme

herbivore pressure—or the character, intensity or frequency of disturbances fundamentally changes as a result of changing conditions—e.g., climate change and invasive species—then ecosystem processes may not be able to secure the persistence of individuals and recovery of populations. This is where ecosystem adaptation comes into play. Under these changed conditions, the ecosystem develops into one with a different composition, structure and functioning [10, 134]. The interplay of absorption and adaptation, as used in this paper, is also called *ecological resilience* [135, 136].

The dichotomy between absorption and adaptation, as proposed in literature [99, 101, 135] naturally connects to the practical discussion on climate change adaptation. Forest managers try to increase forest resilience for securing ecosystem services by optimising ecosystem processes based on the current vegetation composition, thus, relying on the ecosystem’s absorption. When it becomes clear that the current suite of forest tree species face a high risk of losing their vitality due to changing climatic conditions, the forest tree species composition may be adapted in an ongoing and long process by adding new provenances or species, both natives and non-natives [137]. In this way, the absorption capacity is complemented by the adaptation capacity. These two components of resilience, absorption, and adaptation, fit well with the two questions this paper addresses.

- (1) First, what effect has absorption of EITS within forest ecosystems on their components - e.g., abiotic conditions, vegetation structure and species composition - on forest biodiversity and ecosystem services?
- (2) And secondly can EITS, as an absorbed component of tree diversity and redundancy, contribute to absorption and adaptation of climate change effects?

To clarify the connection between the concepts of absorption and adaptation with the development of forest ecosystems under stress, we developed a forest resilience framework inspired mainly by the unified framework for ecological resilience of Falk et al. 2022 [134] (see Fig. 1).




Fig. 1 Framework for forest resilience. Under constant climatic conditions **Complex Forests** will preserve their structure and functions through absorption of stress. **Simplified Forests** under steady state conditions can, through forest succession, develop into **Less Simplified Forest** and when a large tree species pool is present in the landscape, even into **Complex Forests**. When absorption fails, due

to changing climatic conditions, adaptation occurs, involving different species or functional groups. Adaptation can lead to an **Adapted Complex Forest** while maintaining structure and function of the pre-disturbance community, or lead to **Degraded forest** as a result of altered structure or function (modified after Falk et al. (2022) [134])

In our framework we make a distinction between simplified and complex forests. Simplified forests occur where past land use has led to limited species diversity, limited redundancy, and simplified structure; these conditions can facilitate dominance of EITS. Complex forests have a more developed structure and are richer in tree species as a result of a lack of management or more extensive management (e.g., close-to-nature forestry). They are less prone to EITS dominance. Through forest succession and disturbances under stable external conditions, simplified forests develop into less simplified forests, which are more structured and richer in tree species – if propagules are present. In the presence of a diverse tree species pool, simplified and less simplified forests can develop into complex forests, characterised by structural and compositional complexity.

Absorption and adaptation are two different resilience components that may occur simultaneously. Under relatively stable climate conditions, absorption of disturbances dominates in complex forests resulting in continuity of ecosystem structure and functioning. Under changing climate conditions, the complex forest ecosystem continues to absorb disturbances. However, when the absorbing capacity is exceeded, adaptation may occur, and the complex forest develops into an adapted complex forest [138].

As long as the joint processes of absorption and adaptation ensure the preservation of structure and function, we speak of resilient forest ecosystems. However due to changing climatic conditions—reinforcing already existing stressors or introducing new ones [82]—forest structure and function may change, in which case forests may become insufficiently resilient [134, 139–141]. According to the FAO Global Forest Resources Assessment, we refer to these resulting forest ecosystems as degraded forests: forests whose structure and functions are negatively affected resulting in systems with lower capacity to supply products and/or services [142]. Simplified and less simplified forests are more at risk of degradation than complex forests due to limited diversity and species redundancy [137].

Resilience is often expressed in terms of a system's insurance by diversity and redundancy [100, 136, 143, 144]. The functional characteristics of tree species influence their performance in terms of growth, survival or reproduction [79]. Greater diversity in tree species with different functional traits is expected to reduce the impact of disturbance or stressors on the forest [60, 80–82, 145] and provide greater adaptability [83] (Fig. 2a). Besides the variation in tree species, it is also important for the resilience of the forest ecosystem that multiple tree species with similar characteristics occur, in



Fig. 2 Illustration of the concepts of functional diversity and redundancy within two stands. **a.** Although the stand in the upper pane consists of only two tree species, it has high functional diversity because they have very different functional properties: for example, one species is a deciduous broadleaf tree, and the other an evergreen conifer. However, because of the large difference in functional properties between these two species, functional redundancy is weak; if one spe-

cies disappears, several specific functional properties are lost. **b.** The lower stand also has high functional diversity because it consists of six different species, four deciduous broadleaf species and two conifer species with relatively similar characteristics. However, functional redundancy is high in this case because if one species disappears, the variation in functional traits in the stand will be maintained (adapted from Messier et al. 2019 [144])

case one tree species fails due to climate change effects. This is called functional redundancy [82, 84, 85] (Fig. 2b).

EITS challenges and opportunities

Significant negative impacts of EITS on biodiversity, ecosystem processes, and related ecosystem services have been reported [25, 28, 31–35, 146]. Conversely, other studies point at the contribution of EITS to species richness, forest resilience, ecosystem services, and climate change adaptation [47, 50, 52, 56, 57, 67, 69, 78, 147, 148]. Either way, EITS may directly or indirectly influence the availability of resources for other species by causing changes in the biotic or abiotic environment [36–38].

EITS, for example, affect nutrient availability and soil fertility, which varies greatly among tree species. Whereas *P. radiata* may acidify the forest soil, most notably on low-fertility sites [149], *P. serotina* [53], *A. altissima* [37] and *R. pseudoacacia* [150] usually increase nutrient availability. *Eucalyptus globulus* litter degrades slowly in monocultures [151], and the high biomass removal in plantation forestry linked to the high productivity of this tree species seems to reduce nutrient availability [152, 153] (See also Table 6). This high productivity also seems to be the main cause of the sometimes observed high water consumption in *E. globulus* plantations

[154]. However, even less productive EITS can influence the soil moisture availability for other tree species especially when they have a more efficient water uptake strategy [155].

Light availability can also be affected by the presence of EITS. Under pioneer conditions such as in spontaneous forest development on agricultural land, forest recovery after fire or in plantations consisting of tree species which form a translucent canopy, EITS can form dense shrub layers that alter the light environment that can limit the regeneration of native tree species for several years to decades [28, 156, 157]. Also allelopathic effects have been demonstrated in some EITS—e.g., *A. altissima*, *P. serotina*, *R. pseudoacacia*, *E. globulus*—under laboratory and experimental conditions [157–160], albeit with limited impact on tree species regeneration and survival under field conditions [37, 55, 161–164].

Yet, EITS may also positively influence forest recovery. Due to their invasive trait and frequent shade-intolerant nature, they may rapidly colonize deforested areas and accelerate canopy closure. This may benefit survival of many typical forest plants and animal species as they depend on the microclimate that prevails in forests [26, 88]. Through shading and transpiration, the closed crown canopy causes lower maximum temperatures in summer and higher minima in winter, while maintaining levels of air and soil moisture that can be higher than in the open [86, 87, 165].

Ailanthus altissima has often been planted for ornamental reasons in areas climatically similar to the cool northern side of its native range. Warming of the climate offers expansion potential here, which *A. altissima* uses, piggybacking on road and rail transport. Despite its expansive character, the review shows that where *A. altissima* forms mature forests in the presence of native tree species, the latter rejuvenate easily under its canopy.

TRAITS:

- Deciduous broadleaf pioneer tree, 20 – 30 m high
- More climate resilient than competitors
- Positive impact on nutrient availability
- High quality wood, comparable to *Fraxinus* spp.

NATIVE RANGE:

- Eastern China

INTRODUCTION PATHWAYS:


- Horticulture
- Silviculture

INVASION HABITAT:

- Pioneer species in forest margins and open forests, outside forests in urban and rural landscapes, heathlands, dry grasslands, and dunes.

STATUS:

- Naturalized in all continents but Antarctica. Widespread and common in its native Chinese range, Europe, North America, and Australia.



Ailanthus altissima distribution map; colors indicate gradients of occurrence from limited (yellow) to intensive (red) presence (source: <https://www.gbif.org/species/3190653>).



PICTURE: Native forest development under mature *Ailanthus altissima* forest in the Parcul National Portile de Fier, Romania (with *Acer campestre*, *Acer monspessulanum*, *Acer platanoides*, *Acer pseudoplatanus*, *Alnus glutinosa*, *Carpinus betulus*, *Cornus sanguinea*, *Corylus avellana*, *Crataegus monogyna*, *Euonymus europaeus*, *Fagus orientalis*, *Fagus sylvatica*, *Fraxinus excelsior*, *Fraxinus ornus*, *Juglans regia*, *Lonicera periclymenum*, *Prunus avium*, *Prunus spinosa*, *Quercus cerris*, *Quercus robur*, *Sambucus nigra*, *Tilia cordata*, *Tilia platyphyllos*, *Tilia tomentosa*, *Ulmus glabra* and *Ulmus minor*) (Photo Bart Nyssen)

Box. 1 *Ailanthus altissima*. Effective nurse tree with rapid spread facilitated by climate change

Through their influence on resources, EITS also influence forest regeneration [39, 40]. Especially when they grow densely, they can reduce the amount of light reaching the soil surface. Thus, they may often promote late successional tree species at the expense of pioneer tree species, thereby accelerating forest succession. Also, when EITS form the

tree canopy, the influence of their litter on soil development may play a decisive role. EITS with nutrient-rich leaves may increase the nutrient availability in the topsoil, thereby facilitating the establishment of pioneer and late successional tree species [134, 166]. This positive effect can, in fact, compensate for allelopathic effects [55, 165].

Changes due to the presence of EITS in the light environment and in nutrient and moisture availability in the topsoil also influences the herb layer. Research into the effect of EITS on the herb layer also shows conflicting results. EITS can reduce the biodiversity of the herb layer by promoting a higher abundance of generalist species [167]. But the main effects here are a shift from light-demanding to more shade-tolerant tree species and, in the case of EITS with nutrient-rich litter, to species bound to a richer forest floor. These tree species usually promote development towards specialist forest species [168]. To a greater or lesser extent EITS are host tree species for other organisms [54, 87, 169–173]. Co-evolution of species with their host plants has influence on the species richness associated with tree species. Therefore, with longer presence of EITS, their role as host plants increases [169], especially if they are widespread and congeneric native trees are present [171–174].

These impacts of EITS on the availability of resources and species composition can have an impact on ecosystem services and forest biodiversity. Thus, the presence of EITS may have a high potential for conflict with local nature conservation objectives, from which they are mostly excluded [45, 46]. This is especially the case when these nature conservation objectives are derived from strongly human influenced ecosystems, for instance intensively grazed systems or when tree species composition has been reduced to species that form a translucent canopy. In fire-prone regions, EITS can alter fire regimes by increasing fuel availability and flammability [43, 44]. In general, flammability varies strongly between tree species [175]. Some EITS plantations, such as *E. globulus* and *P. radiata*, are highly flammable. But broadleaf EITS mostly limit the fire risk of the—usually coniferous—forests in which they are established. A well-known broadleaved exception to this is *E. globulus*, which increases the fire hazard. If grown under a short-rotation, fire risk of *Eucalyptus* spp. is no different from *Pinus* spp. Older trees develop stringy bark that increases the risk of fires spreading because of a tendency to disperse embers from its bark [176].

Integration of EITS into forest ecosystems

The reason for large-scale planting of many EITS, such as *R. pseudoacacia* [54], *E. globulus* [95, 150] and *P. radiata* [177] was wood provisioning in short to medium rotation plantations. Other EITS, such as *P. serotina* and *A. altissima* have been planted for other reasons. For instance *P. serotina* was mainly planted in Europe as a companion tree species

in reforestation, mostly with *Pinus sylvestris*, besides some planting for wood production in the nineteenth century [178]. *Ailanthus altissima* was and is a popular ornamental tree [157] and, outside China, its country of origin, only small-scale planting for wood production took place in Austria, Hungary, Argentina, Uruguay, India and New Zealand [179–182].

All these EITS are known for their ability to become invasive [58, 183, 184]. The ability of non-native species to invade new communities depends on their traits and the vulnerability (invasibility) of the community [185]. They usually present pioneer traits associated with plants occurring in early successional stages of forests—high seed production at a young age, fast growth, and short juvenile period—enabling them to rapidly colonise open spaces inside and outside the forest [186–189]. EITS usually combine these pioneer traits with a strong light requirement and high light transparency. However, there are exceptions. Some EITS combine pioneer traits—e.g., early seed setting, efficient propagule spreading and rapid juvenile growth—with late successional species traits—e.g., shade-tolerant, shade-casting, and long-lived. The more shade-tolerant and shade-casting a tree species is, the longer it can dominate forest succession. *Acer platanoides* (which is native to Europe and invasive in other temperate regions) typically is such a tree species, combining traits associated with early successional stages [190, 191], with shade tolerance [191, 192], hence the problems with it in North American forests [190, 191].

Evidence from several studies indicates that very few non-native species invade successional advanced plant communities [185, 189, 193–196]. Non-native species are completely missing from undisturbed late successional forests [189, 197, 198], and the high diversity of native tree species limits forest invasion by EITS [199]. Ecosystem invasibility is the result of several factors, including physical environmental characteristics, the competitive ability of resident species, and the disturbance regime of the habitat [185, 193, 200, 201]. However, the availability of resources such as light, nutrients and water facilitates the establishment of EITS [71, 202, 203]. For instance, many EITS have low moisture and nutrient requirements and thus may out-compete many native species on sites with reduced nutrient and moisture availability [15, 58, 94, 204–206]. Ecosystem invasibility often results from anthropogenic or natural degradation of forest ecosystems [207]. Thus, the dominance by EITS is usually a symptom, rather than the cause, of limited forest resilience. Due to climate change, the level of degradation of forest ecosystems is generally expected to increase, and with it, their invasibility [75, 132, 208].

Eucalyptus globulus only regenerates in the immediate vicinity of the mother tree. *E. globulus* owes its reputation as an invasive tree species to massive planting in short rotation plantations impeding opportunities for natural regeneration of other species. But simply increasing rotation length to beyond 10 to 20 years creates opportunities for establishment of native species regeneration.

TRAITS:

- Evergreen broadleaf pioneer tree, growing up to 70 m high
- More climate resilient than most competitors
- Wood: mostly used for pulp and paper, but also high-quality timber applications

NATIVE RANGE:

- Australia

INTRODUCTION PATHWAYS:


- Silviculture.

INVASION HABITAT:

- Low invasion capacity, even in open forests.

STATUS:

- Major occurrence areas globally: Australia (host), Spain, Portugal, California, South Africa and South America. Occurs locally elsewhere in the temperate climatic zone.



Eucalyptus globulus distribution map; colors indicate gradients of occurrence from limited (yellow) to intensive (red) presence (source: <https://www.gbif.org/species/3176787>).



PICTURE: Native oak and sweet chestnut forest succession under 140 years old 70m high *Eucalyptus globulus* plantation. Souto da Retorta, Galicia, Spain (Photo Bart Nyssen)

Box. 2 *Eucalyptus globulus*. Bad reputation due to short rotation plantation forestry

This degradation can be countered by diversifying the tree species composition. Higher species richness results in higher functional diversity as long as the species in the community present different functional traits and therefore different strategies to acquire resources [209]. Higher functional diversity would reduce susceptibility to invasion through the pre-emption of available resources [210–214]. Additionally, functional trait similarities between resident species and introduced species result in overlapping resource requirements and, as a consequence, in competition among species [209, 214, 215]. As resource requirements overlap increases, competition is intense, and no more species can establish [216–218].

Thus, forest invasibility is a dynamic attribute that can be modulated by resource supply, disturbance regimes, and tree species composition of recipient forest ecosystems [71]. These factors acting in synergy, determine a complex scenario to predict the invasibility of forest ecosystems [76]. Nevertheless, the knowledge of the latter can be used as a management tool to prevent or control EITS in forests [77, 219]. Due to the

shade-intolerant nature of the set of EITS addressed here, they are unable to establish in complex forests. Yet, when they are dominant, their translucent crowns enable establishment of late successional species, which eventually makes them disappear from complex forests. However, if the possibility of competition is limited—by lack of seed trees or intensive foraging by a high density of herbivores—resources can be utilized by EITS, thus maintaining their presence [71].

According to this, the possibility of EITS becoming dominant can be reduced by adjusting the abiotic conditions that promote their establishment, such as soil nutrients, soil moisture, or light in the understory by strengthening the vertical forest structure; an approach that fits seamlessly with close-to-nature forest management [220]. At the same time, it is important to strengthen competition from other tree species by promoting the diversity of both pioneer species that can compete in the early stages of succession, and of late-successional species to stimulate forest succession [15, 58, 94, 116, 204–206] (See Table 2).

Table 2 Possibility of reducing EITS dominance in forest ecosystems

Species	Studies on reducing EITS dominance in forest ecosystems
<i>Ailanthus altissima</i>	[55, 121, 221–223]
<i>Eucalyptus globulus</i>	[95, 130, 224–230]
<i>Pinus radiata</i>	[96, 97, 231–242]
<i>Prunus serotina</i>	[94, 164, 165, 205, 243–258]
<i>Robinia pseudoacacia</i>	[54, 70, 124, 259–264]

Contribution of integrated EITS to species diversity and wood production

Adding a tree species—with its species-specific traits—to an existing forest ecosystem will change its biological diversity. This applies equally to native tree species as to EITS [265]. These effects of EITS on species diversity depend very much on EITS abundance [97]. When EITS density is high, these effects, negative or positive, are greater than when EITS occur in lower density [266]. In cultivated forests (i.e., planted or plantation forests), EITS are often intensively managed in short-rotation monoculture plantations. This short forest development period usually prevents the natural regeneration of native woody species comparable to dense, young stands of native trees [267–269]. However, less intensively managed EITS plantations on longer rotations and abandoned EITS plantations in many regions host a high diversity of plants and birds. This phenomenon is well known for pine and eucalypt plantations [95, 97, 230, 237, 270–272].

In forest ecosystems resilient against EITS dominance, both the negative and positive impacts of EITS on the species composition of the forest ecosystems decrease [96, 225, 265]. The negative effects of the presence of EITS, when dominant, can, in resilient forests, turn into positive diversity effects due to increasing diversity of habitats [265]. Meanwhile, a gradual adaptation takes place between species that are already present and EITS [169, 170].

As explained above, due to the pioneer traits of most EITS, their effects on biodiversity, viewed at the time scale of forest development, can be a transitional pioneer phase, and these EITS are likely to decline eventually if sufficient seed sources of successor species are present [94–96]. Due to this process, in complex forests, EITS are expected to be found only locally and temporarily, as they are part of the pioneer stage after disturbance. Also, they can profit from pioneer conditions at forest edges. But, generally speaking, large-scale negative or positive effects of EITS can dissolve over time in forest succession while small-scale pockets of EITS may remain. Therefore, strengthening resilience against dominance by EITS is likely to ensure that potential effects are limited and will not persist for long.

For some EITS, timber production in plantations is an important reason for planting (e.g., *P. radiata*, *E. globulus* and *R. pseudoacacia*). The question arises whether these

Table 3 Examples of high-value applications of EITS wood

Species	Saw logs for construction, furniture, or veneer
<i>Ailanthus altissima</i>	[121, 157, 223, 273–282]
<i>Eucalyptus globulus</i>	[153, 283–289]
<i>Pinus radiata</i>	[290–294]
<i>Prunus serotina</i>	[248, 258, 295–300]
<i>Robinia pseudoacacia</i>	[54, 70, 124, 261, 262, 301–307]

EITS and those whose invasive nature prevents forest managers from using them for timber production outside their area of origin (e.g., *P. serotina* and *A. altissima*) can contribute to wood provisioning in complex forests.

Although in plantation forestry wood production with EITS may be aimed at low-value applications such as firewood, pulp, and paper, EITS are also used to supply wood for high-value applications such as construction framing, furniture, and veneer (see Table 3). However, management of complex forests is characterised by small-scale interventions that limit the amount of light on the forest floor, and the additional growth will be concentrated on a limited number of high-quality trees. Thus, here too, the density of pioneer tree species – and thus of EITS – decreases in favour of late successional species.

The preferred high-end application for the hardwood species *A. altissima*, *E. globulus*, *P. serotina* and *R. pseudoacacia* is in flooring, furniture, and veneer while the softwood *P. radiata* mainly finds valuable applications as structural timber but also for furniture. Experience with silviculture aimed at high-quality wood applications is published for all example EITS (see Table 4). Sometimes, more publications on silviculture for quality wood are available from the area of origin (e.g., *E. globulus* and *P. serotina*) and sometimes from the area of introduction (e.g., *A. altissima*, *P. radiata* and *R. pseudoacacia*). Given the pioneer nature and rapid growth of these EITS, the emphasis in quality wood production is on striving for a large crown with low crown base and, in some cases, timely pruning.

Table 4 Wood production with EITS in complex forest for high-quality wood applications

Species	Experience in area of origin	Experience in introduction area
<i>Ailanthus altissima</i>	[223, 274, 308]	[180, 275–277, 309]
<i>Eucalyptus globulus</i>	[283, 284, 288, 289]	[310]
<i>Pinus radiata</i>	[294]	[293, 294, 311, 312]
<i>Prunus serotina</i>	[295–300, 313–320]	[244, 248–250, 257, 258, 321]
<i>Robinia pseudoacacia</i>	[322]	[54, 70, 261, 262, 302, 304, 306, 323]

Thanks to the wind dispersal of its seeds and its ability to germinate and grow under a range of site conditions, *Pinus radiata* is an efficient pioneer and, according to invasion ecology, an invasive tree species. However, due to its high requirements for light, its invasiveness is usually limited to short vegetation types with potential for forest development. Once established, *P. radiata* becomes like most pine species an efficient nurse tree for native trees, either spontaneous regeneration or enrichment planting.

TRAITS:

- Evergreen coniferous pioneer tree, 15–30 m high
- More climate change tolerant than most competitors
- Negative impact on nutrient availability
- Wood: mostly used for construction timber, pulp and paper.

NATIVE RANGE:

- Endemic to the Californian Monterey Peninsula and several islands off the coast of Baja California (Mexico).

INTRODUCTION PATHWAYS:


- Silviculture.

INVASION HABITAT:

- Pioneer species outside forests in open landscapes, heathlands, dry grassland and dunes.

STATUS:

- Plantation forests mainly along the California coast into southern coastal Oregon, New Zealand (where it is the most common non-native tree), Australia, Chile, SW Europe and South Africa.
- Occurs locally elsewhere in the temperate climatic zone.



Pinus radiata distribution map; colors indicate gradients of occurrence from alimited (yell ow) to intensive (red) presence (source: <https://www.gbif.org/species/5285727>).



PICTURE: *Pinus radiata* plantation stand, with underplanted seedlings of shade-intolerant (*Nothofagus dombeyi*, *Nothofagus obliqua*), semi-tolerant (*Nothofagus alpina*, *Laurelia sempervirens*) and shade tolerant native tree species (*Aextoxicon punctatum*; *Cryptocarya alba*). South-Central Chile (Photo Klaus Kremer)

Box. 3 *Pinus radiata*. Invasive conifer and nurse tree

Contribution of EITS to climate resilience

The effects of climate change on forests can be wide-ranging and complex [324–326]. Warmer temperatures can increase the frequency and intensity of fires, droughts, and insect outbreaks, while changes in precipitation can lead to flooding, soil erosion, and changes in soil moisture. These changes can cause trees to become stressed and more susceptible to disease, pests, and wildfire. Due to the complexity of the mutually interacting effects of climate change, it is still

largely unclear which tree species will face difficulties, leading to changes in tree species diversity, forest composition and vegetation structure, as well which will be favoured [60, 327, 328].

Climate change mainly impacts forests with weak moisture or nutrient supply, or with simplified vegetation structure and restricted tree species diversity and redundancy. These simplified forests often originated from historical and current forest use [134, 137, 329]. If EITS propagules are present, they will often fill the vacant space associated with

Table 5 EITS resilience to climate change: Climatic amplitude, drought, and heat tolerance relative to tree species present in introduction area

Species	Large climatic amplitude	Drought resilience	Heat resilience
<i>Ailanthus altissima</i>	[157, 281, 331]	[157, 281, 332–335]	[157, 332, 333, 335]
<i>Eucalyptus globulus</i>	[150, 154, 336–338]	[283, 338–342]	[336–338]
<i>Pinus radiata</i>	[234, 294, 343–345]	[294, 344, 346, 347]	[294, 344, 346, 347]
<i>Prunus serotina</i>	[244, 348–353]	[244, 335, 349–352, 354]	[244, 335, 349–352]
<i>Robinia pseudoacacia</i>	[54, 70, 322, 355–357]	[54, 159, 335, 356, 358–362]	[54, 262, 335]

the associated disturbances. Given the complexity of climate change effects—changing disease and pest pressures in addition to changing abiotic conditions – it is difficult to predict whether EITS can cope with climate change and contribute to forest climate resilience in a given region. In general, however, it can be stated that most EITS will suffer less from

climate change than most competing native tree species due to their often wider climate amplitude, and drought and heat tolerance. This also applies to our sample EITS, including *P. radiata* which has a very small area of origin, mainly on the Monterey Peninsula in California, but an extensive introduced area [330] (see Table 5).

Prunus serotina is a gap tree species capable of quickly restoring the forest climate after disturbances. Native tree species, with the exception of highly light-demanding pioneer tree species, regenerate without any problem under *P. serotina*, whose nurse tree character is enhanced by its fast-decomposing litter.

TRAITS:

- Deciduous broadleaf pioneer tree, 20 – 30 m high
- More climate resilient than most competitors
- Positive impact on nutrient availability
- High quality furniture wood

NATIVE RANGE:

- Eastern North America

INTRODUCTION PATHWAYS:


- Horticulture
- Silviculture

INVASION HABITAT:

- Pioneer species in forest margins and open forests, outside forests in open landscapes, heathlands, dry grasslands, and dunes.

STATUS:

- Widespread and common in its native range and on the North European sand belt, from northern France to Poland, where it was extensively planted in the past.
- Occurs locally elsewhere in the temperate climatic zone



Prunus serotina distribution map; colors indicate gradients of occurrence from limited (yellow) to intensive (red) presence (source <https://www.gbif.org/species/3021850>)



PICTURE: Native forest succession (*Acer pseudoplatanus*, *Acer platanoides*, *Castanea sativa*, *Fagus sylvatica*, *Quercus robur*, *Carpinus betulus*, *Sorbus aucuparia*, *Rhamnus frangula*, *Ilex aquifolium*, *Taxus baccata*) in *Prunus serotina* forest. Waasmunster, Belgium (Photo Bart Nyssen)

Box. 4 *Prunus serotina*. From herbicide-treated invasive to accelerator of forest succession

Table 7 EITS and forest climate recovery

Species	Fast canopy recovery	Subcanopy and shrub layer
<i>Ailanthus altissima</i>	[121, 157, 281, 428–430]	[121, 281, 428–432]
<i>Eucalyptus globulus</i>	*	*
<i>Pinus radiata</i>	[73, 76, 236, 343]	**
<i>Prunus serotina</i>	[156, 205, 256, 257, 433–442]	[94, 156, 165, 243, 401, 403, 436–448]
<i>Robinia pseudoacacia</i> ***	[54, 70, 124, 150, 163, 264, 335, 449–455]	[54, 70, 150, 163, 456, 457]

* A contribution of *E. globulus* to forest recovery or the formation of a subcanopy or shrub layer is not mentioned. Probably this tree species is limited in this by its high light requirement and low invasive nature

** For *P. radiata*, only a contribution to forest recovery is mentioned, not the formation of a subcanopy or shrub layer of *P. radiata*. Probably this tree species is limited in this by its high light requirement

*** *R. pseudoacacia* also has a high light requirement and a low invasive character. Its ability to form root sprouts means that this EITS does contribute locally to forest recovery and the formation of a subcanopy or shrub layer, although this is limited to open forests such as pine and oak forests

Table 8 Nurse tree effect and regeneration of late successional species under EITS

EITS species	Nurse tree effect and regeneration of native tree species
<i>Ailanthus altissima</i>	[55, 221, 222]
<i>Eucalyptus globulus</i>	[95, 224–230, 462]
<i>Pinus radiata</i>	[96, 97, 231–237, 462]
<i>Prunus serotina</i>	[94, 164, 165, 243–245, 258]
<i>Robinia pseudoacacia</i>	[70, 124, 259, 260]

Especially in forest restoration, nurse tree effects are important to initiate forest succession [78, 96, 97, 227, 231, 458–461]. EITS may contribute to the rapid build-up of a forest canopy, followed by the below-canopy establishment of late successional species (see Table 8).

Conclusions

EITS integrated into forest ecosystems can contribute to climate change adaptation.

Because of the pioneer character of most EITS, spontaneous forest succession limits their dominance in time and space in most situations. EITS may be temporarily and locally dominant in pioneer situations. However, establishment of mid- and late-successional tree species under their translucent canopy should gradually replace them, over several decades, as long as propagules of these species are present [268, 462].

This integration of EITS in the forest ecosystem can be enhanced by measures aiming at strengthening resilience to climate change. In fact, if climate-adaptive forest management is implemented consistently—optimising abiotic conditions, enhancing structural diversity of the canopy, and

increasing tree species diversity—the (dominant) occurrence of EITS in these forests may be reduced and eventually EITS may become regular trees among the other tree species. This may contribute to forest functional diversity and functional redundancy, which helps to secure the continuous provisioning of ecosystem services. For example, *P. radiata* may be able to fulfil functions of native coniferous tree species that are affected by severe pests or disturbances. Likewise, *R. pseudoacacia*, *P. serotina*, *A. altissima* and *E. globulus* can act as alternative fast-growing pioneer deciduous trees if certain native species were to decline.

However, this is only true in regions where climate adaptation is served by a structurally rich forest with high functional diversity and functional redundancy. In (semi)-arid regions where forest resilience to climate change is enhanced by giving trees a large growth space in an open forest [463], the proportion of pioneer tree species, and thus EITS, is likely to increase.

Increased resilience through enhanced tree species functional diversity and functional redundancy will not only reduce dominance by EITS but also reduces the likelihood of any new invasive non-native tree and shrub species to establish in the forest. Reducing forest invasibility is most effective for tree species with high light requirements and light transmission. Reducing forest invasibility is most effective for the majority of EITS (e.g., *P. serotina*, *R. pseudoacacia*, *A. altissima*, *E. globulus* or *P. radiata*), which are light-demanding pioneers. Reducing abundance by forest succession is less effective for the exceptional EITS combining pioneer traits—e.g., early seed setting, efficient propagule spreading, and rapid juvenile growth—with late successional species traits—e.g., shade-tolerant, shade-casting, and long-lived. The higher the shade-tolerance or the shade-casting capacity of EITS, the more difficult their integration into the forest ecosystem without them becoming dominant tree species. Members of the genus *Acer*, and more specifically *Acer platanoides*, are in this category [190, 191].

In resilient forest ecosystems EITS may contribute to further climate change adaptation due to their often higher drought and heat tolerance, and their pioneer character, translating into rapid restoration of a forest microclimate after a stand-replacing disturbance, and a capacity to fulfil the role of nurse tree for establishment of native late-successional species [461]. EITS may further enhance multiple elements of forest structure: abiotic conditions, vegetation structure, and species composition.

The extent and manner in which EITS contribute to climate change adaptation vary depending on the tree species. Nevertheless, EITS are often approached as a homogeneous group in studies of their effects on forest ecosystems [30, 362, 464]. Since the differences between EITS are similar to the differences among native pioneer

tree species, information important for evaluating such effects is then insufficiently considered. Given the significant variations among EITS, it is essential to impartially assess their specific contributions to climate change adaptation, biodiversity, and ecosystem services, without bias towards their origin. There is a great need for further forest research on the autecology, synecology, contribution to associated biodiversity, ecosystem processes, and ecosystem services of the different EITS in specific contexts, just as it has been done in the past for native tree species. EITS have become part of the regional tree species pools of many areas worldwide [465]. They therefore deserve intensified fundamental and applied research into their ecosystem functioning.

Of all example trees in this review, *Robinia pseudoacacia* is probably one of the most heat and drought tolerant temperate deciduous tree species and one of the least invasive. Planted widely over an estimated area of over 30,000 km², in North America, Europe, temperate Asia, temperate South America, northern and southern Africa, Australia, and New Zealand.

TRAITS:

- Deciduous broad-leaved pioneer tree, 20-25 m high
- More climate resilient than competitors
- Nitrogen fixer, positive impact on soil fertility
- Wood: biomass, poles, and quality timber for outdoor use

NATIVE RANGE:

- Eastern North America

PATHWAYS:


- Horticulture
- Erosion control on dikes, embankments
- Silviculture

INVASION HABITAT:

- Pioneer species in forest margins and open forests, outside forests in open landscape, heathlands, dry grassland, and dunes.

STATUS:

- Widespread and common in North America and Europe, but also in South America, South Africa, and China.
- Occurs locally elsewhere in the temperate climatic zone.



Robinia pseudoacacia distribution map; colors indicate gradients of occurrence from limited (yellow) to intensive (red) presence (source <https://www.gbif.org/species/5352251>).



PICTURE: Massive native forest regeneration (*Acer pseudoplatanus*, *Acer platanoides*, *Tilia cordata*, *Fagus sylvatica*, *Quercus robur*, *Carpinus betulus*, *Sorbus aucuparia*, *Rhamnus frangula*, *Ilex aquifolium*, *Taxus baccata*) in adult *Robinia pseudoacacia* forest. Waasmunster, Belgium (Photo Bart Nyssen)

Box. 5 *Robinia pseudoacacia*. A heat and drought tolerant pioneer widely cultivated for wood production

Integration of EITS is promising for forest managers

Preventing dominance of EITS and enhancing forest resilience under changing climatic conditions are mostly seen as separate tasks by forest managers. However, this review shows that these two processes are actually related. Forest management strategies can target increased ecosystem resilience to multiple disturbances [144]. Silvicultural measures aimed at increasing resilience to a changing climate, can simultaneously contribute to increasing resilience to dominance by EITS, especially if such measures restore complex forest structure and tree species diversity [1, 8]. Such an approach also underpins close-to-nature forest management [433].

The contribution that EITS can make to climate change adaptation in forests is promising for forest managers: the large budgets previously reserved for control can be better used to further increase resilience of forests to disturbances exacerbated by global changes. Also, by integrating EITS, forest managers have additional tree species at their disposal to achieve their management objectives.

Here it is essential to emphasise that the integration potential of EITS depends on physical site characteristics, the local forest composition and structure, and on the forest objectives pursued. Strengthening resilience to EITS dominance in conjunction with reinforced resilience to the effects of global warming is not suitable for forests with a natural open canopy structure or where an open canopy structure is aimed for (e.g., to achieve established nature conservation goals or to enhance drought adaptation). Nor is this approach suitable for mono-specific plantation forests, except when these should be changed into mixed structured forests. Resilience – absorption, in this case—towards EITS may be high in plantations of shade-casting tree species but the pursued even-aged monoculture has too little vegetation structure and diversity of tree species to absorb and adapt to climate change impacts.

Yet, notwithstanding all the differences in how the presence of distinct EITS works out in diverse forest ecosystems, society has much to gain from a non-biased approach to these ‘new’ tree species [466]. The sustainable delivery of forest ecosystem services under rapidly changing climatic conditions may partly depend on EITS.

Key References

- Davis MA, Chew MK, Hobbs RJ, Lugo AE, Ewel JJ, Vermeij GJ, et al. Don't judge species on their origins. *Nature*. 2011;474(7350):153-4. doi: <https://doi.org/10.1038/474153a>.

This 2011 comment in *Nature* by renowned ecologists is an early manifestation of a paradigm shift among biological invasion researchers. Replacing a biased anti-exotic stance with a neutral approach to these new species in both research design and implementation as a prerequisite for advancing the understanding how to deal with newcomers in ecosystems.

- Muys B, Messier C. Climate smart forest management caught between a rock and a hard place. *Annals of Forest Science*. 2023;80(43). doi: <https://doi.org/10.1186/s13595-023-01208-5>.

This letter notes the current trend in forest vitality loss that demonstrates the urgent need for forest adaptation. Authors note that measures are insufficiently adopted by foresters in the field and question the reasons for this inaction that cripples climate-smart forest management. A way forward is proposed, using a diversity-based no-regret approach consistent with available knowledge.

- Aquilué N, Messier C, Martins KT, Dumais-Lalonde V, Mina M. A simple-to-use management approach to boost adaptive capacity of forests to global uncertainty. *Forest Ecology and Management*. 2021;481:118692. doi: <https://doi.org/10.1016/j.foreco.2020.118692>.

Using the functional network approach, this paper provides forest practitioners with a well-researched yet easy-to-use tool to evaluate functional diversity, vulnerability and functional connectivity at the landscape level. This tool can be used to inform plans for improving ecosystem adaptability to changing environmental conditions and societal demands.

- Nikinmaa L, Lindner M, Cantarello E, Jump AS, Seidl R, Winkel G, et al. Reviewing the use of resilience concepts in forest sciences. *Current Forestry Reports*. 2020;6(2):61-80. doi: <https://doi.org/10.1007/s40725-020-00110-x>.

This review offers a systematic overview of the recent forest science literature on resilience - a key concept to deal with an uncertain future in forestry - , synthesizing how resilience is defined, assessed and operationalised.

- Brancalion PH, Amazonas NT, Chazdon RL, van Melis J, Rodrigues RR, Silva CC, et al. Exotic eucalypts: From demonized trees to allies of tropical forest restoration? *Journal of Applied Ecology*. 2020;57(1):55-66. doi: <https://doi.org/10.1111/1365-2664.13513>.

This article presents results of experimental studies on the effects of using non-native eucalyptus (*Eucalyptus*

spp.) as a transitional phase in tropical forest restoration on above-ground biomass accumulation, regeneration of native woody species and financial viability. Many of the negative effects attributed to eucalypts on the growth and natural regeneration of native trees depend on characteristics of the production system, landscape structure, soil and climate in which they are grown, rather than on the presence of eucalypts per se.

- Forbes A, Norton D. Transitioning Exotic Plantations to Native Forest: A Report on the State of Knowledge In: Ministry for Primary Industries NZ, editor. Wellington 2021.

This report, prepared for Te Uru Rākau - New Zealand Forestry Service - lays the foundation for national policy in dealing with EITS. It summarises current knowledge on how exotic forest - in New Zealand consisting mainly of radiata pine (*Pinus radiata* D. Don) - can best be converted to native forest.

- Nicolescu V-N, Rédei K, Mason WL, Vor T, Pöetzelsberger E, Bastien J-C, et al. Ecology, growth and management of black locust (*Robinia pseudoacacia* L.), a non-native species integrated into European forests. *Journal of Forestry Research*. 2020;31(4):1081-101. doi: <https://doi.org/10.1007/s11676-020-01116-8>.

This review provides a broad overview of the ecology, adaptability to climate change and contribution to ecosystem services of black locust (*Robinia pseudoacacia* L.), a tree species, due to its pioneer traits, regarded as invasive in large parts of the world where its range is expected to expand under predicted climate changes

- Schilthuizen M, Pimenta LPS, Lammers Y, Steenbergen PJ, Flohil M, Beveridge NG, et al. Incorporation of an invasive plant into a native insect herbivore food web. *PeerJ*. 2016;4:e1954. doi: <https://doi.org/10.7717/peerj.1954>.

This article presents a solid example of unbiased research on ecological incorporation of an EITS, black cherry (*Prunus serotina* Ehrh.) into native food webs. The authors conclude that evolutionary processes can lead to a specialised herbivorous community on an EITS, reducing invasiveness over time.

- Annighöfer P, Kawaletz H, Terwei A, Mölder I, Zerbe S, Ammer C. Managing an invasive tree species—silvicultural recommendations for black cherry (*Prunus*

serotina Ehrh). *Forstarchiv*. 2015;86(5):139-52. doi: <https://doi.org/10.4432/0300411286139>.

This paper analyses different management options to derive recommendations for the future management of black cherry (*Prunus serotina* Ehrh). The options were evaluated in terms of economic profitability and ecological compatibility. The results show that there are promising strategies to integrate these EITS into the forest ecosystem, which are effective in reducing its density and can result in positive income for landowners.

- Brandner R, Schickhofer G. Tree-of-Heaven (*Ailanthus altissima*): enormous and wide potential neglected by the western civilisation. *Proceedings of the 11th World Conference on Timber Engineering Riva del Garda, Italy2010*. p. 1-7.

This article is an early call for an objective approach to ecosystem services by EITS, in this case Tree of Heaven (*Ailanthus altissima* (Mill.) Swingle). Authors make recommendations regarding the usefulness of the wood and point out its wide application range for furniture and construction purposes, reflecting its huge and wide potential.

Acknowledgements We thank the anonymous reviewers for their valuable comments which helped us improve the earlier version of this paper. BN, LR, ES and BM are grateful for the knowledge and network available thanks to the EU Green Deal project SUPERB (www.forest-restoration.eu). BN and LR are grateful for the knowledge and network available thanks to the European Union's LIFE Nature and Biodiversity project RESILIAS (www.resilias.eu/en).

Author Contributions BN and BM conceived the original idea of the research, BN wrote the first draft of the manuscript with support from BM and JdO. All authors made substantial contributions and critically reviewed and approved the final version for submission.

Funding Bart Nyssen, Jan den Ouden, Anja Bindewald, Pedro Brancalion, Klaus Kremer, Katharina Lapin, Lisa Raats, Elisabeth Schatzdorfer, John Stanturf, Kris Verheyen and Bart Muys received no funding for their work on this article.

Data Availability No datasets were generated or analysed during the current study.

Declarations

Conflict of Interest The authors declare no competing interests.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Mina M, Messier C, Duveneck MJ, Fortin MJ, Aquilué N. Managing for the unexpected: building resilient forest landscapes to cope with global change. *Glob Change Biol* 2022. <https://doi.org/10.1111/gcb.16197>.
- Trumbore S, Brando P, Hartmann H. Forest health and global change. *Science*. 2015;349(6250):814–8. <https://doi.org/10.1126/science.aac6759>.
- Elkin C, Gutiérrez AG, Leuzinger S, Manusch C, Temperli C, Rasche L, et al. A 2° C warmer world is not safe for ecosystem services in the European Alps. *Glob Change Biol*. 2013;19(6):1827–40. <https://doi.org/10.1111/gcb.12156>.
- Mina M, Bugmann H, Cordonnier T, Irauschek F, Klopčič M, Pardos M, et al. Future ecosystem services from European mountain forests under climate change. *J Appl Ecol*. 2017;54(2):389–401. <https://doi.org/10.1111/1365-2664.12772>.
- Seidl R, Thom D, Kautz M, Martin-Benito D, Peltoniemi M, Vacchiano G, et al. Forest disturbances under climate change. *Nat Clim Chang*. 2017;7(6):395–402. <https://doi.org/10.1038/nclimate3303>.
- Boulanger Y, Taylor AR, Price DT, Cyr D, McGarrigle E, Rammer W, et al. Climate change impacts on forest landscapes along the Canadian southern boreal forest transition zone. *Landscape Ecol*. 2017;32(7):1415–31. <https://doi.org/10.1007/s10980-016-0421-7>.
- Flory SL, Dillon W, Hiatt D. Interacting global change drivers suppress a foundation tree species. *Ecol Lett*. 2022;25(4):971–80. <https://doi.org/10.1111/ele.13974>.
- Aquilué N, Messier C, Martins KT, Dumais-Lalonde V, Mina M. A simple-to-use management approach to boost adaptive capacity of forests to global uncertainty. *For Ecol Manage*. 2021;481:118692. <https://doi.org/10.1016/j.foreco.2020.118692>.
- Beck PS, Goetz SJ, Mack MC, Alexander HD, Jin Y, Randerston JT, et al. The impacts and implications of an intensifying fire regime on Alaskan boreal forest composition and albedo. *Glob Change Biol*. 2011;17(9):2853–66. <https://doi.org/10.1111/j.1365-2486.2011.02412.x>.
- Millar CI, Stephenson NL. Temperate forest health in an era of emerging megadisturbance. *Science*. 2015;349(6250):823–6. <https://doi.org/10.1126/science.aaa9933>.
- Thom D, Seidl R. Natural disturbance impacts on ecosystem services and biodiversity in temperate and boreal forests. *Biol Rev*. 2016;91(3):760–81. <https://doi.org/10.1111/brv.12193>.
- Musil C. Effect of invasive Australian acacias on the regeneration, growth and nutrient chemistry of South African lowland fynbos. *J Appl Ecol*. 1993;361–72. <https://doi.org/10.2307/2404637>.
- Evans RD, Rimer R, Sperry L, Belnap J. Exotic plant invasion alters nitrogen dynamics in an arid grassland. *Ecol Appl*. 2001;11(5):1301–10. [https://doi.org/10.1890/1051-0761\(2001\)011\[1301:EPIAND\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2001)011[1301:EPIAND]2.0.CO;2).
- Mack MC, D'Antonio CM, Ley RE. Alteration of ecosystem nitrogen dynamics by exotic plants: a case study of C4 grasses in Hawaii. *Ecol Appl*. 2001;11(5):1323–35. [https://doi.org/10.1890/1051-0761\(2001\)011\[1301:EPIAND\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2001)011[1301:EPIAND]2.0.CO;2).
- Forrest Meekins J, McCarthy BC. Effect of environmental variation on the invasive success of a nonindigenous forest herb. *Ecol Appl*. 2001;11(5):1336–48. [https://doi.org/10.1890/1051-0761\(2001\)011\[1323:AOENDB\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2001)011[1323:AOENDB]2.0.CO;2).
- Scott NA, Saggarr S, McIntosh PD. Biogeochemical impact of hieracium invasion in New Zealand's Grazed tussock grasslands: sustainability implications. *Ecol Appl* 2001:1311–22. [https://doi.org/10.1890/1051-0761\(2001\)011\[1311:BIOHII\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2001)011[1311:BIOHII]2.0.CO;2).
- Ehrenfeld JG, Ravit B, Elgersma K. Feedback in the plant-soil system. *Annu Rev Environ Resour*. 2005;30:75–115. <https://doi.org/10.1146/annurev.energy.30.050504.144212>.
- Grant T, Paschke MW, Monaco T, Sheley R. Invasive plant impacts on soil properties nutrient cycling and microbial communities. In: Monaco TA, Sheley RL, editors. *Invasive plant ecology and management, linking processes to practice*. Cambridge: CAB International; 2012. p. 79–104.
- Jo I, Fridley JD, Frank DA. Invasive plants accelerate nitrogen cycling: evidence from experimental woody monocultures. *J Ecol*. 2017;105(4):1105–10. <https://doi.org/10.1111/1365-2745.12732>.
- Fang W, Wang X. A field experimental study on the impact of *Acer platanoides*, an urban tree invader, on forest ecosystem processes in North America. *Ecol Process*. 2020;9(1):1–15. <https://doi.org/10.1186/s13717-020-0213-5>.
- Blackburn TM, Bellard C, Ricciardi A. Alien versus native species as drivers of recent extinctions. *Front Ecol Environ*. 2019;17(4):203–7. <https://doi.org/10.1002/fee.2020>.
- Pyšek P, Jarošík V, Hulme PE, Pergl J, Hejda M, Schaffner U, et al. A global assessment of invasive plant impacts on resident species, communities and ecosystems: the interaction of impact measures, invading species' traits and environment. *Glob Change Biol*. 2012;18(5):1725–37. <https://doi.org/10.1111/j.1365-2486.2011.02636.x>.
- Simberloff D, Martin J-L, Genovesi P, Maris V, Wardle DA, Aronson J, et al. Impacts of biological invasions: what's what and the way forward. *Trends Ecol Evol*. 2013;28(1):58–66.
- Vilà M, Hulme PE. *Impact of Biological Invasions on Ecosystem Services*. 2017.
- Richardson DM. Forestry trees as invasive aliens. *Conserv Biol*. 1998;12(1):18–26. <https://doi.org/10.1111/j.1523-1739.1998.96392.x>.
- Andrés C, Ojeda F. Effects of afforestation with pines on woody plant diversity of Mediterranean heathlands in southern Spain. *Biodivers Conserv*. 2002;11:1511–20. <https://doi.org/10.1023/A:1016850708890>.
- Bond WJ. Ancient grasslands at risk. *Science*. 2016;351(6269):120–2. <https://doi.org/10.1126/science.aad5132>.
- Richardson DM, Rejmánek M. Trees and shrubs as invasive alien species—a global review. *Divers Distrib*. 2011;17(5):788–809. <https://doi.org/10.1111/j.1472-4642.2011.00782.x>.
- Núñez MA, Davis KT, Dimarco RD, Peltzer DA, Paritsis J, Maxwell BD, et al. Should tree invasions be used in treeless ecosystems to mitigate climate change? *Front Ecol Environ*. 2021;19(6):334–41. <https://doi.org/10.1002/fee.2346>.
- Dodet M, Collet C. When should exotic forest plantation tree species be considered as an invasive threat and how should we treat them? *Biol Invasions*. 2012;14(9):1765–78. <https://doi.org/10.1007/s10530-012-0202-4>.
- Richardson DM, Pyšek P, Rejmánek M, Barbour MG, Dane Panetta F, West CJ. Naturalization and invasion of alien plants: Concepts and definitions. *Divers Distrib*. 2000;6:93–107. <https://doi.org/10.1046/j.1472-4642.2000.00083.x>.
- Castro-Díez P, Vaz AS, Silva JS, Van Loo M, Alonso Á, Aponte C, et al. Global effects of non-native tree species on multiple ecosystem services. *Biol Rev*. 2019;94(4):1477–501.
- Brundu G, Pauchard A, Pyšek P, Pergl J, Bindewald A, Brunori A, et al. Global guidelines for the sustainable use of non-native trees to prevent tree invasions and mitigate their negative impacts. *NeoBiota*. 2020;61:65–116. <https://doi.org/10.3897/neobiota.65.58380>.
- Seidl R, Turner MG. Post-disturbance reorganization of forest ecosystems in a changing world. *Proc Natl Acad Sci USA*. 2022;119(28):e2202190119. <https://doi.org/10.1073/pnas.2202190119>.

35. Wohlgemuth T, Gossner MM, Campagnaro T, Marchante H, van Loo M, Vacchiano G, et al. Impact of non-native tree species in Europe on soil properties and biodiversity: a review. *NeoBiota*. 2022;78:45–69. <https://doi.org/10.3897/neobiota.78.87022>.
36. Ayanu Y, Jentsch A, Müller-Mahn D, Rettberg S, Romankiewicz C, Koellner T. Ecosystem engineer unleashed: *Prosopis juliflora* threatening ecosystem services? *Reg Environ Change*. 2015;15(1):155–67. <https://doi.org/10.1007/s10113-014-0616-x>.
37. Medina-Villar S, Rodríguez-Echeverría S, Lorenzo P, Alonso A, Pérez-Corona E, Castro-Díez P. Impacts of the alien trees *Ailanthus altissima* (Mill.) Swingle and *Robinia pseudoacacia* L. on soil nutrients and microbial communities. *Soil Biol Biochem*. 2016;96:65–73. <https://doi.org/10.1016/j.soilbio.2016.01.015>.
38. Gentili R, Ferrè C, Cardarelli E, Montagnani C, Bogliani G, Citterio S, et al. Comparing negative impacts of *Prunus serotina*, *Quercus rubra* and *Robinia pseudoacacia* on native forest ecosystems. *Forests*. 2019;10(10):842. <https://doi.org/10.3390/f10100842>.
39. Gaertner M, Biggs R, Te Beest M, Hui C, Molofsky J, Richardson DM. Invasive plants as drivers of regime shifts: identifying high-priority invaders that alter feedback relationships. *Divers Distrib*. 2014;20(7):733–44. <https://doi.org/10.1111/ddi.12182>.
40. Souza-Alonso P, Rodríguez J, González L, Lorenzo P. Here to stay. Recent advances and perspectives about *Acacia* invasion in Mediterranean areas. *Ann For Sci*. 2017;74(3):1–20. <https://doi.org/10.1007/s13595-017-0651-0>.
41. Zavaleta E, Hulvey K. Realistic variation in species composition affects grassland production, resource use and invasion resistance. *Plant Ecol*. 2007;188(1):39–51. <https://doi.org/10.1007/s11258-006-9146-z>.
42. Matos FA, Edwards DPS, Magnago LF, Heringer G, Viananeri A, Buttschardt T, et al. Invasive alien acacias rapidly stock carbon, but threaten biodiversity recovery in young second-growth forests. *Philos Transact Royal Soc B*. 1867;2023(378):20210072. <https://doi.org/10.1098/rstb.2021.0072>.
43. Paritsis J, Landesmann JB, Kitzberger T, Tiribelli F, Sasal Y, Quintero C, et al. Pine plantations and invasion alter fuel structure and potential fire behavior in a Patagonian forest-steppe ecotone. *Forests*. 2018;9(3):117. <https://doi.org/10.3390/f9030117>.
44. Davis KT, Callaway RM, Fajardo A, Pauchard A, Nuñez MA, Brooker RW, et al. Severity of impacts of an introduced species corresponds with regional eco-evolutionary experience. *Ecography*. 2019;42(1):12–22. <https://doi.org/10.1111/ecog.04014>.
45. Campagnaro T, Brundu G, Sitzia T. Five major invasive alien tree species in European Union forest habitat types of the Alpine and Continental biogeographical regions. *J Nat Conserv*. 2018;43:227–38. <https://doi.org/10.1016/j.jnc.2017.07.007>.
46. Bindewald A, Brundu G, Schueler S, Starfinger U, Bauhus J, Lapin K. Site-specific risk assessment enables trade-off analysis of non-native tree species in European forests. *Ecol Evol*. 2021. <https://doi.org/10.1002/ece3.8407>.
47. Rodewald AD. Spreading messages about invasives. *Divers Distrib*. 2012;18(1):97–9. <https://doi.org/10.1111/j.1472-4642.2011.00817.x>.
48. Davis MA, Chew MK, Hobbs RJ, Lugo AE, Ewel JJ, Vermeij GJ, et al. Don't judge species on their origins. *Nature*. 2011;474(7350):153–4. <https://doi.org/10.1038/474153a>.
49. Chew MK. Ecologists, environmentalists, experts, and the invasion of the 'second greatest threat.' *Int Rev Environ Hist*. 2015;1:7–40.
50. Lugo AE, Carlo T, Wunderle J Jr. Natural mixing of species: novel plant–animal communities on Caribbean Islands. *Anim Conserv*. 2012;15(3):233–41. <https://doi.org/10.1111/j.1469-1795.2012.00523.x>.
51. Lugo AE, Smith JE, Potter KM, Vega HM, Kurtz CM. The contribution of nonnative tree species to the structure and composition of forests in the conterminous United States in comparison with tropical islands in the Pacific and Caribbean. *Gen Tech Rep Int Inst Trop For*. 2022;54:1–81. <https://doi.org/10.2737/IITF-GTR-54>.
52. Schlaepfer MA. Do non-native species contribute to biodiversity? *PLoS Biol*. 2018;16(4):e2005568. <https://doi.org/10.1371/journal.pbio.2005568>.
53. Desie E, Vancampenhout K, Nyssen B, van den Berg L, Weijters M, van Duinen G-J, et al. Litter quality and the law of the most limiting: Opportunities for restoring nutrient cycles in acidified forest soils. *Sci Total Environ*. 2020;699:134383. <https://doi.org/10.1016/j.scitotenv.2019.134383>.
54. Nicolescu V-N, Rédei K, Mason WL, Vor T, Pöetzelsberger E, Bastien J-C, et al. Ecology, growth and management of black locust (*Robinia pseudoacacia* L.), a non-native species integrated into European forests. *J For Res*. 2020;31(4):1081–101. <https://doi.org/10.1007/s11676-020-01116-8>.
55. Gómez-Aparicio L, Canham CD. Neighbourhood analyses of the allelopathic effects of the invasive tree *Ailanthus altissima* in temperate forests. *J Ecol*. 2008;96(3):447–58. <https://doi.org/10.1111/j.1365-2745.2007.01352.x>.
56. Eviner VT, Garbach K, Baty JH, Hoskinson SA. Measuring the effects of invasive plants on ecosystem services: challenges and prospects. *Invasive Plant Sci Manag*. 2012;5(1):125–36. <https://doi.org/10.1614/IPSM-D-11-00095.1>.
57. Rigling A, Gessler A, Feichtinger L, Queloz V, Wohlgemuth T. Introduced or native tree species to maintain forest ecosystem services in a hotter and drier future. In: Krumm F, Vítková L, editors. *Introduced tree species in European forests: opportunities and challenges*. 2016. p. 236–46.
58. Milanović M, Knapp S, Pyšek P, Kühn I. Linking traits of invasive plants with ecosystem services and disservices. *Ecosyst Serv*. 2020;42:101072. <https://doi.org/10.1016/j.ecoser.2020.101072>.
59. Arnet M, Santos B, Brockerhoff EG, Pelsler PB, Ecroyd C, Clemens J. Importance of arboreta for ex situ conservation of threatened trees. *Biodivers Conserv*. 2015;24:3601–20. <https://doi.org/10.1007/s10531-015-1024-3>.
60. IPCC. *Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems*. In: Change. IPoC, editor.: Intergovernmental Panel on Climate Change. 2019.
61. Allen CD, Breshears DD, McDowell NG. On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere*. 2015;6(8):1–55. <https://doi.org/10.1890/ES15-00203.1>.
62. Hartmann H, Moura CF, Anderegg WR, Ruehr NK, Salmon Y, Allen CD, et al. Research frontiers for improving our understanding of drought-induced tree and forest mortality. *New Phytol*. 2018;218(1):15–28. <https://doi.org/10.1111/nph.15048>.
63. Ramsfield T, Bentz B, Faccoli M, Jactel H, Brockerhoff E. Forest health in a changing world: effects of globalization and climate change on forest insect and pathogen impacts. *Forestry*. 2016;89(3):245–52. <https://doi.org/10.1093/forestry/cpw018>.
64. Allen CD, Breshears DD. Drought-induced shift of a forest–woodland ecotone: rapid landscape response to climate variation. *Proc Natl Acad Sci*. 1998;95(25):14839–42. <https://doi.org/10.1073/pnas.95.25.14839>.
65. Esquivel-Muelbert A, Baker TR, Dexter KG, Lewis SL, Brien RJ, Feldpausch TR, et al. Compositional response of Amazon forests to climate change. *Glob Change Biol*. 2019;25(1):39–56. <https://doi.org/10.1111/gcb.14413>.

66. Bauhus J, Pokorny B, van der Meer PJ, Kanowski PJ, Kanninen M. Ecosystem goods and services—the key for sustainable plantations. In: Bauhus J, van der Meer PJ, Kanninen M, editors. *Ecosystem goods and services from plantation forests*. London: Routledge; 2010. p. 221–43.
67. Brockerhoff EG, Jactel H, Parrotta JA, Ferraz SF. Role of eucalypt and other planted forests in biodiversity conservation and the provision of biodiversity-related ecosystem services. *For Ecol Manage.* 2013;301:43–50. <https://doi.org/10.1016/j.foreco.2012.09.018>.
68. Mori AS, Lertzman KP, Gustafsson L. Biodiversity and ecosystem services in forest ecosystems: a research agenda for applied forest ecology. *J Appl Ecol.* 2017;54(1):12–27. <https://doi.org/10.1111/1365-2664.12669>.
69. Tassin J, Kull CA. Facing the broader dimensions of biological invasions. *Land Use Policy.* 2015;42:165–9. <https://doi.org/10.1016/j.landusepol.2014.07.014>.
70. Vítková M, Müllerová J, Sádlo J, Pergl J, Pyšek P. Black locust (*Robinia pseudoacacia*) beloved and despised: A story of an invasive tree in Central Europe. *For Ecol Manage.* 2017;384:287–302. <https://doi.org/10.1016/j.foreco.2016.10.057>.
71. Davis MA, Grime JP, Thompson K. Fluctuating resources in plant communities: a general theory of invasibility. *J Ecol.* 2000;88(3):528–34. <https://doi.org/10.1046/j.1365-2745.2000.00473.x>.
72. Schelhas J, Greenberg R. *Forest patches in tropical landscapes*. Washington, D.C.: Island Press; 1996.
73. Richardson D, Brown P. Invasion of mesic mountain fynbos by *Pinus radiata*. *S Afr J Bot.* 1986;52(6):529–36. [https://doi.org/10.1016/S0254-6299\(16\)31486-7](https://doi.org/10.1016/S0254-6299(16)31486-7).
74. Viana VM, Tabanez A, Batista JLF. Dynamics and restoration of forest fragments in the Brazilian Atlantic moist forest. In: *Tropical forest remnants: ecology, management, and conservation of fragmented communities*. Chicago: University of Chicago Press; 1997.
75. Buckley YM, Bolker BM, Rees M. Disturbance, invasion and re-invasion: managing the weed-shaped hole in disturbed ecosystems. *Ecol Lett.* 2007;10(9):809–17. <https://doi.org/10.1111/j.1461-0248.2007.01067.x>.
76. Gómez P, Murúa M, San Martín J, Goncalves E, Bustamante RO. Maintaining close canopy cover prevents the invasion of *Pinus radiata*: Basic ecology to manage native forest invasibility. *PLoS ONE.* 2019;14(5):e0210849. <https://doi.org/10.1371/journal.pone.0210849>.
77. Gómez P, Espinoza S, Cuadros N, Goncalves E, Bustamante R. Light availability influences the invasion of *Teline monspessulana* (L.) K. Koch in a temperate fragmented forest in Central Chile. *iForest-Biogeosci For.* 2022;15(5):411.
78. Lugo AE. The outcome of alien tree invasions in Puerto Rico. *Front Ecol Environ.* 2004;2(5):265–73. [https://doi.org/10.1890/1540-9295\(2004\)002\[0265:TOOATI\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2004)002[0265:TOOATI]2.0.CO;2).
79. Hobbs RJ, Higgs ES, Hall C. *Novel ecosystems: intervening in the new ecological world order*. John Wiley & Sons; 2013.
80. Evers CR, Wardropper CB, Branoff B, Granek EF, Hirsch SL, Link TE, et al. The ecosystem services and biodiversity of novel ecosystems: A literature review. *Glob Ecol Conserv.* 2018;13:e00362. <https://doi.org/10.1016/j.gecco.2017.e00362>.
81. Hobbs RJ, Higgs E, Harris JA. Novel ecosystems: implications for conservation and restoration. *Trends Ecol Evol.* 2009;24(11):599–605. <https://doi.org/10.1016/j.tree.2009.05.012>.
82. Williams JW, Jackson ST. Novel climates, no-analog communities, and ecological surprises. *Front Ecol Environ.* 2007;5(9):475–82. <https://doi.org/10.1890/070037>.
83. Radeloff VC, Williams JW, Bateman BL, Burke KD, Carter SK, Childress ES, et al. The rise of novelty in ecosystems. *Ecol Appl.* 2015;25(8):2051–68. <https://doi.org/10.1890/14-1781.1>.
84. Gómez-Aparicio L, Canham CD. Neighborhood models of the effects of invasive tree species on ecosystem processes. *Ecol Monogr.* 2008;78(1):69–86. <https://doi.org/10.1890/06-2036.1>.
85. Dyderski MK, Jagodziński AM. Impacts of invasive trees on alpha and beta diversity of temperate forest understories. *Biol Invasions.* 2021:1–18. <https://doi.org/10.1007/s10530-020-02367-6>.
86. Bempah AN, Kyereh B, Ansong M, Asante W. Debarking as a control method for invasive tree species management in tropical forests. *Trop Ecol.* 2022;63(2):200–8. <https://doi.org/10.1007/s42965-021-00198-7>.
87. Pötzelberger E, Spiecker H, Neophytou C, Mohren F, Gazda A, Hasenauer H. Growing non-native trees in European forests brings benefits and opportunities but also has its risks and limits. *Curr For Rep.* 2020:1–15. <https://doi.org/10.1007/s40725-020-00129-0>.
88. Sáenz-Romero C, O'Neill G, Aitken SN, Lindig-Cisneros R. Assisted migration field tests in Canada and Mexico: Lessons, limitations, and challenges. *Forests.* 2020;12(1):9. <https://doi.org/10.3390/f12010009>.
89. Bindewald A, Michiels H-G, Bauhus J. Risk is in the eye of the assessor: comparing risk assessments of four non-native tree species in Germany. *For Int J For Res.* 2019. <https://doi.org/10.1093/forestry/cpz052>.
90. Marinšek A, Bindewald A, Kraxner F, La Porta N, Meisel P, Lapin K. *Management of non-native tree species in forests of Alpine space*. Vienna: BFW; 2022.
91. Pawson SM, Brin A, Brockerhoff EG, Lamb D, Payn TW, Paquette A, et al. *Plantation forests, climate change and biodiversity*. *Biodivers Conserv.* 2013;22:1203–27. <https://doi.org/10.1007/s10531-013-0458-8>.
92. Vor T, Bolte A, Spellmann H, Ammer C. *Potenziale und Risiken eingeführter Baumarten*. Göttinger Forstwissenschaften. Göttingen: Universitätsverlag Göttingen; 2015.
93. Wunder J, Knüsel S, Dorren L, Schwarz M, Bourrier F, Conedera M. *Götterbaum und Paulownie: die «neuen Wilden» im Schweizer Wald? Schweizerische Zeitschrift für Forstwesen.* 2018;169(2):69–76. <https://doi.org/10.3188/szf.2018.0069>.
94. Nysen B, den Ouden J, Verheyen K, Vanhellemont M. Integrating black cherry in forest management in the Netherlands and Belgium. In: Krumm F, Vítková L, editors. *Introduced tree species in European forests: opportunities and challenges*. European Forest Institute; 2016. p. 362–72.
95. Brancalion PH, Amazonas NT, Chazdon RL, van Melis J, Rodrigues RR, Silva CC, et al. Exotic eucalypts: From demonized trees to allies of tropical forest restoration? *J Appl Ecol.* 2020;57(1):55–66. <https://doi.org/10.1111/1365-2664.13513>.
96. Forbes AS, Norton DA, Carswell FE. Opportunities and limitations of exotic *Pinus radiata* as a facilitative nurse for New Zealand indigenous forest restoration. *New Zealand journal of forestry science.* 2019;49. <https://doi.org/10.33494/nzjfs492019x45x>.
97. Brockerhoff EG, Ecroyd CE, Leckie AC, Kimberley MO. Diversity and succession of adventive and indigenous vascular understorey plants in *Pinus radiata* plantation forests in New Zealand. *For Ecol Manage.* 2003;185(3):307–26. [https://doi.org/10.1016/S0378-1127\(03\)00227-5](https://doi.org/10.1016/S0378-1127(03)00227-5).
98. Carpenter S, Walker B, Anderies JM, Abel N. From metaphor to measurement: resilience of what to what? *Ecosystems.* 2001;4(8):765–81. <https://doi.org/10.1007/s10021-001-0045-9>.
99. Nikinmaa L, Lindner M, Cantarello E, Jump AS, Seidl R, Winkel G, et al. Reviewing the use of resilience concepts in forest sciences. *Current Forestry Reports.* 2020;6(2):61–80. <https://doi.org/10.1007/s40725-020-00110-x>.
100. Folke C, Carpenter SR, Walker B, Scheffer M, Chapin T, Rockström J. Resilience thinking: Integrating resilience, adaptability and transformability. *Ecol Soc.* 2010;15(4):20.

101. Gladstone-Gallagher RV, Pilditch CA, Stephenson F, Thrush SF. Linking traits across ecological scales determines functional resilience. *Trends Ecol Evol.* 2019;34(12):1080–91. <https://doi.org/10.1016/j.tree.2019.07.010>.
102. Seidl R, Spies TA, Peterson DL, Stephens SL, Hicke JA. Searching for resilience: addressing the impacts of changing disturbance regimes on forest ecosystem services. *J Appl Ecol.* 2016;53(1):120–9. <https://doi.org/10.1111/1365-2664.12511>.
103. Brockerhoff EG, Barbaro L, Castagneyrol B, Forrester DL, Gardiner B, González-Olabarria JR, et al. Forest biodiversity, ecosystem functioning and the provision of ecosystem services. *Biodivers Conserv.* 2017;26(13):3005–35.
104. Haubrock PJ, Turbelin AJ, Cuthbert RN, Novoa A, Taylor NG, Angulo E, et al. Economic costs of invasive alien species across Europe. *NeoBiota.* 2021;67:153–90. <https://doi.org/10.3897/neobiota.67.58196>.
105. Fantle-Lepczyk JE, Haubrock PJ, Kramer AM, Cuthbert RN, Turbelin AJ, Crystal-Ornelas R, et al. Economic costs of biological invasions in the United States. *Sci Total Environ.* 2022;806:151318. <https://doi.org/10.1016/j.scitotenv.2021.151318>.
106. Diagne C, Catford JA, Essl F, Nuñez MA, Courchamp F. What are the economic costs of biological invasions? A complex topic requiring international and interdisciplinary expertise. *NeoBiota.* 2021. <https://doi.org/10.3897/neobiota.63.55260>.
107. Soto I, Balzani P, Carneiro L, Cuthbert RN, Macêdo R, Serhan Tarkan A, et al. Taming the terminological tempest in invasion science. *Biol Rev.* 2024;99:1357–90. <https://doi.org/10.1111/brv.13071>.
108. Occhipinti-Ambrogi A, Galil BS. A uniform terminology on bioinvasions: a chimera or an operative tool? *Mar Pollut Bull.* 2004;49(9–10):688–94. <https://doi.org/10.1016/j.marpolbul.2004.08.011>.
109. Pyšek P, Richardson DM, Rejmánek M, Webster GL, Williamson M, Kirschner J. Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon.* 2004;53(1):131–43. <https://doi.org/10.2307/4135498>.
110. Richardson DM, Pyšek P, Carlton JT. A compendium of essential concepts and terminology in invasion ecology. In: Richardson DM, editor. *Fifty years of invasion ecology: the legacy of Charles Elton.* Hoboken: John Wiley & Sons; 2011. p. 409–20.
111. Valéry L, Fritz H, Lefeuvre J-C, Simberloff D. In search of a real definition of the biological invasion phenomenon itself. *Biol Invasions.* 2008;10:1345–51. <https://doi.org/10.1007/s10530-007-9209-7>.
112. Branquart E, Fried G. *Espèces envahissantes d'ici et d'ailleurs: Synthèse sur les espèces envahissantes et présentation de 32 espèces.* Paris: Éditions du Gerfaud; 2016.
113. Thompson K, Hodgson JG, Rich TC. Native and alien invasive plants: more of the same? *Ecography.* 1995;18(4):390–402.
114. Ledgard N. The spread of lodgepole pine (*Pinus contorta*, Dougl.) in New Zealand. *For Ecol Manag.* 2001;141(1–2):43–57. [https://doi.org/10.1016/S0378-1127\(00\)00488-6](https://doi.org/10.1016/S0378-1127(00)00488-6).
115. Niinemets Ü, Valladares F. Tolerance to shade, drought, and waterlogging of temperate Northern Hemisphere trees and shrubs. *Ecol Monogr.* 2006;76(4):521–47. [https://doi.org/10.1890/0012-9615\(2006\)076\[0521:TTSDAW\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2006)076[0521:TTSDAW]2.0.CO;2).
116. Jagodziński AM, Dyderski MK, Horodecki P, Knight KS, Rawlik K, Szymt J. Light and propagule pressure affect invasion intensity of *Prunus serotina* in a 14-tree species forest common garden experiment. *NeoBiota.* 2019;46:1. <https://doi.org/10.3897/neobiota.46.30413>.
117. Forte TGW, Brundu G, Celesti-Grappo L, Siniscalco C, Barni E. *Prunus serotina* in Italy a challenging candidate for the national list of priority invasive alien species. *Plant Biosyst.* 2019;153(6):900–4. <https://doi.org/10.1080/11263504.2019.1666173>.
118. Vanhellefont M. Present and future population dynamics of *Prunus serotina* in forests in its introduced range. Gent: Universiteit Gent; 2009.
119. Nemes VEH. Examination of the spread and control possibilities of black cherry (*Prunus serotina* Ehrh.). Sopron: University of Sopron; 2022.
120. Fotiadis G, Kyriazopoulos A, Fraggakis I. The behaviour of *Ailanthus altissima* weed and its effects on natural ecosystems. *J Environ Biol.* 2011;32(6):801.
121. Sladonja B, Sušek M, Guillermic J. Review on invasive tree of heaven (*Ailanthus altissima* (Mill.) Swingle) conflicting values: assessment of its ecosystem services and potential biological threat. *Environ Manag.* 2015;56(4):1009–34. <https://doi.org/10.1007/s00267-015-0546-5>.
122. Iverson LR, Rebbeck J, Peters MP, Hutchinson T, Fox T. Predicting *Ailanthus altissima* presence across a managed forest landscape in southeast Ohio. *For Ecosyst.* 2019;6(1):1–13. <https://doi.org/10.1186/s40663-019-0198-7>.
123. Knüsel S, Conedera M, Bugmann H, Wunder J. Low litter cover, high light availability and rock cover favour the establishment of *Ailanthus altissima* in forests in southern Switzerland. *NeoBiota.* 2019;46:91–116. <https://doi.org/10.3929/ethz-b-000349496>.
124. Meyer-Münzer B, Grotehusmann H, Vor T, Robinie (Robinia pseudoacacia L.). In: Vor T, Spellmann H, Bolte A, Ammer C, editors. *Potenziale und Risiken eingeführter Baumarten, Baumartenportraits mit naturschutzfachlicher Bewertung.* Göttingen: Universitätsverlag Göttingen; 2015.
125. Fernandes P, Máguas C, Correia O, González-Moreno P. What drives *Eucalyptus globulus* natural establishment outside plantations? The relative importance of climate, plantation and site characteristics. *Biol Invasions.* 2018;20(5):1129–46. <https://doi.org/10.1007/s10530-017-1614-y>.
126. Silva JS, Vaz P, Moreira F, Cattry F, Rego FC. Wildfires as a major driver of landscape dynamics in three fire-prone areas of Portugal. *Landsc Urban Plan.* 2011;101(4):349–58. <https://doi.org/10.1016/j.landurbplan.2011.03.001>.
127. Cattry F, Moreira F, Deus E, Silva J, Águas A. Assessing the extent and the environmental drivers of *Eucalyptus globulus* wildling establishment in Portugal: results from a countrywide survey. *Biol Invasions.* 2015;17(11):3163–81. <https://doi.org/10.1007/s10530-015-0943-y>.
128. Yost JM, Wise SL, Love NL, Steane DA, Jones RC, Ritter MK, et al. Origins, diversity and naturalization of *Eucalyptus globulus* (Myrtaceae) in California. *Forests.* 2021;12(8):1129. <https://doi.org/10.3390/f12081129>.
129. Lorentz KA, Minogue PJ. Potential invasiveness for *Eucalyptus* species in Florida. *Invasive Plant Sci Manag.* 2015;8(1):90–7. <https://doi.org/10.1614/IPSM-D-14-00030.1>.
130. El GL. *aucalipto es una especie exótica pero no tiene carácter invasor.* *Foresta.* 2018;72:28–31.
131. Muller da Silva PH, Bouillet J-P, de Paula RC. Assessing the invasive potential of commercial *Eucalyptus* species in Brazil Germination and early establishment. *For Ecol Manag.* 2016;374:129–35. <https://doi.org/10.1016/j.foreco.2016.05.007>.
132. Thompson I, Mackey B, McNulty S, Mosseler A. Forest resilience, biodiversity, and climate change. A synthesis of the biodiversity/resilience/stability relationship in forest ecosystems. *CBD Technical Series.* Montreal: Secretariat of the Convention on Biological Diversity; 2009.
133. Thonicke K, Bahn M, Lavorel S, Bardgett RD, Erb K, Giamberini M, et al. Advancing the understanding of adaptive capacity of social-ecological systems to absorb climate extremes. *Earth's Future.* 2020;8(2):e2019EF001221. <https://doi.org/10.1029/2019EF001221>.

134. Falk DA, van Mantgem PJ, Keeley JE, Gregg RM, Guiterman CH, Tepley AJ, et al. Mechanisms of forest resilience. *For Ecol Manage.* 2022;512:120129. <https://doi.org/10.1016/j.foreco.2022.120129>.
135. Lindner M, Nikinmaa L, Brang P, Cantarello E, Seidl R. Enhancing resilience to address challenges in forest management. In: Krumm F, Schuck A, Rigling A, editors. *How to balance forestry and biodiversity conservation – A view across Europe*. Bonn: European Forest Institute (EFI); 2020.
136. Nikinmaa L, Lindner M, Cantarello E, Jump AS, Seidl R, Winkel G, et al. Reviewing the use of resilience concepts in forest sciences. *Curr For Rep.* 2020;6(2):61–80.
137. Messier C, Bauhus J, Sousa-Silva R, Auge H, Baeten L, Barsoum N, et al. For the sake of resilience and multifunctionality, let's diversify planted forests! *Conserv Lett.* 2022;15(1):e12829. <https://doi.org/10.1111/conl.12829>.
138. Sasaki T, Furukawa T, Iwasaki Y, Seto M, Mori AS. Perspectives for ecosystem management based on ecosystem resilience and ecological thresholds against multiple and stochastic disturbances. *Ecol Ind.* 2015;57:395–408. <https://doi.org/10.1016/j.ecolind.2015.05.019>.
139. Fletcher M-S, Wood SW, Haberle SG. A fire-driven shift from forest to non-forest: evidence for alternative stable states? *Ecology.* 2014;95(9):2504–13. <https://doi.org/10.1890/12-1766.1>.
140. Guiterman CH, Gregg RM, Marshall LA, Beckmann JJ, van Mantgem PJ, Falk DA, et al. Vegetation type conversion in the US Southwest: frontline observations and management responses. *Fire Ecology.* 2022;18(1):1–16. <https://doi.org/10.1186/s42408-022-00131-w>.
141. Miller AD, Thompson JR, Tepley AJ, Anderson-Teixeira KJ. Alternative stable equilibria and critical thresholds created by fire regimes and plant responses in a fire-prone community. *Ecography.* 2019;42(1):55–66. <https://doi.org/10.1111/ecog.03491>.
142. FAO. *Global Forest Resources Assessment 2015. Terms and definition R*. In: FAO, editor. *Forest resources Assessment Working Paper 180*. Rome: FAO; 2015.
143. Walker B, Salt D. *Resilience thinking: sustaining ecosystems and people in a changing world*. Washington: Island Press; 2012.
144. Messier C, Bauhus J, Doyon F, Maure F, Sousa-Silva R, Nolet P, et al. The functional complex network approach to foster forest resilience to global changes. *For Ecosyst.* 2019;6(1):1–16. <https://doi.org/10.1186/s40663-019-0166-2>.
145. Jactel H, Bauhus J, Boberg J, Bonal D, Castagneyrol B, Gardiner B, et al. Tree diversity drives forest stand resistance to natural disturbances. *Curr For Rep.* 2017;3:223–43. <https://doi.org/10.1007/s40725-017-0064-1>.
146. Hazarika R, Lapin K, Bindewald A, Vaz AS, Marinšek A, La Porta N, et al. Balancing Risks and Benefits: Stakeholder Perspective on Managing Non-Native Tree Species in the European Alpine Space. *Mitig Adapt Strateg Glob Chang.* 2024;29(6):55. <https://doi.org/10.1007/s11027-024-10152-2>.
147. Tassin J. *Arbres évadés. Du marronnage au compagnonnage*. Jardins de France. 2019;653:55–7.
148. Goodenough AE. Are the ecological impacts of alien species misrepresented? A review of the “native good, alien bad” philosophy. *Community Ecol.* 2010;11(1):13–21. <https://doi.org/10.1556/comec.11.2010.1.3>.
149. Turner J, Lambert MJ. Soil properties as affected by *Pinus radiata* plantations. *NZ J Forest Sci.* 1988;18(1):77–91.
150. Castro-Díez P, Alonso Á, Saldaña-López A, Granda E. Effects of widespread non-native trees on regulating ecosystem services. *Sci Total Environ.* 2021;146141. <https://doi.org/10.1016/j.scitotenv.2021.146141>.
151. Briones M, Ineson P. Decomposition of eucalyptus leaves in litter mixtures. *Soil Biol Biochem.* 1996;28(10–11):1381–8. [https://doi.org/10.1016/S0038-0717\(96\)00158-7](https://doi.org/10.1016/S0038-0717(96)00158-7).
152. Zerga B, Warkineh B, Teketay D, Woldetsadik M. The sustainability of reforesting landscapes with exotic species: a case study of eucalypts in Ethiopia. *Sustain Earth.* 2021;4(1):1–11. <https://doi.org/10.1186/s42055-021-00044-7>.
153. Tomé M, Almeida MH, Barreiro S, Branco MR, Deus E, Pinto G, et al. Opportunities and challenges of Eucalyptus plantations in Europe: The Iberian Peninsula experience. *Eur J Forest Res.* 2021;140(3):489–510. <https://doi.org/10.1007/s10342-021-01358-z>.
154. Silva JS, Tomé M. Tasmanian blue gum in Portugal – Opportunities and risks of a widely cultivated species. In: Krumm F, Lucie V, editors. *Introduced tree species in European forests: opportunities and challenges*. European Forest Institute, 2016.
155. Haas J, Sass Klaassen U, Akhmetzyanov L, Den Ouden J. Does input of rich litter facilitate tree growth? Growth and climate growth response of oak (*Quercus robur*) growing in the neighborhood of black cherry (*Prunus serotina*). In: Group FEaM, editor. *Report Wageningen University*. Wageningen: Wageningen University and Research, 2020.
156. Muys B, Maddelein D, Lust N. Ecology practice and policy of black cherry (*Prunus serotina* Ehrh.) management in Belgium. *Silva Gandavensis.* 1992;57:28–45.
157. Kowarik I, Säumlé I. Biological flora of central Europe: *Ailanthus altissima* (Mill.) swingle. *Perspect Plant Ecol Evol Syst.* 2007;8(4):207–37. <https://doi.org/10.1016/j.ppees.2007.03.002>.
158. Heisey RM, Kish HT. Herbicidal effects under field conditions of *Ailanthus altissima* bark extract, which contains ailanthone. *Plant Soil.* 2003;256(1):85–99.
159. Boer E. *Risk assessment Robinia pseudoacacia L.* Leiden: Naturalis Biodiversity Center; 2013.
160. Babu RC, Kandasamy O. Allelopathic effect of *Eucalyptus globulus* Labill. on *Cyperus rotundus* L. and *Cynodon dactylon* L. *Pers. J Agron Crop Sci.* 1997;179(2):123–6. <https://doi.org/10.1111/j.1439-037X.1997.tb00507.x>.
161. Castro-Díez P, Fierro-Brunnenmeister N, González-Muñoz N, Gallardo A. Effects of exotic and native tree leaf litter on soil properties of two contrasting sites in the Iberian Peninsula. *Plant Soil.* 2012;350(1):179–91. <https://doi.org/10.1007/s11104-011-0893-9>.
162. Bączek P, Halarewicz A. Effect of black cherry (*Prunus serotina*) litter extracts on germination and growth of Scots pine (*Pinus sylvestris*) seedlings. *Pol J Ecol.* 2019;67(2):137–47. <https://doi.org/10.3161/15052249PJE2019.67.2.004>.
163. Vítková M, Kolbek J. Vegetation classification and synecology of Bohemian *Robinia pseudoacacia* stands in a Central European context. *Phytocoenologia.* 2010;40(2/3):205–41. <https://doi.org/10.1127/0340-269X/2010/0040-0425>.
164. De Vanhessche D. *potentie van Amerikaanse vogelkers als overgangsfase naar veerkrachtig bos*. Faculteit Bio-Ingenieurswetenschappen. Leuven: KULeuven; 2020.
165. Hamm T. *Möglichkeiten und Grenzen einer Integration sukzessionaler Prozesse bei der Renaturierung naturferner und durch die Spätblühende Traubenkirsche (*Prunus serotina* Ehrh.) beeinflusster Kiefernforste mit dem Blick auf die Minimierung der Eingriffsintensität*. Fakultät Umweltwissenschaften Tharandt: Technische Universität Dresden, 2024
166. Xiong S, Nilsson C. The effects of plant litter on vegetation: a meta-analysis. *J Ecol.* 1999;87(6):984–94.
167. Wohlgemuth T, Moser B, Pötzelsberger E, Rigling A, Gossner MM. Über die Invasivität der Douglasie und ihre Auswirkungen auf Boden und Biodiversität. *Schweizerische Zeitschrift Fur Forstwesen.* 2021;172(2):118–27. <https://doi.org/10.3188/szf.2021.0118>.
168. Hermy M, Honnay O, Firbank L, Grashof-Bokdam C, Lawesson JE. An ecological comparison between ancient and other forest plant species of Europe, and the implications for forest

- conservation. *Biol Cons.* 1999;91(1):9–22. [https://doi.org/10.1016/S0006-3207\(99\)00045-2](https://doi.org/10.1016/S0006-3207(99)00045-2).
169. Schilthuizen M, Pimenta LPS, Lammers Y, Steenbergen PJ, Flohil M, Beveridge NG, et al. Incorporation of an invasive plant into a native insect herbivore food web. *PeerJ.* 2016;4:e1954. <https://doi.org/10.7717/peerj.1954>.
 170. Meijer K, Smit C, Beukeboom L, Schilthuizen M. Native insects on non-native plants in The Netherlands: curiosities or common practice? *Entomologische Berichten.* 2012;72(6):288–93.
 171. Bouget C, Brin A, Larrieu L. The use of sentinel logs to assess host shifts in early beetle colonisers of deadwood under climate- and forestry-induced tree species substitutions. *Insect Conserv Divers.* 2021;14(1):117–31. <https://doi.org/10.1111/icad.12434>.
 172. Goßner MM, Chao A, Bailey RI, Prinzing A. Native fauna on exotic trees: phylogenetic conservatism and geographic contingency in two lineages of phytophages on two lineages of trees. *Am Nat.* 2009;173(5):599–614. <https://doi.org/10.1086/597603>.
 173. Goßner M. Diversität und Struktur arborikoler Arthropoden-zönosen fremdländischer und einheimischer Baumarten. München: Technische Universität München; 2004.
 174. Branco M, Brockerhoff EG, Castagneyrol B, Orazio C, Jactel H. Host range expansion of native insects to exotic trees increases with area of introduction and the presence of congeneric native trees. *J Appl Ecol.* 2015;52(1):69–77.
 175. Silva JS, Moreira F, Vaz P, Catry F, Godinho-Ferreira P. Assessing the relative fire proneness of different forest types in Portugal. *Plant Biosyst.* 2009;143(3):597–608. <https://doi.org/10.1080/11263500903233250>.
 176. Goodrick SL, Stanturf JA. Evaluating potential changes in fire risk from Eucalyptus plantings in the Southern United States. *Int J Forestry Res.* 2012; 2012. <https://doi.org/10.1155/2012/680246>.
 177. Borrmann K. Einbürgerung, Ausbreitung und Vorkommen der Späten Traubenkirsche (*Padus serotina* Borkh.) in der Oberförsterei Lütkenhagen (Kreis Neustrelitz). *Botanischer Rundbrief für den Bezirk Neubrandenburg.* 1987;19:13–8.
 178. Nysen B, Schmidt UE, Muys B, van der Lei PB, Pyttel P. The history of introduced tree species in Europe in a nutshell. In: Krumm F, Vítková L, editors. *Introduced tree species in European forests: opportunities and challenges.* European Forest Institute, 2016. p. 44 - 55.
 179. Kowarik I, Böcker R. Zur Verbreitung, Vergesellschaftung und Einbürgerung des Götterbaumes (*Ailanthus altissima* [Mill.] Swingle) in Mitteleuropa. *Tuexenia.* 1984;4:9–29.
 180. Neophytou C, Pötzelsberger E, Curto M, Meimberg H, Hasenauer H. Population bottlenecks have shaped the genetic variation of *Ailanthus altissima* (Mill.) Swingle in an area of early introduction. *Forestry.* 2020;93(4):495–504. <https://doi.org/10.1093/forestry/cpz019>.
 181. Cozzo D. Comportamiento inicial de *Ailanthus altissima* en una plantación experimental. *Rev Forest Argentina.* 1972;16:47–52.
 182. Krüssmann G. *Handbuch der Laubgehölze*: Bd. 1: A-D. Singhofen: Paul Parey, 1976.
 183. Pyšek P, Křivánek M, Jarošík V. Planting intensity, residence time, and species traits determine invasion success of alien woody species. *Ecology.* 2009;90(10):2734–44. <https://doi.org/10.1890/08-0857.1>.
 184. Pyšek P, Manceur AM, Alba C, McGregor KF, Pergl J, Štajerová K, et al. Naturalization of central European plants in North America: species traits, habitats, propagule pressure, residence time. *Ecology.* 2015;96(3):762–74. <https://doi.org/10.1890/14-1005.1.sm>.
 185. Lonsdale WM. Global patterns of plant invasions and the concept of invasibility. *Ecology.* 1999;80(5):1522–36. [https://doi.org/10.1890/0012-9658\(1999\)080\[1522:GPOPIA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1522:GPOPIA]2.0.CO;2).
 186. Rejmánek M. Species richness and resistance to invasions. In: Oriens G, Editor. *Biodiversity and ecosystem processes in tropical forests.* Berlin: Springer; 1996. p. 153–72.
 187. Pyšek P, Richardson DM. Traits associated with invasiveness in alien plants: where do we stand? In: Nentwig W, editor. *Biological invasions.* Springer; 2008. p. 97–125.
 188. Catford JA, Daehler CC, Murphy HT, Sheppard AW, Hardesty BD, Westcott DA, et al. The intermediate disturbance hypothesis and plant invasions: Implications for species richness and management. *Perspect Plant Ecol Evol Syst.* 2012;14(3):231–41. <https://doi.org/10.1016/j.ppees.2011.12.002>.
 189. Rejmánek M, Richardson DM, Pyšek P. Plant invasions and invasibility of plant communities. *Vegetation Ecol.* 2013:387–424. <https://doi.org/10.1002/9781118452592.ch13>.
 190. Webb SL, Dwyer M, Kaunzinger CK, Wyckoff PH. The myth of the resilient forest: case study of the invasive Norway maple (*Acer platanoides*). *Rhodora.* 2000;102(911):332–54.
 191. Martin PH, Marks PL. Intact forests provide only weak resistance to a shade-tolerant invasive Norway maple (*Acer platanoides* L.). *J Ecol.* 2006;94(6):1070–9. <https://doi.org/10.1111/j.1365-2745.2006.01159.x>.
 192. Martin PH, Canham CD, Marks PL. Why forests appear resistant to exotic plant invasions: intentional introductions, stand dynamics, and the role of shade tolerance. *Front Ecol Environ.* 2008;7(3):142–9. <https://doi.org/10.1890/070096>.
 193. Rejmánek M. Invasibility of plant communities In: Drake JA, Mooney HJ, Di Castri F, Groves RH, Kruger FJ, Rejmánek M and Williamson M (eds) *Biological Invasions: a Global Perspective.* John Wiley, Chichester, 1989.
 194. Dechoum M, Castellani T, Zalba S, Rejmánek M, Peroni N, Tamashiro J. Community structure, succession and invasibility in a seasonal deciduous forest in southern Brazil. *Biol Invasions.* 2015;17(6):1697–712. <https://doi.org/10.1007/s10530-014-0827-6>.
 195. Meiners SJ, Pickett ST, Cadenasso ML. Exotic plant invasions over 40 years of old field successions: community patterns and associations. *Ecography.* 2002;25(2):215–23. <https://doi.org/10.1034/j.1600-0587.2002.250209.x>.
 196. Alpert P, Bone E, Holzapfel C. Invasiveness, invasibility and the role of environmental stress in the spread of non-native plants. *Perspect Plant Ecol Evol Syst.* 2000;3(1):52–66. <https://doi.org/10.1078/1433-8319-00004>.
 197. Pyšek P, Danihelka J, Sádlo J, Chrtěk J Jr, Chytrý M, Jarošík V, et al. Catalogue of alien plants of the Czech Republic: checklist update, taxonomic diversity and invasion patterns. *Preslia.* 2012;84(2):155–255.
 198. Pyšek P, Jarošík V, Kučera T. Patterns of invasion in temperate nature reserves. *Biol Conserv.* 2002;104(1):13–24. [https://doi.org/10.1016/S0006-3207\(01\)00150-1](https://doi.org/10.1016/S0006-3207(01)00150-1).
 199. Delavaux CS, Crowther TW, Zohner CM, Robmann NM, Lauber T, van den Hoogen J, et al. Native diversity buffers against severity of non-native tree invasions. *Nature.* 2023;1–9. <https://doi.org/10.1038/s41586-023-06440-7>.
 200. Huston MA. Management strategies for plant invasions: manipulating productivity, disturbance, and competition. *Divers Distrib.* 2004;10(3):167–78. <https://doi.org/10.1111/j.1366-9516.2004.00083.x>.
 201. Whitfield TJ, Lodge AG, Roth AM, Reich PB. Community phylogenetic diversity and abiotic site characteristics influence abundance of the invasive plant *Rhamnus cathartica* L. *J Plant Ecol.* 2014;7(2):202–9. <https://doi.org/10.1093/jpe/rtt020>.
 202. Goldstein LJ, Suding KN. Applying competition theory to invasion: resource impacts indicate invasion mechanisms in California shrublands. *Biol Invasions.* 2014;16(1):191–203. <https://doi.org/10.1007/s10530-013-0513-0>.

203. Symstad AJ. A test of the effects of functional group richness and composition on grassland invasibility. *Ecology*. 2000;81(1):99–109. [https://doi.org/10.1890/0012-9658\(2000\)081\[0099:ATOTEJ\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[0099:ATOTEJ]2.0.CO;2).
204. Dyderski MK, Jagodziński AM. Drivers of invasive tree and shrub natural regeneration in temperate forests. *Biol Invasions*. 2018;1–17. <https://doi.org/10.1007/s10530-018-1706-3>.
205. Nysse B, Muys B, Starfinger U, Conedera M. Spätblühende Traubenkirsche: Waldpest oder Waldbaum, je nach Waldbaukontext. *Schweizerische Zeitschrift für Forstwesen*. 2018;169(2):93–101. <https://doi.org/10.3188/szf.2018.0093>.
206. Koch J, Pearson DE, Huebner CD, Young MK, Sniezko RA. Restoration of landscapes and habitats affected by established invasive species. In: Poland TM, Patel-Weynand T, Finch DM, Ford Miniati C, Hayes DC, Lopez VM, editors. *Invasive species in forests and rangelands of the United States: a comprehensive science synthesis for the United States Forest Sector*. Heidelberg: Springer; 2021. p. 185–202.
207. Daniels MK, Larson ER. Effects of forest windstorm disturbance on invasive plants in protected areas of southern Illinois, USA. *J Ecol*. 2020;108(1):199–211. <https://doi.org/10.1111/1365-2745.13254>.
208. Lemoine RT, Svenning JC. Nativeness is not binary—a graduated terminology for native and non-native species in the Anthropocene. *Restor Ecol*. 2022;30(8):e13636. <https://doi.org/10.1111/rec.13636>.
209. Byun C, de Blois S, Brisson J. Plant functional group identity and diversity determine biotic resistance to invasion by an exotic grass. *J Ecol*. 2013;101(1):128–39. <https://doi.org/10.1111/1365-2745.12016>.
210. Tilman D, Knops J, Wedin D, Reich P, Ritchie M, Siemann E. The influence of functional diversity and composition on ecosystem processes. *Science*. 1997;277(5330):1300–2. <https://doi.org/10.1126/science.277.5330.1300>.
211. Pokorny ML, Sheley RL, Zabinski CA, Engel RE, Svejcar TJ, Borkowski JJ. Plant functional group diversity as a mechanism for invasion resistance. *Restor Ecol*. 2005;13(3):448–59. <https://doi.org/10.1111/j.1526-100X.2005.00056.x>.
212. Brown CS, Rice KJ. Effects of belowground resource use complementarity on invasion of constructed grassland plant communities. *Biol Invasions*. 2010;12(5):1319–34. <https://doi.org/10.1007/s10530-009-9549-6>.
213. Ammond SA, Litton CM. Competition between native Hawaiian plants and the invasive grass *Megathyrus maximus*: implications of functional diversity for ecological restoration. *Restor Ecol*. 2012;20(5):638–46. <https://doi.org/10.1111/j.1526-100X.2011.00806.x>.
214. Nunez-Mir GC, Liebhold AM, Guo Q, Brockerhoff EG, Jo I, Ordonez K, et al. Biotic resistance to exotic invasions: its role in forest ecosystems, confounding artifacts, and future directions. *Biol Invasions*. 2017;19(11):3287–99. <https://doi.org/10.1007/s10530-017-1413-5>.
215. Funk JL, Cleland EE, Suding KN, Zavaleta ES. Restoration through reassembly: plant traits and invasion resistance. *Trends Ecol Evol*. 2008;23(12):695–703. <https://doi.org/10.1016/j.tree.2008.07.013>.
216. Pollnac F, Seipel T, Repath C, Rew LJ. Plant invasion at landscape and local scales along roadways in the mountainous region of the Greater Yellowstone Ecosystem. *Biol Invasions*. 2012;14(8):1753–63. <https://doi.org/10.1007/s10530-012-0188-y>.
217. Relva MA, Nuñez MA. Factores que facilitan y retrasan la invasión de coníferas exóticas en la Isla Victoria. *Ecología Austral*. 2014;24(2):145–53. <https://doi.org/10.25260/EA.14.24.2.0.18>.
218. Riitters K, Potter K, Iannone BV III, Oswalt C, Fei S, Guo Q. Landscape correlates of forest plant invasions: a high-resolution analysis across the eastern United States. *Divers Distrib*. 2018;24(3):274–84. <https://doi.org/10.1111/ddi.12680>.
219. Godefroid S, Koedam N. Identifying indicator plant species of habitat quality and invasibility as a guide for peri-urban forest management. *Biodivers Conserv*. 2003;12(8):1699–713. <https://doi.org/10.1023/A:1023606300039>.
220. Bauhus J, Puettmann K, Kühne C. Close-to-nature forest management in Europe: does it support complexity and adaptability of forest ecosystems. In: Messier C, Puettmann KJ, Coates KD, editors. *Managing Forests as Complex Adaptive Systems*. London and New York: Routledge, The Earthscan Forest Library; 2013. p. 187–213.
221. Csiszár Á, Korda M, Schmidt D, Šporčić D, Süle P, Teleki B, et al. Allelopathic potential of some invasive plant species occurring in Hungary. *Allelopathy J*. 2013;31(2):309–18.
222. Csiszár Á, Korda M. Inváziós növényfajok visszaszorításával kapcsolatos kezelési kísérletek összefoglalása. Csiszár Á, Korda M (szerk) *Özönnövények visszaszorításának gyakorlati tapasztalatai Rosalia kézikönyvek*. 2015; 3:201–35.
223. Brandner R, Schickhofer G. Tree-of-Heaven (*Ailanthus altissima*): enormous and wide potential neglected by the western civilisation. *Proceedings of the 11th World Conference on Timber Engineering Riva del Garda, Italy 2010*. p. 1–7.
224. Woldu K. Natural Regeneration of Juniperous *procera* in E. globulus Plantation. *Skinnskatteberg: Swedish University of Agricultural Sciences*; 1998.
225. Amazonas NT, Forrester DI, Silva CC, Almeida DRA, Rodrigues RR, Brancalion PH. High diversity mixed plantations of Eucalyptus and native trees: An interface between production and restoration for the tropics. *For Ecol Manage*. 2018;417:247–56. <https://doi.org/10.1016/j.foreco.2018.03.015>.
226. Amazonas NT, Forrester DI, Oliveira RS, Brancalion PH. Combining Eucalyptus wood production with the recovery of native tree diversity in mixed plantings: Implications for water use and availability. *For Ecol Manage*. 2018;418:34–40. <https://doi.org/10.1016/j.foreco.2017.12.006>.
227. Senbeta F, Beck E, Lüttge U. Exotic trees as nurse-trees for the regeneration of natural tropical forests. *Trees*. 2002;16(4):245–9. <https://doi.org/10.1007/s00468-002-0161-y>.
228. Senbeta F, Teketay D, Näslund B-Å. Native woody species regeneration in exotic tree plantations at Munessa-Shashemene Forest, southern Ethiopia. *New Forest*. 2002;24(2):131–45. <https://doi.org/10.1023/A:1021201107373>.
229. Senbeta F, Teketay D. Regeneration of indigenous woody species under the canopies of tree plantations in Central Ethiopia. *Trop Ecol*. 2001;42(2):175–85.
230. César RG, Moreno VS, Coletta GD, Chazdon RL, Ferraz SF, de Almeida DR, et al. Early ecological outcomes of natural regeneration and tree plantations for restoring agricultural landscapes. *Ecol Appl*. 2018;28(2):373–84. <https://doi.org/10.1002/eap.1653>.
231. Kremer K, Promis A, Bauhus J. Natural advance regeneration of Native Tree Species in *Pinus radiata* plantations of South-Central Chile suggests potential for a passive restoration approach. *Ecosystems*. 2022;25(5):1096–116. <https://doi.org/10.1007/s10021-021-00704-x>.
232. Onaindia M, Ametzaga-Arregi I, San Sebastián M, Mitxelena A, Rodríguez-Loinaz G, Peña L, et al. Can understorey native woodland plant species regenerate under exotic pine plantations using natural succession? *For Ecol Manage*. 2013;308:136–44. <https://doi.org/10.1016/j.foreco.2013.07.046>.
233. Forbes AS, Allen RB, Herbert JW, Kohiti K, Shaw WB, Taurua L. Determining the balance between active and passive indigenous forest restoration after exotic conifer plantation clear-fell. *For Ecol*

- Manage. 2021;479:118621. <https://doi.org/10.1016/j.foreco.2020.118621>.
234. Forbes A, Norton D. Transitioning Exotic Plantations to Native Forest: A Report on the State of Knowledge In: Ministry for Primary Industries NZ, editor. Wellington 2021.
 235. Kremer Ramirez K. Restoration of native forests from *Pinus radiata* plantations in southern Chile. Faculty of Environment and Natural Resources. Freiburg: Universität Freiburg; 2021. p. 2021.
 236. Becerra PI, Montenegro G. The widely invasive tree *Pinus radiata* facilitates regeneration of native woody species in a semi-arid ecosystem. *Appl Veg Sci*. 2013;16(2):173–83. <https://doi.org/10.1111/j.1654-109X.2012.01221.x>.
 237. Ogden J, Braggins J, Stretton K, Anderson S. Plant species richness under *Pinus radiata* stands on the central North Island volcanic plateau, New Zealand. *New Zealand J Ecol*. 1997;21(1):17–29.
 238. Forbes AS, Wallace KJ, Buckley HL, Case BS, Clarkson BD, Norton DA. Restoring mature-phase forest tree species through enrichment planting in New Zealand's lowland landscapes. *New Zealand J Ecol*. 2020;44(1):1–9. <https://doi.org/10.20417/nzjecol.44.10>.
 239. Forbes AS, Norton DA, Carswell FE. Artificial canopy gaps accelerate restoration within an exotic *Pinus radiata* plantation. *Restor Ecol*. 2016;24(3):336–45. <https://doi.org/10.1111/rec.12313>.
 240. Forbes AS, Norton DA, Carswell FE. Tree fern competition reduces indigenous forest tree seedling growth within exotic *Pinus radiata* plantations. *For Ecol Manage*. 2016;359:1–10. <https://doi.org/10.1016/j.foreco.2015.09.036>.
 241. Forbes AS, Norton DA, Carswell FE. Underplanting degraded exotic *Pinus* with indigenous conifers assists forest restoration. *Ecol Manag Restor*. 2015;16(1):41–9. <https://doi.org/10.1111/emr.12137>.
 242. Norton DA, Forbes A. Can exotic pine trees assist in restoration? *Appl Veg Sci*. 2013;16(2):169–70. <https://doi.org/10.1111/avsc.12014>.
 243. Annighöfer P, Kawaletz H, Terwei A, Mölder I, Zerbe S, Ammer C. Managing an invasive tree species—silvicultural recommendations for black cherry (*Prunus serotina* Ehrh.). *Forstarchiv*. 2015;86(5):139–52. <https://doi.org/10.4432/0300411286139>.
 244. Petersen R, Annighofer P, Spellmann H, Leder B. Spätblühende Traubenkirsche (*Prunus serotina* Ehrh.). In: Vor T, Spellmann H, Bolte A, Ammer C, editors. Potenziale und Risiken eingeführter Baumarten Baumartenportraits mit naturschutzfachlicher Bewertung. Göttingen: Göttinger Forstwissenschaften; 2015.
 245. Hamm T, Huth F, Wagner S. Einfluss der Spätblühenden Traubenkirsche (*Prunus serotina* EHRH.) auf die Verjüngung heimischer Laubbaumarten in Kiefernforsten. Osnabrück: DBU-Stipendienprogramm; 2017. p. 40–6.
 246. Annighöfer P. From short-term activism to long-term pragmatism? Silvicultural recommendations for the regeneration and reforestation of native species under the influence of an invasive species (*Prunus serotina* Ehrh.). 1st International Conference on Reforestation Challenges, 3–6 June 2015, Faculty of Forestry, Belgrade-Serbia. Belgrade Serbia, 2015.
 247. Annighöfer P, Ammer C. Spätblühende Traubenkirsche Pragmatismus statt Aktionismus. *AFZ Der Wald*. 2015;70(4):27–30.
 248. Nicolescu V-N, Vor T. Silviculture and management of black cherry (*Prunus serotina* Ehrh.) in Europe. Unpublished. 2020.
 249. Haag C, Wilhelm U. Arbeiten mit unerwünschter Baumart oder Verschleppung einer Katastrophe. *AFZ*. 1998;53:276–9.
 250. Von Wendorff G. Die *Prunus serotina* in Mitteleuropa: Eine waldbauliche Monographie. Hamburg: Universität Hamburg; 1952. p. 250.
 251. Hamm T, Huth F, Wagner S. Kiefernforsten renaturieren trotz Spätblühender Traubenkirsche? *AFZ Der Wald*. 2015;70(4):22–5.
 252. Hamm T, Huth F, Wagner S. Spätblühende Traubenkirsche—ein Vorteil für das Keimbett? *AFZ Der Wald*. 2015;70(4):26.
 253. Nyssen B, Den Ouden J. Quelle stratégie sylvicole pour le cerisier tardif? Mise au point d'un arbre de décision pour une espèce invasive. *ForêtNature*. 2021;158:29–41.
 254. Nyssen B, Koopmans G, den Ouden J. Beslisboom Amerikaanse vogelkers : Bestrijden, uitfaseren, integreren en bos weerbaar maken. *Vakblad natuur bos landschap*. 2019;16(157):31–5.
 255. Nyssen B, Koopmans G, Den Ouden J: Beslisboom Amerikaanse vogelkers. 2019. https://bosgroepen.nl/wp-content/uploads/2019/09/Beslisboom_Amerikaanse_vogelkers_web.pdf. Accessed 3 Jun 2024.
 256. Le NB. cerisier tardif, nuisance ou espèce au potentiel intéressant ? *Silva Belgica*. 2018;125(2):18–27.
 257. Nyssen B, Den Ouden J, Verheyen K. Amerikaanse vogelkers, van Bospest tot Bosboom. KNNV Uitgeverij: Zeist; 2013.
 258. Engel M, Nyssen B, Desie E, den Ouden J, Raats L, Hagemann U. Managing Black Cherry (*Prunus serotina* Ehrh.) in European Forests: Insights from native and non-native ranges. *Forest Ecol Manag*. 2024;562:11. <https://doi.org/10.1016/j.foreco.2024.121959>.
 259. Kou M, Garcia-Fayos P, Hu S, Jiao J. The effect of *Robinia pseudoacacia* afforestation on soil and vegetation properties in the Loess Plateau (China): A chronosequence approach. *For Ecol Manage*. 2016;375:146–58. <https://doi.org/10.1016/j.foreco.2016.05.025>.
 260. Von Holle B, Joseph K, Largay EF, Lohnes RG. Facilitations between the introduced nitrogen-fixing tree, *Robinia pseudoacacia*, and nonnative plant species in the glacial outwash upland ecosystem of Cape Cod, MA. *Biodivers Conserv*. 2006;15(7):2197–215. <https://doi.org/10.1007/s10531-004-6906-8>.
 261. Nicolescu V-N, Hernea C, Bakti B, Keserű Z, Antal B, Rédei K. Black locust (*Robinia pseudoacacia* L.) as a multi-purpose tree species in Hungary and Romania: a review. *J Forest Res*. 2018;29(6):1449–63. <https://doi.org/10.1007/s11676-018-0626-5>.
 262. Rédei K, Csiha I, Keserű Z, Kamandiné Végh Á, Győri J. The silviculture of black locust (*Robinia pseudoacacia* L.) in Hungary: a review. *South-East Eur Forest*. 2012;2(2):101–7.
 263. Đodan M, Brus R, Eisdold A-M, Nicolescu V-N, ORŠANIĆ M, PRATAŠIENE K, et al. Non-native tree species in the viewpoint of climate change: chances and opportunities – Croatia as a case study. *Šumarski list*. 2018;7–8(CXLII (2018)):391–402. <https://doi.org/10.31298/sl.142.7-8.6>.
 264. Cierjacks A, Kowarik I, Joshi J, Hempel S, Ristow M, von der Lippe M, et al. Biological flora of the British Isles: *Robinia pseudoacacia*. *J Ecol*. 2013;101(6):1623–40. <https://doi.org/10.1111/1365-2745.12162>.
 265. Wohlgenuth T, Hafner J, Höltermann A, Moser B, Nehring S, Rigling A. Impact of Douglas-fir on forests and open land habitats. In: Spiecker H, Lindner M, Schuler J, editors. Douglas-fir - An option for Europe. European Forestry Institute; 2019. p. 57–62.
 266. Kriegel P, Matevski D, Schuldt A. Monoculture and mixture-planting of non-native Douglas fir alters species composition, but promotes the diversity of ground beetles in a temperate forest system. *Biodivers Conserv*. 2021;30(5):1479–99. <https://doi.org/10.1007/s10531-021-02155-1>.
 267. Oliver CD, Larson BC, Oliver C. Forest stand dynamics. New York: John Wiley; 1996.

268. Brockerhoff EG, Jactel H, Parrotta JA, Quine CP, Sayer J. Plantation forests and biodiversity: oxymoron or opportunity? *Biodivers Conserv*. 2008;17:925–51. <https://doi.org/10.1007/s10531-008-9380-x>.
269. Bremer LL, Farley KA. Does plantation forestry restore biodiversity or create green deserts? A synthesis of the effects of land-use transitions on plant species richness. *Biodivers Conserv*. 2010;19:3893–915. <https://doi.org/10.1007/s10531-010-9936-4>.
270. Lopes IT, Gussoni CO, Demarchi LO, De Almeida A, Pizo MA. Diversity of understory birds in old stands of native and Eucalyptus plantations. *Restor Ecol*. 2015;23(5):662–9. <https://doi.org/10.1111/rec.12216>.
271. Marsden SJ, Whiffin M, Galetti M. Bird diversity and abundance in forest fragments and Eucalyptus plantations around an Atlantic forest reserve. *Brazil Biodivers Conserv*. 2001;10:737–51. <https://doi.org/10.1023/A:1016669118956>.
272. Clout M, Gaze P. Effects of plantation forestry on birds in New Zealand. *J Appl Ecol*. 1984;795–815. <https://doi.org/10.2307/2405048>.
273. Forestry VDo. Control and Utilization of Tree-of-heaven: A Guide of Virginia Landowners. Virginia Department of Forestry, 2009.
274. Hu SY. *Ailanthus Arnoldia*. 1979;39(2):29–50.
275. Panayotov P, Kalmukov K, Panayotov M. Biological and wood properties of *Ailanthus altissima* (Mill.) Swingle. *Forestry ideas*. 2011;17(2):42.
276. Berki D. Eigenschaften und Verwendung des Holzes des Götterbaums (*Ailanthus altissima*). Institut für Holztechnologie und nachwachsende. Wien: Universität für Bodenkultur; 2014.
277. Frenes J. Beurteilung von Bekämpfungs- und Förderungsmaßnahmen des Götterbaums in Südtirol. Department für Wald- und Bodenwissenschaften. Wien: Universität für Bodenkultur; 2021.
278. Müller U, Brandner R, Schickhofer G. Der Einsatz des Götterbaums im Baubereich. *Forstwissenschaftliche Tagung2016*.
279. Brandner R, Schickhofer G. The mechanical potential of the Tree-of-Heaven (*Ailanthus altissima*). *Holztechnologie*. 2013;54(5):5–11.
280. Szabolcs K, Varga D. Physical and mechanical properties of wood from invasive tree species. *Maderas Ciencia y tecnología*. 2021;23. <https://doi.org/10.4067/s0718-221x2021000100411>.
281. Wunder J, Knüsel S, Gurtner D, Conedera M. The spread of tree of heaven in Switzerland. In: Krumm F, Vítková L, editors. *Introduced tree species in European forests: opportunities and challenges* Freiburg: European Forest Institute. European Forestry Institute; 2016. p. 374–85.
282. Müller U, Berki D, Freudenthaler M, Kaserer D, Kletzl X, Obernosterer D, et al. *Materialeigenschaften und Nutzungspotential des Götterbaums*. Tulln an der Donau: Universität für Bodenkultur Wien; 2015.
283. Taskhiri MS, Holloway D, Turner P. Non-destructive testing of knot-free timber in a Tasmanian eucalyptus nitens plot European Conference on Non-Destructive Testing. Sweden: European Conference on Non-Destructive Testing; 2018.
284. Washusen R, Harwood C, Morrow A, Northway R, Valencia JC, Volker P, et al. Pruned plantation-grown *Eucalyptus nitens*: Effect of thinning and conventional processing practices on sawn board quality and recovery. *NZ J Forest Sci*. 2009;39(1):39–55.
285. Serrada R, Montero G, Reque JA. *Compendio de selvicultura aplicada en España*. Madrid: INIA; 2008.
286. Nutto L, Touza Vázquez MC. Modelos de producción de madera sólida en plantaciones de *Eucalyptus globulus* de Galicia. *Boletín Informativo CIDEU*. 2006;2:37–50.
287. Sepiarsky F. Nuevos usos industriales de la madera de *Eucalyptus globulus*. *Boletín Informativo CIDEU*. 2006;2:93–102.
288. Wood M, Volker P, Syme M. *Eucalyptus plantations for sawlog production in Tasmania, Australia: optimising thinning regimes*. Proceedings 'Institute of Foresters of Australia and New Zealand Institute of Forestry Conference', 3rd–7th June 2007. Coffs Harbour, New South Wales, Australia: Citeseer; 2007. p. 489–505.
289. Yang J-L, Fife D, Waugh G, Downes G, Blackwell P. The effect of growth strain and other defects on the sawn timber quality of 10-year-old *Eucalyptus globulus* Labill. *Aust For*. 2002;65(1):31–7. <https://doi.org/10.1080/00049158.2002.10674850>.
290. Gavilán E, Alzamora RM, Apiolaza LA, Sáez K, Elissetche JP, Pinto A. Modelling the influence of radiata pine log variables on structural lumber production. *Maderas-Cienc Tecnol*. 2023;25:1–10. <https://doi.org/10.4067/s0718-221x202300100402>.
291. Alzamora RM, Apiolaza LA, Evison DC. Using a production approach to estimate economic weights for structural attributes of *Pinus radiata* wood. *Scand J For Res*. 2013;28(3):282–90. <https://doi.org/10.1080/02827581.2012.734328>.
292. Elzaki OT, Khider TO. Physical and Mechanical Properties of *Pinus radiata* from Jebel Marra Western Sudan. *J For Prod Indust*. 2013;2(3):53–7.
293. Perry C. Economic analysis of a target diameter harvesting system in radiata pine. 2013. <https://doi.org/10.26021/3068>.
294. Mead DJ. Sustainable management of *Pinus radiata* plantations. Food and Agriculture Organization of the United Nations (FAO), 2013.
295. Marquis DA. *Prunus serotina* Ehrh. Black Cherry. In: Burns RM, Honkala BH, editors. *Silvics of North America, Volume 2, Hardwoods*, 1990.
296. Nesom. Black cherry, *Prunus serotina* Ehrh. In: NRCS U, editor. *USDA NRCS Plant Guide*. USDA NRCS, 2008.
297. OMNR. A silvicultural guide for the tolerant hardwood forest in Ontario. In: Resour. OMN, editor. Ontario, 1998.
298. Perkey AW, Wilkins BL. *Crop tree field guide: selecting and managing crop trees in the central Appalachians*. Northeastern Area State and Private Forestry: USDA Forest Service; 2001.
299. Perkey AW, Wilkins BL, Smith HC. *Crop tree management in eastern hardwoods*. Northeastern Area State & Private Forestry: USDA Forest Service; 1994.
300. Hough AF. *Silvical characteristics of Black Cherry (Prunus serotina)*. Upper Darby: Forest Service USDA; 1960.
301. Bijak S, Lachowicz H. Impact of tree age and size on selected properties of black locust (*Robinia pseudoacacia* L.) wood. *Forests*. 2021;12(5):634. <https://doi.org/10.3390/f12050634>.
302. Nicolescu V-N, Buzatu-Goană C, Bartlett D, Iacob N. Regeneration and early tending of black locust (*Robinia pseudoacacia* L.) stands in the north-west of Romania. *South-East Eur Forest*. 2019;10(2):97–105. <https://doi.org/10.15177/see-for.19-14>.
303. Ciuvăț A, Blujdea V, Abrudan I, Nuță I, Negruțiu F. Ecosystem services provided by black locust (*Robinia pseudoacacia* L.) plantations in South-Western Romania. *Proceedings of the Biennial International Symposium Forest and sustainable development, 24-25th October 2014*. Brașov: Transilvania University Press; 2015. p. 151–6.
304. Keresztesi B. Black locust: the tree of agriculture. *Outlook Agric*. 1988;17(2):77–85. <https://doi.org/10.1177/003072708801700207>.
305. Savill PS. *The silviculture of trees used in British forestry*. Boston: CABI; 2019.
306. Sitzia T, Cierjacks A, De Rigo D, Caudullo G. *Robinia pseudoacacia* in Europe: distribution, habitat, usage and threats. *European atlas of forest tree species*. 2016. p. 166–7.
307. Pollet C, Verheyen C, Hebert J, Jourez B. Physical and mechanical properties of black locust (*Robinia pseudoacacia*) wood

- grown in Belgium. *Can J For Res.* 2012;42(5):831–40. <https://doi.org/10.1139/x2012-037>.
308. Zhang J. Coastal saline soil rehabilitation and utilization based on forestry approaches in China. Springer, 2014.
 309. Buffi R. Le specie forestali per la zona castanile insubrica: la crescita giovanile di specie forestali indigene ed esotiche nei rimboschimenti sperimentali di Copera. Eidgenössische Anstalt für das forstliche Versuchswesen, 1987.
 310. Nutto L, Vázquez MCT. Modelos de producción de madera sólida en plantaciones de *Eucalyptus globulus* de Galicia. *Boletín Informativo CIDEU.* 2006;2:37–50.
 311. Whyte A. Radiata pine silviculture in New Zealand: its evolution and future prospects. *Aust For.* 1988;51(3):185–96. <https://doi.org/10.1080/00049158.1988.10676041>.
 312. Woollons R, Manley B. Examining growth dynamics of *Pinus radiata* plantations at old ages in New Zealand. *Forestry.* 2012;85(1):79–86. <https://doi.org/10.1093/forestry/cpr059>.
 313. Marquis DA, Ernst RL, Stout SL. Prescribing silvicultural treatments in hardwood stands of the Alleghenies (revised). General technical report NE-96. 1992.
 314. Hough AF. Preliminary recommendations for the management of black cherry on the northern Allegheny Plateau. *J Forest.* 1953;51(3):184–8.
 315. Carmean WH, Hahn JT, Jacobs RD. Site index curves for forest tree species in the eastern United States. St. Paul: North Central Forest Experiment Station, Forest Service, US Department of Agriculture; 1989.
 316. Carmean WH, Station NCFE, Service USF. Site index curves for northern hardwoods in northern Wisconsin and Upper Michigan. North Central Forest Experiment Station, Forest Service, US Department of Agriculture, 1978.
 317. Smith HC, Miller GW, Lamson NI. Crop-Tree Release Thinning in 65-year-old commercial cherry-maple stands (5-year results). Research paper NE-694. Radnor: USDA Forest Service, Northeastern Forest Experiment station, 1994.
 318. Lamson NI, Smith HC. Stand development and yields of Appalachian hardwood stands managed with single-tree selection for at least 30 years. In: Station NFE, editor. Research Paper NE-655. Radnor, Pennsylvania, 1991.
 319. Lamson NI, Smith HC. Crop-tree release increases growth of 12-year-old yellow-poplar and black cherry. US Dept. of Agriculture, Forest Service, Northeastern Forest Experiment Station; 1989.
 320. Grisez TJ. Pruning black cherry in understocked stands. Forest Service, US Dept. of Agriculture, Northeastern Forest Experiment Station; 1978.
 321. Grundler A. Zu Wachstum von Krone und Schaft der spätblühende Traubenkirsche (*Prunus serotina*) im Käfertaler Wald, Mannheim. Rottenburg: Höhere forstliche Fachschule; 2011. p. 35.
 322. Huntley JC. *Robinia pseudoacacia* L. black locust. *Silvics North Am.* 1990;2:755–61.
 323. Keresztesi B. Breeding and cultivation of black locust, *Robinia pseudoacacia*. Hungary *For Ecol Manag.* 1983;6(3):217–44. [https://doi.org/10.1016/S0378-1127\(83\)80004-8](https://doi.org/10.1016/S0378-1127(83)80004-8).
 324. Anderegg WR, Trugman AT, Badgley G, Anderson CM, Bartuska A, Ciais P, et al. Climate-driven risks to the climate mitigation potential of forests. *Science.* 2020; 368(6497). <https://doi.org/10.1126/science.aaz7005>.
 325. Muys B, Messier C. Climate smart forest management caught between a rock and a hard place. *Ann For Sci* 2023; 80(43). <https://doi.org/10.1186/s13595-023-01208-5>.
 326. Carrasco G, Almeida A, Falvey M, Olmedo G, Taylor P, Santibañez F, et al. Effects of climate change on forest plantation productivity in Chile. *Glob Change Biol.* 2022;28(24):7391–409.
 327. Wang A, Melton AE, Soltis DE, Soltis PS. Potential distributional shifts in North America of allelopathic invasive plant species under climate change models. *Plant Divers.* 2022;44(1):11–9.
 328. Gougherty AV, Keller SR, Fitzpatrick MC. Maladaptation, migration and extirpation fuel climate change risk in a forest tree species. *Nat Clim Chang.* 2021;11(2):166–71. <https://doi.org/10.1038/s41558-020-00968-6>.
 329. Messier C, Puettmann KJ, Coates KD. Managing forests as complex adaptive systems: building resilience to the challenge of global change. New York: Routledge; 2013.
 330. Lavery P, Mead D. *Pinus radiata*: a narrow endemic from North America takes on the world. Ecology and biogeography of *Pinus*. 1998:432–49.
 331. Li H-L. Woody flora of Taiwan. Narberth: Livingston Publishing Company; 1963.
 332. Knüsel S, Conedera M, Zweifel R, Bugmann H, Etzold S, Wunder J. High growth potential of *Ailanthus altissima* in warm and dry weather conditions in novel forests of southern Switzerland. *Trees.* 2018:1–15. <https://doi.org/10.1007/s00468-018-1785-x>.
 333. Pötzelberger E, Hasenauer H. High mortality in tree of heaven plantation experiment in eastern Austria. *Austr J For Sci.* 2015;132(4):241–56.
 334. Trifilo P, Raimondo F, Nardini A, Lo Gullo M, Salleo S. Drought resistance of *Ailanthus altissima*: root hydraulics and water relations. *Tree Physiol.* 2004;24(1):107–14. <https://doi.org/10.1093/treephys/24.1.107>.
 335. Kleinbauer I, Dullinger S, Klingenstein F, May R, Nehring S. Ausbreitungspotenzial ausgewählter neophytischer Gefäßpflanzen unter Klimawandel in Deutschland und Österreich. Berlin: Bundesamt für Naturschutz; 2010.
 336. Butler JB, Harrison PA, Vaillancourt RE, Steane DA, Tibbits JF, Potts BM. Climate adaptation, drought susceptibility, and genomic-informed predictions of future climate refugia for the Australian forest tree *Eucalyptus globulus*. *Forests.* 2022;13(4):575. <https://doi.org/10.3390/f13040575>.
 337. López-Sánchez CA, Castedo-Dorado F, Cámara-Obregón A, Barrio-Anta M. Distribution of *Eucalyptus globulus* Labill. in northern Spain: contemporary cover, suitable habitat and potential expansion under climate change. *For Ecol Manag.* 2021;481:118723. <https://doi.org/10.1016/j.foreco.2020.118723>.
 338. Florence RG. Ecology and silviculture of eucalypt forests. Australia: CSIRO publishing; 2004.
 339. Silva FCE, Shvaleva A, Maroco J, Almeida M, Chaves M, Pereira J. Responses to water stress in two *Eucalyptus globulus* clones differing in drought tolerance. *Tree Physiol.* 2004;24(10):1165–72. <https://doi.org/10.1093/treephys/24.10.1165>.
 340. Saadaoui E, Yahia KB, Dhahri S, Jamaa MLB, Khouja ML. An overview of adaptive responses to drought stress in *Eucalyptus* spp. *For Stud.* 2017;67(1):86. <https://doi.org/10.1515/fsmu-2017-0014>.
 341. Santos EF, Mateus NS, Rosario MO, Garcez TB, Mazzafera P, Lavres J. Enhancing potassium content in leaves and stems improves drought tolerance of eucalyptus clones. *Physiol Plant.* 2021;172(2):552–63. <https://doi.org/10.1111/ppl.13228>.
 342. Pita P, Cañas I, Soria F, Ruiz F, Toval G. Use of physiological traits in tree breeding for improved yield in drought-prone environments. The case of *Eucalyptus globulus*. *For Syst.* 2005;14(3):383–93. <https://doi.org/10.5424/srf/2005143-00931>.
 343. Williams MC, Wardle GM. *Pinus radiata* invasion in Australia: identifying key knowledge gaps and research directions. *Austral Ecol.* 2007;32(7):721–39. <https://doi.org/10.1111/j.1442-9993.2007.01760.x>.
 344. Carnegie A, Kathuria A, Nagel M, Mitchell P, Stone C, Sutton M. Current and future risks of drought-induced mortality in *Pinus radiata* plantations in New South Wales, Australia. *Austr For.* 2022;85(4):161–77. <https://doi.org/10.1080/00049158.2022.2145722>.

345. Espinoza SE, Magni CR, Rubilar RA, Yañez MA, Santelices RE, Cabrera AM, et al. Field performance of various *Pinus radiata* breeding families established on a drought-prone site in central Chile. *NZ J Forest Sci.* 2017;47:1–7. <https://doi.org/10.1186/s40490-017-0093-3>.
346. Ismael A, Xue J, Meason DF, Klápště J, Gallart M, Li Y, et al. Genetic variation in drought-tolerance traits and their relationships to growth in *Pinus radiata* D. Don under water stress. *Front Plant Sci.* 2022;12:3043. <https://doi.org/10.3389/fpls.2021.766803>.
347. Watt MS, Kirschbaum MU, Moore JR, Pearce HG, Bulman LS, Brockerhoff EG, et al. Assessment of multiple climate change effects on plantation forests in New Zealand. *Forestry.* 2019;92(1):1–15. <https://doi.org/10.1093/forestry/cpy024>.
348. Guzmán FA, Segura-Ledesma SD, Almáguera-Vargas G. El capulín (*Prunus serotina* Ehrh.): árbol multipropósito con potencial forestal en México. *Madera y Bosques.* 2020;26(1). <https://doi.org/10.21829/myb.2020.2611866>.
349. Segura S, Guzmán-Díaz F, López-Upton J, Mathuriau C, López-Medina J. Distribution of *Prunus serotina* Ehrh. in North America and Its Invasion in Europe. *J Geosci Environ Protect.* 2018;6(09):111. <https://doi.org/10.4236/gep.2018.69009>.
350. Prasad A, Iverson L, Peters M, Matthews S: Climate change tree atlas. Northern Research Station, US Forest Service, Delaware, OH. 2014. <http://www.nrs.fs.fed.us/atlas>. Accessed 3 Jun 2024.
351. Prasad A, Pedlar J, Peters M, McKenney D, Iverson L, Matthews S, et al. Combining US and Canadian forest inventories to assess habitat suitability and migration potential of 25 tree species under climate change. *Divers Distrib.* 2020;26(9):1142–59. <https://doi.org/10.1111/ddi.13078>.
352. Leites LP, Rehfeldt GE, Steiner KC. Adaptation to climate in five eastern North America broadleaf deciduous species: growth clines and evidence of the growth-cold tolerance trade-off. *Perspect Plant Ecol Evol Syst.* 2019;37:64–72. <https://doi.org/10.1016/j.ppees.2019.02.002>.
353. Starfinger U. Die Einbürgerung der Spätblühenden Traubenkirsche (*Prunus serotina* Ehrh.) in Mitteleuropa. Berlin: Technische Universität Berlin; 1990.
354. Van Duinhoven P. Drought response of *Prunus serotina* in the Netherlands. Forest Ecology and Forest Management Group. WUR: Wageningen; 2023.
355. Rosu C. Black locust (*Robinia pseudoacacia* L.) between legend and reality. *Revista Padurilor.* 1996;4:17–20.
356. Mantovani D, Veste M, Freese D. Black locust (*Robinia pseudoacacia* L.) ecophysiological and morphological adaptations to drought and their consequence on biomass production and water-use efficiency. *New Zealand J For Sci.* 2014;44(1):1–11. <https://doi.org/10.1186/s40490-014-0029-0>.
357. Engel J, Knoche D, Lange C. Bewirtschaftung von Robinien-Beständen in Brandenburg-Ergebnisse aus dem FNR-Projekt FastWOOD. Informationen für Waldbesitzer Landesbetrieb Forst Brandenburg. 2014.
358. Claessens H, Alderweireld M, Thibaut A. Potentialités du robinier en Wallonie. *Forêt Nat.* 2006;84:30–9.
359. Grünewald H, Böhm C, Quinkenstein A, Grundmann P, Eberts J, von Wühlisch G. *Robinia pseudoacacia* L.: a lesser known tree species for biomass production. *BioEnergy Res.* 2009;2(3):123–33. <https://doi.org/10.1007/s12155-009-9038-x>.
360. Kutnar L, Kobler A. The current distribution of black locust (*Robinia pseudoacacia* L.) in Slovenia and predictions for the future. *Acta Silvae et Ligni.* 2013;102:21–30.
361. Giuliani C, Lazzaro L, Mariotti Lippi M, Calamassi R, Foggi B. Temperature-related effects on the germination capacity of black locust (*Robinia pseudoacacia* L., Fabaceae) seeds. *Folia Geobotanica.* 2015;50(3):275–82. <https://doi.org/10.1007/s12224-015-9224-x>.
362. Dyderski MK, Paź S, Frelich LE, Jagodziński AM. How much does climate change threaten European forest tree species distributions? *Glob Change Biol.* 2018;24(3):1150–63. <https://doi.org/10.1111/gcb.13925>.
363. Lavelle P, Dugdale R, Scholes R, Berhe A, Carpenter E, Codispoti L, et al. Nutrient cycling. *Ecosystems and Human Well-Being: Current State and Trends: Findings of the Condition and Trends Working Group*, Island Press, Washington, Covelo, London. 2005.
364. Carnol M, Bazgir M. Nutrient return to the forest floor through litter and throughfall under 7 forest species after conversion from Norway spruce. *For Ecol Manage.* 2013;309:66–75. <https://doi.org/10.1016/j.foreco.2013.04.008>.
365. Likens GE, Driscoll CT, Buso DC. Long-term effects of acid rain: response and recovery of a forest ecosystem. *Science.* 1996;272(5259):244–6. <https://doi.org/10.1126/science.272.5259.244>.
366. Schaberg PG, DeHayes DH, Hawley GJ. Anthropogenic calcium depletion: a unique threat to forest ecosystem health? *Ecosyst Health.* 2001;7(4):214–28. <https://doi.org/10.1046/j.1526-0992.2001.01046.x>.
367. Bowman WD, Cleveland CC, Halada L, Hřeško J, Baron JS. Negative impact of nitrogen deposition on soil buffering capacity. *Nat Geosci.* 2008;1(11):767–70. <https://doi.org/10.1038/ngeo339>.
368. Sumner M, Fey M, Noble A. Nutrient status and toxicity problems in acid soils. Berlin: Springer; 1991.
369. Galloway JN. Acidification of the world: natural and anthropogenic. *Water Air Soil Pollut.* 2001;130(1–4):17–24. <https://doi.org/10.1023/A:1012272431583>.
370. Muys B, Lust N, Granval PH. Effects of grassland afforestation with different tree species on earthworm communities, litter decomposition and nutrient status. *Soil Biol Biochem.* 1992;24(12):1459–66. [https://doi.org/10.1016/0038-0717\(92\)90133-I](https://doi.org/10.1016/0038-0717(92)90133-I).
371. Finzi AC, Van Breemen N, Canham CD. Canopy tree-soil interactions within temperate forests: species effects on soil carbon and nitrogen. *Ecol Appl.* 1998;8(2):440–6. [https://doi.org/10.1890/1051-0761\(1998\)008\[0440:CTSIWT\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1998)008[0440:CTSIWT]2.0.CO;2).
372. Hommel P, Waal R, Muys B, Ouden J, Spek T. Terug naar het lindewoud, Strooiselkwaliteit als basis voor ecologisch bosbeheer. KNNV Uitgeverij: Zeist; 2007.
373. Mueller KE, Hobbie SE, Chorover J, Reich PB, Eisenhauer N, Castellano MJ, et al. Effects of litter traits, soil biota, and soil chemistry on soil carbon stocks at a common garden with 14 tree species. *Biogeochemistry.* 2015;123(3):313–27. <https://doi.org/10.1007/s10533-015-0083-6>.
374. Reich PB, Oleksyn J, Modrzynski J, Mrozinski P, Hobbie SE, Eissenstat DM, et al. Linking litter calcium, earthworms and soil properties: a common garden test with 14 tree species. *Ecol Lett.* 2005;8(8):811–8. <https://doi.org/10.1111/j.1461-0248.2005.00779.x>.
375. Castro-Díez P, González-Muñoz N, Alonso A, Gallardo A, Poorter L. Effects of exotic invasive trees on nitrogen cycling: a case study in Central Spain. *Biol Invasions.* 2009;11(8):1973–86. <https://doi.org/10.1007/s10530-008-9374-3>.
376. González-Muñoz N, Castro-Díez P, Parker I. Differences in nitrogen use strategies between native and exotic tree species: predicting impacts on invaded ecosystems. *Plant Soil.* 2013;363(1):319–29. <https://doi.org/10.1007/s11104-012-1329-x>.
377. Gutiérrez-López M, Ranera E, Novo M, Fernández R, Trigo D. Does the invasion of the exotic tree *Ailanthus altissima* affect the soil arthropod community? The case of a riparian forest of the Henares River (Madrid). *Eur J Soil Biol.* 2014;62:39–48. <https://doi.org/10.1016/j.ejsobi.2014.02.010>.

378. Motard E, Dusz S, Geslin B, Akpa-Vinceslas M, Hignard C, Babiar O, et al. How invasion by *Ailanthus altissima* transforms soil and litter communities in a temperate forest ecosystem. *Biol Invasions*. 2015;17(6):1817–32. <https://doi.org/10.1007/s10530-014-0838-3>.
379. Canhoto C, Graça MA. Decomposition of *Eucalyptus globulus* leaves and three native leaf species (*Alnus glutinosa*, *Castanea sativa* and *Quercus faginea*) in a Portuguese low order stream. *Hydrobiologia*. 1996;333(2):79–85. <https://doi.org/10.1007/BF00017570>.
380. Ribeiro C, Madeira M, Araújo M. Decomposition and nutrient release from leaf litter of *Eucalyptus globulus* grown under different water and nutrient regimes. *For Ecol Manage*. 2002;171(1–2):31–41. [https://doi.org/10.1016/S0378-1127\(02\)00459-0](https://doi.org/10.1016/S0378-1127(02)00459-0).
381. Madeira M. Influência dos Povoamentos de Eucalipto (*Eucalyptus globulus* Labill.) no Solo, Comparativamente aos Povoamentos de Sobreiro (*Quercus suber* L.) e de Pinheiro (*Pinus pinaster* Ait.). Lisbon: Instituto Superior de Agronomia; 1986.
382. Madeira M. Changes in soil properties under *Eucalyptus* plantations in Portugal. In: Pereira JS, Landsberg JJ, editors. *Biomass Production by Fast-growing trees*. Dordrecht: Springer; 1989. p. 81–99.
383. Madeira M, Pereira J. Productivity, nutrient immobilization and soil chemical properties in an *Eucalyptus globulus* plantation under different irrigation and fertilization regimes. *Water Air Soil Pollut*. 1990;54:621–34. <https://doi.org/10.1007/BF02385159>.
384. Madeira M, Araújo M, Pereira J. Effects of water and nutrient supply on amount and on nutrient concentration of litterfall and forest floor litter in *Eucalyptus globulus* plantations. *Plant Soil*. 1995;168:287–95. <https://doi.org/10.1007/BF00029340>.
385. Serralheiro F, Madeira M. Acari colonization of *Quercus suber* and *Eucalyptus globulus* litter. In: Dusbicsek F, Bukva V, editors. *Modern Acarology*. Prague: Academic Publishing; 1991. p. 353–8.
386. Jones H, Madeira M, Herraes L, Dighton J, Fabião A, González-Río F, et al. The effect of organic-matter management on the productivity of *Eucalyptus globulus* stands in Spain and Portugal: tree growth and harvest residue decomposition in relation to site and treatment. *For Ecol Manage*. 1999;122(1–2):73–86. [https://doi.org/10.1016/S0378-1127\(99\)00033-X](https://doi.org/10.1016/S0378-1127(99)00033-X).
387. Cortez N. Compartimentos e Ciclos de Nutrientes em Plantações de *Eucalyptus globulus* Labill. ssp. *globulus* e *Pinus pinaster* Aiton. Lisbon: Universidade Técnica de Lisboa; 1996.
388. Le Maitre DC, van Wilgen BW, Gelderblom C, Bailey C, Chapman RA, Nel J. Invasive alien trees and water resources in South Africa: case studies of the costs and benefits of management. *For Ecol Manage*. 2002;160(1–3):143–59. [https://doi.org/10.1016/S0378-1127\(01\)00474-1](https://doi.org/10.1016/S0378-1127(01)00474-1).
389. Velarde S, Paul T, Monge J, Yao R. Cost benefit analysis of wilding conifer management in New Zealand; Part I—Important Impacts under current management. In: Scion, editor. *Report S0013*. Bloxham: New Zealand Forest Research Institute; 2015.
390. Sapsford SJ, Brandt AJ, Davis KT, Peralta G, Dickie IA, Gibson RD, et al. Towards a framework for understanding the context dependence of impacts of non-native tree species. *Funct Ecol*. 2020;34(5):944–55. <https://doi.org/10.1111/1365-2435.13544>.
391. Maddelein D. Driving Forces and limiting factors in long-term dynamics of forest ecosystems on sandy soils. Ghent: Laboratory of Forestry, University of Ghent; 1991.
392. Adams M, Angradi T. Decomposition and nutrient dynamics of hardwood leaf litter in the Fernow whole-watershed acidification experiment. *For Ecol Manage*. 1996;83(1–2):61–9. [https://doi.org/10.1016/0378-1127\(95\)03695-4](https://doi.org/10.1016/0378-1127(95)03695-4).
393. Lorenz K, Preston CM, Krumrei S, Feger KH. Decomposition of needle/leaf litter from Scots pine, black cherry, common oak and European beech at a conurbation forest site. *Eur J Forest Res*. 2004;123(3):177–88. <https://doi.org/10.1007/s10342-004-0025-7>.
394. Dassonville N, Vanderhoeven S, Vanparys V, Hayez M, Gruber W, Meerts P. Impacts of alien invasive plants on soil nutrients are correlated with initial site conditions in NW Europe. *Oecologia*. 2008;157(1):131–40. <https://doi.org/10.1007/s00442-008-1054-6>.
395. Vanderhoeven S, Dassonville N, Meerts P. Increased topsoil mineral nutrient concentrations under exotic invasive plants in Belgium. *Plant Soil*. 2005;275(1):169–79. <https://doi.org/10.1007/s11104-005-1257-0>.
396. Horodecki P, Jagodziński AM. Tree species effects on litter decomposition in pure stands on afforested post-mining sites. *For Ecol Manage*. 2017;406:1–11. <https://doi.org/10.1016/j.foreco.2017.09.059>.
397. Horodecki P, Nowiński M, Jagodziński AM. Advantages of mixed tree stands in restoration of upper soil layers on postmining sites: A five-year leaf litter decomposition experiment. *Land Degrad Dev*. 2019;30(1):3–13. <https://doi.org/10.1002/ldr.3194>.
398. Hobbie SE, Reich PB, Oleksyn J, Ogdahl M, Zytowski R, Hale C, et al. Tree species effects on decomposition and forest floor dynamics in a common garden. *Ecology*. 2006;87(9):2288–97. [https://doi.org/10.1890/0012-9658\(2006\)87\[2288:TSEODA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2288:TSEODA]2.0.CO;2).
399. Desie E, Vancampenhout K, Van den Berg L, Nyssen B, Weijters M, Den Ouden J, et al. Litter share and clay content determine soil restoration effects of rich litter tree species in forests on acidified sandy soils. *For Ecol Manage* 2020. <https://doi.org/10.1016/j.foreco.2020.118377>.
400. De AN. invloed van *Prunus serotina* (Amerikaanse vogelkers) op duinbodemeigenschappen in de Amsterdamse waterleidingduinen. Hoorn: Universiteit van Amsterdam; 2005.
401. Chabrierie O, Verheyen K, Saguez R, Decocq G. Disentangling relationships between habitat conditions, disturbance history, plant diversity, and American black cherry (*Prunus serotina* Ehrh.) invasion in a European temperate forest. *Divers Distrib*. 2008;14(2):204–12. <https://doi.org/10.1111/j.1472-4642.2007.00453.x>.
402. Plichta W, Kuczynska I, Rutkowski L. The effect of american cherry [*Prunus serotina* Erh.] plantations on organic and humus horizons of cambic arenosols in a pine forest. *Ekologia Polska*. 1997;2(45):385–94.
403. Aerts R, Ewald M, Nicolas M, Piat J, Skowronek S, Lenoir J, et al. Invasion by the Alien Tree *Prunus serotina* alters ecosystem functions in a temperate deciduous forest. *Front Plant Sci*. 2017;8:179. <https://doi.org/10.3389/fpls.2017.00179>.
404. Liao C, Peng R, Luo Y, Zhou X, Wu X, Fang C, et al. Altered ecosystem carbon and nitrogen cycles by plant invasion: a meta-analysis. *New Phytol*. 2008;177(3):706–14. <https://doi.org/10.1111/j.1469-8137.2007.02290.x>.
405. Vilà M, Espinar JL, Hejda M, Hulme PE, Jarošík V, Maron JL, et al. Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol Lett*. 2011;14(7):702–8. <https://doi.org/10.1111/j.1461-0248.2011.01628.x>.
406. Alfredsson H, Condron LM, Clarholm M, Davis MR. Changes in soil acidity and organic matter following the establishment of conifers on former grassland in New Zealand. *For Ecol Manage*. 1998;112(3):245–52. [https://doi.org/10.1016/S0378-1127\(98\)00346-6](https://doi.org/10.1016/S0378-1127(98)00346-6).
407. Ntshuxeko VE, Ruwanza S. Physical properties of soil in *Pine elliottii* and *Eucalyptus cloeziana* plantations in the Vhembe

- biosphere, Limpopo Province of South Africa. *J For Res.* 2020;31:625–35. <https://doi.org/10.1007/s11676-018-0830-3>.
408. Ewers RM, Banks-Leite C. Fragmentation impairs the microclimate buffering effect of tropical forests. *PLoS ONE.* 2013;8(3):e58093. <https://doi.org/10.1371/journal.pone.0058093>.
409. Von Arx G, Graf Pannatier E, Thimonier A, Rebetez M. Microclimate in forests with varying leaf area index and soil moisture: potential implications for seedling establishment in a changing climate. *J Ecol.* 2013;101(5):1201–13. <https://doi.org/10.1111/1365-2745.12121>.
410. De Frenne P, Zellweger F, Rodríguez-Sánchez F, Scheffers BR, Hylander K, Luoto M, et al. Global buffering of temperatures under forest canopies. *Nat Ecol Evol.* 2019;3(5):744–9. <https://doi.org/10.1038/s41559-019-0842-1>.
411. Geiger R, Aron RH, Todhunter P. *The climate near the ground.* Plymouth: Rowman & Littlefield; 2009.
412. Zellweger F, Coomes D, Lenoir J, Depauw L, Maes SL, Wulf M, et al. Seasonal drivers of understorey temperature buffering in temperate deciduous forests across Europe. *Glob Ecol Biogeogr.* 2019;28(12):1774–86. <https://doi.org/10.1111/geb.12991>.
413. Chen J, Saunders SC, Crow TR, Naiman RJ, Brososke KD, Mroz GD, et al. Microclimate in forest ecosystem and landscape ecology: variations in local climate can be used to monitor and compare the effects of different management regimes. *Bioscience.* 1999;49(4):288–97. <https://doi.org/10.2307/1313612>.
414. Jensen AM, Löf M, Gardiner ES. Effects of above- and below-ground competition from shrubs on photosynthesis, transpiration and growth in *Quercus robur* L. seedlings. *Environ Exper Botany.* 2011;71(3):367–75.
415. Johnstone JF, Allen CD, Franklin JF, Frelich LE, Harvey BJ, Higuera PE, et al. Changing disturbance regimes, ecological memory, and forest resilience. *Front Ecol Environ.* 2016;14(7):369–78. <https://doi.org/10.1002/fee.1311>.
416. Stevens JT, Haffey CM, Coop JD, Fornwalt PJ, Yocom L, Allen CD, et al. Tamm Review: Postfire landscape management in frequent-fire conifer forests of the southwestern United States. *For Ecol Manage.* 2021;502:119678. <https://doi.org/10.1016/j.foreco.2021.119678>.
417. Lenoir J, Hattab T, Pierre G. Climatic microrefugia under anthropogenic climate change: implications for species redistribution. *Ecography.* 2017;40(2):253–66. <https://doi.org/10.1111/ecog.02788>.
418. De Frenne P, Lenoir J, Luoto M, Scheffers BR, Zellweger F, Aalto J, et al. Forest microclimates and climate change: Importance, drivers and future research agenda. *Glob Change Biol.* 2021;27(11):2279–97. <https://doi.org/10.1111/gcb.15569>.
419. Keppel G, Van Niel KP, Wardell-Johnson GW, Yates CJ, Byrne M, Mucina L, et al. Refugia: identifying and understanding safe havens for biodiversity under climate change. *Glob Ecol Biogeogr.* 2012;21(4):393–404. <https://doi.org/10.1111/j.1466-8238.2011.00686.x>.
420. Ashcroft MB, Gollan JR, Warton DI, Ramp D. A novel approach to quantify and locate potential microrefugia using topoclimate, climate stability, and isolation from the matrix. *Glob Change Biol.* 2012;18(6):1866–79. <https://doi.org/10.1111/j.1365-2486.2012.02661.x>.
421. Hampe A, Jump AS. Climate relicts: past, present, future. 2011. <https://doi.org/10.1146/annurev-ecolsys-102710-145015>.
422. Greiser C, Ehrlén J, Meineri E, Hylander K. Hiding from the climate: Characterizing microrefugia for boreal forest understorey species. *Glob Change Biol.* 2020;26(2):471–83. <https://doi.org/10.1111/gcb.14874>.
423. Kovács B, Tinya F, Ódor P. Stand structural drivers of microclimate in mature temperate mixed forests. *Agric For Meteorol.* 2017;234:11–21. <https://doi.org/10.1016/j.agrformet.2016.11.268>.
424. Gehlhausen SM, Schwartz MW, Augspurger CK. Vegetation and microclimatic edge effects in two mixed-mesophytic forest fragments. *Plant Ecol.* 2000;147(1):21–35. <https://doi.org/10.1023/A:1009846507652>.
425. De Frenne P, Rodríguez-Sánchez F, Coomes DA, Baeten L, Verstraeten G, Vellend M, et al. Microclimate moderates plant responses to macroclimate warming. *Proc Natl Acad Sci.* 2013;110(46):18561–5. <https://doi.org/10.1073/pnas.1311190110>.
426. Frey SJ, Hadley AS, Johnson SL, Schulze M, Jones JA, Betts MG. Spatial models reveal the microclimatic buffering capacity of old-growth forests. *Sci Adv.* 2016;2(4):e1501392. <https://doi.org/10.1126/sciadv.150139>.
427. Latimer CE, Zuckerberg B. Forest fragmentation alters winter microclimates and microrefugia in human-modified landscapes. *Ecography.* 2017;40(1):158–70. <https://doi.org/10.1111/ecog.02551>.
428. Niculescu M, Olaru AL, Cojoacă FD. Study of phytosociology and ecology of *Ailanthus altissima* (Miller) swingle–invasive species in the south-western of Romania. *Ann Univ Craiova-Agric Montanol Cadastre Ser.* 2018;48(1):280–6.
429. Montecchiari S, Tesei G, Allegranza M. *Ailanthus altissima* forests determine a shift in herbaceous layer richness: a paired comparison with hardwood native forests in sub-Mediterranean Europe. *Plants.* 2020;9(10):1404. <https://doi.org/10.3390/plant9101404>.
430. Dinucă NC. *Ailanthus altissima* (Miller) swingle invasive species in the forest habitats in the upper basin of the Motru River. *Curr Trends Nat Sci.* 2021;10(19):175–80. <https://doi.org/10.47068/ctns.2021.v10i19.023>.
431. Spiess U. Wachstumsanalyse von juvenilen Götterbäumen in der Südschweiz. In: *Umweltsystemwissenschaften D*, editor. Zürich: ETH; 2014.
432. Maringer J, Wohlgemuth T, Neff C, Pezzatti GB, Conedera M. Post-fire spread of alien plant species in a mixed broad-leaved forest of the Insubric region. *Flora-Morphol Distrib Funct Ecol Plants.* 2012;207(1):19–29. <https://doi.org/10.1016/j.flora.2011.07.016>.
433. Brang P, Spathelf P, Larsen JB, Bauhus J, Boncčina A, Chauvin C, et al. Suitability of close-to-nature silviculture for adapting temperate European forests to climate change. *Forestry.* 2014;87(4):492–503. <https://doi.org/10.1093/forestry/cpu018>.
434. Wołkowycki D, Próchnicki P. Spatial expansion pattern of black cherry *Padus serotina* Ehrh. in suburban zone of Białystok (NE Poland). *Biodivers Res Conserv.* 2015;40(1):59–67. <https://doi.org/10.1515/biorc-2015-0030>.
435. Bułaj B, Okpiasz K, Rutkowski P, Tomczak A. Occurrence of invasive black cherry (*Prunus serotina* Ehrh.) on abandoned farmland in west-central Poland. *Forestry Letters.* 2017;110(12):26–31.
436. Starfinger U. Population Biology of an Invading Tree Species *Prunus serotina* In: Seitz A, Loeschke V. (Eds), *Species Conservation: A Population Biology Approach.* Basel, Birkhäuser Verlag; 1991.
437. Starfinger U. Introduction and naturalization of *Prunus serotina* in central Europe. In: Brock JH, Wade M, Pysek P, Green D, editors. *Plant invasions: studies from North America and Europe.* Leiden: Backhuys; 1997.
438. Olsthoorn A, van Hees A. 40 years of black cherry (*Prunus serotina*) control in the Netherlands: lessons for management of invasive tree species. In: Kowarik I, Starfinger U, editors. *Biological invasions in Germany—a challenge to act?; contributions and results of a conference in Berlin, October 4th–7th, 2000.* Berlin: BfN-Skripten; 2001. p. 43–4.
439. Deckers B, Verheyen K, Hermy M, Muys B. Effects of landscape structure on the invasive spread of black cherry *Prunus serotina* in an agricultural landscape in Flanders, Belgium. *Belgium Ecograp.* 2005;28:99–109.

440. Godefroid S, Phartyal SS, Weyembergh G, Koedam N. Ecological factors controlling the abundance of non-native invasive black cherry (*Prunus serotina*) in deciduous forest understory in Belgium. *For Ecol Manage.* 2005;210(1–3):91–105. <https://doi.org/10.1016/j.foreco.2005.02.024>.
441. Verheyen K, Vanhellefont M, Stock T, Hermy M. Predicting patterns of invasion by black cherry (*Prunus serotina* Ehrh.) in Flanders (Belgium) and its impact on the forest understory community. *Divers Distrib.* 2007;13(5):487–97. <https://doi.org/10.1111/j.1472-4642.2007.00334.x>.
442. Vanhellefont M, Verheyen K, Hermy M. Long-term population dynamics of *Prunus serotina* in natural forests on sandy soils 2nd European Congress of Conservation Biology (ECCB 2009): conservation biology and beyond: from science to practice. Prague: Czech Republic; 2009. p. 44.
443. Rode M, Kowarik I, Müller T, Wendebourg T. the effects of *Prunus serotina* on pine forest ecosystems in lower saxony. In: Starfinger U, Kowarik I, editors. *Biological Invasions – a challenge to act? contributions and results of a conference in Berlin, 4–7 October 2000*. Bonn: Federal Nature Conservation Agency; 2001. p. 91–2.
444. Decocq G. Dynamique invasive du cerisier tardif, *Prunus serotina* L., en système forestier tempéré: déterminants, mécanismes, impacts écologiques, économiques et socioanthropologiques. In: Verne UdPJ, editor. *Amiens: Université de Picardie Jules Verne; 2007*.
445. Chabrierie O, Loinard J, Perrin S, Saguez R, Decocq G. Impact of *Prunus serotina* invasion on understory functional diversity in a European temperate forest. *Biol Invasions.* 2010;12(6):1891–907. <https://doi.org/10.1007/s10530-009-9599-9>.
446. Halarewicz A, Pruchniewicz D. Vegetation and environmental changes in a Scots pine forest invaded by *Prunus serotina*: what is the threat to terricolous bryophytes? *Eur J Forest Res.* 2015;134(5):793–801. <https://doi.org/10.1007/s10342-015-0890-2>.
447. Halarewicz A, Zolniercz L. Changes in the understory of mixed coniferous forest plant communities dominated by the American black cherry (*Prunus serotina* Ehrh.). *For Ecol Manage.* 2014;313:91–7. <https://doi.org/10.1016/j.foreco.2013.11.006>.
448. Dyderski MK, Jagodziński AM. How do invasive trees impact shrub layer diversity and productivity in temperate forests? *Ann For Sci.* 2021;78(1):1–14. <https://doi.org/10.1007/s13595-021-01033-8>.
449. Swearingen J, Barger C. *Invasive plant atlas of the United States*. Shepherdstown: National Conservation Training Center; 2017.
450. Kowarik I. Zur einföhrung und ausbreitung der Robinie (*Robinia pseudoacacia* L.) in Brandenburg und zur gehölsukzession ruderaler robinienbestände in Berlin. *Verh Berliner Bot Ver.* 1990;8:33–67.
451. Kowarik I, Lippe M, Cierjacks A. Prevalence of alien versus native species of woody plants in Berlin differs between habitats and at different scales. *Preslia.* 2013;85(2):113–32.
452. Botta-Dukát Z. Invasion of alien species to Hungarian (semi-) natural habitats. *Acta Bot Hung.* 2008;50(1):219–27. <https://doi.org/10.1556/abot.50.2008.suppl.11>.
453. Essl F, Hauser E. Verbreitung, Lebensraumbindung und Managementkonzept ausgewählter invasiver Neophyten im Nationalpark Thayatal und Umgebung (Österreich). *Linzer Biol Beitr.* 2003;35:75–101.
454. Wojda T, Klisz M, Jastrzębowski S, Mionskowski M, Szyp-Borowska I, Szczygieł K. The geographical distribution of the black locust (*Robinia pseudoacacia* L.) in Poland and its role on non-forested land. *Papers on Global Change IGBP.* 2015(22). <https://doi.org/10.1515/igbp-2015-0018>
455. Essl F, Nehring S, Klingenstein F, Milasowszky N, Nowack C, Rabitsch W. Review of risk assessment systems of IAS in Europe and introducing the German-Austrian Black List Information System (GABLIS). *J Nat Conserv.* 2011;19(6):339–50. <https://doi.org/10.1016/j.jnc.2011.08.005>.
456. Cseresnyes I, Csontos P. Soil seed bank of the invasive *Robinia pseudoacacia* in planted *Pinus nigra* stands. *Acta Bot Croat.* 2012;71(2):249–60. <https://doi.org/10.2478/v10184-011-0065-2>.
457. Dyderski MK, Jagodziński AM. Seedling survival of *Prunus serotina* Ehrh., *Quercus rubra* L. and *Robinia pseudoacacia* L. in temperate forests of Western Poland. *For Ecol Manage.* 2019;450:117498. <https://doi.org/10.1016/j.foreco.2019.117498>.
458. Parrotta JA. The role of plantation forests in rehabilitating degraded tropical ecosystems. *Agr Ecosyst Environ.* 1992;41(2):115–33. [https://doi.org/10.1016/0167-8809\(92\)90105-K](https://doi.org/10.1016/0167-8809(92)90105-K).
459. Parrotta JA, Turnbull JW, Jones N. Catalyzing native forest regeneration on degraded tropical lands. *For Ecol Manage.* 1997;99(1–2):1–7. [https://doi.org/10.1016/S0378-1127\(97\)00190-4](https://doi.org/10.1016/S0378-1127(97)00190-4).
460. Geldenhuys CJ. Native forest regeneration in pine and eucalypt plantations in Northern Province. *South Africa Forest Ecol Manage.* 1997;99(1–2):101–15. [https://doi.org/10.1016/S0378-1127\(97\)00197-7](https://doi.org/10.1016/S0378-1127(97)00197-7).
461. Lamb D. Large-scale ecological restoration of degraded tropical forest lands: the potential role of timber plantations. *Restor Ecol.* 1998;6(3):271–9. <https://doi.org/10.1046/j.1526-100X.1998.00632.x>.
462. Simões LH, Guillemot J, Ronquim CC, Weidlich EW, Muys B, Fuza MS, et al. Green deserts, but not always: A global synthesis of native woody species regeneration under tropical tree monocultures. *Glob Change Biol.* 2024;30(4):e17269. <https://doi.org/10.1111/gcb.17269>.
463. del Campo AD, Otsuki K, Serengil Y, Blanco JA, Yousefpour R, Wei X. A global synthesis on the effects of thinning on hydrological processes: Implications for forest management. *For Ecol Manage.* 2022;519:120324. <https://doi.org/10.1016/j.foreco.2022.120324>.
464. Linders TEW, Schaffner U, Eschen R, Abebe A, Choge SK, Nigatu L, et al. Direct and indirect effects of invasive species: Biodiversity loss is a major mechanism by which an invasive tree affects ecosystem functioning. *J Ecol.* 2019;107(6):2660–72. <https://doi.org/10.1111/1365-2745.13268>.
465. Brus R, Pötzelsberger E, Lapin K, Brundu G, Orazio C, Straigyte L, et al. Extent, distribution and origin of non-native forest tree species in Europe. *Scand J For Res.* 2019;34(7):533–44. <https://doi.org/10.1080/02827581.2019.1676464>.
466. Konic J, Heiling C, Haeler E, Chakraborty D, Lapin K, Schueler S. The potential of non-native tree species to provide major ecosystem services in Austrian forests. *Front Plant Sci.* 2024;15. <https://doi.org/10.3389/fpls.2024.1402601>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.