

























Clarifying the trophic state concept to advance macroscale freshwater science and management

Michael F. Meyer^{1,2}  | Benjamin M. Kraemer³  | Carolina C. Barbosa⁴  |
 Davi G. F. Cunha⁵  | Walter K. Dodds⁶  | Stephanie E. Hampton⁷  |
 César Ordóñez⁸  | Rachel M. Pilla⁹  | Amina I. Pollard¹⁰  |
 Joshua A. Culpepper¹¹  | Alexander K. Fremier¹²  | Tyler V. King¹³  |
 Robert Ladwig^{2,14}  | Dina M. Leech¹⁵  | Shin-Ichiro S. Matsuzaki¹⁶  |
 Isabella A. Oleksy¹⁷  | Simon N. Topp¹⁸  | R. Iestyn Woolway¹⁹  |
 Ludmila S. Brighenti²⁰  | Kate C. Fickas^{21,22}  | Brian P. Lanouette²³  |
 Jianning Ren²⁴  | Mortimer Werther²⁵  | Xiao Yang²⁶ 

Correspondence

Michael F. Meyer

Email: mfmeyer@usgs.gov

Funding information

Conselho Nacional de Desenvolvimento Científico e Tecnológico, Grant/Award Number: #310844/2020-7; Apoio Institucional a Pesquisa from the Universidade do Estado de Minas Gerais, Grant/Award Number: PAPq-UEMG 11/2022; Natural Environment Research Council, Grant/Award Number: NE/T011246/1; National Science Foundation, Grant/Award Numbers: DEB-2306895, EPS-2019528; Oak Ridge National Laboratory; U.S. Geological Survey, Mendenhall Fellowship

Handling Editor: Wyatt F. Cross

Abstract

For over a century, ecologists have used the concept of trophic state (TS) to characterize an aquatic ecosystem's biological productivity. However, multiple TS classification schemes, each relying on a variety of measurable parameters as proxies for productivity, have emerged to meet use-specific needs. Frequently, chlorophyll a, phosphorus, and Secchi depth are used to classify TS based on autotrophic production, whereas phosphorus, dissolved organic carbon, and true color are used to classify TS based on both autotrophic and heterotrophic production. Both classification approaches aim to characterize an ecosystem's function broadly, but with varying degrees of autotrophic and heterotrophic processes considered in those characterizations. Moreover, differing classification schemes can create inconsistent interpretations of ecosystem integrity. For example, the US Clean Water Act focuses exclusively on algal threats to water quality, framed in terms of eutrophication in response to nutrient loading. This usage lacks information about non-algal threats to water quality, such as dystrophication in response to dissolved organic carbon loading. Consequently, the TS classification schemes used to identify eutrophication and dystrophication may refer to ecosystems similarly (e.g., oligotrophic and eutrophic), yet these categories are derived from different proxies. These inconsistencies in TS classification schemes may be compounded when interdisciplinary projects employ varied TS frameworks. Even with these shortcomings, TS can still be used to distill information on complex aquatic ecosystem

For affiliations refer to page 16

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 Oak Ridge National Laboratory and The Author(s). *Ecosphere* published by Wiley Periodicals LLC on behalf of The Ecological Society of America. This article has been contributed to by U.S. Government employees and their work is in the public domain in the USA.

function into a set of generalizable expectations. The usefulness of distilling complex information into a TS index is substantial such that usage inconsistencies should be explicitly addressed and resolved. To emphasize the consequences of diverging TS classification schemes, we present three case studies for which an improved understanding of the TS concept advances freshwater research, management efforts, and interdisciplinary collaboration. To increase clarity in TS, the aquatic sciences could benefit from including information about the proxy variables, ecosystem type, as well as the spatiotemporal domains used to classify TS. As the field of aquatic sciences expands and climatic irregularity increases, we highlight the importance of re-evaluating fundamental concepts, such as TS, to ensure their compatibility with evolving science.

KEYWORDS

classification, ecosystem function, limnology, metabolism, water quality

INTRODUCTION

Trophic state (TS) is a fundamental concept in the aquatic sciences that describes an ecosystem's characteristic biological productivity. Operationally, productivity can be challenging to estimate because spatial and temporal heterogeneities within ecosystems can substantially influence waterbody-wide productivity. Therefore, several proxies and formulations for classifying TS have emerged since the beginning of the 20th century (Box 1; Table 1). Nutrients, chlorophyll, and Secchi disk depth are common proxy variables for evaluating TS, whereas alternative formulations rely on true color, organic carbon, biomass estimates, and even microbial community composition (Table 1). Depending on data availability, TS may be expressed as a discrete group, a continuous index, or even a probability of being assigned to a discrete group, where each formulation communicates information about uncertainty in the classification (Nojvan et al., 2019). The proxies measured and consequent classifications of TS can also differ between ecosystem types and regions (Table 1). In lakes and reservoirs, Hutchinson (1957) focused on hypolimnetic oxygen depletion rates. Lindeman (1942) and Horne and Goldman (1983) focused on TS as phases of a waterbody's ontogeny. Carlson (1977) focused solely on autochthony (i.e., primary production), whereas Naumann (1917), Thienemann (1921), and Wetzel (2001) focused on both autochthony and allochthony. In rivers and streams, the TS concept has likewise focused on productivity as a function of autochthony and allochthony, but more specifically, the ratio of photosynthesis to respiration at the scale of river reach (Dodds & Cole, 2007; Odum, 1956). Dodds (2006) classified TS on the

probability of observing a given phosphorus, nitrogen, or chlorophyll concentration in a river reach, where benthic and pelagic algae can be independently considered.

Regardless of the exact TS classification scheme, scientists, managers, and the public rely upon these simplified discretizations to characterize complex ecosystem processes, thereby allowing for scientific progress when detailed data are lacking (Kraemer, 2020). In this sense, TS is a fundamental ecosystem characteristic that can be inferred from basic water quality data. Such simplifications are important for generalizing our understanding of ecosystem function, when often only discrete observations of ecosystems are available (*sensu* Palmer & Febria, 2012). Wetzel (2001) demonstrates this point by using TS to make broad predictions about how and why water quality constituent depth profiles, such as dissolved oxygen and pH, differ between stratified oligotrophic and eutrophic lakes (Figure 1). Although these generalities may not always apply to every ecosystem, the TS concept is a useful, conceptual framework that can predict water quality conditions when direct measurements are sparse. For example, depth profiles are uncommon relative to grab samples from surface waters, yet consistent depth-profile patterns for a given TS allow us to infer profile dynamics from available depth-profile datasets (Figure 1). These uses of TS have even extended to policy, where the language of TS is included in Sections 106 and 314 of the U.S. Clean Water Act (33 U.S.C. 1252 et seq.) for the "identification and classification [of lakes] according to eutrophic condition," and eutrophic conditions can trigger "procedures, processes, and methods ... to control sources of pollution and ... to restore [water quality]." In each of these instances, proxy variables are

BOX 1 A comparison of select lake TS classification schemes.

Several classification schemes have been adopted to classify trophic states, including those mentioned in Table 1 and others (e.g., Dodds & Whiles, 2019). Although we cannot compare all these approaches, we have compared three approaches to illustrate how differences in the definition of TS can lead to divergent interpretations if considered independently and without context. Together, the TS classifications offer broader insights into the trophic state concept.

Trophic State Index (TSI), developed by Carlson (1977) and subsequently refined, has been used as a descriptor of water quality in lentic waterbodies and has been frequently adopted by management agencies, including the U.S. Environmental Protection Agency (1990). TSI provides both a continuous metric and a categorical grouping but only indicates autotrophic productivity. Furthermore, TSI has been adapted to accommodate values typical to a given location. For example, in Brazil, TSI relationships have been adapted to classify tropical reservoirs to consider the overall greater productivity of tropical ecosystems compared to other climate zones (Cunha et al., 2013). Proxies of this scheme are Secchi disk depth, total phosphorus, and chlorophyll *a*; classifications include oligotrophic, mesotrophic, eutrophic, and hypereutrophic.

Nutrient Color Paradigm (NCP) groups lakes based on water clarity (measured as carbon concentration, water color, or absorption coefficient) and autotrophic capacity. Rohde (1969) first arranged the four quadrants of the NCP, placing autochthony on the horizontal axis and allochthony on the vertical axis. This second dimension discriminates “oligotrophic” (low autochthony, low allochthony) and “eutrophic” (high autochthony, low allochthony) lakes from “dystrophic” (low autochthony, high allochthony) lakes and “mixotrophic” (high autochthony, high allochthony) lakes. Proxies of this scheme are total phosphorus and carbon concentration, water color, or absorption coefficient; classifications include oligotrophic, eutrophic, dystrophic, and mixotrophic.

Ecological State (ES) is a component of the European Union’s Water Framework Directive (WFD; Directive 2000/60/EC of the European Parliament and of the Council (EU), 2000), which introduces a planning process and assessment schema to manage, protect, and improve the surface and subsurface water environment. ES is an assessment of the structure and function of surface waters. ES accounts for the abundance of aquatic flora and fish fauna, the availability of nutrients, and aspects like salinity, temperature, and presence of chemical pollutants. Notably, ES includes benthic variables as well as water column conditions. As defined in the WFD, ES refers not to a specific level of a variable or a characteristic of an ecosystem but rather to a change from the baseline undisturbed state. Proxies of this scheme include but are not limited to total phosphorus, dissolved oxygen, water temperature, and macrophyte density; classifications include bad, poor, moderate, good, and high.

related to a TS classification, which is then used to predict ecosystem productivity, function, and integrity.

Over the last half century, the proxies used to classify TS have become synonymous with productivity, potentially leading to TS being, as Hutchinson (1957) warned, “[a] terminology that is so widely and often so inaccurately employed in discussing productivity.” For example, oligotrophic lakes and rivers are associated with low biological productivity, which is associated with low phosphorus and nitrogen concentrations. Consequently, nutrient concentrations become the defining feature of an ecosystem’s TS, rather than the biological productivity itself. This conceptual merger of biological productivity with its measured proxies can be beneficial for projecting ecological information across landscapes. However, it can also lead to confusion, where, for example, reference to a TS category may actually translate to value ranges of the proxy variable. In these instances, the same word in

reference to a given TS classification can create miscommunication, where multiple individuals may refer to the same TS classification but through the lenses of different proxies. Confusion can be further compounded when different classification schemes suggest diverging expectations for ecosystem function, such as characteristic oxyclines across various TS classification schemes throughout the contiguous United States (Figure 1). Regardless of how “oligotrophic” is defined, oxycline and thermocline profiles produce anticipated orthograde curves. In contrast, eutrophic profiles are visually similar, yet the most distinctive clinograde curves are observed in dystrophic and mixotrophic lakes (Figure 1), systems that are less well studied than eutrophic lakes (Leech et al., 2018). These incongruences between expected ecosystem functioning call into question the extent to which autotrophic-focused metrics, such as Trophic State Index (TSI), might discourage consideration of heterotrophic

TABLE 1 Aggregation of select trophic state classification schemes.

Index name	Data type	Geography	Season	Type	Zone sampled	TS group	Variables used and value limit	Source
TSI	Cont	Midwest United States	Summer	Lakes	Epi	Oligo	SDD, >4 TP, <12 Chl, <2.6	1
						Meso	SDD, 2–4 TP, 12–24 Chl, 2.6–7.3	
						Eu	SDD, 0.5–2 TP, 24–96 Chl, 7.3–56	
						Hypereu	SDD, <0.5 TP, >96 Chl, >56	
TSI	Cat Cont	Brazil	All	Reservoirs	Epi	Ultraoligo	TP, <15.9 Chl, <2	2
						Oligo	TP, 16–23.8 Chl, 2.1–3.9	
						Meso	TP, 23.9–36.7 Chl, 4–10	
						Eu	TP, 36.8–63.7 Chl, 4–10	
						Supereu	TP, 63.8–77.6 Chl, 20.3–27.1	
						Hupereu	TP, >77.6 Chl, >27.2	
TS	Cat Cont Prob	Worldwide	NA	Lakes and Reservoirs	Epi	Ultraoligo	TP, 0–4 Chl, < 1 Max Chl, <2.5 SDD, >12 Min SDD, >6	3
						Oligo	TP, 4–10 Chl, 1–2.5 Max Chl, 2.5–8 SDD, 6–12 Min SDD, 3–6	
						Meso	TP, 10–35 Chl, 2.5–8 Max Chl, 8–25 SDD, 3–6 Min SDD, 1.5–3	
						Eu	TP, 35–100 Chl, 8–25 Max Chl, 25–75 SDD, 1.5–3 Min SDD, 0.7–1.5	
						Hupereu	TP, >100 Chl, >25 Max Chl, >75 SDD, <1.5 Min SDD, <0.7	

TABLE 1 (Continued)

Index name	Data type	Geography	Season	Type	Zone sampled	TS group	Variables used and value limit	Source
TS	Cat Cont	Brazil	All	Reservoirs	Epi	Ultraoligo	TP, < 8 Chl, < 1.2	4
						Oligo	TP, 8.1–19 Chl, 1.3–3.2	
						Meso	TP, 19.1–52 Chl, 3.3–11	
						Eu	TP, 52.1–120 Chl, 11.1–30.6	
						Supereu	TP, 120.1–233 Chl, 30.7–69.1	
					Hupereu	TP, >233 Chl, >69.2		
TS	Cat Cont	Japan	All	Lakes	Epi	Oligo	TP, 2–20 TN, 20–200	5
						Meso	TP, 10–30 TN, 100–700	
						Eu	TP, 10–90 TN, 500–1300	
NCP	Cat	Maine, New Hampshire, Michigan, Wisconsin (United States)	Summer	Lakes	NA	Oligo	TP, <30 + Color, <20	6
						Eu	TP, >30 + Color, <20	
						Dys	TP, <30 + Color, >20	
						Mixo	TP, >30 + Color, >20	
NCP	Cat	United States, Argentina	Summer	Lakes	Epi	Oligo	TP, <10 + CDOC, <25	7
						Eu	TP, >10 + CDOC, <25	
						Dys	TP, <10 + CDOC, >25	
						Mixo	TP, >10 + CDOC, >25	
QI	Cat Cont	Hungary	All	Lakes	Epi	Excellent Good Medium Tolerable	PB, <1 PB, 1–4 PB, 4–8 PB, >8	8
PTSI	Cat Cont	Germany	All	Lakes and Reservoirs	Epi	Oligo	PTSI, <1.5	9
						Meso	PTSI, 1.5–2.5	
						Eu	PTSI, 2.5–3.5	
						Poly	PTSI, 3.5–4.5	
						Hyper	PTSI, >4.5	
TLI	Cat Cont	New Zealand	NA	Lakes	Epi	Ultramicro	Chl, <0.33 SDD, >25 TP, <1.8 TN, <34	10
						Micro	Chl, 0.33–0.82 SDD, 15–25 TP, 1.8–4.1 TN, 34–73	
						Oligo	Chl, 0.82–2.0	

(Continues)

TABLE 1 (Continued)

Index name	Data type	Geography	Season	Type	Zone sampled	TS group	Variables used and value limit	Source
							SDD, 15–7 TP, 4.1–9.0 TN, 73–157	
						Meso	Chl, 2–5 SDD, 2.8–7 TP, 9.0–20 TN, 157–337	
						Eu	Chl, 5–12 SDD, 1.1–2.8 TP, 20–43 TN, 337–725	
						Supereu	Chl, 12–31 SDD, 0.4–1.1 TP, 43–96 TN, 725–1558	
						Hypereu	Chl, >31 SDD, <0.4 TP, >96 TN, >1558	
SBTI	Cat Cont	New Zealand	NA	Lakes	Sed	Micro	SBTI, <2	11
						Oligo	SBTI, 2–3	
						Meso	SBTI, 3–4	
						Eu	SBTI, 4–5	
						Supereu	SBTI, 5–6	
						Hypereu	SBTI, >6	
TS	Cat Cont	Worldwide	Summer	Lakes	Epi Hypo	Oligo	TP, <10 TN, <350 Chl, <3.5 SDD, >4 AF, <20 AHOD, <250	12, 13
						Meso	TP, 10–30 TN, 350–650 Chl, 3.5–9 AF, 20–40 AHOD, 250–400	
						Eu	TP, 30–100 TN, 650–1200 Chl, 9–25 AF, 40–60 AHOD, 400–550	
						Hypereu	TP, >100 TN, >1200 Chl, >25 AF, >60 AHOD, >550	

TABLE 1 (Continued)

Index name	Data type	Geography	Season	Type	Zone sampled	TS group	Variables used and value limit	Source
TS	Cat Prob	Worldwide	All	Lotic	Ben	Oligo	TP, <25 TN, <325	14
						Meso	TP, 25–60 TN, 325–700	
						Eu	TP, >60 TN, >700	
TS	Cat Cont Prob	United States	Summer	Lotic	Ben	Oligo	GPP, <6.25 R, <119 GPP/R, <0.19	15
						Meso	GPP, 6.25–37.5 R, 119–243 GPP/R, 0.19–0.63	
						Eu	GPP, >37.5 R, >243 GPP/R, >0.63	
TS	Cat Cont Prob	United States	Summer	Lakes	Epi	Oligo	GPP, <13 R, <35 GPP/R, <0.68	15
						Meso	GPP, 13–34 R, 35–46 GPP/R, 0.67–0.9	
						Eu	GPP, >34 R, >46 GPP/R, >0.9	
TS	Cat Cont Prob	Worldwide	All	Lotic	Ben Ses	Oligo	Mean Benthic Chl, <20 Max Benthic Chl, <60 Sestonic Chl, <10 TP, <25 TN, <700	16
						Meso	Mean Benthic Chl, 20–70 Max Benthic Chl, 60–200 Sestonic Chl, 10–30 TP, 25–75 TN, 700–1500	
						Eu	Mean Benthic Chl, >70 Max Benthic Chl, >200 Sestonic Chl, >30 TP, >75 TN, >1500	
TS	Cat Cont	Argentina		Lakes	Epi	Ultraoligo	cDOM, <3	17
						Oligo	cDOM, 3–40	
						Meso	cDOM, 40–45	
						Eu	cDOM, >45	
TS	Cat Cont Prob	China		Lakes	Epi	Oligo	cDOM, <4	18
						Meso	cDOM, 4–10	

(Continues)

TABLE 1 (Continued)

Index name	Data type	Geography	Season	Type	Zone sampled	TS group	Variables used and value limit	Source
						Eu	cDOM, 10–23	
						Hypereu	cDOM, >23	

Note: Nürnberg (1996) likewise contains an extensive list of trophic state classification schemes, a portion of which are included in this list. Additionally, some trophic state indices mentioned in this table contain uncommon metrics, such as sediment bacterial abundance, but have complex formulations that do not necessarily fit the format of this table. For completeness, we added their reference and ranges to this table, but future users are encouraged to refer to the primary literature for those indices' calculations. Abbreviations in the "Index name" column are TSI, Trophic State Index; TS, trophic state; NCP, Nutrient Color Paradigm; QI, Q Index; PTSI, Phyto-See-Index; TLI, Trophic Level Index; SBTI, Sediment Bacterial Trophic Index. Abbreviations in the "Data type" column are Cont, continuous; Cat, categorical; Prob, probabilistic. Abbreviations in the "Zone sampled" column are Epi, epilimnion; Sed, sediment; Hypo, hypolimnion; Ben, benthic; Ses, seston. Abbreviations in the "TS group" column are Oligo, oligotrophic; Meso, mesotrophic; Eu, eutrophic; Hypereu, hypereutrophic; Ultraoligo, ultraoligotrophic; Supereu, supereutrophic; Poly, polytrophic; Ultramicro, ultramicrotrophic; Micro, microtrophic. Abbreviations in the "Variables used and value limits" column are TP, total phosphorus ($\mu\text{g/L}$); SDD, Secchi disk depth (m); Chl, chlorophyll *a* ($\mu\text{g/L}$); Max Chl, maximum chlorophyll *a* ($\mu\text{g/L}$); Min SDD, minimum Secchi disk depth (m); TN, total nitrogen ($\mu\text{g/L}$); Color, true color (platinum-cobalt units); CDOC, colored dissolved organic carbon (m^{-1}); PB, phytoplankton biomass (mg/L); PTSI, Phyto-See-Index; SBTI, Sediment Bacterial Trophic Index; AF, anoxic factor (no. days/summer); AHOD, areal hypolimnetic oxygen depletion ($\text{mg} \cdot \text{m}^{-2} \cdot \text{d}^{-1}$); GPP, gross primary production ($\text{mmol O}_2 \cdot \text{m}^{-2} \cdot \text{d}^{-1}$); R, respiration ($\text{mmol O}_2 \cdot \text{m}^{-2} \cdot \text{d}^{-1}$); Mean Benthic Chl, mean benthic chlorophyll *a* (mg/m^2); Max Benthic Chl, maximum benthic chlorophyll *a* (mg/m^2); Sestonic Chl, sestonic chlorophyll *a* ($\mu\text{g/L}$); cDOM, chromophoric dissolved organic matter (a_{254}). Numbers in the "Source" column are 1, Carlson (1977); 2, Cunha et al. (2013); 3, Vollenweider and Kerekes (1982); 4, Lamparelli (2004); 5, Sakamoto (1966); 6, Webster et al. (2008); 7, Williamson et al. (1999); 8, Padišák et al. (2006); 9, Mischke (2015); 10, Burns et al. (1999); 11, Wood et al. (2023); 12, Nürnberg (1996); 13, Nürnberg and Shaw (1998); 14, Dodds (2007); 15, Dodds and Cole (2007); 16, Dodds et al. (1998); 17, Gianello et al. (2024); 18, Zhang et al. (2018).

processes that likewise influence ecosystem patterns and processes. In this vein, clarifying language in TS classification has potential consequences for how both water quality conditions and ecosystem function are perceived.

Without an epistemological evolution, the fields of aquatic ecology and water quality management run the risk of developing divergent understandings of ecosystem function. Considering the pace and magnitude of climatic uncertainty, clarifying existing TS categories can allow for more standardized approaches to understanding the structure and function of aquatic ecosystems over past and future decades. To illustrate the usefulness of a clear and consistent, yet dynamic, conceptual framework for advancing the aquatic sciences, we detail three instances where clarifying the TS concept can guide aquatic research and management. In each case study, we highlight the clarity gained by combining emerging scientific themes, data streams, and technologies with the TS concept. Moreover, we demonstrate that conveying (1) the proxy variables and classification scheme employed, (2) the spatial and temporal domains of the proxy data, and (3) the ecosystem type considered are three concrete steps toward clarifying the TS concept. Communicating these pieces of information is an initial step in ensuring scientific reproducibility, thereby furthering the aquatic sciences, water resource management, and interdisciplinary collaboration.

CLARIFYING THE TS CONCEPT CAN ENHANCE OUR UNDERSTANDING OF AQUATIC ECOSYSTEMS ACROSS SEASONS AND BIOMES

Especially in northern, temperate regions, the rise of under-ice studies has expanded our understanding of biological productivity beyond open-water seasons (Hampton et al., 2015, 2017). Oligotrophic lakes, such as Lake Baikal (Kozhova & Izmet'eva, 1998), can experience multi-week, under-ice algal blooms that attain biomass comparable to eutrophic systems (Popovskaya, 2000). Conversely, eutrophic lakes may experience decreased primary production when light-blocking snow is located on ice or as ice becomes opaque (Garcia et al., 2019; Kivilä et al., 2023), which can drive wintertime chlorophyll concentrations towards levels typically observed in oligotrophic lakes (Kalinowska & Karpowicz, 2020). In both cases, TS classifications based on samples collected under-ice may differ from those based on samples collected during open-water conditions, which can alter interpretations of the dominant processes in the waterbody. Given surface waters' rapid warming (Huang et al., 2024; O'Reilly et al., 2015) and declining ice cover worldwide (Sharma et al., 2019), an emphasis on how wintertime dynamics influence TS classifications can help refine our expectations of ecosystem change. For example, diminishing ice cover over coming decades could reduce algal productivity by both hindering ice-obligate algal communities and suppressing

growth of open-water taxa, even though psychrophilic and psychrotolerant blooms can occur (Reinl et al., 2023). Therefore, ice loss may homogenize the anticipated

behaviors of eutrophic and oligotrophic water bodies during winter, with both ecosystems being similarly productive during ice-free winters and only diverging in




FIGURE 1 Legend on next page.

summer. Thus, clarifying the TS concept can present a conceptual framework upon which seasonal investigations of waterbody productivity lead to new hypotheses.

In contrast to temperate ecosystems, tropical aquatic ecosystems have less pronounced seasonal variation in temperature and photosynthetically available radiation but are highly driven by hydrological variation in the dry and wet seasons (Cunha et al., 2021). In this case, TS schemes from temperate lakes are inadequate in tropical lakes. The productivity of Brazilian lakes, for example, is influenced by lake water level, water column stability changes, allochthonous nutrient loading, and turbidity related to dry–wet seasonal shifts (Brighenti et al., 2024; Cunha et al., 2021; Gagliardi et al., 2019). This alternative framing of seasonality based on dry–wet periods rather than cold–warm periods diverges from anticipated conceptual models of how eutrophic and oligotrophic lakes function, such as via the buildup and breakdown of thermoclines and oxyclines (Figure 1). Consequently, these contrasting frameworks for characterizing intra-annual variation in productivity further complicate the comparability of TS assessments made across biomes when using a single classification system.

To clarify the TS concept, the spatial and temporal domains of classifications are especially important. If we assume that TS is based on cyclical degrees of autotrophy and heterotrophy (Wetzel, 2001), characterizing TS relative to an annual baseline is necessary. While estimates of TS solely based on summertime productivity in temperate lakes can be useful for water quality management, such estimates cannot always be extrapolated across seasons or ecosystems. As most TS assessments are currently based on summertime productivity, our current understanding of characteristic productivity is likely biased toward summertime conditions and temperate regions. Yet, the growth of our understanding of wintertime and tropical productivity highlights how important seasonality can be for holistically understanding and inferring waterbody productivity and function. As the field of aquatic sciences continues to expand into seasons and geographic locations that are less well represented in the

literature (Barbosa et al., 2023; Mejia et al., 2018; Rogers et al., 2023), we can build on decades of research using the TS concept to create opportunities for scaling the TS concept beyond the specific periods and biomes used to lay its foundations. Thus, we could further assess how aquatic ecosystems function, how this functionality varies regionally and seasonally, and how to contextualize regional ecosystems' functioning within global patterns.

CLARIFYING THE TS CONCEPT CAN INFORM FRESHWATER MANAGEMENT ACROSS AQUATIC ECOSYSTEMS

For management purposes, the language of TS has been used to classify water quality characteristics to represent ecosystem functions and services. For example, eutrophic conditions may be desirable for increasing fish production (Rast & Thornton, 1996). Conversely, hypereutrophic conditions may promote widespread anoxia, which can lead to fish kills. Even though TS is classified via a subset of proxy variables, the TS classification can imply a suite of generalizable expectations for ecosystem function and integrity. These expectations may not empower precise characterization across ecosystems or prescribe specific actions but can be used for flagging ecosystems for targeted investigation, thereby aiding managers to prioritize ecosystems for restoration efforts. When applied across landscapes and geopolitical boundaries, TS categories can aid managers as a high-level comparative and contextual tool to communicate ecosystem integrity without the need for relying on specific water quality parameters.

Because various TS classification schemes rely on different proxy variables (Table 1), categories detectable by each scheme can narrow the focus of how water quality is perceived and communicated. For example, managers may use Secchi disk depth to calculate TSI (Carlson, 1977), and then use TSI to identify waterbodies that are hypereutrophic and at greater risk of cyanobacterial

FIGURE 1 Daytime temperature and dissolved oxygen depth profiles for all lakes in the 2017 U.S. Environmental Protection Agency (EPA) National Lakes Assessment (NLA) sampling campaign across four trophic state (TS) classification schemes. Dissolved oxygen concentrations have been scaled relative to temperature values so that profiles approximately overlap, similar to those portrayed in Wetzel (2001). Lines are drawn using locally estimated scatterplot smoothing (loess) fits across all lakes within each TS classification, where confidence envelopes (gray bands) signify increasing uncertainty in the model fit. Broadly, oligotrophic lakes all demonstrate anticipated orthograde curves, whereas eutrophic lakes do not demonstrate distinctive clinograde curves relative to those observed in explicitly dystrophic, mixotrophic, or hypereutrophic lakes. Data for TS classifications come from the 2017 EPA NLA sampling campaign (U.S. Environmental Protection Agency, 2017a, 2017b). Further details on how this analysis was conducted, including information on the EPA NLA's sampling design, are included in Appendix S1 as well as the R scripts “strat_check.R” and “depth_profile_plots_condensed.R” in the companion software release (Meyer et al., 2025).

blooms. However, Secchi disk depth can also indicate high concentrations of suspended inorganic sediments or highly colored waters (Cunha et al., 2021). Such highly colored dystrophic lakes are not necessarily at a similar risk of cyanobacterial blooms and present a different suite of management implications despite having overlapping Secchi disk depth ranges with eutrophic lakes. This incongruence can be consequential for water quality estimates because mixotrophic lakes, as defined by the Nutrient Color Paradigm (NCP), also have an elevated risk of cyanobacterial blooms (Leech et al., 2018). Continental-scale data demonstrate that Secchi disk depth can fail to differentiate eutrophic, mixotrophic, and dystrophic lakes (Figure 2).

Translation across TS schemes can be challenging. For example, TSI-derived classifications do not identify dystrophic and mixotrophic states, unlike NCP, creating different interpretations of TS class occurrences at landscape scales (Figure 3). The Upper Midwest and Northern Appalachians (United States; Omernik, 1987) are two ecoregions that highlight extremes in TS classification differences (Figure 3). Among TSI-derived metrics using total phosphorus, Secchi disk depth, chlorophyll,

rotifer abundance, and crustacean zooplankton abundance, 41% and 35.4% of lakes in the Upper Midwest and Northern Appalachians would be classified as eutrophic, respectively, whereas NCP suggests 10.4% and 3.7% of lakes as eutrophic, respectively. Conversely, NCP suggests that 51.5% and 58.3% of lakes in the Upper Midwest and Northern Appalachians would be classified as dystrophic or mixotrophic, underscoring both the prevalence of high dissolved organic carbon (DOC) in these ecosystems and the potential for management to overlook a widespread water quality concern (Solomon et al., 2015).

Even when TS classification schemes are conceptually comparable, irregularities in sample collection may limit the extent to which TS classifications are interchangeable. In particular, samples necessary for linking classification schemes may not be co-located or collected contemporaneously. For example, TSI and ecological state (ES) can rely upon total phosphorus, Secchi disk depth, and chlorophyll measurements — measurements that may be more commonly collected relative to the true color or dissolved organic matter/carbon (DOM/DOC) measurements needed for the NCP (Box 1). TSI and ES, then, might allow for finer scale assessments, but less




FIGURE 2 Boxplots representing characteristic Secchi disk depths for lakes from the U.S. Environmental Protection Agency (EPA) 2012 and 2017 National Lakes Assessment (NLA) and their associated trophic categories as determined by Nutrient Color Paradigm (NCP) and Trophic State Index (TSI). Boxplots representing NCP-based categories have diagonal hatches, whereas boxplots lacking diagonal hatches represent TSI-based categories. Boxplot lines represent quartile statistics, and points outside lines represent outliers. Secchi disk depth, total phosphorus, and true color data come from the EPA NLA (U.S. Environmental Protection Agency, 2011, 2012, 2017a, 2017b). TSI delineations were made following guidelines in Carlson (1977). NCP delineations were made following thresholds established in Webster et al. (2008) and Leech et al. (2018). Further details on the EPA NLA sampling design are included in Appendix S1. This figure was created using the R script “secchi_depth_boxplots.R” provided in the companion software release (Meyer et al., 2025).




FIGURE 3 (A) Map and (B) aggregated percentages in Omernik Level III aggregated ecoregions of lake trophic state (TS) using various classification schemes. Classification schemes shown here are not necessarily the most common but are intended to reflect the diversity of potential schemes. Notably, TS classifications can produce dramatic differences in anticipated frequency and abundance of a given TS. For example, the Nutrient Color Paradigm (NCP) tends to identify far fewer eutrophic lakes in the Northern Appalachian (3.7% of lakes) and Upper Midwest (10.7%) ecoregions, relative to Trophic State Index (TSI)-based metrics (35.4% and 41% on average, respectively). TS classifications are based on chlorophyll a (TSI_{CHLa}), total phosphorus (TSI_{TP}), and Secchi disk depth (TSI_{SDD}), rotifer abundance (TSI_{ROT} ; Ejsmont-Karabin, 2012), and crustacean zooplankton abundance (TSI_{CR1} ; Ejsmont-Karabin & Karabin, 2013), as well as the NCP. Data for trophic state (TS) classifications come from the 2017 U.S. Environmental Protection Agency (EPA) National Lakes Assessment (NLA) sampling campaign (U.S. Environmental Protection Agency, 2017a, 2017b). Further details on the EPA NLA sampling design are included in Appendix S1. This figure was created using the R script “lts_refinement_map.R” provided in the companion software release (Meyer et al., 2025).

frequent true color or DOM data collection may be too coarse for comparative analyses. Given potential problems with translating across TS classification schemes,

clarifying the proxy data used to assign a TS classification ensures the interpretability of TS classifications. Further, ensuring clear and consistent proxy data across

classification schemes will improve our ability to quantify expected variability in TS over space and time, thereby optimizing successive management decisions.

CLARIFYING THE TS CONCEPT CAN BENEFIT INTERDISCIPLINARY COLLABORATION

Progress in the aquatic sciences has benefited from a suite of disciplines, and other disciplines reciprocally benefit from the aquatic sciences. Scientists and managers can maximize these benefits by providing greater consistency and clarity to avoid confusion in their applications across disciplines. In particular, a limitation for field-based methods is the capacity to upscale in situ observations across landscapes due to limited time, funding, and personnel. Emerging technologies and tools, such as remote sensing and machine learning techniques, have enabled progress in extending localized insights across continental and interannual scales. Although these techniques have historically focused on understanding structural components of aquatic ecosystems (e.g., mixing), they are beginning to address water quality and ecosystem functional change (Calamita et al., 2024). Consequently, these approaches may provide the most tractable paths for understanding macroscale water quality patterns if grounded in clear and sound limnology that guides interpretation.

Remote sensing is a pertinent example of a discipline that has implemented the TS concept to understand aquatic ecosystem change from local-to-global and monthly-to-decadal scales. For example, Wang et al. (2018), Gilarranz et al. (2022), and Sillen et al. (2024) quantified TSI based on chlorophyll *a* (TSI_{CHLA}) and its associated variability from remotely sensed surface reflectance for hundreds of lakes worldwide. Werther et al. (2021) and Meyer et al. (2024) also developed models to classify TS from remote sensing surface reflectance across broad spatial scales, but Meyer et al. (2024) used the NCP to classify TS, whereas Werther et al. (2021) used TSI_{CHLA} . Together, the range and number of remote sensing studies aimed at evaluating ecosystem productivity indicate the incorporation of the TS concept into remote sensing-based approaches, yet few studies consider how different TS classification schemes may hinder communication and comparison of results.

This lack of clarity could lead to diverging perceptions of ecosystem integrity across scales that are otherwise not feasible to manually sample, thereby undermining the strength of remote sensing's capacity to sample large spatial scales. For example, in situ data may indicate oligotrophic status for both a river and a lake, but the

characteristics of TS in those ecosystem types are different. In lakes, oligotrophic conditions are associated with low pelagic productivity in the water column, resulting in high reflectance values in the blue portion of the spectrum and lower values in the green portion. In rivers, oligotrophic conditions often are associated with high benthic productivity, resulting in high reflectance values in the green portion of the spectrum and lower values in the blue portion. Even within a given ecosystem type, varying TS formulations may mask water quality change. For example, dominant wavelength is a powerful remotely sensed metric that can suggest autotrophic productivity in lakes (Sillen et al., 2024; Topp, Pavelsky, Dugan, et al., 2021; Topp, Pavelsky, Stanley, et al., 2021; Yang et al., 2022) and rivers (Gardner et al., 2021). However, dominant wavelength can fail to capture organic matter concentrations in lakes, making dystrophic and mixotrophic lakes indistinguishable from oligotrophic and eutrophic lakes, respectively (Figure 4). Thus, insights made from dominant wavelength may be limited to assessing remotely sensed TSI-based classifications as opposed to NCP-based classifications. Because remote sensing of inland water quality has historically had weak connections with limnology, hydrology, and ecology (Bukata, 2013; Topp et al., 2020), there is potential for remote sensing science to obfuscate distinct ecological states, especially when both states may be referred to by the same word (e.g., "eutrophic"). If limnological expertise is not included in the modeling framework or considered when applying models across ecosystems, remotely sensed water quality observations may offer spurious conclusions about water quality distribution and trends.

Beyond remote sensing, the proliferation of data collection methods in aquatic ecosystems has substantially expanded, including automated buoys, platforms, and in situ sensors. This wealth of information from multiple sources has enabled the creation of diverse machine learning methods for better understanding complex aquatic ecosystem dynamics. However, in instances where TS classification schemes vary across training datasets or only focus on specific dimensions of water quality, machine learning methods may not robustly form generalizable models capable of classifying diverse aquatic ecosystems. When limnological knowledge is integrated into machine learning models, such as in knowledge-guided machine learning (Appling et al., 2022; Karpatne et al., 2024), consistent and well-documented classification schemes can aid in reflecting anticipated ecosystem dynamics. For example, Hanson et al. (2020) used knowledge-guided machine learning to model phosphorus dynamics in a lake over 20 years, where the integrated model replicated a downward trend in lake total phosphorus (TP) concentrations and, by




FIGURE 4 Legend on next page.

extension, reduction in eutrophy. Regardless of the exact implementation, consistent and clear communication of TS classification schemes used to develop training datasets would maximize the predictive accuracy of these data-driven models.

New information gathered via emerging technologies may deepen our understanding of aquatic ecosystem properties across scales but will also demand periodic re-evaluation of how TS classification is operationalized. Remote sensing and data-driven modeling can expand spatial and temporal domains that may be impractical to manually sample. Clarification of the TS concept can enable researchers using remote sensing and data-intensive modeling to take advantage of the rich history of limnological principles (Topp et al., 2020). Effectively integrating limnology and technical aspects of remote sensing and machine learning will require investment and cross-pollination of communities. This blending of communities will need to overcome disciplinary barriers and norms, including jargon, concepts, and even data formats. Many grassroots efforts, such as the Community for Data Science and Open Science in the Aquatic Sciences and “Hacking Limnology” Workshop Series (Meyer et al., 2021; Meyer & Zwart, 2020), are breaking down these divides. Given the rapid growth of remote sensing and machine learning techniques for water quality assessment, further clarification of TS can benefit the increasing interdisciplinarity of limnology by disentangling concepts for non-limnologists, thereby enabling broader uptake and insights.

MOVING FORWARD: CLARIFYING THE TS CONCEPT TO ADVANCE THE FRESHWATER SCIENCES

For many ecologists, TS is often the first conceptual framework for understanding aquatic ecosystems. Similarly, for many policymakers and water managers, TS is often a guiding paradigm to qualitatively evaluate

water quality and prioritize ecosystems for restoration (e.g., 33 U.S.C. 1252 et seq.; Carlson, 1977). Even though TS may not be as specific or prescriptive as individual ecosystem characteristics, TS provides a generalized conceptual framework to compress complex, interconnected processes into a single metric. TS then becomes a tool to infer ecosystem processes when data may be limited; however, there may be nuanced inconsistencies across applications. We highlight how TS can be a useful conceptual framework for evaluating ecosystem function, yet we demonstrate a clear need to re-evaluate and expand this conceptual framework to include less well sampled ecosystems, water quality parameters, biomes, and seasons. Without this clarification, miscommunication among limnologists and collaborators has the potential to offer spurious conclusions of water quality and ecosystem change and to limit scientific reproducibility.

Given divergent TS schemes and end-user needs, convergent understanding of TS will require at least three pieces of metadata: (1) the ecosystem type, (2) the proxy data, and (3) the spatial and temporal domains of the proxy data. Each component provides detail on how TS can be understood in a given ecosystem. The ecosystem type details the dominant processes at play within an ecosystem. The proxies used for a classification scheme detail how a TS classification is determined, its comparability to other ecosystems, and the balance of autotrophy and heterotrophy considered. Defining the spatial and temporal domains of the proxy data allows for nuanced understanding of a classification, where inferences can be conveyed based on the temporal scales, geographic regions, and biomes considered. When provided, these metadata could empower robust assertions of landscape-level patterns (Figure 3), consistency in usage (Figure 4), and even re-evaluation of core expectations for each TS (Figure 1). In cases when metadata are thoroughly documented, such as for the U.S. Environmental Protection Agency National Lakes Assessment, multiple TS classification schemes may be applied to more holistically understand implications of divergent definitions

FIGURE 4 Mean and standard deviation of dominant wavelengths and true color (platinum-cobalt units [PCU]) ranges across trophic state (TS) classifications. When assessed with Trophic State Index (TSI) groupings (A), mean dominant wavelength largely separates oligotrophic, mesotrophic, eutrophic, and hypereutrophic lakes, although substantial variation within and overlap between classifications exist. Unlike dominant wavelength, true color values for TSI classifications substantially overlap between classifications. When assessed with Nutrient Color Paradigm groupings (B), dominant wavelength separates autotrophic groups but masks dystrophic and mixotrophic lakes with oligotrophic and eutrophic lakes, respectively. When viewed across a gradient (C), dominant wavelength is a powerful metric for assessing autotrophic productivity, yet variation within groups can likely contribute noise to classification models trained on solely remotely sensed surface reflectance data. Data for TS classifications come from the 2017 U.S. Environmental Protection Agency (EPA) National Lakes Assessment (NLA) sampling campaign (U.S. Environmental Protection Agency, 2017a, 2017b). Further details on how this analysis was conducted, including information on the EPA NLA sampling design are included in Appendix S1 as well as the “rs_ts_differences.R” R script in the companion software release (Meyer et al., 2025).

across macroscales. Regardless of the level of detail given to each metadata criterion, communicating these pieces of information is an initial step forward in improving clarity among TS concepts.

Ultimately, how the TS concept is implemented will likely stem from the task at hand. In the case of management, TS may be used to characterize water quality, prioritize water bodies for additional investigation, and communicate those water quality characteristics to decision-makers. In the case of scientific investigations, varying classification schemes may be applied to fully characterize ecosystems occurring at a particular spatial or temporal scale. As a foundational concept, understanding trophic states expected over macroscales is necessary for a global understanding of freshwater systems that moves beyond that derived from northwestern Europe and northeastern North America (Dodds et al., 2019). A wealth of existing literature has already underscored the importance of including understudied regions (Santoso & Toruan, 2020) and seasons (Hampton et al., 2015), and increased data sharing and publication practices will likely expedite the pace and scope of understanding limnological function worldwide (sensu Wulder et al., 2012). Global change is pushing ecosystems into novel states, and understanding the characteristics, particularly trophic state, of fresh waters across a range of climate conditions is necessary to predict ecosystem trajectories.

AUTHOR CONTRIBUTIONS

Michael F. Meyer conceived the idea for the manuscript and provided leadership throughout its development. Michael F. Meyer, Benjamin M. Kraemer, Stephanie E. Hampton, Amina I. Pollard, Alexander K. Fremier, Tyler V. King, Robert Ladwig, Isabella A. Oleksy, Simon N. Topp, and Ludmila S. Brighenti contributed to the design of the manuscript. Michael F. Meyer and Benjamin M. Kraemer wrangled and harmonized data for the manuscript. Rachel M. Pilla, Benjamin M. Kraemer, and Michael F. Meyer contributed to table and figure development. All coauthors contributed either to writing or critically editing the manuscript.

AFFILIATIONS

¹U.S. Geological Survey, Oregon Water Science Center, Portland, Oregon, USA

²Center for Limnology, University of Wisconsin, Madison, Wisconsin, USA

³Department of Hydrology, University of Freiburg, Freiburg, Germany

⁴Department of Ecosystem Science and Sustainability, Colorado State University, Fort Collins, Colorado, USA

⁵Departamento de Hidráulica e Saneamento, Universidade de São Paulo, São Carlos, São Paulo, Brazil

⁶Division of Biology, Kansas State University, Manhattan, Kansas, USA

⁷Tahoe Environmental Research Center, University of California Davis, Davis, California, USA

⁸Department F.-A. Forel for Aquatic and Environmental Sciences, University of Geneva, Geneva, Switzerland

⁹Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

¹⁰Office of Water, U.S. Environmental Protection Agency, Washington, DC, USA

¹¹Department of Biology, York University, Toronto, Ontario, Canada

¹²School of the Environment, Washington State University, Pullman, Washington, USA

¹³U.S. Geological Survey, Observing Systems Division, Boise, Idaho, USA

¹⁴Department of Ecoscience, Aarhus University, Aarhus, Denmark

¹⁵Biological and Environmental Sciences Department, Longwood University, Farmville, Virginia, USA

¹⁶Biodiversity Division, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

¹⁷Institute for Arctic and Alpine Research, Colorado University–Boulder, Boulder, Colorado, USA

¹⁸Upstream Tech, Carrboro, North Carolina, USA

¹⁹School of Ocean Sciences, Bangor University, Anglesey, UK

²⁰Departamento de Ciências Naturais e da Terra, Universidade do Estado de Minas Gerais, Divinópolis, Minas Gerais, Brazil

²¹Esri, Redlands, California, USA

²²Bren School of Environment, University of California, Santa Barbara, California, USA

²³Natural Resources Department, Upper Skagit Indian Tribe, Sedro Woolley, Washington, USA

²⁴Department of Civil and Environmental Engineering, National University of Singapore, Singapore, Singapore

²⁵Surface Waters - Research and Management, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

²⁶Earth Sciences Department, Southern Methodist University, Dallas, Texas, USA

ACKNOWLEDGMENTS

We thank Matthew R. Brousil, Jack R. Eggleston, Matthew R.V. Ross, Dustin W. Kincaid, and Jacob A. Zwart for diverse creative support during the formation of this manuscript. We are very grateful to Craig E. Williamson, Linnea A. Rock, Bryan M. Maitland, and Paul C. Hanson for commenting on a previous version of this manuscript. Michael F. Meyer, Simon N. Topp, and Kate C. Fickas were supported by a Mendenhall Fellowship from the US Geological Survey Water Mission

Area. Rachel M. Pilla was supported by the US Department of Energy (DOE), Office of Energy Efficiency and Renewable Energy, Water Power Technologies Office, and Environmental Sciences Division at Oak Ridge National Laboratory (ORNL). ORNL is managed by UT-Battelle, LLC, for the US DOE under contract DE-AC05-00OR22725. IAO was supported by awards EPS-2019528 and DEB-2306895. DGFC thanks Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the research productivity grant (number 310844/2020-7). LSB thanks the Programa de Apoio Institucional a Pesquisa from the Universidade do Estado de Minas Gerais for the research productivity grant (PAPq-UEMG 11/2022). RIW was supported by a UKRI Natural Environment Research Council (NERC) Independent Research Fellowship (grant number NE/T011246/1). Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the US Government.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

All U.S. Environmental Protection Agency National Lakes Assessment data are available at <https://www.epa.gov/national-aquatic-resource-surveys/data-national-aquatic-resource-surveys>. R scripts necessary for producing figures (Meyer et al., 2025) are available from the US Geological Survey: <https://doi.org/10.5066/P14KNKHD>.

ORCID

Michael F. Meyer  <https://orcid.org/0000-0002-8034-9434>

Benjamin M. Kraemer  <https://orcid.org/0000-0002-3390-9005>

Carolina C. Barbosa  <https://orcid.org/0000-0002-6393-5730>

Davi G. F. Cunha  <https://orcid.org/0000-0003-1876-3623>

Walter K. Dodds  <https://orcid.org/0000-0002-6666-8930>

Stephanie E. Hampton  <https://orcid.org/0000-0003-2389-4249>

César Ordóñez  <https://orcid.org/0000-0001-6295-7690>

Rachel M. Pilla  <https://orcid.org/0000-0001-9156-9486>

Amina I. Pollard  <https://orcid.org/0000-0002-5010-0961>

Joshua A. Culpepper  <https://orcid.org/0000-0002-0468-0539>

Alexander K. Fremier  <https://orcid.org/0000-0002-4104-6633>

Tyler V. King  <https://orcid.org/0000-0002-5785-3077>

Robert Ladwig  <https://orcid.org/0000-0001-8443-1999>

Dina M. Leech  <https://orcid.org/0000-0002-0674-3433>

Shin-Ichiro S. Matsuzaki  <https://orcid.org/0000-0003-2744-1343>

Isabella A. Oleksy  <https://orcid.org/0000-0003-2572-5457>

Simon N. Topp  <https://orcid.org/0000-0001-7741-5982>

R. Iestyn Woolway  <https://orcid.org/0000-0003-0498-7968>

Ludmila S. Brighenti  <https://orcid.org/0000-0003-1305-2689>

Kate C. Fickas  <https://orcid.org/0000-0002-6617-2441>

Jianning Ren  <https://orcid.org/0000-0002-5849-2189>

Mortimer Werther  <https://orcid.org/0000-0002-0775-9285>

Xiao Yang  <https://orcid.org/0000-0002-0046-832X>

REFERENCES

- Appling, A. P., S. K. Oliver, J. S. Read, J. M. Sadler, and J. A. Zwart. 2022. "Machine Learning for Understanding Inland Water Quantity, Quality, and Ecology." In *Encyclopedia of Inland Waters*, 2nd ed., edited by T. Mehner and K. Tockner, 585–606. Oxford: Elsevier.
- Barbosa, C. C., L. A. Rock, A. N. Curtis, C. A. Sharitt, and P. Rogers. 2023. "Conducting Communication, Research, and Education from Climate Change Perspectives." *Limnology and Oceanography Bulletin* 32: 52–53.
- Brighenti, L. S., E. A. P. Viana, D. G. F. Pujoni, F. A. R. Barbosa, and J. F. Bezerra-Neto. 2024. "Post-Drought Leads to Increasing Metabolic Rates in the Surface Waters of a Natural Tropical Lake." *Frontiers in Geochemistry* 2: 1393444.
- Bukata, R. P. 2013. "Retrospection and Introspection on Remote Sensing of Inland Water Quality: 'Like Déjà Vu All Over Again'." *Journal of Great Lakes Research* 39: 2–5.
- Burns, N. M., J. C. Rutherford, and J. S. Clayton. 1999. "A Monitoring and Classification System for New Zealand Lakes and Reservoirs." *Lake and Reservoir Management* 15: 255–271.
- Calamita, E., J. J. Lever, C. Albergel, R. I. Woolway, and D. Odermatt. 2024. "Detecting Climate-Related Shifts in Lakes: A Review of the Use of Satellite Earth Observation." *Limnology and Oceanography* 69: 723–741.
- Carlson, R. E. 1977. "A Trophic State Index for Lakes." *Limnology and Oceanography* 22: 361–69.
- Cunha, D. G. F., M. d. C. Calijuri, and M. C. Lamparelli. 2013. "A Trophic State Index for Tropical/Subtropical Reservoirs (TSI_{tr})." *Ecological Engineering* 60: 126–134.
- Cunha, D. G. F., N. R. Finkler, M. C. Lamparelli, M. d. C. Calijuri, W. K. Dodds, and R. E. Carlson. 2021. "Characterizing Trophic State in Tropical/Subtropical Reservoirs: Deviations among Indexes in the Lower Latitudes." *Environmental Management* 68: 491–504.
- Directive 2000/60/EC of the European Parliament and of the Council (EU). 2000. *The EU Water Framework Directive*. Publications Office.
- Dodds, W. K. 2006. "Eutrophication and Trophic State in Rivers and Streams." *Limnology and Oceanography* 51: 671–680.
- Dodds, W. K. 2007. "Trophic State, Eutrophication and Nutrient Criteria in Streams." *Trends in Ecology & Evolution* 22: 669–676.

- Dodds, W. K., L. Bruckerhoff, D. Batzer, A. Schechner, C. Pennock, E. Renner, F. Tromboni, K. Bigham, and S. Grieger. 2019. "The Freshwater Biome Gradient Framework: Predicting Macroscale Properties Based on Latitude, Altitude, and Precipitation." *Ecosphere* 10: e02786.
- Dodds, W. K., and J. J. Cole. 2007. "Expanding the Concept of Trophic State in Aquatic Ecosystems: It's Not Just the Autotrophs." *Aquatic Sciences* 69: 427–439.
- Dodds, W. K., J. R. Jones, and E. B. Welch. 1998. "Suggested Classification of Stream Trophic State: Distributions of Temperate Stream Types by Chlorophyll, Total Nitrogen, and Phosphorus." *Water Research* 32: 1455–62.
- Dodds, W. K., and M. R. Whiles. 2019. *Freshwater Ecology: Concepts and Environmental Applications of Limnology*, 3rd ed. Burlington: Academic Press.
- Ejsmont-Karabin, J. 2012. "The Usefulness of Zooplankton as Lake Ecosystem Indicators: Rotifer Trophic State Index." *Polish Journal of Ecology* 60: 339–350.
- Ejsmont-Karabin, J., and A. Karabin. 2013. "The Suitability of Zooplankton as Lake Ecosystem Indicator: Crustacean Trophic State Index." *Polish Journal of Ecology* 61: 561–573.
- Gagliardi, L. M., L. S. Brighenti, P. A. Staehr, F. A. R. Barbosa, and J. F. Bezerra-Neto. 2019. "Reduced Rainfall Increases Metabolic Rates in Upper Mixed Layers of Tropical Lakes." *Ecosystems* 22: 1406–23.
- Garcia, S. L., A. J. Szekely, C. Bergvall, M. Schattener, and S. Peura. 2019. "Decreased Snow Cover Stimulates Under-Ice Primary Producers but Impairs Methanotrophic Capacity." *mSphere* 4. <https://doi.org/10.1128/msphere.00626-18>.
- Gardner, J. R., X. Yang, S. N. Topp, M. R. V. Ross, E. H. Altenau, and T. M. Pavelsky. 2021. "The Color of Rivers." *Geophysical Research Letters* 48: e2020GL088946.
- Gianello, D., M. Reissig, G. L. Pérez, M. Rechencq, L. G. De Stefano, and C. Queimaliños. 2024. "Effects of Water Level Fluctuations on the Trophic State of a Connected Lacustrine System (Southern Andes, Patagonia, Argentina): Applicability of the CDOM Method as a Tool for Monitoring of Eutrophication." *Aquatic Sciences* 86: 1–25.
- Gilarranz, L. J., A. Narwani, D. Odermatt, R. Siber, and V. Dakos. 2022. "Regime Shifts, Trends, and Variability of Lake Productivity at a Global Scale." *Proceedings of the National Academy of Sciences of the United States of America* 119: e2116413119.
- Hampton, S. E., A. W. E. Galloway, S. M. Powers, T. Ozersky, K. H. Woo, R. D. Batt, S. G. Labou, et al. 2017. "Ecology under Lake Ice." *Ecology Letters* 20: 98–111.
- Hampton, S. E., M. V. Moore, T. Ozersky, E. H. Stanley, C. M. Polashenski, and A. W. E. Galloway. 2015. "Heating up a Cold Subject: Prospects for under-Ice Plankton Research in Lakes." *Journal of Plankton Research* 37: 277–284.
- Hanson, P. C., A. B. Stillman, X. Jia, A. Karpatne, H. A. Dugan, C. C. Carey, J. Stachelek, et al. 2020. "Predicting Lake Surface Water Phosphorus Dynamics Using Process-Guided Machine Learning." *Ecological Modelling* 430: 109136.
- Horne, A. J., and C. R. Goldman. 1983. *Limnology*, 1st ed. New York: McGraw-Hill.
- Huang, L., R. I. Woolway, A. Timmermann, S.-S. Lee, K. B. Rodgers, and R. Yamaguchi. 2024. "Emergence of Lake Conditions that Exceed Natural Temperature Variability." *Nature Geoscience* 17: 763–69.
- Hutchinson, G. E. 1957. *A Treatise on Limnology*, Vol. 1. New York: Wiley.
- Kalinowska, K., and M. Karpowicz. 2020. "Ice-on and Ice-off Dynamics of Ciliates and Metazooplankton in the Łuczański Canal (Poland)." *Aquatic Ecology* 54: 1121–34.
- Karpatne, A., X. Jia, and V. Kumar. 2024. "Knowledge-Guided Machine Learning: Current Trends and Future Prospects." arXiv.
- Kivilä, E. H., V. Prėskienis, N. Gaudreault, C. Girard, and M. Rautio. 2023. "Variability in Lake Bacterial Growth and Primary Production under Lake Ice: Evidence from Early Winter to Spring Melt." *Limnology and Oceanography* 68: 2603–16.
- Kozhova, O. M., and L. R. Izmes'teva. 1998. *Lake Baikal: Evolution and Biodiversity*, 2nd ed. Leiden: Backhuys Publishers.
- Kraemer, B. M. 2020. "Rethinking Discretization to Advance Limnology amid the Ongoing Information Explosion." *Water Research* 178: 115801.
- Lamparelli, M. C. 2004. "Graus de trofia em corpos d'água do estado de São Paulo: avaliação dos métodos de monitoramento." Dissertation, Universidade de São Paulo.
- Leech, D. M., A. I. Pollard, S. G. Labou, and S. E. Hampton. 2018. "Fewer Blue Lakes and more Murky Lakes across the Continental U.S.: Implications for Planktonic Food Webs." *Limnology and Oceanography* 63: 2661–80.
- Lindeman, R. L. 1942. "The Trophic-Dynamic Aspect of Ecology." *Ecology* 23: 399–417.
- Mejia, F. H., A. K. Fremier, J. R. Benjamin, J. R. Bellmore, A. Z. Grimm, G. A. Watson, and M. Newsom. 2018. "Stream Metabolism Increases with Drainage Area and Peaks Asynchronously across a Stream Network." *Aquatic Sciences* 81: 9.
- Meyer, M. F., R. Ladwig, J. P. Mesman, I. A. Oleksy, C. C. Barbosa, K. M. Cawley, A. N. Cramer, et al. 2021. "The AEMON-J 'Hacking Limnology' Workshop Series & Virtual Summit: Incorporating Data Science and Open Science in Aquatic Research." *Limnology and Oceanography Bulletin* 30: 140–43.
- Meyer, M. F., R. L. Ladwig, and R. M. Pilla. 2025. *Clarifying the Trophic State Concept to Advance Macroscale Freshwater Science and Management*. Reston, VA: U.S. Geological Survey Software Release. <https://doi.org/10.5066/P14KNKHD>.
- Meyer, M. F., S. N. Topp, T. V. King, R. Ladwig, R. M. Pilla, H. A. Dugan, J. R. Egleston, et al. 2024. "National-Scale Remotely Sensed Lake Trophic State from 1984 through 2020." *Scientific Data* 11: 77.
- Meyer, M. F., and J. A. Zwart. 2020. "Virtual Summit: Incorporating Data Science and Open Science in Aquatic Research." *Limnology and Oceanography Bulletin* 29: 144–46.
- Mischke, U. 2015. "Handbuch Für Die Seenbewertung Mittels Plankton – Phyto-See-Index (Teil A) Und Phytoloss-Modul Zooplankton (TEIL B) Stand 27.02.2015." Leibniz-Institut für Gewässerökologie und Binnenfischerei.
- Naumann, E. 1917. "Undersökningar över fytoplankton och under den pelagiska regionen fösiggäende gyttje-och dybildningar inom vissa syd- och mellansvenska urbergsvatten." *Kungliga Svenska Vetenskapsakademiens Handlingar* 56: 1–165.
- Nojvan, A. F., B. J. Kreakie, J. W. Hollister, and S. S. Qian. 2019. "Rethinking the Lake Trophic State Index." *PeerJ* 7: e7936.
- Nürnberg, G. K. 1996. "Trophic State of Clear and Colored, Soft-and Hardwater Lakes with Special Consideration of Nutrients, Anoxia, Phytoplankton and Fish." *Lake and Reservoir Management* 12: 432–447.

- Nürnberg, G. K., and M. Shaw. 1998. "Productivity of Clear and Humic Lakes: Nutrients, Phytoplankton, Bacteria." *Hydrobiologia* 382: 97–112.
- Odum, H. T. 1956. "Primary Production in Flowing Waters." *Limnology and Oceanography* 1: 102–117.
- Omernik, J. M. 1987. "Ecoregions of the Conterminous United States." *Annals of the Association of American Geographers* 77: 118–125.
- O'Reilly, C. M., S. Sharma, D. K. Gray, S. E. Hampton, J. S. Read, R. J. Rowley, P. Schneider, et al. 2015. "Rapid and Highly Variable Warming of Lake Surface Waters around the Globe." *Geophysical Research Letters* 42: 10,773–10,781.
- Padisák, J., G. Borics, I. Grigorczyk, and É. Soróczki-Pintér. 2006. "Use of Phytoplankton Assemblages for Monitoring Ecological Status of Lakes within the Water Framework Directive: The Assemblage Index." *Hydrobiologia* 553: 1–14.
- Palmer, M. A., and C. M. Febria. 2012. "The Heartbeat of Ecosystems." *Science* 336: 1393–94.
- Popovskaya, G. I. 2000. "Ecological Monitoring of Phytoplankton in Lake Baikal." *Aquatic Ecosystem Health and Management* 3: 215–225.
- Rast, W., and J. A. Thornton. 1996. "Trends in Eutrophication Research and Control." *Hydrological Processes* 10: 295–313.
- Reinl, K. L., T. D. Harris, R. L. North, P. Almela, S. A. Berger, M. Bizic, S. H. Burnet, et al. 2023. "Blooms Also like it Cold." *Limnology and Oceanography Letters* 8: 546–564.
- Rogers, P., C. C. Barbosa, A. N. Curtis, L. A. Rock, and C. A. Sharitt. 2023. "Uncited Science in the Global South: Striving toward Inclusivity and Bettering our Understanding of Global Climate Change." *Limnology and Oceanography Bulletin* 32: 54–55.
- Rohde, W. 1969. "Crystallization of Eutrophication Concepts in Northern Europe." In *Eutrophication: Causes, Consequences, Correctives* 20256. Washington, DC: National Academies Press.
- Sakamoto, M. 1966. "Primary Production by Phytoplankton Community in some Japanese Lakes and Its Dependence on Lake Depth." *Archiv für Hydrobiologie* 62: 1–28.
- Santoso, A. B., and R. L. Toruan. 2020. "Limnology in the Tropics: Next Steps?" *IOP Conference Series: Earth and Environmental Science* 535: 012007.
- Sharma, S., K. Blaggrave, J. J. Magnuson, C. M. O'Reilly, S. Oliver, R. D. Batt, M. R. Magee, et al. 2019. "Widespread Loss of Lake Ice around the Northern Hemisphere in a Warming World." *Nature Climate Change* 9: 227–231.
- Sillen, S. J., M. R. V. Ross, and S. M. Collins. 2024. "Long-Term Trends in Productivity across Intermountain West Lakes Provide No Evidence of Widespread Eutrophication." *Water Resources Research* 60: e2023WR034997.
- Solomon, C. T., S. E. Jones, B. C. Weidel, I. Buffam, M. L. Fork, J. Karlsson, S. Larsen, et al. 2015. "Ecosystem Consequences of Changing Inputs of Terrestrial Dissolved Organic Matter to Lakes: Current Knowledge and Future Challenges." *Ecosystems* 18: 376–389.
- Thienemann, A. 1921. "Seetypen." *Naturwissenschaften* 9: 343–46.
- Topp, S. N., T. M. Pavelsky, H. A. Dugan, X. Yang, J. Gardner, and M. R. V. Ross. 2021. "Shifting Patterns of Summer Lake Color Phenology in over 26,000 US Lakes." *Water Resources Research* 57: e2020WR029123.
- Topp, S. N., T. M. Pavelsky, D. Jensen, M. Simard, and M. R. V. Ross. 2020. "Research Trends in the Use of Remote Sensing for Inland Water Quality Science: Moving Towards Multidisciplinary Applications." *Water* 12: 169.
- Topp, S. N., T. M. Pavelsky, E. H. Stanley, X. Yang, C. G. Griffin, and M. R. V. Ross. 2021. "Multi-Decadal Improvement in US Lake Water Clarity." *Environmental Research Letters* 16: 055025.
- U.S. Environmental Protection Agency. 1990. *Lake and Reservoir Restoration Guidance Manual*. Washington, DC: U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2011. *2012 National Lakes Assessment. Field Operations Manual*. Washington, DC: U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2012. *National Lakes Assessment. Laboratory Operations Manual*. Washington, DC: U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2017a. *National Lakes Assessment 2017. Laboratory Operations Manual. V.1.1*. Washington, DC: U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2017b. *National Lakes Assessment 2017. Field Operations Manual*. Washington, DC: U.S. Environmental Protection Agency.
- Vollenweider, R. A., and J. Kerekes. 1982. *Eutrophication of Waters: Monitoring, Assessment, and Control* 155. Paris: Environment Directorate, OECD.
- Wang, S., J. Li, B. Zhang, E. Spyarakos, A. N. Tyler, Q. Shen, F. Zhang, et al. 2018. "Trophic State Assessment of Global Inland Waters Using a MODIS-Derived Forel-Ule Index." *Remote Sensing of Environment* 217: 444–460.
- Webster, K. E., P. A. Soranno, K. S. Cheruvellil, M. T. Bremigan, J. A. Downing, P. D. Vaux, T. R. Asplund, L. C. Bacon, and J. Connor. 2008. "An Empirical Evaluation of the Nutrient-Color Paradigm for Lakes." *Limnology and Oceanography* 53: 1137–48.
- Werther, M., E. Spyarakos, S. G. H. Simis, D. Odermatt, K. Stelzer, H. Krawczyk, O. Berlage, P. Hunter, and A. Tyler. 2021. "Meta-Classification of Remote Sensing Reflectance to Estimate Trophic Status of Inland and Nearshore Waters." *ISPRS Journal of Photogrammetry and Remote Sensing* 176: 109–126.
- Wetzel, R. G. 2001. *Limnology: Lake and River Ecosystems*, 3rd ed. San Diego: Academic Press.
- Williamson, C. E., D. P. Morris, M. L. Pace, and O. G. Olson. 1999. "Dissolved Organic Carbon and Nutrients as Regulators of Lake Ecosystems: Resurrection of a more Integrated Paradigm." *Limnology and Oceanography* 44: 795–803.
- Wood, S. A., M. J. Vandergoes, J. Atalah, J. D. Howarth, S. Waters, G. Thomson-Laing, L. Thompson, et al. 2023. "A National-Scale Trophic State Analysis to Prioritize Lakes for Restoration in Aotearoa New Zealand: Kilham Memorial Lecture on Occasion of the 100th Anniversary of SIL." *Inland Waters* 13(3): 303–315.
- Wulder, M. A., J. G. Masek, W. B. Cohen, T. R. Loveland, and C. E. Woodcock. 2012. "Opening the Archive: How Free Data Has Enabled the Science and Monitoring Promise of Landsat." *Remote Sensing of Environment* 122: 2–10.
- Yang, X., C. M. O'Reilly, J. R. Gardner, M. R. V. Ross, S. N. Topp, J. Wang, and T. M. Pavelsky. 2022. "The Color of Earth's Lakes." *Geophysical Research Letters* 49: e2022GL098925.

Zhang, Y., Y. Zhou, K. Shi, B. Qin, X. Yao, and Y. Zhang. 2018. "Optical Properties and Composition Changes in Chromophoric Dissolved Organic Matter along Trophic Gradients: Implications for Monitoring and Assessing Lake Eutrophication." *Water Research* 131: 255–263.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Meyer, Michael F., Benjamin M. Kraemer, Carolina C. Barbosa, Davi G. F. Cunha, Walter K. Dodds, Stephanie E. Hampton, César Ordóñez, et al. 2025. "Clarifying the Trophic State Concept to Advance Macroscale Freshwater Science and Management." *Ecosphere* 16(9): e70392. <https://doi.org/10.1002/ecs2.70392>