



Antifungal activity of *Rhopalurus crassicauda* venom against *Candida* spp.

Umberto Zottich^{a,*,1}, Isadora Sousa de Oliveira^{b,1}, Isabela Gobbo Ferreira^b, Felipe Augusto Cerni^c, Bordon Karla de Castro Figueiredo^b, Eliane Candiani Arantes^b, Valdirene Moreira Gomes^d, Germana Bueno Dias^c, Manuela Berto Pucca^{a,c,*}

^a Medical School, Federal University of Roraima, Boa Vista, Roraima, Brazil

^b Department of BioMolecular Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, São Paulo, Brazil

^c Health Sciences Postgraduate Program, Federal University of Roraima, Boa Vista, Brazil

^d Centro de Biociências e Biotecnologia, Universidade Estadual Do Norte Fluminense, 28013-602, Campos Dos Goytacazes, Rio de Janeiro, Brazil

ARTICLE INFO

Handling Editor: Glenn King

Keywords:

Rhopalurus crassicauda
scorpion Toxin
Candida albicans
Candida parapsilosis
Antifungal

ABSTRACT

Fungal infections are becoming a serious problem of human diseases, being one of the most important fungal pathogens the yeast of the genus *Candida*. So far, fungal infection treatment faces different challenges, including the limited number of therapeutic drugs. Scorpions are known to be a valuable source of biologically active molecules, especially of peptide-derived molecules with a variety of biological effects and useful, lead compounds for drugs development. Here, we pioneer described the antifungal effect of venom, mucus, and the major toxin (Rc1) from *Rhopalurus crassicauda* scorpion. These results support the potential for Rc1 to be further investigated as a novel antifungal therapeutic to treat *Candida* infections.

Credit author statement

UZ performed the experiments and participated in the data interpretation. ISO interpreted the data and drafted the manuscript. IGF, FAC, KCFB, ECA, VMG and GBD participated in the data interpretation and wrote the manuscript. MBP coordinated and designed all the experiments, analyzed and interpreted the data. All authors read, corrected, and approved the final manuscript.

Candida albicans as well as other emerging non albicans species, including *Candida glabrata*, *Candida krusei*, *Candida tropicalis*, and *Candida parapsilosis*, represent an important source of systemic infections throughout the globe. The body sites that *Candida* spp. preferably colonizes are the oropharynx, vagina, and skin, especially skin folds (axilla, groin, perineum). It can produce proteolytic enzymes (such as aspartyl proteinases and phospholipases) and toxins (such as mycotoxin) that affect host defenses and enhance their chances of colonizing and invading the host (De Bernardis et al., 1990; Schaller et al., 2003). Also, under favorable conditions, they can enter the bloodstream leading to deep-tissue infections (Conti et al., 2014; Turner and Butler, 2014; Whaley et al., 2016).

Fungal pathogens are a major cause of morbidity and mortality among several patient groups, including those hospitalized in the intensive care unit, solid organ and stem cell transplant recipients, and human immunodeficiency virus (HIV) infected patients. Also, there is a deepening appreciation of genetic risk factors, which may predispose to complications from fungal pathogens that occur among otherwise normal hosts. Most researches regarding novel antifungals have focused on patients who are known to be at greatest risk of invasive fungal infection, since the opportunistic mycoses are a cause of great concern to both clinicians and investigators (Pappas, 2010). The traditional pathogens in this group include *Aspergillus* species, *Cryptococcus* species, the zygomycetes, *Fusarium* species, and *Candida* species (Pfaller and Diekema, 2010).

Fungal pathogens cause life-threatening invasive diseases (e.g. fungaemia, meningitis, pneumonia), severe chronic conditions (e.g. chronic pulmonary aspergillosis, allergic bronchopulmonary aspergillosis), complex chronic respiratory conditions (e.g. asthma, chronic obstructive pulmonary disease) and recurrent infections, such as oral and vaginal candidiasis (Brown et al., 2012). In general, these infections are associated with high mortality, and successful clinical outcome requires

* Corresponding author. Medical School, Federal University of Roraima, Boa Vista, Roraima, Brazil.

** Corresponding author.

E-mail addresses: umberto.zottich@ufr.br (U. Zottich), manu.pucca@ufr.br (M.B. Pucca).

¹ These authors also contributed equally to this study.




Fig. 1. Inhibition of *C. albicans*. Kinetic growth inhibition by (A) *R. crassicauda* venom (100 µg/mL), (B) Rc1 (100 µg/mL, corresponding to 14 µM), and (C) *R. crassicauda* mucus (100 µg/mL). (D) Growth inhibition after 24 h. Non-treated cells were used as controls. All experiments were performed in triplicate and data are expressed as the mean $\pm$ standard deviation. *** $p < 0.001$ when compared to control.

early diagnosis and effective antifungal therapy. However, antimycotic drugs are few, with chemical classes for invasive disease treatment limited to azoles, echinocandins, polyenes, and flucytosine (Odds et al., 2003).

Scorpion venoms are known source of components able to inhibit microorganism growing. In fact, many antibacterial and antifungal compounds have been isolated from scorpion venoms (Almaaytah et al., 2012; Machado et al., 2016). In a previous work, authors evaluated 11 scorpion-venom derived non-disulfide-bridged peptides which demonstrated activity against *Cryptococcus neoformans*, *Candida* spp., and analogous activity against *C. albicans* biofilms (Guilhelmelli et al., 2016).

From *Tityus* species, *Tityus serrulatus* venom (Tsv) was capable of inhibiting filamentous fungi growth of *Aspergillus nidulans*, *Aspergillus terreus*, *Penicillium corylophilum*, and *Penicillium verrucosum* (Santussi et al., 2017). Ts1, the major toxin from Tsv, showed a dose-dependent inhibition against *A. nidulans* (Santussi et al., 2017). From *Tityus stigmurus*, two peptides present activity against fungi strains, stigmurin which has the ability to inhibit growth of *C. albicans*, *C. krusei* and *C. glabrata* (de Melo et al., 2015), and TistH was analyzed towards its antifungal properties against different strains of *C. albicans*, *C. tropicalis*, *A. flavus* and the filamentous fungus *Trichophyton rubrum* (Machado et al., 2016).

Inhabiting Amazonian savannah in the Brazilian northern and Guyana Southern (di Caporiacco, 1947; Martins et al., 2021), the scorpion *Rhopalurus crassicauda* Lourenço, 2002 (synonym *Rhopalurus amazonicus* Lourenço, 1986 and *Rhopalurus laticauda* Thorell, 1876) (Esposito et al., 2017; Martins et al., 2021) is a poorly studied species. There is only one study developed by Abreu et al. (2020) revealing the structure and function of the major toxin (Rc1 with ~ 7 kDa) and the isolation of a hyaluronidase. The same study also explored some activities of the whole venom (e.g. nociception and enzymatic activities) (Abreu et al., 2020). However, no antifungal activity has been explored so far. Based on these facts, this study aimed to describe a novel antifungal activity for

R. crassicauda scorpion venom.

R. crassicauda scorpions were collected in Boa Vista - RR (Brazil, 2°49'14.88" N and 60°40'19.20" W). The species were kept at Medical School of Federal University of Roraima, received water daily and were fed with cockroaches or crickets twice a month, normally, 5 days prior to venom milking. Venom milking was performed by electrical stimulation of venom gland using electrical pulses (18 V) (Abreu et al., 2020). Pooled venom was stored at -20°C until its use.

All the procedures involving the scorpions were in accordance with the ethical principles in animal research adopted by Sistema de Autorização e Informação em Biodiversidade (SISBIO), under registration number 57491/9.

The soluble crude venom (supernatant without mucus - 2 mg of proteins) was obtained, applied on a reversed-phase C18 column (10 mm $\times$ 250 mm) and the major peak was rechromatographed on another C18 column (2.1 mm $\times$ 250 mm) to obtain Rc1, as described by Abreu et al. (2020). The soluble venom and the toxin Rc1 (which represents 24% of the total protein of the soluble venom) were submitted to antifungal assays.

Yeast cells *C. albicans* (CE022) and *C. parapsilosis* (CE002) were preserved in the Laboratório de Fisiologia e Bioquímica de Microrganismos (LFBM), Universidade Estadual do Norte Fluminense Darcy Ribeiro (UENF), Campos dos Goytacazes, RJ, Brazil, were cultured in Sabouraud broth and quantified in a Neubauer chamber. To monitor the effect of proteins on the growth of yeasts, cells (10^4 in 1 mL Sabouraud broth) were incubated in the presence of the venom (10, 50 and 100 µg/mL), toxin Rc1 or mucus (100 µg/mL), at 28°C in 200 µL microplates (Broekaert et al., 1990). The positive control was performed with fluconazole 100 µg/mL, getting 100% inhibition (data not shown).

Through the growth kinetics, it was observed that 100 µg/mL of *R. crassicauda* venom or Rc1 venom (representing 14 µM) were statistically able to inhibit the growth of *C. albicans* after 24 h, differently from mucus, which did not show the ability to inhibit the growth of the microorganism (Fig. 1A–D).




Fig. 2. Inhibition of *C. parapsilosis*. Kinetic growth inhibition by (A) *R. crassicauda* venom (100 µg/mL), (B) Rc1 (100 µg/mL, corresponding to 14 µM), and (C) *R. crassicauda* mucus (100 µg/mL). (D) Growth inhibition after 24 h. Non-treated cells were used as controls. All experiments were performed in triplicate and data are expressed as the mean $\pm$ standard deviation. *** $p < 0.001$ when compared to control.

When the same treatment was performed on *C. parapsilosis*, it was observed that *R. crassicauda* venom and mucus, as well as Rc1, were statistically able to inhibit its growth in 24 h, except for the 10 µg/mL of venom (Fig. 2A–D). When at a concentration of 50 µg/mL, *C. albicans* has a higher growth than the control (Fig. 1A). Dimorphic transitions with the development of pseudo hyphae could explain our failure to detect growth inhibition in this yeast species: pseudo hyphae formation might disrupt absorption measurements (Zottich et al., 2011).

Rc1 showed inhibitory activity in the order of magnitude of micromolar concentration, such as those reported for scorpion venom toxins. Serrulin, a glycine-rich native peptide of the *T. serrulatus* scorpion hemolymph, showed the minimal inhibitory concentration (MIC) of 1.5–3 µM and 3–6 µM against *C. albicans* and *Aspergillus niger*, respectively (Oliveira et al., 2019). Ts1, the major toxin from *T. serrulatus* venom, showed 100% inhibition against *A. nidulans* from 4.36 µM (Santussi et al., 2017). Stigmurin from *T. stigmurus* venom showed MIC of 37.5 µM against *C. albicans*, and its analogues of 4.69–9.38 µM (Parente et al., 2018). Other manuscripts report that the antifungal activity depends on fungal strain (de Melo et al., 2015; Guilhaumelli et al., 2016; Machado et al., 2016) and antifungal agent (Ahmadi et al., 2020). The scorpion venom D-amino acid analogue dKn2–7 from *Mesobuthus martensii* showed an amphotericin B-like killing kinetics for *C. albicans* with rapid onset of antifungal activity (Snyder et al., 2021). It is interesting to note that *R. crassicauda* venom showed inhibitory activity against *C. albicans* and *C. parapsilosis* at a concentration 10–50 times lower than that reported for *T. serrulatus* venom (1 and 5 mg/mL) against the growth of *A. nidulans*, *A. terreus*, *P. corylophilum*, and *P. verrucosum* (Santussi et al., 2017).

In conclusion, this study pioneer investigated the venom, mucus, and the major toxin of *R. crassicauda* scorpion for antifungal potential against *C. albicans* and *C. parapsilosis* strains. The isolated toxin, Rc1, exhibited the most potent antifungal efficacy against both strains tested. Future studies involving a wider range of *R. crassicauda* venom-derived toxins and analysis of peptide effects on host tissues are required to better determine if the scorpion venom could be a source of potential

and novel antifungal drugs.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, São Paulo Research Foundation; scholarship to ISO no. 2020/13176-3), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, The National Council for Scientific and Technological Development, scholarship to MBP no. 307184/2020-0), and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES, Finance Code 001, scholarship to IGF). MBP acknowledges Prospecção PRPPG-UFRR (Edital 16/2020).

References

- Abreu, C.B., Bordon, K.C.F., Cerni, F.A., Oliveira, I.S., Balenzuela, C., Alexandre-Silva, G. M., Zoccal, K.F., Reis, M.B., Wiesel, G.A., Peigneur, S., Pinheiro-Júnior, E.L., Tytgat, J., Cunha, T.M., Quinton, L., Faccioli, L.H., Arantes, E.C., Zottich, U., Pucca, M.B., 2020. Pioneering study on *Rhopalurus crassicauda* scorpion venom: isolation and characterization of the major toxin and hyaluronidase. *Front. Immunol.* 11 <https://doi.org/10.3389/fimmu.2020.02011>.
- Ahmadi, S., Knerr, J.M., Argemi, L., Bordon, K.C.F., Pucca, M.B., Cerni, F.A., Arantes, E. C., Çalışkan, F., Laustsen, A.H., 2020. Scorpion venom: detriments and benefits. *Biomedicines* 8, 118. <https://doi.org/10.3390/biomedicines8050118>.
- Almaaytah, A., Zhou, M., Wang, L., Chen, T., Walker, B., Shaw, C., 2012. Antimicrobial/cytolytic peptides from the venom of the North African scorpion, *Androctonus amoreuxi*: biochemical and functional characterization of natural peptides and a single site-substituted analog. *Peptides* 35, 291–299. <https://doi.org/10.1016/j.peptides.2012.03.016>.
- Broekaert, W.F., Terras, F.R.G., Cammue, B.P.A., Vanderleyden, J., 1990. An automated quantitative assay for fungal growth inhibition. *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Lett.* 69, 55–59. <https://doi.org/10.1111/j.1574-6968.1990.tb04174.x>.

- Brown, G.D., Denning, D.W., Gow, N.A.R., Levitz, S.M., Netea, M.G., White, T.C., 2012. Hidden killers: human fungal infections. *Sci. Transl. Med.* 4, 165rv13. <https://doi.org/10.1126/scitranslmed.3004404>.
- Conti, H.R., Huppler, A.R., Whibley, N., Gaffen, S.L., 2014. Animal models for candidiasis. *Curr. Protoc. Im.* 105, 19. <https://doi.org/10.1002/0471142735.im1906s105>.
- De Bernardis, F., Agatensi, L., Ross, I.K., Emerson, G.W., Lorenzini, R., Sullivan, P.A., Cassone, A., 1990. Evidence for a role for secreted aspartate proteinase of *Candida albicans* in vulvovaginal candidiasis. *J. Infect. Dis.* 161, 1276–1283. <https://doi.org/10.1093/infdis/161.6.1276>.
- de Melo, E.T., Estrela, A.B., Santos, E.C.G., Machado, P.R.L., Farias, K.J.S., Torres, T.M., Carvalho, E., Lima, J.P.M.S., Silva-Júnior, A.A., Barbosa, E.G., Fernandes-Pedrosa, M. de F., 2015. Structural characterization of a novel peptide with antimicrobial activity from the venom gland of the scorpion *Tityus stigmurus*: Stigmurin. *Peptides* 68, 3–10. <https://doi.org/10.1016/j.peptides.2015.03.003>.
- di Caporiacco, L., 1947. Diagnosi preliminari di specie nuove di Aracnidi della Guiana britannica. *Monit. Zool. Ital.* 56, 20–34.
- Espósito, L.A., Yamaguti, H.Y., Souza, C.A., Pinto-da-Rocha, R., Prendini, L., 2017. Systematic Revision of the Neotropical Club-Tailed Scorpions, *Physoctonus*, *Rhopalurus*, and *Troglorhopalurus*, Revalidation of *Heteroctenus*, and Descriptions of Two New Genera and Three New Species (Buthidae, Rhopalurinae). *Neotropical club-tailed scorpions*, p. 415.
- Guilhelmelli, F., Vilela, N., Smidt, K.S., de Oliveira, M.A., da Cunha Moraes Álvares, A., Rigonatto, M.C.L., da Silva Costa, P.H., Tavares, A.H., Freitas, S.M. de, Nicola, A.M., Franco, O.L., Derengowski, L. da S., Schwartz, E.F., Mortari, M.R., Bocca, A.L., Albuquerque, P., Silva-Pereira, I., 2016. Activity of scorpion venom-derived antifungal peptides against planktonic cells of *Candida* spp. and *Cryptococcus neoformans* and *Candida albicans* biofilms. *Front. Microbiol.* 7 <https://doi.org/10.3389/fmicb.2016.01844>.
- Machado, R.J.A., Estrela, A.B., Nascimento, A.K.L., Melo, M.M.A., Torres-Rêgo, M., Lima, E.O., Rocha, H.A.O., Carvalho, E., Silva-Junior, A.A., Fernandes-Pedrosa, M.F., 2016. Characterization of TistH, a multifunctional peptide from the scorpion *Tityus stigmurus*: structure, cytotoxicity and antimicrobial activity. *Toxicon* 119, 362–370. <https://doi.org/10.1016/j.toxicon.2016.06.002>.
- Martins, J.G., Santos, G.C., Procópio, R.E. de L., Arantes, E.C., Bordon, K. de C.F., 2021. Scorpion species of medical importance in the Brazilian Amazon: a review to identify knowledge gaps. *J. Venom. Anim. Toxins Incl. Trop. Dis.* 27 <https://doi.org/10.1590/1678-9199-JVATITD-2021-0012>.
- Odds, F.C., Brown, A.J.P., Gow, N.A.R., 2003. Antifungal agents: mechanisms of action. *Trends Microbiol.* 11, 272–279. [https://doi.org/10.1016/s0966-842x\(03\)00117-3](https://doi.org/10.1016/s0966-842x(03)00117-3).
- Oliveira, T. de J., Oliveira, U.C. de, da Silva Junior, P.I., 2019. Serrulin: a glycine-rich bioactive peptide from the hemolymph of the yellow *Tityus serrulatus* scorpion. *Toxins* 11, 517. <https://doi.org/10.3390/toxins11090517>.
- Pappas, P.G., 2010. Opportunistic fungi: a view to the future. *Am. J. Med. Sci.* 340, 253–257. <https://doi.org/10.1097/MAJ.0b013e3181e99c88>.
- Parente, A.M.S., Daniele-Silva, A., Furtado, A.A., Melo, M.A., Lacerda, A.F., Queiroz, M., Moreno, C., Santos, E., Rocha, H.A.O., Barbosa, E.G., Carvalho, E., Silva-Júnior, A.A., Silva, M.S., Fernandes-Pedrosa, M.D.F., 2018. Analogs of the scorpion venom peptide stigmurin: structural assessment, toxicity, and increased antimicrobial activity. *Toxins* 10, 161. <https://doi.org/10.3390/toxins10040161>.
- Pfaller, M.A., Diekema, D.J., 2010. Epidemiology of invasive mycoses in North America. *Crit. Rev. Microbiol.* 36, 1–53. <https://doi.org/10.3109/10408410903241444>.
- Santussi, W.M., Bordon, K.C.F., Rodrigues Alves, A.P.N., Cologna, C.T., Said, S., Arantes, E.C., 2017. Antifungal activity against filamentous fungi of Ts1, a multifunctional toxin from *Tityus serrulatus* scorpion venom. *Front. Microbiol.* 8, 984. <https://doi.org/10.3389/fmicb.2017.00984>.
- Schaller, M., Bein, M., Kortling, H.C., Baur, S., Hamm, G., Monod, M., Beinhauer, S., Hube, B., 2003. The secreted aspartyl proteinases Sap 1 and Sap 2 cause tissue damage in an in vitro model of vaginal candidiasis based on reconstituted human vaginal epithelium. *Infect. Immun.* 71, 3227–3234. <https://doi.org/10.1128/IAI.71.6.3227-3234.2003>.
- Snyder, S.S., Gleaton, J.W., Kirui, D., Chen, W., Millenbaugh, N.J., 2021. Antifungal activity of synthetic scorpion venom-derived peptide analogues against *Candida albicans*. *Int. J. Pept. Res. Therapeut.* 27, 281–291. <https://doi.org/10.1007/s10989-020-10084-w>.
- Turner, S.A., Butler, G., 2014. The *Candida* pathogenic species complex. *Cold Spring Harb. Perspect. Med.* 4, a019778. <https://doi.org/10.1101/cshperspect.a019778>.
- Whaley, S.G., Berkow, E.L., Rybak, J.M., Nishimoto, A.T., Barker, K.S., Rogers, P.D., 2016. Azole antifungal resistance in *Candida albicans* and emerging non-*albicans* *Candida* species. *Front. Microbiol.* 7, 2173. <https://doi.org/10.3389/fmicb.2016.02173>.
- Zottich, U., Da Cunha, M., Carvalho, A.O., Dias, G.B., Silva, N.C.M., Santos, I.S., do Nascimento, V.V., Miguel, E.C., Machado, O.L.T., Gomes, V.M., 2011. Purification, biochemical characterization and antifungal activity of a new lipid transfer protein (LTP) from *Coffea canephora* seeds with α -amylase inhibitor properties. *Biochim. Biophys. Acta Gen. Subj.* 1810, 375–383. <https://doi.org/10.1016/j.bbagen.2010.12.002>.