

Chapter 09

The Role of Mast Cell Specific Chymases and Trypsases in Tumor Angiogenesis

Devandir Antonio de Souza Junior, Ana Carolina Santana, Elaine Zayas Marcelino da Silva, Constance Oliver and Maria Celia Jamur*

Department of Cell and Molecular Biology and Pathogenic Bioagents,
Ribeirão Preto Medical School, University of São Paulo, Brazil

***Corresponding Author:** Maria Celia Jamur, Department of Cell and Molecular Biology and Pathogenic Bioagents, Ribeirão Preto Medical School, University of São Paulo, Avenida Bandeirantes, 3900, 14049-900 Ribeirão Preto, SP, Brazil, Tel: 55-16-3315-3217; Fax: 55-16-3315-0728

First Published **January 21, 2019**

This Book Chapter is an excerpt from an article published by Maria Celia Jamur, et al. at BioMed Research International in February 2015. (Devandir Antonio de Souza Junior, Ana Carolina Santana, Elaine Zayas Marcelino da Silva, Constance Oliver, and Maria Celia Jamur, "The Role of Mast Cell Specific Chymases and Trypsases in Tumor Angiogenesis," BioMed Research International, vol. 2015, Article ID 142359, 13 pages, 2015. <https://doi.org/10.1155/2015/142359>.)

Copyright: © 2019 Maria Celia Jamur, et al.

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source.

Abstract

An association between mast cells and tumor angiogenesis is known to exist, but the exact role that mast cells play in this process is still unclear. It is thought that the mediators released by mast cells are important in neovascularization. However, it is not known how individual mediators are involved in this process. The major constituents of mast cell secretory granules are the mast cell specific proteases chymase, tryptase and carboxypeptidase A3. Several previous studies aimed to understand the way in which specific mast cell granule constituents act to induce tumor angiogenesis. A body of evidence indicates that mast cell proteases are the pivotal players in inducing tumor angiogenesis. In this review, the likely mechanisms by which tryptase and chymase can act directly or indirectly to induce tumor angiogenesis are discussed. Finally, information presented here in this review indicates that mast cell proteases significantly influence angiogenesis and thus affecting tumor growth and progression. This also suggests that these proteases could serve as novel therapeutic targets for the treatment of various types of cancer.

Key Words

Mast Cell; Tryptase; Chymase; Tumor Angiogenesis

Introduction

Angiogenesis is a dynamic process mediated by endothelial cells whereby new blood vessels are formed from existing ones [1,2]. Angiogenesis is crucial during physiological processes such as embryonic development and corpus luteum formation, and it is also involved in the development of pathological conditions such as tumorigenesis and chronic inflammation [3,4]. This process is highly regulated by the balance between proangiogenic and antiangiogenic factors within the vascular microenvironment and involves the participation of extracellular matrix (ECM) proteins, adhesion molecules and proteolytic enzymes [5,6]. The main proangiogenic factors include, Vascular Endothelial Growth Factor (VEGF), Fibroblast Growth Factor (FGF), Transforming Growth Factor-Beta (TGF- β) Platelet-Derived Growth

Factor (PDGF), Interleukin-8 (IL-8), and angiopoietin 1 [7,8]. A change in the balance between proangiogenic and antiangiogenic factors towards the activation of usually quiescent endothelial cells initiates a series of sequential events that characterize the angiogenic process, i.e.: ECM degradation; migration and proliferation of endothelial cells; formation of tube-like structures on 2D matrices and sprouting in 3D matrices; spatial distribution of newly formed vessels; deposition of new ECM; mobilization of pericytes for vessel stabilization; and finally, a shift in the balance towards inhibition of angiogenesis returns the endothelial cells to their quiescent state [9-12].

The importance of angiogenesis for tumor development has long been investigated. In the beginning of the 20th century, Goldman [13] had already envisioned the importance of the vascular system for tumor growth. However, it was not until 1971 that Folkman [14] showed that tumor growth was dependent on angiogenesis and proposed the use of antiangiogenic therapy against cancer. Today, tumor angiogenesis is considered crucial for cancer development through induction of tumor growth, invasion, and metastasis [15,16]. Moreover, the use of antiangiogenic agents as adjuvants for cancer treatment has become a reality [17].

Tumor angiogenesis relies on the abnormal switch in the balance between positive and negative regulators, usually triggered by the release of angiostimulatory mediators from neoplastic and inflammatory cells. The binding of these mediators to their receptors on endothelial cells activates the angiogenesis cascade [18-20]. The pro and anti-tumorigenic signals provided by tumor cells, resident and recruited immune cells are ultimately responsible for the outcome of tumor progression. These signals act in processes such as proliferation and survival of tumor cells, angiogenesis, immune system modulation, tissue remodeling and metastasis [21]. Therefore, the interaction between tumor, stromal, and immune cells collaborate to establish a pro angiogenic/tumorigenic microenvironment favoring the survival, development and spreading of tumor cells [22].

The view that immune cells are critical in providing the angiogenic stimulus required for tumor growth and progression is supported by a number of studies assigning proangiogenic roles to various innate immune cell types, in particular macrophages and mast cells [23]. In this review article, the importance of mast cells in tumor angiogenesis with special reference on the role of mast cell proteases in this process will be discussed.

Mast Cell Biology

Mast cells are bone-marrow derived multifunctional immune cells implicated in both physiological and pathological settings (reviewed in [24]). Mast cell progenitors leave the blood circulation to mature and reside in the connective tissue, at the interface between the host and the outside environment, predominantly in sub-epithelial regions and in the connective tissue surrounding blood vessels, nerves, lymphatic vessels, and mucus glands [25,26]. Mast cell distribution relies on mechanisms of constitutive homing, enhanced recruitment, survival, and local maturation of mast cell progenitors. In physiological conditions, their numbers remain relatively constant; however, proliferation and the enhancement of the mechanisms described above lead to mast cell hyperplasia during hypersensitivity reactions, infections, and in response to various disease processes [27-29].

Mast cell functions are a direct consequence of the diversity of biologically active compounds that they are able to produce and secrete. Upon activation, mast cells can release a wide range of mediators belonging to three distinct classes: preformed mediators, which are stored in mast cell cytoplasmic granules; neoformed or lipid mediators, which are products of phospholipase A2 activation; and neo-synthesized mediators, which are produced and released upon transcriptional activation [25].

Mast cells may be activated by distinct stimuli acting on the various surface receptors expressed by these cells. Moreover, the variety of mast cell responses to different stimuli, i.e. the nature of the mediators

released, is influenced by mast cell heterogeneity. This heterogeneity is ultimately dictated by the intrinsic and microenvironmental factors that these cells encounter in normal and pathological conditions [25,30].

A distinct feature of mature mast cells is the high content of metachromatic, electron dense secretory granules present in their cytoplasm. These secretory granules store an extensive variety of pre-formed bioactive mediators, which are all stored in the active form and are promptly released by degranulation. Mast cell degranulation can be triggered by various mechanisms such as: the crosslinking of FcεRI-bound IgE by specific antigens, crosslinking of FcγRIII by IgG immune complexes, complement receptor engagement by complement components such as C3a and C5a, c-Kit receptor binding by Stem Cell Factor (SCF), TLR2 (Toll-like Receptor 2) stimulation, neuropeptides and other peptides [31-37]. The preformed mediators include, but are not restricted to: biogenic amines, lysosomal enzymes, proteases, proteoglycans, cytokines, chemokines, growth factors, and peptides [31,33,36-38].

The major constituents of mast cell secretory granules are the mast cell specific proteases chymase, tryptase and carboxypeptidase A3 (CPA3) (Table 1) [39]. These proteases are exclusively expressed by mast cells and they represent approximately 25% of the cell protein content. They are stored in their active form within the secretory granules [40-42]. The tryptases and chymases are serine-proteases while CPA3 is a zinc-dependent metalloprotease [39,43].

Like other positively charged preformed mediators, mast cell proteases are efficiently packed into mast cell secretory granules owing to their interaction with negatively charged serglycin proteoglycans [43,44]. However, not all subtypes of mast cell proteases are dependent on serglycin for storage. A study using knockout mice for serglycin core protein showed that although the protease mRNA levels were similar between knockout and wild type mice the interaction of mast cell proteases with serglycin was necessary to regulate the storage of mouse mast cell protease (mMCP)-4, -5, -6, and CPA in the gran-

ules [45]. On the other hand, Braga et al. [46] showed that the storage of mMCP-7 and mMCP-1 was independent of serglycin. Moreover, Mello et al. [44] have recently shown that serglycin dependent storage of mast cell proteases is critical in the induction of apoptosis induced by permeabilization of the granule membrane.

The mast cell protease content varies according to the tissue distribution of mast cells as well as from species to species, and these differences are used to phenotypically classify mast cells (Table 1) [47]. Human mast cells are divided into MC_T that express tryptases α and β and MC_{TC} that express chymases, tryptases and CPA3. A third phenotype expressing tryptases and CPA3 was recently described in airway epithelium and esophageal samples of patients with asthma and eosinophilic esophagitis [48,49]. In rodents, mast cells are classified according to their distribution as connective tissue mast cells (CTMCs), which express chymases (α and β), tryptases and CPA; and mucosal mast cells (MMC), which express the β chymases, mMCP-1 and 2. It is noteworthy to mention that these mouse mast cell phenotypes can vary significantly according to mast cell location, animal strain, and whether or not the tissue is inflamed [50-53]. Mast cell proteases have been implicated in a number of pathological states including arthritis, allergic airway inflammation, and tumor angiogenesis [54-59].

Mast Cells, Tumors and Angiogenesis

The association between mast cells, inflammation, and cancer is conflicting and involves both promotion of and protection against tumor progression. The first association of mast cells with tumors dates from the initial description of mast cells by Paul Ehrlich in 1878, when he reported that mast cells were numerous in some tumors Ehrlich [60]. Since then, interest in the contribution of mast cells to tumor development has increased progressively. Mast cells have been shown to accumulate around several types of tumors and are generally the first inflammatory cells to infiltrate developing tumors such as malignant melanoma, and breast and colorectal tumors [61-64]. This accumulation typically occurs around blood vessels within the tumor environment and correlates with both good and poor prognosis in

different cancers demonstrating the paradoxical involvement of mast cells in tumor progression [65-67]. Mast cells are recruited by several tumor-derived factors including the angiogenic factors VEGF, PDGF, and FGF-2 [68]. Notably, Huang et al. [69] using an hepatocarcinoma model, showed that mast cells were unable to migrate in SCF-knock-out tumors or in the presence of anti-c-kit antibodies, ultimately resulting in decreased tumor growth. In addition SCF stimulation lead to the release of matrix metalloproteinase 9 (MMP-9), IL-6, TNF- α , VEGF, Cox2, i-NOS, and chemokine (C-C motif) ligand 2 (CCL2). These results suggest that mast cell recruitment and activation, with the consequent release of inflammatory mediators that participate in tissue remodeling and immune suppression, are mainly mediated by SCF and its receptor c-kit expressed on the mast cell surface. The contribution of mast cells to the promotion of cancer includes their action on immunosuppression, the release of pro-angiogenic and mitogenic factors, and degradation of the ECM [61].

Mast cells have been shown to be associated with angiogenesis and tumor progression in numerous types of cancer [65,70-80]. Stimulation of angiogenesis is undoubtedly the most significant role of mast cells in the promotion of tumor growth and development [81]. In fact, mast cell deficient mice showed a reduced angiogenesis and metastatic capacity during tumor induction [82]. The involvement of mast cells in these processes is most certainly related to the release of a plethora of potent pro-angiogenic mediators which include, histamine, angiopoietin-1, VEGF, FGF-2, TGF- β , Tumor Necrosis Factor alpha (TNF- α), and IL-18 [81,83,84]. Furthermore, a role for mast cell-specific proteases has gained increasing support [74,85-89]. For instance, mast cell specific proteases are involved in the initial phases of tumor growth and also modulate vascular growth in the later stages of tumor progression [59]. Mast cell mediators can induce angiogenesis through their action at different stages of angiogenesis i.e.: degradation of the ECM, migration and proliferation of endothelial cells, formation and distribution of new vessels, synthesis of ECM and pericyte mobilization [10,90].

Most angiogenic mediators released by mast cells are not exclusive to mast cells, so it is difficult to attribute their biological actions to mast cells alone. Therefore, the majority of recent investigations on the role of mast cells in tumor angiogenesis have focused on the ability of mast cells to synthesize, store and release mast cell specific chymases and tryptases. The role of these proteases in tumor angiogenesis is the subject of this review.

Table 1: Characteristics of major human and murine mast cell proteases.

Protein designation	Mast cell subtype	Chomosomal location	Gene Name	Primary substrate specificity	Subunit structure
CHYMASE					
Human					
Chymase (CMA1)	MC _{TC}	14q11.2	<i>CMA1</i>	Chymotrypsin-like (aromatic amino acid)	Monomer
Mouse					
mMCP-1	MMC	14C1/2	<i>Mcpt1</i>	Chymotrypsin-like (aromatic amino acid)	Monomer
mMCP-2	MMC	14C1/2	<i>Mcpt2</i>	Unknown (low enzymatic activity)	Monomer
mMCP-4	CTMC	14C1/2	<i>Mcpt4</i>	Chymotrypsin-like (aromatic amino acid)	Monomer
mMCP-5	CTMC	14C1/2	<i>Cma1/Mcpt5</i>	Elastase-like (Val/Ala/Ile)	Monomer
TRYPTASE					
Human					
Tryptase β II/ β III (human)	MC _{TC} /MC _T	16p13.3	<i>TPSB2</i>	Trypsin-like (Arg/Lys)	tetramer
Tryptase α I/ β I (human)	MC _{TC} /MC _T	16p13.3	<i>TPSAB1</i>	Trypsin-like (Arg/Lys)	tetramer
Mouse					
mMCP-6	CTMC	17A3.3	<i>Mcpt6</i>	Trypsin-like (Arg/Lys)	tetramer
mMCP-7	CTMC	17A3.3	<i>Tpsab1</i>	Trypsin-like (Arg/Lys)	tetramer
CARBOXYPEPTIDASE A3					
Human					
MC-CPA	MC _{TC}	3q24	<i>CPA3</i>	CPA-like (C-terminal aromatic/aliphatic amino acids)	Monomer
Mouse					
MC-CPA	CTMC	3A3	<i>Cpa3</i>	CPA- CPA-like (C-terminal aromatic/aliphatic amino acids)like	Monomer

Tryptases in Tumor Angiogenesis

Mast cell tryptase is a tetrameric neutral serine protease with a molecular weight of 134 kDa [54,91]. Mast cells contain 10-35 pg tryptase per cell [92]. The enzyme consists of four non-covalently bound subunits. Each subunit has one active catalytic site and disruption of the tetramer into monomers causes inactivation of the tryptase [40, 93]. Tryptases were named because of their trypsin-like substrate specificity (Table 1). They cleave preferentially after Lys/Arg residues [39,94]. Humans express two classes of tryptase (alpha and beta). Alpha-tryptase is the major circulating isoform and β -tryptase appears to be the major type stored in secretory granules [54,94-96]. Three similar β -tryptases (β I-III), and one α -tryptase have been identified. The β -tryptases are 98-99% identical in amino acid sequence, while the α -tryptase is less closely related. The mast cell tryptases (MCP-6 and MCP-7) expressed in rodents most likely represent the counterparts to human α -tryptase and β -tryptases respectively. Additionally, both subtypes are initially expressed as monomers in humans and rodents [95,96] and then combine to form tetramers.

The first reports on the association between tryptase and tumor angiogenesis were focused on the use of tryptase as a specific marker for mast cells. Kankkunen and colleagues [97] were among the first to study this relationship. They analyzed the number and distribution of tryptase and chymase-positive mast cells in benign and malignant breast lesions [97]. Their results show that the number of tryptase positive mast cells in malignant tumors was increased four times in relation to the benign lesions. Moreover, in malignant lesions, tryptase positive mast cells were concentrated at the tumor edge within the invasion zone. Since then several other studies have corroborated this correlation between tryptase and malignant tumors in various types of tumor, such as: lung adenocarcinoma [98]; B-cell non-Hodgkin's lymphomas [80]; brain [99], cervical cancer [70]; myelodysplastic syndrome [4]; B cell chronic lymphocytic leukemia [100]; melanoma [101]; hepatocellular carcinoma [102]; gastrointestinal cancer [103];

early breast cancer [104]; and colorectal cancer [105]. During the last two decades, several studies, using tryptase as a mast cell marker, have shown that mast cells are strategically located in proximity to blood vessels where they have been associated with tumor angiogenesis. These studies on the correlation between tryptase positive mast cells and angiogenesis also showed a positive association between tumor aggressiveness and poor prognosis [106-108].

Tryptases act on a variety of substrates [39,96]. Several of these substrates are known to be key factors in angiogenesis, such as: degradation of extracellular matrix [109,110], plasminogen activation [111], fibrinogen degradation [112,113], activation of latent collagenase and matrix metalloproteinases [114, 115], and tryptase-induced endothelial barrier dysfunction [116,117]. The correlation between the presence of tryptase positive mast cells and angiogenesis as well as the action of tryptase on angiogenesis related substrates supports the hypothesis that the presence of tryptase is critical for the initiation of angiogenesis. In spite of the fact that these studies only showed the presence of mast cells at angiogenic sites, they provided a basis for further investigation on the direct relationship between tryptase and angiogenesis.

Blair et al. [118] were the first to investigate the potential of tryptase to induce in vitro angiogenesis using a tube formation angiogenesis assay. This assay involves plating endothelial cells on a gel matrix and assaying adhesion, migration, and formation of capillary or tube-like structures. These capillary-like structures consist of tubes and loops. Tubes are formed by endothelial cell sheets and go from one branch point to another branch point or to a loose end. Loops are the enclosed (or almost enclosed) spaces inside the tubes. In these experiments, they incubated human dermal microvascular endothelial cells (HDMEC), plated in matrigel-coated wells, in the presence of purified human lung tryptase for 16 hours. The tryptase induced an increase in capillary-like structures and proliferation of HDMEC cells. This effect was suppressed by the use of protease inhibitors. These results lead to the conclusion that tryptase interacts directly with endothelial cells via an unidentified mechanism to induce angiogenesis.

Compton et al. [119] also investigated the ability of tryptase to induce in vitro angiogenesis. However, in this study tryptase did not stimulate growth of human umbilical vein endothelial cells (HUVEC). They observed that tryptase induced the release of IL-8 in a dose-dependent manner. They also confirmed that this effect on HUVEC required enzymatically active tryptase. Release of IL-8 did not occur in the presence of heat inactivated enzyme, or when the tryptase was preincubated with the protease inhibitors.

Using a mouse model of epithelial carcinogenesis, Coussens et al. [120] observed that during the dysplasia stage, vascular changes indicative of an angiogenic switch from vascular quiescence to modest neovascularization occurred in early modest to low-grade lesions [120]. At this stage, the researchers observed a significant increase in tryptase activity and mMCP-6 mRNA expression. In this same study, mMCP-6 did not directly affect the extracellular matrix, but mMCP-6 acted indirectly through the stimulation of procollagen synthesis by fibroblasts. The results of this elegant study indicate that tryptase can affect the extracellular matrix by stimulating the cellular components, thus altering its composition that then leads to neovascularization. Additionally, other studies have demonstrated the ability of tryptase to digest isolated extracellular matrix [109,121,122]. Further studies have shown that tryptase can also activate metalloproteinases thus inducing degradation of the extracellular matrix [114,123]. Taken together, these results suggest that tryptase may have both a direct and indirect action in the remodeling and degradation of the extracellular matrix during angiogenesis. The role of tryptase in degradation/remodeling of the extracellular matrix may vary depending on the type of tumor and phase of tumor progression. This process of degradation/remodeling could increase the space available for neovascularization as well as contribute to the release of angiogenic factors present in the matrix such as VEGF and FGF-2 [113,124].

Another direct effect of tryptase in various cell types, including endothelial cells, is stimulation of Protease Activated Receptor-2 (PAR-2) [125,126]. Yoshii and coworkers [74] studying the direct

effects of tryptase on the proliferation and intracellular signaling in DLD-1 colorectal adenocarcinoma cells in vitro demonstrated that tryptase induces proliferation in DLD-1 cells through its proteolytic activity on PAR-2. The increase in proliferation was mediated in a cyclooxygenase- and MAP kinase-dependent manner. Recently, Zhi and colleagues [127] demonstrated in vitro that tryptase increased bEnd.3 mouse brain endothelial cell proliferation, migration, and tube formation, suggesting a role for tryptase in microvessel formation. They also observed that tryptase increased tissue Plasminogen Activator (tPA) and inhibited Plasminogen Activator Inhibitor-1 (PAI-1) expression at both the mRNA and protein levels in bEnd.3 cells.

Therefore, tryptase, through the activation of PAR-2, may have a role in modulating the balance between expression of urokinase Plasminogen Activator (uPA) and PAI-1. Expression of uPA and PAI-1 is PAR-dependent [128,129]. Additionally, PAR-2 activation in vascular cells promotes angiogenesis during tissue repair as well as during neovascularization of the retina [130,131]. PAR-2 also induces endothelial cell proliferation [132,133] and cytokine expression in endothelial cells and other cell types [134-136]. Together these studies indicate that the activation of PAR-2 by tryptase is directly related to the role of mast cells in angiogenesis. The exact mechanism by which PAR-2 contributes to angiogenesis remains to be elucidated. Nevertheless, the involvement of tryptase in PAR-2 activation and angiogenesis is clear.

Tryptase (mMCP-6 and mMCP-7) expression and activity were increased during progression of skin tumors in mice [59]. Furthermore, the increase in tryptase activity was correlated with the tumor angiogenesis. mMCP-7 was not detectable in control animals, but became measurable concomitantly with the initiation of angiogenesis during tumor progression. The ability of these two subtypes of tryptase to induce angiogenesis in vitro was compared. mMCP-6 and mMCP-7 were both able to induce tube formation in SVEC4-10 endothelial cells. However, mMCP-7 was more efficient in inducing cell spreading and tube formation than mMCP-6 (Figure 1). These data

show that both subtypes of mast cell specific tryptases may interact with endothelial cells to promote neovascularization of tumors.




Figure 1: rmMCP-6 and 7 induce tube formation by mouse endothelial cells.

When endothelial cells were incubated in absence of tryptases a few SVEC4-10 cell are spread and an occasional tube is seen (arrow). When the SVEC4-10 cells were incubated with rmMCP -6, cells are spread and tubes (arrows) and loops (L) are present. When the SVEC4-10 cells were incubated with rmMCP-7 tubes (arrows) and loops (L) are more prevalent. Endothelial cells (SVEC4-10) were cultivated in wells of μ -slides angiogenesis (Ibidi GmbH, Martinsried, DE) with Geltrex® (Thermo Fisher Scientific Inc. Waltham, MA) for 5 hours at 37°C in the presence or absence of the subtypes of tryptase. Following incubation, the samples were fixed, viewed and photographed using a Nikon Eclipse TE2000-U (Nikon Instruments Inc. Mellville, NY) phase contrast inverted microscope equipped with a Nikon DXM-1200 digital camera.

Recently, Wang et al. [137] examined the involvement of mast cells in angiogenesis in vitro using by co-culturing myocardial microvascular endothelial cells (MMVEC) with rat peritoneal mast cells (activated or nonactivated) or with only the granules from rat peritoneal mast cells. Mast cells activated by 48/80 induced in vitro angiogenesis, but this was not detected with nonactivated mast cells. To confirm if the mast cell granules (MCG) alone can induce in vitro angiogenesis, MMVEC were cultivated with MCG. There was significantly higher migration, proliferation, and capillary tube-like formation in the presence of MCG than without MCG. In addition, in the presence of MCG a decrease in Angiopoietin-2 (Ang-2) protein expression and mRNA by endothelial cells was seen, indicating that the MCG had a suppressive effect on Ang-2 expression. In order to determine if tryptase or chymase was responsible for the induction

of angiogenesis via Ang-Tie-2, myocardial microvascular endothelial cells were cultivated with MCG in the presence of tryptase or chymase inhibitors. The ability of MCG to induce angiogenesis could be reversed by either tryptase or chymase inhibitors, although the tryptase inhibitor was more effective. Therefore, tryptase may be more important than chymase in inducing angiogenesis. Also tryptase may be the major factor released by mast cell that promotes neovessel formation. Taken together the results indicate that tryptase affects angiogenesis in MMVEC via Ang-Tie-2.

Since tryptase is known to be involved in tumor angiogenesis (Figure 2), recently it has been suggested that tryptase may serve not only as a biomarker in different pathologies but also as a promising therapeutic target. Ammendola et al. [103] review the possible molecular mechanisms of three drugs targeting tryptase, and discusses their possible role in cancer therapy. Furthermore, the authors suggest mast cell tryptase as potential antiangiogenic strategy.




Figure 2: Trypsin can act directly or indirectly to induce tumor angiogenesis.

Chymases and Tumor Angiogenesis

Chymases are mast cell specific proteases that belongs to the family of serine-proteases. Mast cell chymases are monomeric endopeptidases [138] that have chymotrypsin-like specificities (Table 1). They preferentially cleave proteins after aromatic amino acid residues, except rat and mouse chymase 5 (rMCP-5 and mMCP-5), which have elastase-like rather than chymotrypsin-like substrate specificity [139]. Different chymases have been identified in various species, and have been divided into two groups (α - and β -chymases) according to their structure and substrate specificity [140,141]. Both subgroups of chymases convert Angiotensin I to Angiotensin II, but only β -chymases degrade Angiotensin II. The α -chymases cleave the Phe⁸-His⁹ bond of Angiotensin I [141] and are widely expressed in mammalian mast cells. The β -chymases can convert Angiotensin I to Angiotensin II but

preferentially hydrolyze the Tyr⁴-Ile⁵ bond of Ang I and Ang II, yielding inactive peptide fragments [142].

Many other chymase substrates have been reported such as procollagen [143], pro-matrix metalloproteinase-9 (pro-MMP-9) [144,145], tissue inhibitor of metalloproteinase1 (TIMP-1) [146], big-endothelin-1 and big-endothelin-2 [147,148], TGF- β 1 [149], and thrombin [150]. However, chymases are relatively promiscuous in terms of cleavage specificity, and they have the capacity to cleave a multitude of substrates and can even cleave the same substrate at multiple positions.

In mast cells, chymases are stored as active enzymes in a macromolecular complex together with heparin in secretory granules. Prior to secretion, mast cell chymases are activated by dipeptidyl peptidase I [151]. Once secreted into the extracellular fluid, chymases remain in a complex with heparin proteoglycan, which protects the chymases from inhibition by endogenous inhibitors of chymase present in the extracellular fluid [152].

In man, chymases are found in mast cells around blood vessels, in the heart, and many other tissues [153], but not in plasma [138]. The fact that chymases are present in different tissues and that they cleave numerous substrates indicates that chymases can play many different roles in physiological and pathological conditions, including tumor angiogenesis, but these roles are not well characterized. Additionally, the distribution of mast cells and mast cell subtypes is variable depending on the type of tumor. In colorectal cancer, hepatocellular carcinoma and intrahepatic cholangiocarcinoma, both MC_{TC} and MC_T proliferate or infiltrate in the tumor [154,155]. In contrast, in some malignant tumors, a significant increase in mast cell number was found. This observation was due to an increase in MC_T at the invasion zone, whereas the number of MC_{TC} remained constant in this area [70,156].

Muramatsu et al. [157], using a hamster sponge implant model of angiogenesis, suggested a possible involvement of chymase in angiogenesis. Both transfection of human pro-chymase cDNA and injections of purified chymase into implanted sponges demonstrated that chymase was a powerful angiogenic factor. The authors then went on to investigate chymase as an alternative angiotensin II-generating enzyme in angiogenesis. Chymase inhibitors reduced mast cell mediated angiogenesis in sponge hamster [86, 158]. Furthermore, chymostatin (a chymase inhibitor) and TCV-116 (an angiotensin II (Ang II) type 1 receptor agonist) completely inhibited bFGF induced angiogenesis [85]. These results suggest that hamster chymase generates Angiotensin II from Angiotensin I, triggering bFGF induced angiogenesis (Figure 3). Additionally, chymase mRNA expression was detected in bFGF-induced angiogenesis, but not in the control hamsters treated with saline. The increased angiogenesis was accompanied by a considerable increase in chymase activity in the granuloma tissues [86].

Local injection of compound 48/80, a mast cell activator, also promoted angiogenesis in hamster sponge granulomas through a chymase-dependent mechanism [85] and enhanced VEGF mRNA expression [159]. Treatment with neutralizing antibody against VEGF or an antisense oligodeoxynucleotide against VEGF suppressed Angiotensin II-induced angiogenesis, showing that VEGF is a downstream factor in Angiotensin II-induced angiogenesis [159]. Therefore, the chymase-Ang II-VEGF pathway in granulation tissue may be the primary functional mediator of angiogenesis.

Arbeit et al. [160] developed a mouse model of epithelial carcinogenesis, by targeting the early region of human papillomavirus type 16 (HPV16) to the basal cells of the squamous epithelium using a human keratin 14 (K14) enhancer/promoter. With time after birth, these mice develop epidermal hyperplasia, which advances to angiogenic dysplasias. After one year, these dysplasias can progress to invasive squamous cell carcinomas [120]. In this transgenic mouse model

an increase in mast cell number was observed during progression of the disease, especially in dysplasia [120], which is characterized by increased angiogenesis. At the same time, there was a significant increase in chymase activity and mMCP-4 mRNA was detected in dysplastic skin. Finally, hyperplastic ear skin from the transgenic mice was incubated in medium containing mMCP-4 and subsequently embedded in a collagen gel containing randomly dispersed bovine capillary endothelial cells (BCE). The BCEs showed a dramatic response to the presence of skin that had been preincubated with mMCP-4. The endothelial cells became radially aligned, had increased rates of proliferation and migration, and exhibited tube formation toward the dermal surface, but not the epidermal surface of the skin pieces. However, the addition of mMCP-4 directly to BCEs did not induce a response, indicating that the angiogenic activity came from sequestered factors in the skin pieces used in this assay. Furthermore, canine and human chymases can cleave pro-MMP-9 producing active MMP-9. This matrix metalloproteinase participates in extracellular matrix remodeling [161] and regulation of angiogenesis [162]. In order to verify that mMCP-4 contributes to angiogenesis by activating MMP-9, hyperplastic skin lysates (containing predominantly pro-MMP-9) were incubated with mMCP-4 and then subjected to gelatin-substrate zymography. A substantially increased amount of active MMP-9 was detected, confirming that mMCP-4 plays a role in angiogenesis by activating MMP-9 thus inducing extracellular matrix remodeling [120]. These results, taken together, suggest that mMCP-4 acts indirectly to induce angiogenesis by activating pro-MMP-9, thus releasing sequestered angiogenic factors (Figure 3).

MCP-5 is emerging as a key player in chymase induced angiogenesis. The level of rMCP-5 mRNA in the granulomatous tissue of λ -carrageenin-soaked sponge implants was significantly increased in comparison to control granulomas [163]. Furthermore, angiogenesis during granuloma formation could be blocked by chymostatin, and by blocking rMCP-5 with a specific antisense oligonucleotide. Recently, de Souza et al. [59] reported increased mast cell and blood

vessel numbers as well as blood vessel diameter during mouse skin tumor progression. mMCP-5 expression was not detected in the control animals, but was present in all stages of tumor progression. Total chymase activity and mMCP-5 expression increased during tumor progression and was correlated with tumor angiogenesis. In contrast, although mMCP-4 expression could be detected in control animals and in all phases of tumor progression, there was no difference in expression at the various phases.

The majority of the studies attempting to define the role of specific mast cell proteases, such as chymase in tumor angiogenesis in humans have been focused on a pathological examination of tumor biopsies. The stage of the tumor, mast cell numbers and mast cell subtype i.e., MC_{TC} and MC_T were all analyzed [78,80,156,164-167]. These investigations have suggested a relation between chymase and angiogenesis in tumors. However, research into the chymase-containing mast cell subpopulations has been hampered by lack of suitable reagents.

In small sized adenocarcinomas of the lung, the mast cell number is increased in comparison with normal lung tissue and ratios of MC_{TC} to total mast cells in the adenocarcinomas is significantly higher than in normal lung. Furthermore, the accumulation of MC_{TC}, but not MC_T correlates with a poor prognosis for patients with small sized lung adenocarcinoma. However, the numbers of both mast cell types correlated significantly with microvessel counts in small sized lung adenocarcinomas [168].

Mast cells are present in all three developmental phases of hemangiomas, which are benign, usually self-involuting tumors of the microvasculature [169]. In the initial proliferative phase, angiogenesis in hemangiomas is excessive, but is followed by spontaneous regression, and the cellular parenchyma is gradually replaced with fibrofatty tissue. The mast cell number is highest in the involuting phase, reduced in the involuted phase and least in the proliferative phase. However, the proportion of MC_{TC} decreases from the proliferative

phase through the involuting phase to the involuted phase, indicating a proangiogenic role for chymase in this tumor. At the later stages, when the proportion of MC_T are increasing, these mast cells may be releasing tumor inhibitors.

In conclusion, the studies presented here support a relationship between mast cell chymases and tumor angiogenesis (Figure 3). However, additional studies are needed in order to elucidate the precise role of mast cells.




Figure 3: Chymase acts indirectly to induce angiogenesis. Chymase can active MMP-9 or convert Angiotensin I to Angiotensin II.

Conclusion

This review has presented evidence that tryptase and chymase act directly (on endothelial cells) or indirectly (on the extracellular

matrix) to induce tumor angiogenesis. Most importantly, tryptase appears to play a more significant role in tumor progression than chymase. This evidence supports the use of mast cell proteases as novel tumor biomarkers as well as targets for antiangiogenic therapy. Nevertheless, studies using chymase or tryptase knockout animals are necessary to demonstrate the functional impact of either chymase or tryptase on tumor angiogenesis.

References

1. J Folkman, Y Shing. Angiogenesis. *J Biol Chem.* 1992; 267: 10931-10934.
2. TO Daniel, D Abrahamson. Endothelial signal integration in vascular assembly, *Annu Rev Physiol.* 2000; 62: 649-671.
3. D Ribatti. Genetic and epigenetic mechanisms in the early development of the vascular system. *J Anat.* 2006; 208: 139-152.
4. D Ribatti, E Crivellato. Mast cells, angiogenesis, and tumour growth. *Biochim Biophys Acta.* 2012; 1822: 2-8.
5. D Hanahan, J Folkman. Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell.* 1996; 86: 353-364.
6. ML Iruela-Arispe, HF Dvorak. Angiogenesis: a dynamic balance of stimulators and inhibitors. *Thromb Haemost.* 1997; 78: 672-677.
7. M Sawatsubashi, T Yamada, N Fukushima, H Mizokami, O Tokunaga, T Shin. Association of vascular endothelial growth factor and mast cells with angiogenesis in laryngeal squamous cell carcinoma. *Virchows Arch.* 2000; 436: 243-248.
8. J Talreja, MH Kabir, BFM, DJ Stechschulte, KN Dileepan.

Histamine induces Toll-like receptor 2 and 4 expression in endothelial cells and enhances sensitivity to Gram-positive and Gram-negative bacterial cell wall components. *Immunology*. 2004; 113: 224-233.

9. B Hobson, J Denekamp. Endothelial proliferation in tumours and normal tissues: continuous labelling studies. *Br J Cancer*. 1984; 49: 405-413.
10. PA D'Amore, RW Thompson. Mechanisms of angiogenesis. *Annu Rev Physiol*. 1987; 49: 453-464.
11. A Le Querrec, D Duval, G Tobelem. Tumour angiogenesis. *Baillieres Clin Haematol*. 1993; 6: 711-730.
12. M Potente, H Gerhardt, P Carmeliet. Basic and therapeutic aspects of angiogenesis. *Cell*. 2011; 146: 873-887.
13. E Goldmann. The Growth of Malignant Disease in Man and the Lower Animals, with special reference to the Vascular System. *Proc R Soc Med*. 1908; 1: 1-13.
14. J Folkman. Tumor angiogenesis: therapeutic implications. *N Engl J Med*. 1971; 285: 1182-1186.
15. G Ranieri, CD Gadaleta, R Patruno, N Zizzo, MG Daidone, et al. A model of study for human cancer: Spontaneous occurring tumors in dogs. Biological features and translation for new anticancer therapies. *Crit Rev Oncol Hematol*. 2013; 88: 187-197.
16. G Ranieri, M Pantaleo, M Piccinno, M Roncetti, M Mutinati, et al. Tyrosine kinase inhibitors (TKIs) in human and pet tumours with special reference to breast cancer: a comparative review. *Crit Rev Oncol Hematol*. 2013; 88: 293-308.
17. B Al-Husein, M Abdalla, M Trepte, DL Deremer, PR Somnath. Antiangiogenic therapy for cancer: an update. *Pharmacotherapy*. 2012; 32: 1095-1111.

Top 10 Contributions on Cell Biology

18. AW Griffioen, G Molema. Angiogenesis: potentials for pharmacologic intervention in the treatment of cancer, cardiovascular diseases, and chronic inflammation. *Pharmacol Rev.* 2000; 52: 237-268.
19. P Carmeliet, RK Jain. Molecular mechanisms and clinical applications of angiogenesis. *Nature.* 2011; 473: 298-307.
20. F Moccia, S Dragoni, V Poletto, V Rosti, F Tanzi, et al. Orai1 and transient receptor potential channels as novel molecular targets to impair tumor neovascularization in renal cell carcinoma and other malignancies. *Anticancer Agents Med Chem.* 2014; 14: 296-312.
21. T Marichal, M Tsai, SJ Galli. Mast cells: potential positive and negative roles in tumor biology. *Cancer Immunol Res.* 2013; 1: 269-279.
22. D Hanahan, RA Weinberg. Hallmarks of cancer: the next generation. *Cell.* 2011; 144: 646-674.
23. D Ribatti, E Crivellato. Immune cells and angiogenesis. *J Cell Mol Med.* 2009; 13: 2822-2833.
24. EZ da Silva, MC Jamur, C Oliver. Mast cell function: a new vision of an old cell. *J Histochem Cytochem.* 2014; 62: 698-738.
25. SJ Galli, J Kalesnikoff, MA Grimbaldston, AM Piliponsky, CM Williams, et al. Mast cells as tunable effector and immunoregulatory cells: recent advances. *Annu Rev Immunol.* 2005; 23: 749-786.
26. MF Gurish, KF Austen. Developmental origin and functional specialization of mast cell subsets. *Immunity.* 2012; 37: 25-33.
27. MF Gurish, JA Boyce. Mast cells: ontogeny, homing, and recruitment of a unique innate effector cell. *J Allergy Clin Immunol.* 2006; 117: 1285-1291.

28. KD Stone, C Prussin, DD Metcalfe. IgE, mast cells, basophils, and eosinophils. *J Allergy Clin Immunol.* 2010; 125: S73-80.
29. MC Jamur, C Oliver. Origin, maturation and recruitment of mast cell precursors. *Front Biosci (Schol Ed.* 2011; 3: 1390-1406.
30. DD Metcalfe, RD Peavy, AM Gilfillan. Mechanisms of mast cell signaling in anaphylaxis. *J Allergy Clin Immunol.* 2009; 124: 639-646.
31. HR Katz, MB Raizman, CS Gartner, HC Scott, AC Benson, et al. Secretory granule mediator release and generation of oxidative metabolites of arachidonic acid via Fc-IgG receptor bridging in mouse mast cells. *J Immunol.* 1992; 148: 868-871.
32. JW Coleman, MR Holliday, I Kimber, KM Zsebo, SJ Galli. Regulation of mouse peritoneal mast cell secretory function by stem cell factor. IL-3 or IL-4, *J Immunol.* 1993; 150: 556-562.
33. SJ Galli, A Iemura, DS Garlick, C Gamba-Vitalo, KM Zsebo, et al. Reversible expansion of primate mast cell populations in vivo by stem cell factor. *J Clin Invest.* 1993; 91: 148-152.
34. V Supajatura, H Ushio, A Nakao, S Akira, K Okumura, et al. Differential responses of mast cell Toll-like receptors 2 and 4 in allergy and innate immunity. *J Clin Invest.* 2002; 109: 1351-1359.
35. SJ Galli, S Nakae, M Tsai. Mast cells in the development of adaptive immune responses. *Nat Immunol.* 2005; 6: 135-142.
36. RT Venkatesha, E Berla Thangam, AK Zaidi, H Ali. Distinct regulation of C3a-induced MCP-1/CCL2 and RANTES/CCL5 production in human mast cells by extracellular sig-

Top 10 Contributions on Cell Biology

- nal regulated kinase and PI3 kinase. *Mol Immunol.* 2005; 42: 581-587.
37. SJ Galli, M Tsai. Mast cells: versatile regulators of inflammation, tissue remodeling, host defense and homeostasis. *J Dermatol Sci.* 2008; 49: 7-19.
 38. HR Katz, JP Arm, AC Benson, KF Austen. Maturation-related changes in the expression of Fc gamma RII and Fc gamma RIII on mouse mast cells derived in vitro and in vivo. *J Immunol.* 1990; 145: 3412-3417.
 39. G Pejler, M Abrink, M Ringvall, S Wernersson. Mast cell proteases. *Adv Immunol.* 2007; 95: 167-255.
 40. LB Schwartz, TR Bradford. Regulation of tryptase from human lung mast cells by heparin. Stabilization of the active tetramer. *J Biol Chem.* 1986; 261: 7372-7379.
 41. LB Schwartz, AM Irani, K Roller, MC Castells, NM Schechter. Quantitation of histamine, tryptase, and chymase in dispersed human T and TC mast cells. *J Immunol.* 1987; 138: 2611-2615.
 42. C Huang, G Morales, A Vagi, K Chanasyk, M Ferrazzi, et al. Formation of enzymatically active, homotypic, and heterotypic tetramers of mouse mast cell tryptases. Dependence on a conserved Trp-rich domain on the surface. *J Biol Chem.* 2000; 275: 351-358.
 43. G Pejler, SD Knight, F Henningsson, S Wernersson. Novel insights into the biological function of mast cell carboxypeptidase A. *Trends Immunol.* 2009; 30: 401-408.
 44. FR Melo, I Waern, E Ronnberg, M Abrink, DM Lee, et al. A role for serglycin proteoglycan in mast cell apoptosis induced by a secretory granule-mediated pathway. *J Biol Chem.* 2011; 286: 5423-5433.

Top 10 Contributions on Cell Biology

45. M Abrink, M Grujic, G Pejler. Serglycin is essential for maturation of mast cell secretory granule. *J Biol Chem.* 2004; 279: 40897-40905.
46. T Braga, M Grujic, A Lukinius, L Hellman, M Abrink et al. Serglycin proteoglycan is required for secretory granule integrity in mucosal mast cells. *Biochem J.* 2007; 403: 49-57.
47. SJ Galli, N Borregaard, TA Wynn. Phenotypic and functional plasticity of cells of innate immunity: macrophages, mast cells and neutrophils. *Nat Immunol.* 2011; 12: 1035-1044.
48. JP Abonia, C Blanchard, BB Butz, HF Rainey, MH Collins, et al. Involvement of mast cells in eosinophilic esophagitis. *J Allergy Clin Immunol.* 2010; 126: 140-149.
49. RH Dougherty, SS Sidhu, K Raman, M Solon, OD Solberg, et al. Accumulation of intraepithelial mast cells with a unique protease phenotype in T(H)2-high asthma. *J Allergy Clin Immunol.* 2010; 125: 1046-1053 e1048.
50. RL Stevens, DS Friend, HP McNeil, V Schiller, N Ghildyal, et al. Strain-specific and tissue-specific expression of mouse mast cell secretory granule proteases. *Proc Natl Acad Sci U S A.* 1994; 91: 128-132.
51. MF Gurish, WS Pear, RL Stevens, ML Scott, K Sokol, et al. Tissue-regulated differentiation and maturation of a v-abl-immortalized mast cell-committed progenitor. *Immunity.* 1995; 3: 175-186.
52. DS Friend, N Ghildyal, KF Austen, MF Gurish, R Matsumoto et al. Mast cells that reside at different locations in the jejunum of mice infected with *Trichinella spiralis* exhibit sequential changes in their granule ultrastructure and chymase phenotype. *J Cell Biol.* 1996; 135: 279-290.
53. W Xing, KF Austen, MF Gurish, TG Jones. Protease phenotype of constitutive connective tissue and of induced mu-

- cosal mast cells in mice is regulated by the tissue. *Proc Natl Acad Sci U S A*. 2011; 108: 14210-14215.
54. HP McNeil, K Shin, IK Campbell, IP Wicks, R Adachi, et al. The mouse mast cell-restricted tetramer-forming tryptases mouse mast cell protease 6 and mouse mast cell protease 7 are critical mediators in inflammatory arthritis. *Arthritis Rheum*. 2008; 58: 2338-2346.
 55. K Shin, GF Watts, HC Oettgen, DS Friend, AD Pemberton, et al. Mouse mast cell tryptase mMCP-6 is a critical link between adaptive and innate immunity in the chronic phase of *Trichinella spiralis* infection. *J Immunol*. 2008; 180: 4885-4891.
 56. Q Sun, W Li, R She, D Wang, D Han, et al. Evidence for a role of mast cells in the mucosal injury induced by Newcastle disease virus. *Poult Sci*. 2009; 88: 554-561.
 57. L Scandiuizzi, W Beghdadi, E Daugas, M Abrink, N Tiwari, et al. Mouse mast cell protease-4 deteriorates renal function by contributing to inflammation and fibrosis in immune complex-mediated glomerulonephritis. *J Immunol*. 2010; 185: 624-633.
 58. A Lambiase, A Micera, S Bonini. Multiple action agents and the eye: do they really stabilize mast cells? *Curr Opin Allergy Clin Immunol*. 2009; 9: 454-465.
 59. DA de Souza, VD Toso, MR Campos, VS Lara, C Oliver et al. Expression of mast cell proteases correlates with mast cell maturation and angiogenesis during tumor progression. *PLoS One*. 2012; 7: e40790.
 60. EP. Beiträge zur Theorie und Praxis der histologischen Färbung, edn. 1878.

61. S Ch'ng, RA Wallis, L Yuan, PF Davis, ST Tan. Mast cells and cutaneous malignancies. *Mod Pathol.* 2006; 19: 149-159.
62. M Gulubova, T Vlaykova. Prognostic significance of mast cell number and microvascular density for the survival of patients with primary colorectal cancer. *J Gastroenterol Hepatol.* 2009; 24: 1265-1275.
63. G Ranieri, M Ammendola, R Patruno, G Celano, FA Zito, et al. Tryptase-positive mast cells correlate with angiogenesis in early breast cancer patients. *Int J Oncol.* 2009; 35: 115-120.
64. A Mangia, A Malfettone, R Rossi, A Paradiso, G Ranieri, et al. Tissue remodelling in breast cancer: human mast cell tryptase as an initiator of myofibroblast differentiation. *Histopathology.* 2011; 58: 1096-1106.
65. I Takanami, K Takeuchi, M Naruke. Mast cell density is associated with angiogenesis and poor prognosis in pulmonary adenocarcinoma. *Cancer.* 2000; 88: 2686-2692.
66. S Dabiri, D Huntsman, N Makretsov, M Cheang, B Gilks, et al. The presence of stromal mast cells identifies a subset of invasive breast cancers with a favorable prognosis, *Mod Pathol.* 2004; 17: 690-695.
67. P Conti, ML Castellani, D Kempuraj, V Salini, J Vecchiet, et al. Role of mast cells in tumor growth. *Ann Clin Lab Sci.* 2007; 37: 315-322.
68. BL Gruber, MJ Marchese, R Kew. Angiogenic factors stimulate mast-cell migration. *Blood.* 1995; 86: 2488-2493.
69. B Huang, Z Lei, GM Zhang, D Li, C Song, et al. SCF-mediated mast cell infiltration and activation exacerbate the inflammation and immunosuppression in tumor microenvironment. *Blood.* 2008; 112: 1269-1279.

70. L Benitez-Bribiesca, A Wong, D Utrera, E Castellanos. The role of mast cell tryptase in neoangiogenesis of premalignant and malignant lesions of the uterine cervix. *J Histochem Cytochem.* 2001; 49: 1061-1062.
71. MP Wymann, K Bjorklof, R Calvez, P Finan, M Thomast, et al. Phosphoinositide 3-kinase gamma: a key modulator in inflammation and allergy. *Biochem Soc Trans.* 2003; 31: 275-280.
72. MA Grimbaldston, AL Pearce, BO Robertson, BJ Coventry, G Marshman, et al. Association between melanoma and dermal mast cell prevalence in sun-unexposed skin. *Br J Dermatol.* 2004; 150: 895-903.
73. D Ribatti, N Finato, E Crivellato, A Marzullo, D Mangieri, et al. Neovascularization and mast cells with tryptase activity increase simultaneously with pathologic progression in human endometrial cancer. *Am J Obstet Gynecol.* 2005; 193: 1961-1965.
74. M Yoshii, A Jikuhara, S Mori, H Iwagaki, HK Takahashi, et al. Mast cell tryptase stimulates DLD-1 carcinoma through prostaglandin- and MAP kinase-dependent manners. *J Pharmacol Sci.* 2005; 98: 450-458.
75. NC Diaconu, R Kaminska, A Naukkarinen, RJ Harvima, IT Harvima. The increase in tryptase- and chymase-positive mast cells is associated with partial inactivation of chymase and increase in protease inhibitors in basal cell carcinoma. *J Eur Acad Dermatol Venereol.* 2007; 21: 908-915.
76. N Nonomura, H Takayama, K Nishimura, D Oka, Y Nakai, et al. Decreased number of mast cells infiltrating into needle biopsy specimens leads to a better prognosis of prostate cancer. *Br J Cancer.* 2007; 97: 952-956.
77. A Fleischmann, T Schlomm, J Kollermann, N Sekulic, H Huland, et al. Immunological microenvironment in prostate

cancer: high mast cell densities are associated with favorable tumor characteristics and good prognosis. *Prostate*. 2009; 69: 976-981.

78. MJ Carlini, MC Dalurzo, JM Lastiri, DE Smith, BC Vasallo, et al. Mast cell phenotypes and microvessels in non-small cell lung cancer and its prognostic significance. *Hum Pathol*. 2010; 41: 697-705.
79. A Johansson, S Rudolfsson, P Hammarsten, S Halin, K Pietras, et al. Mast cells are novel independent prognostic markers in prostate cancer and represent a target for therapy. *Am J Pathol*. 2010; 177: 1031-1041.
80. D Ribatti, D Guidolin, A Marzullo, B Nico, T Annese, et al. Mast cells and angiogenesis in gastric carcinoma. *Int J Exp Pathol*. 2010; 91: 350-356.
81. G Dyduch, K Kaczmarczyk, K Okon. Mast cells and cancer: enemies or allies? *Pol J Pathol*. 2012; 63: 1-7.
82. D Ribatti, A Vacca, B Nico, E Crivellato, L Roncali et al. The role of mast cells in tumour angiogenesis. *Br J Haematol*. 2001; 115: 514-521.
83. E Crivellato, CA Beltrami, F Mallardi, D Ribatti. The mast cell: an active participant or an innocent bystander? *Histol Histopathol*. 2004; 19: 259-270.
84. X Tan, S Essengue, J Talreja, J Reese, DJ Stechschulte, et al. Histamine directly and synergistically with lipopolysaccharide stimulates cyclooxygenase-2 expression and prostaglandin I(2) and E(2) production in human coronary artery endothelial cells. *J Immunol*. 2007; 179: 7899-7906.
85. M Muramatsu, J Katada, M Hattori, I Hayashi, M Majima. Chymase mediates mast cell-induced angiogenesis in hamster sponge granulomas. *Eur J Pharmacol*. 2000; 402: 181-191.

86. M Muramatsu, J Katada, I Hayashi, M Majima. Chymase as a proangiogenic factor. A possible involvement of chymase-angiotensin-dependent pathway in the hamster sponge angiogenesis model. *J Biol Chem.* 2000; 275: 5545-5552.
87. R Toth-Jakatics, S Jimi, S Takebayashi, N Kawamoto. Cutaneous malignant melanoma: correlation between neovascularization and peritumor accumulation of mast cells overexpressing vascular endothelial growth factor. *Hum Pathol.* 2000; 31: 955-960.
88. K Norrby. Mast cells and angiogenesis. *APMIS.* 2002; 110: 355-371.
89. I Feoktistov, S Ryzhov, AE Goldstein, I Biaggioni. Mast cell-mediated stimulation of angiogenesis: cooperative interaction between A2B and A3 adenosine receptors. *Circ Res.* 2003; 92: 485-492.
90. M Juczevska, L Chyczewski. Angiogenesis in cancer. *Rocz Akad Med Bialymst.* 1997; 42: 86-100.
91. LB Schwartz, RA Lewis, KF Austen. Tryptase from human pulmonary mast cells. Purification and characterization. *J Biol Chem.* 1981; 256: 11939-11943.
92. LB Schwartz. Tryptase, a mediator of human mast cells. *J Allergy Clin Immunol.* 1990; 86: 594-598.
93. V Payne, PC Kam. Mast cell tryptase: a review of its physiology and clinical significance. *Anaesthesia.* 2004; 59: 695-703.
94. G Pejler, E Ronnberg, I Waern, S Wernersson. Mast cell proteases: multifaceted regulators of inflammatory disease. *Blood.* 2010; 115: 4981-4990.
95. HZ Xia, CL Kepley, K Sakai, J Chelliah, AM Irani, et al. Quantitation of tryptase, chymase, Fc epsilon RI alpha, and Fc epsilon RI gamma mRNAs in human mast cells and ba-

- sophils by competitive reverse transcription-polymerase chain reaction. *J Immunol.* 1995; 154: 5472-5480.
96. S Wernersson, G Pejler. Mast cell secretory granules: armed for battle. *Nat Rev Immunol.* 2014; 14: 478-494.
97. JP Kankkunen, IT Harvima, A Naukkarinen. Quantitative analysis of tryptase and chymase containing mast cells in benign and malignant breast lesions. *Int J Cancer.* 1997; 72: 385-388.
98. A Imada, N Shijubo, H Kojima, S Abe. Mast cells correlate with angiogenesis and poor outcome in stage I lung adenocarcinoma. *Eur Respir J.* 2000; 15: 1087-1093.
99. B Nico, A Marzullo, P Corsi, A Vacca, L Roncali, et al. A possible role of tryptase in angiogenesis in the brain of mdx mouse, a model of Duchenne muscular dystrophy. *Neuroscience.* 2004; 123: 585-588.
100. D Ribatti, S Molica, A Vacca, B Nico, E Crivellato, et al. Tryptase-positive mast cells correlate positively with bone marrow angiogenesis in B-cell chronic lymphocytic leukemia. *Leukemia.* 2003; 17: 1428-1430.
101. D Ribatti, MG Ennas, A Vacca, F Ferreli, B Nico, et al. Tumor vascularity and tryptase-positive mast cells correlate with a poor prognosis in melanoma. *Eur J Clin Invest.* 2003; 33: 420-425.
102. V Goffredo, CD Gadaleta, A Laterza, A Vacca, G Ranieri. Tryptase serum levels in patients suffering from hepatocellular carcinoma undergoing intra-arterial chemoembolization: Possible predictive role of response to treatment. *Mol Clin Oncol.* 2013; 1: 385-389.
103. M Ammendola, R Sacco, G Donato, V Zuccala, E Russo, et al. Mast cell positivity to tryptase correlates with metastatic lymph nodes in gastrointestinal cancer patients treated surgically. *Oncology.* 2013; 85: 111-116.

104. I Marech, M Ammendola, R Sacco, GS Capriuolo, R Patruno, et al. Serum tryptase, mast cells positive to tryptase and microvascular density evaluation in early breast cancer patients: possible translational significance. *BMC Cancer*. 2014; 14: 534.
105. I Marech, M Ammendola, C Gadaleta, N Zizzo, C Oakley, et al. Possible biological and translational significance of mast cells density in colorectal cancer. *World J Gastroenterol*. 2014; 20: 8910-8920.
106. TC Theoharides, KD Alysandratos, A Angelidou, DA Delivanis, N Sismanopoulos, et al. Mast cells and inflammation. *Biochim Biophys Acta*. 2012; 1822: 21-33.
107. A Malfettone, N Silvestris, C Saponaro, G Ranieri, A Russo, et al. High density of tryptase-positive mast cells in human colorectal cancer: a poor prognostic factor related to protease-activated receptor 2 expression. *J Cell Mol Med*. 2013; 17: 1025-1037.
108. M Ammendola, C Leporini, I Marech, CD Gadaleta, G Scognamillo, et al. Targeting mast cells tryptase in tumor microenvironment: a potential antiangiogenetic strategy. *Biomed Res Int*. 2014; 2014: 154702.
109. CM Kielty, M Lees, CA Shuttleworth, D Woolley. Catabolism of intact type VI collagen microfibrils: susceptibility to degradation by serine proteinases. *Biochem Biophys Res Commun*. 1993; 191: 1230-1236.
110. D Ribatti, G Ranieri. Tryptase, a novel angiogenic factor stored in mast cell granules. *Exp Cell Res*. 2015; 332: 157-162.
111. MS Stack, DA Johnson. Human mast cell tryptase activates single-chain urinary-type plasminogen activator (pro-urokinase). *J Biol Chem*. 1994; 269: 9416-9419.

112. LB Schwartz, TR Bradford, BH Littman, BU Wintroub. The fibrinogenolytic activity of purified tryptase from human lung mast cells. *J Immunol.* 1985; 135: 2762-2767.
113. C Huang, GW Wong, N Ghildyal, MF Gurish, A Sali, et al. The tryptase, mouse mast cell protease 7, exhibits anticoagulant activity in vivo and in vitro due to its ability to degrade fibrinogen in the presence of the diverse array of protease inhibitors in plasma. *J Biol Chem.* 1997; 272: 31885-31893.
114. BL Gruber, MJ Marchese, K Suzuki, LB Schwartz, Y Okada, et al. Synovial procollagenase activation by human mast cell tryptase dependence upon matrix metalloproteinase 3 activation. *J Clin Invest.* 1989; 84: 1657-1662.
115. KC Fang, PJ Wolters, M Steinhoff, A Bidgol, JL Blount et al. Mast cell expression of gelatinases A and B is regulated by kit ligand and TGF-beta. *J Immunol.* 1999; 162: 5528-5535.
116. T Sendo, T Sumimura, Y Itoh, T Goromaru, K Aki, et al. Involvement of proteinase-activated receptor-2 in mast cell tryptase-induced barrier dysfunction in bovine aortic endothelial cells. *Cell Signal.* 2003; 15: 773-781.
117. Y Itoh, T Sendo, T Hirakawa, T Goromaru, S Takasaki, et al. Role of sensory nerve peptides rather than mast cell histamine in paclitaxel hypersensitivity. *Am J Respir Crit Care Med.* 2004; 169: 113-119.
118. RJ Blair, H Meng, MJ Marchese, S Ren, LB Schwartz, et al. Human mast cells stimulate vascular tube formation. Tryptase is a novel, potent angiogenic factor. *J Clin Invest.* 1997; 99: 2691-2700.
119. SJ Compton, JA Cairns, ST Holgate, AF Walls. The role of mast cell tryptase in regulating endothelial cell proliferation, cytokine release, and adhesion molecule expression: tryptase induces expression of mRNA for IL-1 beta and IL-8

- and stimulates the selective release of IL-8 from human umbilical vein endothelial cells. *J Immunol.* 1998; 161: 1939-1946.
120. LM Coussens, WW Raymond, G Bergers, M Laig-Webster, O Behrendtsen, et al. Inflammatory mast cells up-regulate angiogenesis during squamous epithelial carcinogenesis. *Genes Dev.* 1999; 13: 1382-1397.
121. J Lohi, I Harvima, J Keski-Oja. Pericellular substrates of human mast cell tryptase: 72,000 dalton gelatinase and fibronectin. *J Cell Biochem.* 1992; 50: 337-349.
122. R Kaminska, P Helisalmi, RJ Harvima, A Naukkarinen, M Horsmanheimo, et al. Focal dermal-epidermal separation and fibronectin cleavage in basement membrane by human mast cell tryptase. *J Invest Dermatol.* 1999; 113: 567-573.
123. M Lees, DJ Taylor, DE Woolley. Mast cell proteinases activate precursor forms of collagenase and stromelysin, but not of gelatinases A and B. *Eur J Biochem.* 1994; 223: 171-177.
124. W Zhang, G Stoica, SI Tasca, KA Kelly, CJ Meininger. Modulation of tumor angiogenesis by stem cell factor. *Cancer Res.* 2000; 60: 6757-6762.
125. IA Akers, M Parsons, MR Hill, MD Hollenberg, S Sanjar, et al. Mast cell tryptase stimulates human lung fibroblast proliferation via protease-activated receptor-2. *Am J Physiol Lung Cell Mol Physiol.* 2000; 278: L193-201.
126. C Lu, FD Zhao, XB Li, LH Yin. Up regulation of interleukin-8 expressions induced by mast cell tryptase via protease activated receptor-2 in endothelial cell line. *Chin Med J (Engl.)* 2005; 118: 1900-1906.
127. X Zhi, C Xu, H Zhang, D Tian, X Li, et al. Tryptase promotes atherosclerotic plaque haemorrhage in ApoE^{-/-} mice. *PLoS One.* 2013; 8: e60960.

128. TA McEachron, FC Church, N Mackman. Regulation of thrombin-induced plasminogen activator inhibitor-1 in 4T1 murine breast cancer cells. *Blood Coagul Fibrinolysis*. 2011; 22: 576-582.
129. TA McEachron, R Pawlinski, KL Richards, FC Church, N Mackman. Protease-activated receptors mediate crosstalk between coagulation and fibrinolysis. *Blood*. 2010; 116: 5037-5044.
130. AF Milia, MB Salis, T Stacca, A Pinna, P Madeddu, et al. Protease-activated receptor-2 stimulates angiogenesis and accelerates hemodynamic recovery in a mouse model of hindlimb ischemia. *Circ Res*. 2002; 91: 346-352.
131. M Belting, MI Dorrell, S Sandgren, E Aguilar, J Ahamed, et al. Regulation of angiogenesis by tissue factor cytoplasmic domain signaling. *Nat Med*. 2004; 10: 502-509.
132. H Mirza, V Yatsula, WF Bahou. The proteinase activated receptor-2 (PAR-2) mediates mitogenic responses in human vascular endothelial cells. *J Clin Invest*. 1996; 97: 1705-1714.
133. C Cicala. Protease activated receptor 2 and the cardiovascular system. *Br J Pharmacol*. 2002; 135: 14-20.
134. GM Hjortoe, LC Petersen, T Albrechtsen, BB Sorensen, PL Norby, et al. Tissue factor-factor VIIa-specific up-regulation of IL-8 expression in MDA-MB-231 cells is mediated by PAR-2 and results in increased cell migration. *Blood*. 2004; 103: 3029-3037.
135. VM Shpacovitch, T Brzoska, J Buddenkotte, C Stroh, CP Sommerhoff, et al. Agonists of proteinase-activated receptor 2 induce cytokine release and activation of nuclear transcription factor kappaB in human dermal microvascular endothelial cells. *J Invest Dermatol*. 2002; 118: 380-385.
136. N Asokanathan, PT Graham, J Fink, DA Knight, AJ Bakker, et al. Activation of protease-activated receptor (PAR)-1,

- PAR-2, and PAR-4 stimulates IL-6, IL-8, and prostaglandin E2 release from human respiratory epithelial cells. *J Immunol.* 2002; 168: 3577-3585.
137. ZH Wang, W Zhu, JP Tao, QY Zhang, M Wei. Stimulated mast cells promote maturation of myocardial microvascular endothelial cell neovessels by modulating the angiopoietin-Tie-2 signaling pathway. *Braz J Med Biol Res.* 2013; 46: 920-928.
138. H Urata, A Kinoshita, KS Misono, FM Bumpus, A Husain. Identification of a highly specific chymase as the major angiotensin II-forming enzyme in the human heart. *J Biol Chem.* 1990; 265: 22348-22357.
139. Y Kunori, M Koizumi, T Masegi, H Kasai, H Kawabata, et al. Rodent alpha-chymases are elastase-like proteases. *Eur J Biochem.* 2002; 269: 5921-5930.
140. UM Chandrasekharan, S Sanker, MJ Glynias, SS Karnik, A Husain. Angiotensin II-forming activity in a reconstructed ancestral chymase. *Science.* 1996; 271: 502-505.
141. GH Caughey, WW Raymond, PJ Wolters. Angiotensin II generation by mast cell alpha- and beta-chymases. *Biochim Biophys Acta.* 2000; 1480: 245-257.
142. S Sanker, UM Chandrasekharan, D Wilk, MJ Glynias, SS Karnik et al. Distinct multisite synergistic interactions determine substrate specificities of human chymase and rat chymase-1 for angiotensin II formation and degradation. *J Biol Chem.* 1997; 272: 2963-2968.
143. MW Kofford, LB Schwartz, NM Schechter, DR Yager, RF Diegelmann, et al. Cleavage of type I procollagen by human mast cell chymase initiates collagen fibril formation and generates a unique carboxyl-terminal propeptide. *J Biol Chem.* 1997; 272: 7127-7131.

144. KC Fang, WW Raymond, SC Lazarus, GH Caughey. Dog mastocytoma cells secrete a 92-kD gelatinase activated extracellularly by mast cell chymase. *J Clin Invest.* 1996; 97: 1589-1596.
145. E Tchougounova, A Lundequist, I Fajardo, JO Winberg, M Abrink, et al. A key role for mast cell chymase in the activation of pro-matrix metalloprotease-9 and pro-matrix metalloprotease-2. *J Biol Chem.* 2005; 280: 9291-9296.
146. BT Frank, JC Rossall, GH Caughey, KC Fang. Mast cell tissue inhibitor of metalloproteinase-1 is cleaved and inactivated extracellularly by alpha-chymase. *J Immunol.* 2001; 166: 2783-2792.
147. A Nakano, F Kishi, K Minami, H Wakabayashi, Y Nakaya, et al. Selective conversion of big endothelins to tracheal smooth muscle-constricting 31-amino acid-length endothelins by chymase from human mast cells. *J Immunol.* 1997; 159: 1987-1992.
148. S Takai, N Shiota, D Jin, M Miyazaki. Chymase processes big-endothelin-2 to endothelin-2-(1-31) that induces contractile responses in the isolated monkey trachea. *Eur J Pharmacol.* 1998; 358: 229-233.
149. J Taipale, J Lohi, J Saarinen, PT Kovanen, J Keski-Oja. Human mast cell chymase and leukocyte elastase release latent transforming growth factor-beta 1 from the extracellular matrix of cultured human epithelial and endothelial cells. *J Biol Chem.* 1995; 270: 4689-4696.
150. G Pejler, A Karlstrom. Thrombin is inactivated by mast cell secretory granule chymase. *J Biol Chem.* 1993; 268: 11817-11822.
151. GH Caughey. New developments in the genetics and activation of mast cell proteases. *Mol Immunol.* 2002; 38: 1353-1357.

152. L Lindstedt, M Lee, PT Kovanen. Chymase bound to heparin is resistant to its natural inhibitors and capable of proteolyzing high density lipoproteins in aortic intimal fluid, Atherosclerosis. 2001; 155: 87-97.
153. H Urata, S Hoffmann, D Ganten. Tissue angiotensin II system in the human heart. Eur Heart J. 1994; 15: 68-78.
154. T Terada, Y Matsunaga. Increased mast cells in hepatocellular carcinoma and intrahepatic cholangiocarcinoma. J Hepatol. 2000; 33: 961-966.
155. SY Tan, Y Fan, HS Luo, ZX Shen, Y Guo, et al. Prognostic significance of cell infiltrations of immunosurveillance in colorectal cancer. World J Gastroenterol. 2005; 11: 1210-1214.
156. A Cabanillas-Saez, JA Schalper, SM Nicovani, MI Rudolph. Characterization of mast cells according to their content of tryptase and chymase in normal and neoplastic human uterine cervix. Int J Gynecol Cancer. 2002; 12: 92-98.
157. M Muramatsu, J Katada, I Hayashi, M Majima. [Involvement of chymase in angiogenesis in hamster sponge granulomas]. Nihon Yakurigaku Zasshi. 1999; 114: 48P-54P.
158. M Muramatsu, M Yamada, S Takai, M Miyazaki. Suppression of basic fibroblast growth factor-induced angiogenesis by a specific chymase inhibitor, BCEAB, through the chymase-angiotensin-dependent pathway in hamster sponge granulomas. Br J Pharmacol. 2002; 137: 554-560.
159. J Katada, M Muramatsu, I Hayashi, M Tsutsumi, Y Konishi et al. Significance of vascular endothelial cell growth factor up-regulation mediated via a chymase-angiotensin-dependent pathway during angiogenesis in hamster sponge granulomas. J Pharmacol Exp Ther. 2002; 302: 949-956.
160. JM Arbeit, K Munger, PM Howley, D Hanahan. Progressive squamous epithelial neoplasia in K14-human papillomavirus type 16 transgenic mice. J Virol. 1994; 68: 4358-4368.

161. LM Coussens, Z Werb. Matrix metalloproteinases and the development of cancer. *Chem Biol.* 1996; 3: 895-904.
162. EI Deryugina, JP Quigley. Pleiotropic roles of matrix metalloproteinases in tumor angiogenesis: contrasting, overlapping and compensatory functions. *Biochim Biophys Acta.* 2010; 1803: 103-120.
163. A Russo, G Russo, M Peticca, C Pietropaolo, M Di Rosa, et al. Inhibition of granuloma-associated angiogenesis by controlling mast cell mediator release: role of mast cell protease-5. *Br J Pharmacol.* 2005; 145: 24-33.
164. IG Rojas, ML Spencer, A Martinez, MA Maurelia, MI Rudolph. Characterization of mast cell subpopulations in lip cancer. *J Oral Pathol Med.* 2005; 34: 268-273.
165. M Bialas, G Dyduch, J Szpor, S Demczuk, K Okon. Microvascular density and mast cells in benign and malignant pheochromocytomas. *Pol J Pathol.* 2012; 63: 235-242.
166. T Globa, L Saptefriti, RA Ceausu, P Gaje, AM Cimpean, et al. Mast cell phenotype in benign and malignant tumors of the prostate. *Pol J Pathol.* 2014; 65: 147-153.
167. M Wilk, L Liszka, P Palen, A Gabriel, P Laudanski. Intensity of angiogenesis and mast cell infiltration in cervical intraepithelial and invasive lesions - are they correlated? *Pathol Res Pract.* 2010; 206: 217-222.
168. M Nagata, N Shijubo, AF Walls, S Ichimiya, S Abe, et al. Chymase-positive mast cells in small sized adenocarcinoma of the lung. *Virchows Arch.* 2003; 443: 565-573.
169. ST Tan, RA Wallis, Y He, PF Davis. Mast cells and hemangioma. *Plast Reconstr Surg.* 2004; 113: 999-1011.