

## 20 a 23 de outubro de 2025

Majestic Hotel - Águas de Lindóia - SP - Brasil

Nanocomposite-modified screen-printed electrode based on f-MWCNT/AgNPs for diuron detection in Brazilian agricultural products

Adriana C. da Silva<sup>1</sup>, Thiago S. da Sena<sup>1</sup>, Igor G. S. Oliveira<sup>1</sup>, Fausto E. B. Junior<sup>2</sup>, Robson S. Souto<sup>2</sup>, Oswaldo C. Junior<sup>2</sup>, Marcos R. V. Lanza<sup>2</sup>, <u>Adriana E. de Carvalho<sup>1</sup></u>\*, Willyam R. P. Barros <sup>1</sup>

<sup>1</sup> Federal University of Grande Dourados, <sup>2</sup>Institute of Chemistry of São Carlos - University of São Paulo \*e-mail: adrianacarvalho@ufgd.edu.br

In this work, a screen-printed electrode (SPE) modified with a nanocomposite based on functionalized multi-walled carbon nanotubes (f-MWCNTs), chitosan (CS) and silver nanoparticles (AgNPs) was developed for the electrochemical detection of diuron (DIU), a widely used herbicide in agricultural practices. The AgNPs were synthesized by a greener route using lemon leaf extract as a natural reducing and stabilizing agent. The composite f-MWCNT@CS-AgNPs was drop-cast onto the working area of the SPE, resulting the sensing platform SPE/f-MWCNT@CS-AgNPs. Physicochemical and electrochemical characterizations revealed a significant increase in the electroactive area and heterogeneous electron transfer constant, especially when 0.70% CS was used. DFT calculations showed that modifying f-MWCNT@CS with AgNPs lowered its Gibbs free energy. The electrochemical behavior of DIU was studied by DPV, revealing a pHdependent irreversible oxidation consistent with a 2e<sup>-</sup>/2H<sup>+</sup> mechanism<sup>1</sup> after optimization. The analytical curve was linear in the range of 0.02 to 50.0 µmol L-1, with a limit of detection (LOD) of 5.0 nmol L<sup>-1</sup>. The platform showed high selectivity in the presence of common interferents and excellent recovery values (up to 115.6%) in real agricultural matrices - tomato, orange, tangerine and sugarcane. These results demonstrate the potential of the proposed platform as a simple, low-cost, and reproducible sensor for monitoring pesticide residues in food matrices.

## **Acknowledgments:**

The authors are grateful for the financial support provided by CAPES, CAPES and FAPESP.

**Reference:** [1] R.S. Oliveira, H.B. da Silva, C.C. de Souza, W.B. Veríssimo, M.A.C. Matos, T.P. Lisboa, R.C. Matos, Development of an electrochemical sensor utilizing recycled ABS filaments for 3D printing in the determination of diuron, **Microchemical Journal** 201 (2024) 110454.