

LETTER • OPEN ACCESS

Carbon dioxide emissions under different soil tillage systems in mechanically harvested sugarcane

To cite this article: A M Silva-Olaya *et al* 2013 *Environ. Res. Lett.* **8** 015014

View the [article online](#) for updates and enhancements.

You may also like

- [Soil properties and fenugreek \(*Trigonella foenum-graecum* L\) growth, yield and quality as affected by tillage systems, crop rotation and nitrogen fertilization under semi-arid climatic conditions](#)
Mouna Mechri, Afef Othmani, Taqi Raza et al.
- [Satellites reveal a small positive yield effect from conservation tillage across the US Corn Belt](#)
Jillian M Deines, Sherrie Wang and David B Lobell
- [Impacts of tillage practices on soil carbon stocks in the US corn-soybean cropping system during 1998 to 2016](#)
Zhen Yu, Chaoqun Lu, David A Hennessy et al.



The poster features a dark blue background with a large, stylized graphic of two hands shaking in a circle, rendered in shades of green and blue. The text is white and green. The ECS logo is in the top right, followed by the society's name and tagline. The meeting details are in the center right. A green circle in the bottom right contains a registration incentive. The bottom left has a call to action.

ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

ECS UNITED

247th ECS Meeting
Montréal, Canada
May 18-22, 2025
Palais des Congrès de Montréal

**Register to
save \$\$
before
May 17**

Unite with the ECS Community

Carbon dioxide emissions under different soil tillage systems in mechanically harvested sugarcane

A M Silva-Olaya¹, C E P Cerri¹, N La Scala Jr³, C T S Dias¹ and C C Cerri²

¹ ESALQ-USP, Avenida Pádua Dias, 11, Piracicaba, SP, Brazil

² CENA-USP, Avenida Centenário, 303, Piracicaba, SP, Brazil

³ FCAV-UNESP, Via de Acesso Prof. Paulo Donato Castellane s/n. 14884-900, Jaboticabal, SP, Brazil

E-mail: adrianasilva1221@gmail.com

Received 31 August 2012

Accepted for publication 10 December 2012

Published 12 February 2013

Online at stacks.iop.org/ERL/8/015014

Abstract

Soil tillage and other methods of soil management may influence CO₂ emissions because they accelerate the mineralization of organic carbon in the soil. This study aimed to quantify the CO₂ emissions under conventional tillage (CT), minimum tillage (MT) and reduced tillage (RT) during the renovation of sugarcane fields in southern Brazil. The experiment was performed on an Oxisol in the sugarcane-planting area with mechanical harvesting. An undisturbed or no-till (NT) plot was left as a control treatment. The CO₂ emissions results indicated a significant interaction ($p < 0.001$) between tillage method and time after tillage. By quantifying the accumulated emissions over the 44 days after soil tillage, we observed that tillage-induced emissions were higher after the CT system than the RT and MT systems, reaching 350.09 g m⁻² of CO₂ in CT, and 51.7 and 5.5 g m⁻² of CO₂ in RT and MT respectively. The amount of C lost in the form of CO₂ due to soil tillage practices was significant and comparable to the estimated value of potential annual C accumulation resulting from changes in the harvesting system in Brazil from burning of plant residues to the adoption of green cane harvesting. The CO₂ emissions in the CT system could respond to a loss of 80% of the potential soil C accumulated over one year as result of the adoption of mechanized sugarcane harvesting. Meanwhile, soil tillage during the renewal of the sugar plantation using RT and MT methods would result in low impact, with losses of 12% and 2% of the C that could potentially be accumulated during a one year period.

Keywords: emission after tillage, soil carbon, greenhouse gas, Brazil, ethanol, C footprint, CO₂ emissions

1. Introduction

The atmospheric concentration of greenhouse gases (GHGs) has been increased considerably in recent years as a result of

human activities. Carbon dioxide (CO₂) is the most important anthropogenic GHG; its annual emissions increased by about 80% between 1970 and 2004. Activities such as the burning of fossil fuels, the burning of forests and the loss of soil organic matter (Houghton *et al* 1992) have been recognized as the main causes of that increase.

The replacement of fossil fuels by fuels produced from biomass is an alternative approach that could reduce GHG emissions. Life-cycle analyses have found that the substitution



Content from this work may be used under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 licence](http://creativecommons.org/licenses/by-nc-sa/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

of ethanol produced from corn for gasoline effectively reduces emissions. However, the reduction in emissions may be even greater when sugarcane is used to produce ethanol from grazing land (Searchinger *et al* 2008). Brazil is the largest producer of sugarcane in the world and is consequently the primary exporter of ethanol. However, regional estimates indicate that Brazilian agriculture and land use changes are responsible for 75% of CO₂ emissions, 91% of CH₄ emissions and 94% of N₂O emissions (Cerri and Cerri 2007). These data create a need to develop agricultural practices that will decrease the environmental impact of sugarcane cultivation, ensure the sustainability of ethanol production and minimize the environmental impact.

Among mitigation processes, soil is an important natural reservoir of carbon (C). The emission of CO₂ from the soil is the second largest component of the global carbon cycle and is important in climatic variation (Reth *et al* 2005). Cultivation of the soil by tillage increases the mineralization of soil organic carbon (SOC) and the emission of CO₂ (Reicosky 1999). The decomposition of soil organic matter is increased by the physical disturbance caused by soil tillage, breaking down macroaggregates and exposing the carbon protected in their interiors to microbial processes (Cambardella and Elliott 1992). The amount of C lost in the form of CO₂ due to soil tillage practices is strongly correlated with the intensity of the disruption and the volume of soil disturbed by the implements used (Dao 1998, Rastogi *et al* 2002, La Scala *et al* 2006, Teixeira 2007, Reicosky and Archer 2007). Thus, depending on the soil type and management system, soils may be important sources or sinks of atmospheric carbon (Lal *et al* 1995, Bernoux *et al* 2005).

Thus, it is necessary to understand the role of the soil in the production of sugarcane. Soil used in sugarcane production may be either a mitigator or a contributor to global warming. Therefore, we seek to identify and quantify the impact of agricultural practices on GHG emissions and to measure the carbon footprint of ethanol derived from sugarcane production as a source of energy.

In Brazil, the area planted with sugarcane is currently about 8.4 million hectares (Conab 2011), of which approximately 50% is harvested mechanically, resulting in straw deposition on the soil surface (Galdos *et al* 2009, Razafimbelo *et al* 2006, Blair *et al* 1998), activity that should increase soil C stocks over time. In this sense, this study aimed to quantify the CO₂ emissions resulting from three systems of soil tillage used during the renovation of sugarcane plantations in the southeast region of Brazil. Significant soil CO₂ efflux due to tillage has been demonstrated in short and intermediate-term experiments (Reicosky and Lindstrom 1993, Reicosky *et al* 1997, Rochette and Angers 1999, Alvarez *et al* 2001, La Scala *et al* 2001, La Scala *et al* 2006); however, there is little information about the impact of that agricultural practice on the tropical soils that are mainly used for sugarcane production. Most of these studies quantified CO₂ fluxes resulting from one tillage operation, whereas during the renovation of sugarcane fields several tillage operations are conducted, which can result in a greater impact than reported in previous work. This research was a

qualitative evaluation of different types of soil tillage systems widely used this agricultural system.

2. Materials and methods

2.1. Study area

The study was carried out in the region of Piracicaba, in northeastern São Paulo State, Brazil (22° 33' S; 47° 30' W), in an area under sugarcane variety SP91-1049, which had been mechanically harvested (without burning) for the past six years, resulting in deposition of 14 Mg ha⁻¹ of dry matter from straw on the soil surface.

The soil is a Typic Hapludox (Soil Survey Staff 1999), with a pH of 5.5 and SOC stocks of 94.6 Mg ha⁻¹ (0–30 cm). The average particle sizes in the surface layer (0–30 cm) were 37% clay, 22% silt and 41% sand.

The climate of the region is classified according to the Koppen Climate Classification (1936) as Aw: tropical with a wet summer and a dry winter. The mean annual temperature and precipitation were 21.45 °C and 1271 mm, respectively.

2.2. Evaluated systems of soil tillage

We established four experimental plots with dimensions of 12 m × 35 m. Each plot had the same soil type and was located at the same topography. The treatments were as follows. (i) *Conventional tillage* (CT), which consisted of two heavy offset disk harrowing operations and a subsoiling operation. The first heavy offset disk harrowing was performed using a tool with 18 disks × 28" working at a speed of 6.5 km h⁻¹ perturbing the first 15–20 cm depth. The second heavy harrowing was conducted seven days after the first and used a tool with 24 disks × 29" working at 5.2 km h⁻¹ (20–25 cm depth). Twelve days after that operation row subsoiling was carried out, which employed a subsoiler with five shanks reaching nearly 45 cm depth. (ii) *Minimum tillage* (MT), in which two applications of herbicide were performed to chemically eliminate sugarcane ratoon (new shoots of sugarcane growing after harvest) followed by a subsoiling operation in the row planting. The second herbicide application was executed sixteen days after the first application; meanwhile, the subsoiling was performed 28 days after the same operation, once the plants' death had been verified. The tool used for that subsoiling was a subsoiler with seven shanks, with a difference of 50 cm between shanks, reaching a depth of 40 cm. (iii) *Reduced tillage* (RT), which involved two phases of mechanical elimination of the ratoon and two subsoiling operations. The mechanical eliminator of the ratoon was an implement working in two rows of planting simultaneously; composed of two plows which pull out the plants, lifting them to expose them to a set of mounted knives in a rotative spiral system that destroy the vegetal material and leave them on the soil surface. Seven days after the first mechanical elimination operation, the second elimination was performed through every other two planting rows. The time between these two operations was established to guarantee the death of a pest (*Sphenophorus levis*) that




Figure 1. The timeline of tillage events for each treatment: CT—conventional tillage; MT—minimum tillage; RT—reduced tillage and NT—no-tillage.

stays in the remaining root system of the old ratoon. The mechanical eliminator worked at a speed of 3.0 km h^{-1} , perturbing the soil to 20 cm depth. To complete the tillage operation system row subsoiling was executed 12 days after the last ratoon elimination; for that a subsoiler with five shanks working at 3.9 km h^{-1} , reaching between 20 and 30 cm of depth, was used. Finally, nine days after the last operation a cross-subsoiling was carried out at an angle of 45° using an implement with seven shanks reaching nearly 50 cm of depth and working at 4.4 km h^{-1} . (iv) *No-tillage* (NT), where the soil and sugarcane trash were left from the last harvesting procedure, which was performed about two weeks before starting the experiment.

The first operations in all treatments were performed simultaneously on the same day (7 October 2009). Figure 1 illustrates the tillage events over time for each treatment.

2.3. Measurement of soil CO_2 emissions, temperature and moisture

After the tillage procedures were finished, eight PVC soil collars (10 cm in diameter) were inserted into the soil in each experimental plot at a depth of 3 cm. The CO_2 flux measurements were initiated 24 h after this procedure and were repeated over a period of 44 days, once a day, in the early morning, by the end of which the CO_2 emissions in the plots with the experimental soil tillage systems reached values equivalent to those observed in the control treatment ($p < 0.05$).

The emission of CO_2 was monitored using a portable LI-COR system (model LI-8100; Lincoln, NE, USA), which monitors the changes in CO_2 concentration inside the chamber using an infrared gas analyzer (IRGA) (La Scala *et al* 2001, Panosso *et al* 2009).

Soil temperature was monitored by using a dial probe soil thermometer, which was inserted vertically to 12 cm from the soil surface close to the collars. Soil moisture was registered with a portable hydrosense system (TDR Probe, Campbell, USA) that estimated the percentage of moisture in the soil volume.




Figure 2. Soil temperature and precipitation under different systems of tillage during the study period. CT—conventional tillage; MT—minimum tillage; RT—reduced tillage and NT—no-tillage treatment.

2.4. Statistical analyses

The CO_2 flux was expressed on a mass basis ($\text{g m}^{-2} \text{ h}^{-1}$) and illustrated based on the time after soil tillage. This research was a qualitative evaluation of different types of soil tillage systems used in a sugarcane crop. In this way the approach was based on pseudo-replicates (8) for each treatment. A repeated-measure experimental design was used, meaning that the replicated data comprised measurements on the same experimental unit (Kepner and Robinson 1988). To compare treatments, we performed an *F*-test of analysis of variance using a GLM model and then compared the average values of the variables among treatments using Tukey's test at a significance level of $p < 0.05$.

The multiple correlation among moisture, temperature and CO_2 fluxes from each tillage system was evaluated through the Gauss–Newton method. All statistical analyses were performed using the Statistical Analysis Systems program (SAS Institute 1987).

The emissions accumulated over the study period were calculated by integration of the area under the emissions curve using the Origin 7 program (Origin Lab Corporation 2002).

3. Results and discussion

3.1. Soil temperature and moisture

The soil temperature trends over the study period are presented in figure 2. Significant differences among the plots ($p < 0.05$) were observed on days 7, 19, 27, 29, 30 and 31 and on the last four days of the study. Some of these days coincided with tillage events or with precipitation in the area. The average soil temperature over the entire study period was significantly different among the different soil tillage systems. In the CT system, the average temperature was 23.6°C , with maximum and minimum values of 28.6 and 19.8°C . In RT and MT the average soil temperatures were 22.9 and 22.7°C respectively, ranging between 26.7 and 19.4°C in RT and 26.9 and 20.0°C in MT. In the NT system the average soil temperature was 23.2 (ranging from 27.8 to 20.4°C).




Figure 3. Volumetric soil moisture under different systems of soil tillage and precipitation during the study period. CT—conventional tillage; MT—minimum tillage; RT—reduced tillage and NT—no-tillage treatment.

Soil disturbance due to tillage operations may modify the relationship between the volume of air and the volume of soil particles inducing changes in the soil thermal conductivity and affecting the soil temperature. Soil tillage exhibits lower conductivity compared to soil non-tillage (Hillel 1998).

Soil moisture was more differentiated than temperature among the tillage systems, with wider variations over the study period (figure 3). Differences in soil moisture were observed during 79% of the study period. The MT and NT treatments generally exhibited greater soil moisture compared to RT and CT.

The average soil moisture was highest ($p < 0.05$) in the NT (37.0%) and MT systems (31.3%), followed by the RT (24.3%) and CT (20.8%) systems. This pattern indicates that tillage systems that involve less physical disturbance help to maintain or conserve soil moisture. The tillage activity exposed more soil surface area to the atmosphere, providing a greater evaporative area and, consequently, a greater water loss. This effect depends on the depth, degree and frequency of soil tillage activities, the subsequent climatic conditions and the restructuring of the prepared layer (Salton and Mielniczuk 1995).

According to Smith *et al* (2003) soil temperature and moisture generate a direct effect on CO₂ emissions. Based on these findings, we determined the Pearson correlation coefficients (r) between the CO₂ flux from each plot and the soil temperature and soil moisture.

Our results indicate that these variables only had an influence on CO₂ flux in the CT ($r^2 = 0.44$). Multiple linear regression between emissions (FCO₂), volumetric moisture and soil temperature in the CT was established according to the following relationship:

$$\text{FCO}_2 = -0.087 + 0.024 \times \text{moisture} + 0.015 \times \text{temperature.} \quad (1)$$

Equation (1) suggests that increases in soil moisture and temperature would favor increase in the CO₂ emission process in the CT treatment. Several studies have indicated soil moisture as a controlling factor in CO₂ emissions and that the sensitivity of CO₂ flux to soil moisture is greater under the conventional system of tillage compared to no-till and minimum tillage treatments (Jabro *et al* 2008, La Scala *et al* 2006, Ussiri and Lal 2009). As the percolation of water is facilitated by soil tillage, the microbial activity is stimulated, which resulting in increases on CO₂ emissions (Jabro *et al* 2008, La Scala *et al* 2006).

During the time period assessed, the soil temperature exhibited only minor variations and remained close to the optimal temperature for microbial activity. The same trend has been reported in previous tests conducted in soil under sugarcane (Panosso *et al* 2009, La Scala *et al* 2006).

3.2. Evolution of CO₂ emissions

The average CO₂ emissions from the soil were higher in the CT plot than in the RT, MT and NT plots, with values of 0.82, 0.56, 0.51 and 0.49 g m⁻² h⁻¹, respectively (table 1). The analysis of variance indicated a highly significant interaction ($p < 0.001$) between the tillage method and the time after tillage, showing that time must be considered when evaluating the effect of soil tillage on emissions. Similar average emissions (but lower than those found in this study) were observed by La Scala *et al* (2001).

One day after performing the first tillage operations, we found no significant differences between treatments ($p < 0.05$) in soil CO₂ emissions. Alvarez *et al* (2001), in an Argentine study, reported similar trends to ours, where the CO₂ emissions one day after tillage operations were not significantly different from the previous day. Several studies have indicated that the effect of soil tillage on CO₂ emissions




Figure 4. CO₂ emissions after soil tillage using CT—conventional tillage; MT—minimum tillage; RT—reduced tillage and NT—no-tillage treatment.

Table 1. Mean $\pm$ standard error, maximum, minimum and total CO₂ emissions during the study period for conventional tillage (CT), reduced tillage (RT), minimum tillage (MT) and no-tillage (NT) treatments. (Note: means followed by the same capital letter within a row are not significantly different according to Tukey's test at the 5% level. $N = 232$.)

Soil tillage system	CO ₂ emissions			
	Mean	Maximum	Minimum	Total
	g m ⁻² h ⁻¹			g m ²
CT	0.82 A $\pm$ 0.37	1.89	0.17	881.6
RT	0.56 B $\pm$ 0.22	1.37	0.22	583.2
MT	0.51 B $\pm$ 0.17	0.95	0.22	537.0
NT	0.49 B $\pm$ 0.19	0.87	0.17	531.5

is immediate, with CO₂ flux decreasing considerably just a few hours after tillage operations (Omonode *et al* 2007, Al-Kaisi and Yin 2005, Reicosky *et al* 1997, Alvaro-Fuentes *et al* 2007).

The changes in CO₂ flux due to soil tillage over the study period may be better observed in figure 4. Emissions increased during the days when tillage operations occurred and after precipitation events. Significant differences ($p < 0.05$) in CO₂ flux between tillage systems were dependent on the day observed. CO₂ emissions were only found to be different between the treatments on the study days 7, 12, 13, 15, 19 and 20 (table 2).

On day 7 the CO₂ flux immediately after tillage in the RT system differed significantly from that observed on the previous day ($p < 0.05$). The emissions observed in the CT system did not differ from the average emission of the RT plot (table 2). The mechanical eliminator of the ratoon disturbed the soil at a shallower depth compared to harrowing; however, it caused a greater breakdown of the soil structure into smaller aggregates, exposing labile organic matter more quickly and releasing the CO₂ accumulated in the soil pores. Immediate increase in CO₂ emissions after soil tillage may be due to a rapid increase in microbial activity for decomposition of labile organic matter caused by the increased soil aeration, as was exposed by Al-Kaisi and Yin (2005), or may instead be due

to the phenomenon described by Rochette and Angers (1999) as 'degassing', which involves the physical expulsion of CO₂ from the soil at the time of tillage.

Teixeira *et al* (2011) assessed the CO₂ emissions due to the incorporation of sugarcane crop residues and the aggregate breaking after rotary tillage, and reported high emissions immediately after tillage operations with exponential decay over time. The behavior of exponential decay in the emission over time after tillage has been reported for other research performed in Brazil (La Scala *et al* 2009, 2008).

The highest CO₂ emissions from the soil in the CT system were observed 12 and 13 days after the first tillage operation (figure 4). Increases in soil moisture during this period due to precipitation events can be associated with the highest soil CO₂ emissions in this treatment. In studies performed on the Korean peninsula, the flow of CO₂ significantly increased around ten hours after precipitation and then strongly decreased until it had returned to the level observed before precipitation after four days (Kim *et al* 2010). The same trend was observed by La Scala *et al* (2005) in sugarcane fields in Brazil, with increases in the CO₂ flow during precipitation periods.

On the other hand, the in-row subsoiling operations on the CT and RT plots during day 19 of the study were the apparent source of the differences ($p < 0.05$) observed in the CO₂

Table 2. Mean values of CO₂ emissions at the measured time points for each of the soil tillage systems evaluated over the studied period. (Note: means followed by the same capital letter within a row are not significantly different according to Tukey's test at the 5% level.)

Day of evaluation	CO ₂ emissions (g m ⁻² h ⁻¹)			
	Conventional tillage	Minimum tillage	Reduced tillage	No-till treatment
1	0.57 (0.45) A	0.48 (0.19) A	0.64 (0.17) A	0.33 (0.11) A
2	0.36 (0.24) A	0.40 (0.14) A	0.45 (0.21) A	0.30 (0.12) A
3	0.45 (0.29) A	0.36 (0.11) A	0.40 (0.15) A	0.25 (0.11) A
4	0.53 (0.22) A	0.36 (0.12) A	0.44 (0.12) A	0.23 (0.05) A
5	0.17 (0.11) A	0.35 (0.17) A	0.29 (0.38) A	0.18 (0.07) A
6	0.75 (0.27) A	0.49 (0.21) A	0.64 (0.15) A	0.42 (0.11) A
7	1.13 (0.24) AB	0.53 (0.99) BC	1.37 (0.20) A	0.47 (0.11) C
8	0.79 (0.16) A	0.51 (0.26) A	0.48 (0.22) A	0.73 (0.12) A
9	0.97 (0.44) A	0.63 (0.01) A	0.42 (0.20) A	0.66 (0.12) A
10	0.98 (0.26) A	0.59 (0.13) A	0.40 (0.26) A	0.58 (0.20) A
12	1.41 (0.54) A	0.73 (0.14) B	0.45 (0.16) B	0.77 (0.22) B
13	1.89 (0.96) A	0.95 (0.22) B	0.48 (0.28) B	0.87 (0.12) B
15	1.05 (0.84) A	0.55 (0.20) AB	0.36 (0.17) B	0.57 (0.08) AB
16	1.14 (0.28) A	0.77 (0.19) A	0.54 (0.25) A	0.60 (0.17) A
19	1.52 (0.60) A	0.58 (0.32) B	0.84 (0.17) B	0.45 (0.13) B
20	1.25 (0.84) A	0.65 (0.26) AB	0.69 (0.20) AB	0.60 (0.08) B
21	0.52 (0.16) A	0.47 (0.11) A	0.59 (0.21) A	0.47 (0.13) A
22	0.90 (0.45) A	0.44 (0.31) A	0.65 (0.15) A	0.39 (0.13) A
24	0.88 (0.30) A	0.45 (0.24) A	0.60 (0.16) A	0.41 (0.13) A
27	0.80 (0.12) A	0.48 (0.20) A	0.51 (0.13) A	0.41 (0.15) A
28	0.65 (0.16) A	0.76 (0.47) A	0.90 (0.14) A	0.37 (0.17) A
29	0.47 (0.13) A	0.38 (0.08) A	0.22 (0.23) A	0.27 (0.12) A
30	0.44 (0.11) A	0.27 (0.11) A	0.42 (0.09) A	0.30 (0.13) A
31	0.45 (0.28) A	0.21 (0.13) A	0.35 (0.10) A	0.33 (0.08) A
35	0.79 (0.19) A	0.70 (0.17) A	0.72 (0.37) A	0.79 (0.23) A
37	0.76 (0.14) A	0.40 (0.05) A	0.66 (0.15) A	0.56 (0.13) A
40	0.93 (0.12) A	0.69 (0.13) A	0.73 (0.22) A	0.82 (0.14) A
42	0.61 (0.26) A	0.26 (0.19) A	0.44 (0.10) A	0.38 (0.10) A
44	0.60 (0.16) A	0.35 (0.08) A	0.50 (0.16) A	0.60 (0.17) A
Means	0.82 (0.37) A	0.51 (0.22) B	0.56 (0.17) B	0.49 (0.19) B

emissions from the soil on that day and on day 20 (table 2). The interaction between the incorporation of organic residues and the greater soil aeration may have positively affected oxidation, influencing CO₂ flow. A detailed observation of the soil structure after tillage activities allows us to estimate that the subsoiling operation performed in the CT affected a greater depth (approximately 45 cm), which is more important than the incorporation of residues in the magnitude of immediate CO₂ emissions (Reicosky and Lindstrom 1993). In an experiment conducted in São Paulo to evaluate the influence of the depth of the scarifier on CO₂ emissions, Teixeira (2007) found that the greater the working depth of this implement, the higher the emissions.

Although there were few significant differences in CO₂ flux between tillage systems on certain days, the tendency observed over the study period was for higher emissions in the CT system compared to the other tillage systems (figure 4). The heavy harrowing combined with subsoiling in the CT system in areas with straw generated a greater impact on CO₂ losses to the atmosphere during the study period.

3.3. Cumulated emissions

On quantifying the total emissions during the study period we found that they were 34%, 39% and 40% higher ($p < 0.05$) in the CT than the RT, MT and NT plots respectively

(table 1). The cumulative emissions in the RT, MT and NT did not differ significantly from each other. When comparing the accumulated emission in the CT plot with the NT plot, we observed that 350.09 g m⁻² of CO₂ was released to the atmosphere due to soil tillage and straw decomposition, amount equivalent to 954.79 kg ha⁻¹ of C-CO₂.

Meanwhile, the impact of the RT and MT systems was smaller; the increases in emissions in these treatments when compared to NT were 51.7 and 5.5 g m⁻², respectively, suggesting losses of 141 and 15.2 kg ha⁻¹ of C-CO₂.

The adoption of mechanical harvesting systems in sugarcane affect the dynamics of SOC, resulting in an annual increase of 1.2 Mg ha⁻¹ in C stocks in the 0–20 cm layer of the soil (Galdos *et al* 2009). Thus, soil tillage under conventional practices (CT) for sugarcane reformation would generate a loss equivalent to 80% of the C that could potentially be accumulated in this soil layer during one year of mechanical harvesting over a period of only 44 days.

The RT and MT systems have a smaller effect on CO₂ emissions; these would generate losses of 12% and 2% of the accumulated C in one year, respectively.

Several studies have suggested that tillage systems involving plowing and harrowing generate a greater impact on CO₂ emissions than systems using scarification and/or subsoiling as the sole method of soil tillage (Reicosky *et al* 1997, Dao 1998, Rochette and Angers 1999, Alvarez *et al*

2001, Al-Kaisi and Yin 2005, La Scala et al 2006, Jabro et al 2008). The area planted with sugarcane in Brazil is currently 8.4 million hectares (Conab 2011). Today, approximately 50% of the total area is harvested mechanically, which should increase soil C stocks over time.

However, it is estimated that about 60% of the sugarcane plantation area is under Oxisols (the same type of soil as in this study) and that between 15% and 20% of the plantations are renewed annually (0.75 to 1 million hectares). Losses of SOC may be caused as a function of the tillage system adopted during such renovation. Thus, the use of the CT system will result in an annual loss of C-CO₂ of about 0.72–0.96 Tg of C. However, with soil tillage using RT or MT methods, the loss of C-CO₂ would be reduced to approximately 0.10–0.14 Tg or 0.01 Tg, respectively.

The current production of Brazilian ethanol is based primarily on sugarcane, and this biofuel has both an energy and a GHG balance that makes it one of the most sustainable biofuels currently produced at commercial scales (Galdos et al 2010, Macedo et al 2008). Additional efforts to reduce GHG emissions in sugarcane production are necessary to guarantee the environmental sustainability of the ethanol produced from this crop.

When comparing C loss from tillage practices with C loss caused by other agricultural activities such as mineral and organic fertilization, vinasse application and liming in sugarcane fields we verified that tillage is an important source of CO₂ to the atmosphere. The intensity of soil tillage should be minimized to reduce CO₂ emissions from the soil, aiming for a sustainable condition.

Figueiredo et al (2010) quantified the GHG emission sources related to the sugarcane production in the agricultural sector in Brazil and found that mineral and organic N fertilization cause a loss of 162.9 kg C_{eq} ha⁻¹ year⁻¹; meanwhile, liming accounts for 49.8 kg C_{eq} ha⁻¹ year⁻¹. Similar values were reported by Galdos et al (2010) who estimated C loss due to mineral N fertilization and liming of the order of 110.7 and 74.7 kg C_{eq} ha⁻¹ year⁻¹ respectively.

In addition to soil C loss due to tillage, the use of fossil fuel to perform that agricultural operation causes CO₂ emissions. The selection of a tillage system involving fewer operations could reduce CO₂ emissions as a consequence of fuel savings. Koga et al (2003) found that a change from conventional tillage to minimum tillage could reduce the total CO₂ emissions by between 15% and 29% in northern Japan.

In this sense, despite soil tillage in the sugarcane crop being an activity performed every 5–6 years, the intensity of soil tillage should be minimized to reduce CO₂ emissions from the soil. It is greatly beneficial to select sustainable management practices that allow for increased carbon accumulation, improve soil quality and minimize CO₂ emissions from agricultural soils. These practices can help to reduce the carbon footprint of ethanol, increasing the environmental benefits of replacing fossil fuels with this biofuel.

4. Conclusions

Soil temperature was not a controlling factor in CO₂ emissions among the tillage systems evaluated. Soil moisture, however, showed a positive correlation with the CO₂ flux from the conventional tillage plot, suggesting that, in this tillage system, increased moisture favors the emission processes.

Soil tillage under conventional operations (CT) causes emissions of 350.09 g m⁻² of CO₂ to the atmosphere, a value that suggests the loss of 80% of the C that can potentially be stocked in the 0–20 cm soil layer as a result of adoption of harvesting of green cane. The impact of RT and MT on SOC loss was lower than that observed for CT, suggesting losses of 12% and 2% of the C stored in the 0–20 cm soil layer under those harvesting systems.

The adoption of a soil tillage system in sugarcane crop involving the RT and MT methods could contribute a decrease of the rate of GHG emissions from the agricultural sector in Brazil, enhancing the sustainability of Brazilian ethanol.

References

- Al-Kaisi M M and Yin X H 2005 Tillage and crop residue effects on soil carbon and carbon dioxide emission in corn-soybean rotations *J. Environ. Qual.* **34** 437–45
- Alvarez R, Alvarez C R and Lorenzo G 2001 Carbon dioxide fluxes following tillage from a mollisol in the Argentine rolling Pampa *Eur. J. Soil Biol.* **37** 161–6
- Alvaro-Fuentes J, Cantero-Martinez C, Lopez M V and Arrue J L 2007 Soil carbon dioxide fluxes following tillage in semiarid Mediterranean agroecosystems *Soil Tillage Res.* **96** 331–41
- Bernoux M, Cerri C C, Volkoff B, Carvalho M C S, Feller C, Cerri C E P, Eschenbrenner V, Piccolo M C and Feigl B J 2005 Gases do efeito estufa e estoque de carbono nos solos: inventário do Brasil *Cadernos de Ciência e Tecnologia* **22** 235–46
- Blair G J, Chapman L, Whitbread A M, Ball-Coelho B, Larsen P and Tiessen H 1998 Soil carbon changes resulting from sugarcane trash management at two locations in Queensland, Australia, and in North-East Brazil *Aust. J. Soil Res.* **36** 873–81
- Cambardella C A and Elliott E T 1992 Particulate soil organic-matter changes across a grassland cultivation sequence *Soil Sci. Soc. Am. J.* **56** 777–83
- Cerri C C and Cerri C E P 2007 Agricultura e aquecimento global *Boletim Informativo de Sociedade Brasileira da Ciência do Solo* **32** 40–4
- Conab 2011 *Acompanhamento de Safra Brasileira: Cana-de-Açúcar, Primeiro Levantamento, Abril/2011* (Brasília: Companhia Nacional de Abastecimento) p 19
- Dao T H 1998 Tillage and crop residue effects on carbon dioxide evolution and carbon storage in a Paleustoll *Soil Sci. Soc. Am. J.* **62** 250–6
- Figueiredo E B, Panosso A R, Romão R and La Scala N 2010 Greenhouse gas emission associated with sugar production in southern Brazil *Carbon Balance Manag.* **5** 1–7
- Galdos M, Cerri C and Cerri C 2009 Soil carbon stocks under burned and unburned sugarcane in Brazil *Geoderma* **153** 347–52
- Galdos M V, Cerri C C, Lal R, Bernoux M, Feigl B and Cerri C E P 2010 Net greenhouse gas fluxes in Brazilian ethanol production systems *GCB Bioenergy* **2** 37–44
- Hillel D 1998 *Environmental Soil Physics* (San Diego, CA: Academic)

- Houghton J, Callander B and Varney S 1992 *Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment* (Cambridge: Cambridge University Press)
- Jabro J D, Sainju U, Stevens W B and Evans R G 2008 Carbon dioxide flux as affected by tillage and irrigation in soil converted from perennial forages to annual crops *J. Environ. Manag.* **88** 1478–84
- Kepner J L and Robinson D H 1988 Nonparametric methods for detecting treatment effects in repeated-measures designs *J. Am. Stat. Assoc.* **456**–61
- Kim D G, Mu S, Kang S and Lee D 2010 Factors controlling soil CO₂ effluxes and the effects of rewetting on effluxes in adjacent deciduous, coniferous, and mixed forests in Korea *Soil Biol. Biochem.* **42** 576–85
- Koga N, Tsuruta H, Tsuji H and Nakano H 2003 Fuel consumption-derived CO₂ emissions under conventional and reduced tillage cropping systems in northern Japan *Agric. Ecosyst. Environ.* **99** 213–9
- Köppen W P 1936 *Das Geographische System der Klimate (Handbuch der Klimatologie Bd. 1 Teil C)* (Berlin: Gebrüder Borntraeger)
- Lal R, Fausey N and Eckert D 1995 Land use and soil management effects on emissions of radiatively active gases from two soils in Ohio *Soil Management and Greenhouse Effect (Advances in Soil Science Series)* ed R Lal, J Kimble, E Levine and B A Stewart (Boca Raton, FL: CRC Press) pp 41–59
- La Scala N, Bolonhezi D and Pereira G T 2006 Short-term soil CO₂ emission after conventional and reduced tillage of a no-till sugar cane area in southern Brazil *Soil Tillage Res.* **91** 244–8
- La Scala N, Lopes A, Marques J and Pereira G T 2001 Carbon dioxide emissions after application of tillage systems for a dark red latosol in southern Brazil *Soil Tillage Res.* **62** 163–6
- La Scala N, Lopes A, Panosso A R, Camara F T and Pereira G T 2005 Soil CO₂ efflux following rotary tillage of a tropical soil *Soil Tillage Res.* **84** 222–5
- La Scala N, Lopes A, Spokas K, Archer D and Reicosky D C 2009 Short-term temporal changes of bare soil CO₂ fluxes after tillage described by first-order decay models *Eur. J. Soil Sci.* **60** 258–64
- La Scala N, Lopes A, Spokas K, Bolonhezi D, Archer D and Reicosky D C 2008 Short-term temporal changes of soil carbon losses after tillage described by a first-order decay model *Soil Tillage Res.* **99** 108–18
- Macedo I C, Seabra J E A and Silva J 2008 Green house gases emissions in the production and use of ethanol from sugarcane in Brazil: the 2005/2006 averages and a prediction for 2020 *Biomass Bioenergy* **32** 582–95
- Omonode R A, Vyn T J, Smith D R, Hegymegi P and Gal A 2007 Soil carbon dioxide and methane fluxes from long-term tillage systems in continuous corn and corn-soybean rotations *Soil Tillage Res.* **95** 182–95
- Panosso A R, Marques J, Pereira G T and La Scala N 2009 Spatial and temporal variability of soil CO₂ emission in a sugarcane area under green and slash-and-burn managements *Soil Tillage Res.* **105** 275–82
- Rastogi M, Singh S and Pathak H 2002 Emission of carbon dioxide from soil *Curr. Sci.* **82** 510–7
- Razafimbelo T, Barthes B, Larre-Larrouy M C, De Luca E F, Laurent J Y, Cerri C C and Feller C 2006 Effect of sugarcane residue management (mulching versus burning) on organic matter in a clayey Oxisol from southern Brazil *Agric. Ecosyst. Environ.* **115** 285–9
- Reicosky D 1999 Effects of conservation tillage on soil organic carbon dynamics: field experiments in the US corn belt *Selected Paper from the 10th International Soil Conservation Organization (Purdue University and the USDA-ARS National Soil Erosion Research Laboratory, 24–29 May 1999)*
- Reicosky D C and Archer D W 2007 Moldboard plow tillage depth and short-term carbon dioxide release *Soil Tillage Res.* **94** 109–21
- Reicosky D C and Lindstrom M J 1993 Fall tillage method—effect on short-term carbon-dioxide flux from soil *Agron. J.* **85** 1237–43
- Reicosky D C, Dugas W A and Torbert H A 1997 Tillage-induced soil carbon dioxide loss from different cropping systems *Soil Tillage Res.* **41** 105–18
- Reth S, Reichstein M and Falge E 2005 The effect of soil water content, soil temperature, soil pH-value and the root mass on soil CO₂ efflux—a modified model *Plant Soil* **268** 21–33
- Rochette P and Angers D A 1999 Soil surface carbon dioxide fluxes induced by spring, summer, and fall moldboard plowing in a sandy loam *Soil Sci. Soc. Am. J.* **63** 621–8
- Salton J and Mielniczuk J 1995 Relações entre sistema de preparo, temperatura e umidade de um podzólico vermelho-escuro de Eldorado do Sul *Revista Brasileira de Ciencia do Solo* **19** 313–9
- Searchinger T, Heimlich R, Houghton R A, Dong F X, Elobeid A, Fabiosa J, Tokgoz S, Hayes D and Yu T H 2008 Use of US croplands for biofuels increases greenhouse gases through emissions from land-use change *Science* **319** 1238–40
- Smith K, Ball T, Conen F, Dobbie K, Massheder J and Rey A 2003 Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes *Eur. J. Soil Sci.* **54** 779–91
- Soil Survey Staff 1999 *Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys* 2nd edn (Agricultural Handbook no 436) (Washington, DC: US Government Printing Office)
- Teixeira L G 2007 Emissão de CO₂ de um Latossolo vermelho após preparo mecânico *Undergraduate Thesis Universidade Estadual Paulista Escola ‘Júlio de Mesquita Filho’ (Jaboticabal)* p 98
- Teixeira L G, Fukuda A, Panosso A R, Lopes A and La Scala N 2011 Soil CO₂ emission as related to the sugarcane crop residues incorporation and aggregates break after rotary tillage *Eng. Agríc.* **31** 1075–84
- Ussiri D A N and Lal R 2009 Long-term tillage effects on soil carbon storage and carbon dioxide emissions in continuous corn cropping system from an alfisol in Ohio *Soil Tillage Res.* **104** 39–47