

Essential oils for dairy calves: effects on performance, scours, rumen fermentation and intestinal fauna

F. H. R. Santos, M. R. De Paula, D. Lezier, J. T. Silva, G. Santos and C. M. M. Bittar[†]

Animal Science Department, University of São Paulo – USP/ESALQ, Avenida Pádua Dias, 11-13418-900, Piracicaba, SP, Brazil

(Received 26 November 2013; Accepted 15 January 2015; First published online 18 February 2015)

The first cause of death of dairy calves is often diarrhea which is mainly caused by pathogenic bacteria, which can result in excessive use of antibiotics. However, facing the increase concern by the industry and consumers, the use of antibiotics not only to control pathogens, but also to manipulate growth, has become a challenge. Alternative additives, such essential oils, have the potential to decrease antibiotic use, without reducing performance or increasing mortality of dairy calves. The objective of this study was to evaluate the use of a commercial blend of essential oils, incorporated into the calf starter and/or milk replacer to monitor the effect on overall calf performance, fecal scores and rumen fermentation parameters. A total of 30 Holstein calves received 6 l/day of a liquid diet, consisting of a commercial milk replacer containing 20% CP: 15% fat (EE). Calves had free choice access to water and calf starter. Weaning occurred at week 8, and calves were followed until the 10th week of age. Calves were assigned to one of the three treatment groups in a randomized block design. Treatments: (1) control without essential oils supplementation (C); (2) essential oils blend in the milk replacer at 400 mg/kg (MR) and (3) essential oils blend in the milk replacer (200 mg/kg) and starter feed (200 mg/kg) (MRS). From the 2nd week, calves were weighed and body measurements were taken, while concentrate intake and fecal scores were monitored daily. Blood samples were drawn weekly for determination of glucose and β-hydroxybutyrate. Fecal samples were collected weekly and analyzed for lactic acid bacteria and Enterobacteria; and ruminal fluid for determination of pH, short chain fatty acids, ammonia-N and counts of amylolytic and cellulolytic bacteria, and protozoa. Performance, fecal scores and intestines microorganisms were not affected by the essential oils supplementation. Ruminal and blood parameters were also not affected, with the exception the rumen ammonia-N concentration, with higher values when essential oils were supplemented in a combination of milk replacer and starter feed. Most of the evaluated parameters were affected by age of calves, mainly as a response to the increase in concentrate intake as animals' aged. Essential oils are promising substitutes for antibiotics. However, the dose and routes of administration deserve further studies, allowing a better animal performance and health to be achieved.

Keywords: ammonia nitrogen, dairy calf performance, diarrhea, intestinal fauna, rumen fermentation

Implications

This study was conducted to evaluate the inclusion of essential oils in starter feed and/or milk replacer for dairy calves as alternatives to the use of antibiotics for disease control and/or growth promotion in response to their prohibited use in animal feed in the European Union (Official Journal of the European Union, 2003). These antimicrobial effects may result in neonatal diarrhea reduction, improved rumen development, enhanced calf performance and overall a decrease in calf mortality rates.

Introduction

Antibiotics are often used in ruminant nutrition aiming to manipulate rumen fermentation and minimize the growth of microorganisms that are known to reduce performance and increase morbidity and mortality rates. Various antibiotics are forbidden in the European Union, owing to the allegation of possible bacteria development of resistance and food security concerns. Currently, there is no definitive scientific data that supports bacterial resistance as a result of the use of antibiotics in animal production. However, alternatives are needed for animal production to be optimized, maintaining the animal health and welfare, minimizing environmental risks and ensuring food safety. Faced with these impasses,

[†] E-mail: carlabittar@usp.br

the scientific community has been looking for new compounds that may be used as alternatives to antibiotics.

Recent literature shows that plant secondary metabolites are alternatives to modulate ruminal fermentation (Calsamiglia *et al.*, 2007; Macheboeuf *et al.*, 2008); control microorganisms present in the intestine (Bampidis *et al.*, 2006) and improve overall animal performance (Hill *et al.*, 2007; Chester-Jones *et al.*, 2010). These compounds are considered Generally Recognized as Safe according to the Food and Drug Administration, by virtue of being safe for human consumption.

The most common health problem in dairy calf rearing systems is the occurrence of diarrhea and the consequent high mortality, morbidity and poor performance. Therefore, the reduction of intestinal pathogenic microorganisms may improve animal health, well-being and animal performance. Etiological microorganisms responsible for diarrhea in calves may be modulated by essential oils (EOs), as it was observed for piglets (Manzanilla *et al.*, 2004). Among the effects observed for supplemented animals are improved feed digestion and efficiency, maintenance of beneficial intestinal microflora and inhibition of pathogenic microorganisms (Durmic and Blache, 2012). Hill *et al.* (2007) studied the effect of EOs in milk replacer and observed increases in starter feed intake and better feed efficiency in dairy calves. Similar results were observed by Chester-Jones *et al.* (2010), including the reduction of intestinal disorders.

Moreover, EOs may effect rumen development due to their effect on microorganism populations and subsequent change in the rumen fermentation profiles. However, changes in short chain fatty acids (SCFA) profile are variable, depending on dose and specific EO evaluated (Busquet *et al.*, 2006; Castillejos *et al.*, 2007). On the other hand, ammonia-N concentration decreased in response to EOs supplementation *in vitro* (Busquet *et al.*, 2006; Macheboeuf *et al.*, 2008) or *in vivo* (Benchaar *et al.*, 2007; Giannenas *et al.*, 2011). These changes in rumen SCFA and ammonia-N insert are a consequence of modulation of the rumen microorganism

populations. Giannenas *et al.* (2011) observed a decrease in counts of ammonia-hyper producing bacteria (AHP), while counts of cellulolytic bacteria and protozoa were not affected by EOs supplementation.

The objectives of the current study were to evaluate the use of a commercial blend of EOs, incorporated in milk replacer and starter feed in relation to improvements in fecal scores and the overall performance of the young dairy calves, as well as the effects of these oils on ruminal fermentation and microbial population.

Material and methods

A total of 30 newborn and colostrum-fed Holstein calves were individually housed at the experimental calf facility of the Department of Animal Science of the 'Luiz de Queiroz' College of Agriculture. Calves received 6 l of milk replacer daily (Sprayfo Violet® 20% CP : 15%EE, 12.5% solids; Sloten of Brazil Ltd, Santos, SP, Brazil), divided into two meals (0700 and 1800 h), and had free access to water and starter feed. Calves were randomly assigned by birth weight to one of the three treatments as follows: (1) control with no EOs supplementation (C); (2) supplementation of 400 mg/kg EO blend in the milk replacer (MR) and (3) supplementation of 200 mg/kg EO blend in the milk replacer and 200 mg/kg EOs blend the starter feed (MRS). The composition of the milk replacer and starter feed are summarized in Table 1. The starter feed was formulated according to National Research Council (2001) guidelines. EOs were blended into the starter feed using a 'Y' mixer (Lucato, Limeira, SP, Brazil). Individual doses of EO blend were mixed in the milk replacer at feeding time. The EO blend (Activo®, GRASP, PR, Brazil) was composed of a combination of several microencapsulated EO: carvacrol, cineole, cinnamaldehyde, pepper oil resin and mannanoligosaccharide as a vehicle.

The starter feed was supplied *ad libitum* and every morning the refusals were weighed so as to obtain the daily starter

Table 1 Chemical composition of starter feed and milk replacer

	Feeds		
	Concentrate with EO	Concentrate without EO	Milk replacer ¹
DM (g/kg)	868	871	956
Mineral matter (g/kg DM)	55	58	86
CP (g/kg DM)	231	230	204
Ether extract (g/kg DM)	35	37	145
NDF (g/kg DM)	146	148	0.3
ADF (g/kg DM)	72	73	0.0
N-NDF (g/kg total N)	103	89	0.0
N-ADF (g/kg total N)	52	42	0.0
Lignin (g/kg DM)	11	10	0.0
Non-fibrous carbohydrates (g/kg DM)	53	53	56
Total digestible nutrients (g/kg DM)	815	823	—

EO = essential oils; DM = dry matter.

¹Sprayfo Violeta Sloten of Brazil Ltda., Santos, São Paulo, Brazil.

feed intake. After abruptly weaning calves at 8 weeks of age, calves received starter concentrate and coastcross hay (*Cynodon* spp.) *ad libitum* until the 10th week of age.

Calves were weighed weekly, before the morning milk feeding, on a mechanical balance (ICS-300; Coimma Ltd, Dracena, SP, Brazil) and measurements of withers height, heart girth and hip width were taken. Every morning feces were scored for fluidity, color, consistency and odor, as described by Larson *et al.* (1977).

From the 2nd week of age fecal samples were collected for lactic acid bacteria (LAB) and *Enterobacteria* counting, after serial dilution with distilled water (10⁻¹ to 10⁻¹⁰). LAB were counted using Petri plates with nutrient agar Man Rogosa Sharper (Becton, Dickson and Company, MD, USA) (Kawakami *et al.*, 2011). Samples were placed in closed jars containing anaerobic condition generator (ANAEROBAC; Probac Bacteriological Products of Brazil Ltda., São Paulo, SP, Brazil) in a biochemical oxygen demand (BOD) incubator Model TE-402 (Tecnal, Piracicaba, SP, Brazil) at 32 ± 1°C for 48 ± 3 h. For *Enterobacteria* counts, Petrifilm plates (3M® Brazil; Sumaré, SP, Brazil) were used and samples were incubated at 32.5 ± 1°C in BOD incubator Model TE-402 for 24 ± 2 h. After the incubation period, counting proceeded by considering the areas with red colonies with yellow and/or red colonies with gas bubbles, with or without yellow borders positive for *Enterobacteria*.

From the 2nd week of life, ruminal fluid samples were collected weekly, using an oro-esophageal tube and a vacuum pump, with care to avoid saliva contamination. Samples (50 ml) were filtered through an appropriate cloth tissue and pH was immediately measured (TEC-5; Tecnal, Piracicaba, SP, Brazil). Samples were frozen for later analyses of SCFA and ammonia-N. Samples were centrifuged at 15 000 × *g* (Universal 320R; Hettich, Tuttlinger, German) for determination of SCFA, carried out as described by Palmquist and Conrad (1971); and ammonia-N nitrogen concentration by the method of Chaney and Marbach (1962), adapted for a biochemistry automatic reader (SBA-200; CELM, Barueri, SP, Brazil) using a 540 nm absorbance filter.

The quantification of cellulolytic and amylolytic bacteria were performed using mineral solution (Goshe, 1987) supplemented with 1% carboxymethylcellulose and soluble starch (Cromaline Fine Chemicals Ltd, Diadema, SP, Brazil). Plates were placed in anaerobic jars and incubated in BOD incubator (TE-402) at 38°C until the appearance of colonies. The methodology for quantification of protozoa used was an adaptation of the proposal by Dehority (1977). After sample filtration, 500 µl of rumen contents was added to 500 µl of 18.5% formaldehyde solution and then 2 ml of dye solution (formaldehyde 35% NaCl, methyl green and water) to give a solution of 1 : 5 in the final product. Counts were done using MacMaster-like chamber in a binocular microscope (Bioval, Jiangbei, China). Blood samples were taken weekly, 2 h after morning feeding, via jugular venipuncture using vacuum tubes containing sodium fluoride and potassium ethylenediaminetetraacetic acid. Samples were centrifuged at 2000 × *g* (20 min at 4°C) and plasma stored until analysis.

Specific enzymatic kits were used to analyze plasma concentrations of glucose (Glicose HK Liquiform – Ref.: 85; Labtest Diagnóstica S.A., Belo Horizonte, MG, Brazil) and β-hydroxybutyrate (BHBA, RANBUT – Ref.: RB1007, Randox Laboratories, Crumlin, UK) in an automatic biochemistry system (SBA-200, CELM, São Paulo SP, Brazil).

Starter feed and hay were periodically sampled, dried at 55°C and ground through a 1-mm mesh for dry matter (DM), ash, EE and CP determination according to Association of Official Analytical Chemists (2000). CP was determined by combustion (Dumas method), using a N analyzer by LECO, model FP-528 (St. Joseph, MI, USA) and CP was calculated by multiplying results by 6.25; free-ash NDF and ADF and lignin were determined, according to the method of Van Soest *et al.* (1991), using sodium sulfite and thermo stable amylase when required; and N-NDF and N-ADF were determined according to Licitra *et al.* (1996). The total digestible nutrients values were calculated according to the equation proposed by Weiss (1993) and non-fiber carbohydrates as suggested by Hall *et al.* (2000).

Statistics

Performance and plasma metabolites data were analyzed as repeated measures by the PROC MIXED of SAS according to the model (1) below. Data for fecal or rumen microorganisms counts were also performed by the model (1) after transformation to log 10. Significance was adopted for values of *P* < 0.05 for all parameters:

$$Y_{ijk} = \mu + T_i + B_j + W_k + T_i W_k + E_{ijk} \quad (1)$$

where Y_{ijk} is the response variable, μ the overall mean, T_i the treatment effect, B_j the block effect, W_k the age effect and E_{ijk} the residual effect.

Results and discussion

The starter feed intake, BW and daily weight gain were not affected by treatments (*P* > 0.05). There was an interaction between treatment and age for the starter feed intake (*P* < 0.05); however, this interaction was not observed for the treatments in the same week of life of the calves (Table 2). As might be expected, the age effect was significant (*P* < 0.0001) for the starter feed intake, since there is an increased need to consume more nutrients as animals age (Figure 1). Manzanilla *et al.* (2004) supplemented piglets with EO blend (0, 150 and 300 mg/kg) and observed similar results, with no difference for BW or daily weight gain.

Starter feed intake is negatively related to the liquid diet volume (milk or milk replacer) fed (Davis and Drackley, 1998). The starter feed intake increased over the weeks (Figure 1), but from the 6th week the shape of the curve changed with a dramatic increase from the 8th week. Data shows that the EO blend had no effect on acceptance of liquid or solid diet, suggesting that, if there was a change in palatability, it was not deleterious to feed intake. Even though supplementation with EO blend did not affect (*P* > 0.05)

Table 2 Average starter intake, BW, average daily gain and gain of body measurements of calves supplemented with an essential oils (EO) blend

	Treatments ¹			s.e.m.	P		
	C	MR	MRS		T	A	T × A
Starter Intake (g/day)	589.1	450.9	545.7	56.0	0.21	<0.0001	0.04
BW (kg)	50.7	46.6	49.3	1.5	0.18	<0.0001	0.82
Average daily gain (g)	395.5	378.3	392.1	36.1	0.91	<0.0001	0.77
Withers height gain (cm/week)	0.12	0.16	0.11	0.02	0.06	0.15	0.07
Heart girth gain (cm/week)	0.20	0.22	0.22	0.02	0.89	0.004	0.09
Hip width gain (cm/week)	0.05	0.05	0.05	0.01	0.78	0.04	0.15

C = control; MR = 400 mg/kg of EO in milk replacer; MRS = 200 mg/kg of EO in milk replacer and 200 mg/kg of EO in starter feed; T = treatment effect; A = age (week) effect; T × A = treatment and age interaction effect.

¹A total of 10 animals/treatment during 70 days of life.




Figure 1 Starter feed intake (g/day), according to week of age, of calves supplemented with an essential oils (EOs) blend. C = control; MR = 400 mg/kg EO in milk replacer, MRS = 200 mg/kg EO in milk replacer and 200 mg/kg in the starter feed.

BW and average daily gain (Figure 2), there was a significant effect of age for these variables ($P < 0.0001$; Table 2), which progressively increased until the 10th week of life. The weight gain values are similar to those found by Ferreira and Bittar (2011), in similar housing and climate conditions, when additives to improve the overall performance of dairy calves were tested.

Body measurement gains (withers height, heart girth and hip width) were also not affected by supplementation with the EO blend (Table 2). However, since calves were growing, there was a significant age effect ($P < 0.0001$) for all measures. Observed body measurements are slightly lower than those found by Ferreira and Bittar (2011), with sodium propionate supplementation as an alternative to antibiotics in similar experimental conditions.

The average fecal scores (fluidity, color, consistency and odor) were not affected by EO supplementation ($P > 0.05$), which was unexpected. However, the age of the calves had a significant effect ($P < 0.0001$) in changes of fecal scores of calves (Table 3), with decreasing scores after the 4th week of age. According to Larson *et al.* (1977), when the animal has diarrhea, score for fluidity is higher than 2.5 (considering a 1 to 4 scale). During week 4, calves in group MR (400 mg/kg of EO blend on milk replacer) exceeded the score of 2.5 (Figure 3). Bampidis *et al.* (2006) observed no differences for




Figure 2 BW (kg) and average daily gain (g/day), according to week of age, of calves supplemented with an essential oils (EOs) blend. C = control; MR = 400 mg/kg EO in milk replacer, MRS = 200 mg/kg EO in milk replacer and 200 mg/kg in the starter feed.

fecal scores of dairy calves supplemented with neomycin (2.58) or with oregano EO (2.67). According to Durmic and Blache (2012), the EO may assist in the maintenance of the intestinal flora balance, minimizing cases of diarrhea in animal husbandry. The supplementations of EO blend by different routes were not associated to any score of color, odor or consistency. The number of days that calves showed fluidity fecal score above 2.5 was not affected by EO blend supplementation ($P > 0.5$). However, the age effect was observed ($P < 0.0001$), with a reducing number of days as calves' aged, until reaching score 0 in weeks 9 and 10. The average total days with diarrhea was also not different among treatments ($P > 0.5$), with 13, 15 and 14 days for the

Table 3 Mean values of fecal scores and counts of intestinal microorganisms of calves supplemented with an essential oils (EO) blend

	Treatments ¹			s.e.m.	P		
	C	MR	MRS		T	A	T × A
Score²							
Fluidity	1.81	1.77	1.67	0.09	0.53	<0.0001	0.48
Color	3.84	4.01	4.23	0.22	0.45	<0.0001	0.93
Consistency	1.52	1.51	1.45	0.08	0.80	<0.0001	0.95
Odor	1.52	1.52	1.56	0.08	0.88	<0.0001	0.79
Microorganisms (log 10 CFU/gDM)							
Enterobacteria	4.45	4.53	4.28	0.24	0.65	0.0001	0.64
Lactic acid bacteria	7.01	7.13	7.01	0.26	0.86	0.0059	0.74

C = control; MR = 400 mg/kg of EO in milk replacer; MRS = 200 mg/kg of EO in milk replacer and 200 mg/kg of EO in starter feed; T = treatment effect; A = age (week) effect; T × A = treatment and age interaction effect.

¹A total of 10 animals/treatment during 70 days of life.

²Scale of score and fecal components by Larson *et al.* (1977): (1) normal-firm but not hard, (2) soft-does not hold form, (3) runny-spreads easily and (4) devoid of solid matter.




Figure 3 Fecal scores of fluidity, according to week of age, of calves supplemented with an essential oils (EOs) blend. C = control; MR = 400 mg/kg EO in milk replacer, MRS = 200 mg/kg EO in milk replacer and 200 mg/kg in the starter feed.

control, EO on replacer milk and EO in the total diet, respectively.

Control treatment had lower number of calves with score higher than 2.5 (67%) as compared with calves supplemented via liquid diet (90%) or via solid and liquid diet (80%); in addition, the fecal score and the scores of severity of diarrhea tended to be very low compared with non-supplemented calves. One possible explanation for these results is an adverse effect caused by EO in the calves' intestinal tract. Durmic and Blache (2012) reported that EO may be associated with reduced enzyme activity and changes in the anatomy and physiology of the intestinal lining epithelium, contributing therefore, to the development of intestinal problems.

The evaluated intestinal flora (Table 3) showed no significant difference in response to supplementation of EO in either route of supply ($P > 0.05$). The comparison between the beneficial microorganisms (LAB) and the harmful ones (*Enterobacteria*) helped in the conclusion that the intestinal flora of the calves was predominantly populated by bacteria that helps maintain balance and normal functioning of the

intestine. The count of LAB was about 7.0 log 10 colony-forming units (CFU/gDM), while the count of *Enterobacteria* was 4.5 log 10 CFU/gDM (Table 4). These results confirm the fecal scores results discussed, since only the calves that received EO in a liquid diet had fecal score (fluidity) above 2.5 on the 4th week of age (Figure 3). Similar to the results of the fecal scores, the *Enterobacteria* counts were affected only by the age effect ($P < 0.0001$), with decreasing counts as calves' aged.

Despite the decreased counts of LAB over the weeks, this reduction was much smaller when compared with the decrease in the *Enterobacteria* counts. As a whole, the microbial flora tends to decrease or stabilize with the immune system and animal development. It was observed that the LAB counts were almost constant up to 6 weeks and decreased thereafter while at this point *Enterobacteria* were in higher concentration. That occurred probably because of the control action that *Lactobacillus* strains have on *Escherichia coli*. From week 6, both groups of microorganisms had reduced counts, indicating stability of the intestinal tract flora, which may be confirmed by fecal characteristics, since at the 6th week fluidity scores were lower than 2 for supplemented calves.

There were no differences in ruminal parameters that might have been reflected in calf performance or rumen development. Rumen pH were not affected by treatments ($P > 0.7$) or the age of the calves ($P > 0.1$; Table 4). SCFA, the fermentation end products, are highly correlated with starter intake. After weaning, calves showed a significant increase in starter feed intake, which resulted in similar response for the SCFA concentrations, as well as for the acetate to propionate ratio (C2 : C3; Table 4). The main SCFA (acetate, propionate and butyrate) concentrations were significantly higher ($P < 0.0001$) after weaning. However, the SCFA concentrations were not affected by EO supplementation ($P > 0.05$). Although all SCFA increased as calves' aged, with increasing starter intake, and consequent rumen development, propionate had a higher increase in concentration,

Table 4 Mean values of pH, ammonia nitrogen (ammonia-N) and short chain fatty acids (SCFA) of calves supplemented with an essential oils (EO) blend

	Treatments ¹			s.e.m.	P		
	C	MR	MRS		T	A	T × A
pH	5.81	5.92	5.82	0.12	0.81	0.11	0.06
Ammonia-N (mg/dl)	11.5b	12.7b	16.7a	1.5	0.04	0.09	0.92
SCFA (μmol/ml)							
Acetic (C2)	57.37	56.06	62.63	3.42	0.34	<0.0001	0.16
Propionic (C3)	23.78	20.74	26.51	2.58	0.28	<0.0001	0.11
Butyric (C4)	8.92	7.86	8.78	0.88	0.66	<0.0001	0.25
Total	93.58	88.09	101.29	6.91	0.39	<0.0001	0.09
C2 : C3	2.65	2.86	2.69	0.21	0.76	0.0030	0.002

C = control; MR = 400 mg/kg of EO in milk replacer; MRS = 200 mg/kg of EO in milk replacer and 200 mg/kg of EO in starter feed; T = treatment effect; A = age (week) effect; T × A = treatment and age interaction effect.

¹A total of 10 animals/treatment during 70 days of life.

Table 5 Mean values of rumen amylolytic bacteria, cellulolytic bacteria and protozoa counts of calves supplemented with an essential oils (EO) blend

	Treatments ¹			s.e.m.	P		
	C	MR	MRS		T	A	T × A
Amylolytic (log 10 CFU/ml)	3.92	3.98	3.94	0.16	0.96	0.14	0.28
Cellulolytic (log 10 CFU/ml)	4.11	3.95	3.98	0.11	0.45	0.0055	0.12
Protozoa (n/ml)	4.91	5.15	4.62	2.56	0.32	<0.0001	0.82

C = control; MR = 400 mg/kg of EO in milk replacer; MRS = 200 mg/kg of EO in milk replacer and 200 mg/kg of EO in starter feed; T = treatment effect; A = age (week) effect; T × A = treatment and age interaction effect.

¹A total of 10 animals/treatment during 70 days of life.

which reduced the C2 : C3 ratio at weaning. Similar results were found by Benchaar *et al.* (2006) when dairy cows supplemented with EO blend (2 g/day) containing thymol, eugenol and limonene and total SCFA, propionate, butyrate or C2 : C3 ratio were not affected. In subsequent studies, Benchaar *et al.* (2007) supplemented dairy cows with a smaller dose (0.75 g/day) of the same EO blend, with no significant effects on rumen fermentation profile. Apparently, EO at the doses tested in the present study were not able to modulate the rumen fermentation.

The ammonia-N concentrations were altered with the EO blend supplementation ($P < 0.05$). Calves receiving EO in the total diet (MRS) presenting significantly higher ($P < 0.05$) concentrations than calves not supplemented or supplemented only via liquid diet (Table 4). These results disagree with the results found in dairy sheep by Giannenas *et al.* (2011), where the concentrations of ammonia-N decreased with EO supplementation. The EO are recognized as a modulator of the nitrogen metabolism in the ruminal environment, reducing the deamination and also the population of AHP. Busquet *et al.* (2006) observed significant decrease of ammonia-N concentration with increasing dose of EO tested *in vitro* (32.2 mg/dl in the control group and 11.6 mg N/100 ml at the dose of 3.000 mg/l). Macheboeuf *et al.* (2008) conducted *in vitro* studies with different EO doses in sheep ruminal fluid and the ammonia-N

concentration varied depending on the EO doses. The cinnamaldehyde, cinnamon EO, was the only one responsible for raising the concentrations of ammonia-N in a 3 mM EO dose. In the current study, the concentration of ammonia-N for the treatment with EO supply on milk replacer and starter feed (MRS) was high from week 2 and has remained high until the end of the experiment at week 10, as compared with the other treatments.

Counts of protozoa, cellulolytic and amylolytic bacteria were not affected by the addition of EO in the diet of dairy calves (Table 5). In studies conducted by Benchaar *et al.* (2007), high producing dairy cows supplemented with EO had counts of ruminal microorganisms similar to the data observed in this present study. The cellulolytic bacteria were not affected by supplementation and maintained count values similar to the control group. The same was observed for protozoa. However, cellulolytic bacteria and protozoa counts were affected as calves' aged ($P < 0.05$). Protozoa counts increased progressively as animals' aged and ate more starter feed. After weaning, at the 8th week of life, when there is a sharp increase in starter intake (Figure 1), there is also an increase in protozoa counts. Protozoa use bacteria as the main source of amino acids and nucleic acids, and engulfment occurs more intensively in diets rich in grains (Russell, 2002).

The counts of cellulolytic bacteria showed fluctuation, and with the increasing consumption of starter feed, there was a




Figure 4 Plasma glucose (mg/dL) and β -hydroxybutyrate concentration (mmol/l), according to week of age, of calves supplemented with an essential oils (EOs) blend. C = control; MR = 400 mg/kg EO in milk replacer, MRS = 200 mg/kg EO in milk replacer and 200 mg/kg in the starter feed.

decrease in counts of these microorganisms. This occurred probably due to the reduced pH that affects the growth viability of microorganisms that degrade fiber. Even though the lowest average pH was higher than 5.0, it is possible that even lower pH have occurred during other periods of the day. The growth of amylolytic bacteria was constant over the weeks, and after weaning there was a slight drop in growth. These data do not agree with the study of Anderson *et al.* (1987), where the average proportion of amylolytic bacteria from total anaerobic bacteria tended to increase over the weeks. According to Beharka *et al.* (1998), amylolytic microorganisms are more tolerant to variations in pH and therefore do not undergo oscillations like the cellulolytic bacteria.

The low and very variable (SEM = 56 g) concentrate intake during the 1st weeks of life, associated with the EO dose tested, may have decreased the opportunity to show its effect on microbial rumen population. Data suggest that a higher dose should be used for dairy calves so a change in rumen flora and consequent change in fermentation profile, which may affect positively rumen development, should be tested.

Glucose plasma concentrations were not affected ($P > 0.5$) by EO supplementation (107.21, 105.74 and 102.93, respectively, for C, MR and MRS), and observed values are in agreement with the literature (Shingu *et al.*, 2007). During the first 2 to 3 weeks of age calves exhibit a high capacity for glucose utilization, because the rumen is not developed, so the SCFA can not be used as the primary energy source.

According to Quigley *et al.* (1991) plasma glucose values are higher during the period when the calf has the liquid diet as the main source of nutrients, as compared with weaned calves. Glucose concentrations were high during the 1st weeks (100 to 140 mg/dL), but decreased progressively until reaching values similar to those observed in adult ruminants (70 to 80 mg/dL; Figure 4). After weaning the main source of energy comes from SCFA produced by ruminal microbiota stimulated by the consumption of solid feed.

Another plasma metabolite highly correlated to starter intake and consequently to rumen development is the BHBA (Quigley *et al.*, 1991). Plasma BHBA concentrations were also not altered ($P > 0.5$) in response to EO supplementation (0.14, 0.14 and 0.19, respectively, for C, MR and MRS). According to Quigley and Bernard (1992), BHBA plasma concentrations before weaning are lower than 0.2 mmol/l, as observed in this study (Figure 4). However, BHBA concentrations were affected by age ($P < 0.0001$), with increasing values from 0.15 mmol/l at week 8 to 0.50 mmol/l at week 10, featuring rumen development of these calves at weaning time.

Conclusion

The EO blend supplementation, in doses and routes used, had no effect on performance or improved health of dairy calves, with no decrease on the incidence of diarrhea. However, the intestinal flora of calves contained a higher proportion of beneficial microorganisms, leading to the conclusion that EO did not adversely affect the intestinal microbial population. The ruminal and blood parameters were not negatively affected by supplementation; yet, caused no beneficial effect on the performance or rumen development. EO are considered promising replacements for antibiotics. However, dose and the routes of EO supply deserve more studies.

Acknowledgments

The authors wish to express their appreciation for the financial support provided by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) as scholarship; and GRASP® Company for the tested essential oils supply.

References

- Anderson KL, Nagaraja TG, Morrill JL, Avery TB, Galitzer SJ and Boyer JE 1987. Ruminal microbial development in conventionally or early-weaned calves. *Journal of Animal Science* 64, 1215–1226.
- Association of Official Analytical Chemists 2000. *Official methods of analysis*, 17th edition. AOAC, Arlington, VA, USA.
- Bampidis VA, Christodoulou V, Florou-Paneri P and Christaki E 2006. Effect of dried oregano leaves versus neomycin in treating newborn calves with colibacillosis. *Journal of Veterinary Medicine* 53, 154–156.
- Beharka AA, Nagaraja TG, Morrill JL, Kennedy GA and Klemm RD 1998. Effects of form of the diet on anatomical, microbial, and fermentative development of the rumen of neonatal calves. *Journal of Dairy Science* 81, 1946–1955.
- Benchaar C, Petit HV, Berthiaume RT, Whyte D and Chouinard PY 2006. Effects of dietary addition of essential oils and monensin premix on digestion, ruminal

fermentation characteristics, milk production, and milk composition in dairy cows. *Journal of Dairy Science* 89, 4352–4364.

Benchaar C, Petit HV, Berthiaume R, Ouellet DR, Chiquette J and Chouinard PY 2007. Effects of essential oils on digestion, ruminal fermentation, rumen microbial populations, milk production, and milk composition in dairy cows fed alfalfa silage or corn silage. *Journal of Dairy Science* 90, 886–897.

Busquet M, Calsamiglia S, Ferret A and Kamel C 2006. Screening for effects of plant extracts and secondary plant metabolites on rumen microbial fermentation. *Animal Feed Science and Technology* 123/124, 597–613.

Calsamiglia S, Busquet M, Cardozo PW, Castillejos L, Ferret A 2007. Invited review: essential oils as modifiers of rumen microbial fermentation. *Journal of Dairy Science* 90, 2580–2595.

Castillejos L, Calsamiglia S, Ferret A, Losa R 2007. Effects of dose and adaptation time of a specific blend of essential oil compounds on rumen fermentation. *Animal Feed Science and Technology* 132, 186–201.

Chaney AL and Marbach EP 1962. Modified reagents for determination of urea and ammonia. *Clinical Chemistry* 8, 130–132.

Chester-Jones H, Steiner T, Watkins M, Taylor D, Ziegler D, Raeth-Knight M and Golombeski G 2010. Pre- and post-weaning performance and health of calves fed milk replacers and calf starters with or without essential oils. *Journal of Animal Science* 88, 421.

Davis CL and Drackley JK 1998. The development nutrition and management of the young calf, 1st edition. Iowa State University Press, Ames, IA, USA.

Dehority BA 1977. Classification and morphology of rumen protozoa, 1st edition. University of Ohio, Columbus, OH, USA.

Durmic Z and Blache D 2012. Bioactive plants and plant products: effects on animal function, health and welfare. *Animal Feed Science and Technology* 176, 150–162.

Ferreira LS and Bittar CMM 2011. Performance and plasma metabolites of dairy calves fed starter containing sodium butyrate, calcium propionate or sodium monensina. *Animal Journal* 5, 239–245.

Giannenas I, Skoufos J, Giannakopoulos C, Wiemann M, Gortzi O, Lalas S and Kyriazakis I 2011. Effects of essential oils on milk production, milk composition, and rumen microbiota in Chios dairy ewes. *Journal of Dairy Science* 94, 5569–5577.

Goshe TK 1987. Measurement of cellulase activities. *Pure and Applied Chemistry* 59, 257–268.

Hall MB, Jennings JP, Lewis BA and Robertson JB 2000. Evaluation of starch analysis methods for feed samples. *Journal of the Science of Food and Agriculture* 81, 17–21.

Hill TM, Aldrich JM, Schlotterbeck RL and Bateman HG 2007. Apex plant botanicals for neonatal calf milk replacers and starters. *The Professional Animal Scientist* 23, 521–526.

Kawakami SI, Yamada T, Nakanishi N and Cai Y 2011. Feeding of lactic acid bacteria and yeast affects fecal flora of Holstein calves. *Journal Animal and Veterinary Advances* 10, 269–271.

Larson LL, Owen FG, Albright JL, Appleman RD, Lamb RC and Muller LD 1977. Guidelines toward more uniformity in measuring and reporting calf experimental data. *Journal of Dairy Science* 60, 989–991.

Licitra G, Hernandez TM and Van Soest PJ 1996. Standardization of procedures for nitrogen fractionation of ruminant feeds. *Animal Feed Science and Technology* 57, 347–358.

Macheboeuf D, Morgavi DP, Papon Y, Mousset JL and Arturo-Schaan M 2008. Dose-response effects of essential oils on in vitro fermentation activity of the rumen microbial population. *Animal Feed Science and Technology* 145, 335–350.

Manzanilla EG, Perez JF, Martin M, Kamel C, Baucells F and Gasa J 2004. Effect of plant extracts and formic acid on the intestinal equilibrium of early-weaned pigs. *Journal of Animal Science* 82, 3210–3218.

National Research Council 2001. Nutrient requirement of dairy cattle, 7th revised edition. NRC and National Academy of Science, Washington, DC, USA.

Official Journal of the European Union 2003. Regulation (EC) No1831/2003 of the European Parliament and Council of 22 September 2003 on additives for use in animal nutrition. OJEU. Retrieved 31 November 2014, from <http://eurex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L2003:268:0029:0043:EN:pdf>.

Palmquist D and Conrad H 1971. Origin of plasma fatty acid in lactating dairy cows fed high fat diets. *Journal of Dairy Science* 54, 1025–1033.

Quigley JD and Bernard JK 1992. Effects of nutrient source and time of feeding on changes in blood metabolites in young calves. *Journal of Animal Science* 70, 1543–1549.

Quigley JD, Caldwell LA and Sinks GD 1991. Changes in plasma volatile fatty acids in response to weaning and feed intake in young calves. *Journal of Dairy Science* 74, 258–263.

Russel JB 2002. Rumen microbiology and its role in ruminant nutrition, 1st edition. Blackwell Publishing Ltd, Oxford, UK.

Shingu H, Hayashi H, Touno E, Oshibe A, Kushibiki S, Oda S, Katoh K and Obara Y 2007. Characteristics of developmental changes in the kinetics of glucose and urea in Japanese Black calves: comparison with Holstein calves. *Journal of Animal Science* 85, 2910–2915.

Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber neutral detergent fiber, and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3597.

Weiss WP 1993. Predicting energy values of feeds. In symposium: prevailing concepts in energy utilization by ruminants. *Journal of Dairy Science* 76, 1802–1811.