



Maintaining grass coverage increases methane uptake in Amazonian pastures, with a reduction of methanogenic archaea in the rhizosphere

Leandro Fonseca de Souza^{a,b,*}, Dasiel Obregon Alvarez^{a,c}, Luiz A. Domeignoz-Horta^{b,d}, Fabio Vitorino Gomes^a, Cassio de Souza Almeida^a, Luis Fernando Merloti^a, Lucas William Mendes^a, Fernando Dini Andreote^e, Brendan J.M. Bohannan^f, Jorge L. Mazza Rodrigues^g, Klaus Nüsslein^b, Siu Mui Tsai^a

^a Center for Nuclear Energy in Agriculture, University of Sao Paulo, Piracicaba, SP, Brazil

^b Department of Microbiology, University of Massachusetts, Amherst, MA, USA

^c Applied Soil Ecology Lab, School of Environmental Sciences, University of Guelph, Guelph, ON, Canada

^d Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland

^e Luiz de Queiroz College of Agriculture, University of São Paulo, Piracicaba, SP, Brazil


^f Institute of Ecology and Evolution, University of Oregon, Eugene, OR, USA

^g Department of Land, Air and Water Resources, University of California Davis, CA, USA

HIGHLIGHTS

- Grass coverage increases methane uptake in pasture soils compared to bare soils.
- Methanogens were reduced by 10 fold in the grass rhizosphere compared to bulk soil.
- Soil liming can compromise the capacity of forest and pasture soils to sink methane.
- Pasture management strategies have potential to mitigate soil methane emissions.

GRAPHICAL ABSTRACT



ARTICLE INFO

Editor: Jay Gan

Keywords:

Climate change
Soil methane cycling
Methanogenic archaea
Methanotrophy
Environmental microbiology

ABSTRACT

Cattle ranching is the largest driver of deforestation in the Brazilian Amazon. The rainforest-to-pasture conversion affects the methane cycle in upland soils, changing it from sink to source of atmospheric methane. However, it remains unknown if management practices could reduce the impact of land-use on methane cycling. In this work, we evaluated how pasture management can regulate the soil methane cycle either by maintaining continuous grass coverage on pasture soils, or by liming the soil to amend acidity. Methane fluxes from forest and pasture soils were evaluated in moisture-controlled greenhouse experiments with and without grass cover (*Urochloa brizantha* cv. Marandu) or liming. We also assessed changes in the soil microbial community structure of both bare (bulk) and rhizospheric pasture soils through high throughput sequencing of the 16S rRNA gene, and quantified the methane cycling microbiota by their respective marker genes related to methane generation (*mcrA*) or oxidation (*pmoA*). The experiments used soils from eastern and western Amazonia, and concurrent field studies allowed us to confirm greenhouse data. The presence of a grass cover not only increased methane uptake by up to 35% in pasture soils, but also reduced the abundance of the methane-producing community. In the grass rhizosphere this reduction was up to 10-fold. Methane-producing archaea belonged to the genera *Methanosarcina* sp., *Methanocella* sp., *Methanobacterium* sp., and Rice Cluster I. Further, we showed that soil liming to increasing pH compromised the capacity of forest and pasture soils to be a sink for methane,

* Corresponding author at: Center for Nuclear Energy in Agriculture, University of Sao Paulo, Piracicaba, SP, Brazil.

E-mail address: leandro_fonseca@alumni.usp.br (L. Fonseca de Souza).

and instead converted formerly methane-consuming forest soils to become methane sources in only 40–80 days. Liming reduced the relative abundance of Beijerinckiaceae family in forest soils, which account for many known methanotrophs. Our results demonstrate that pasture management that maintains grass coverage can mitigate soil methane emissions, compared to bare (bulk) pasture soil.

1. Introduction

The establishment of pasture lands has been the main cause of deforestation in the Amazon region since the 1970s (Dias et al., 2016; Margulis, 2003). This transformation of rainforest into pastures has led to a net increased emission of the powerful greenhouse gas methane, turning a methane consuming forest soil into a methane producing pasture soil (Meyer et al., 2020; Fernandes et al., 2002; Steudler et al., 1996; Goreau and De Mello, 1988). The impact of land-use conversion on the annual balance of gas fluxes is noticeable considering pastures in western Amazonia can emit up to 270 mg C-CH₄/m², while nearby forest soils can consume up to 470 mg C-CH₄/m² (Steudler et al., 1996).

Methane (CH₄) gas has an 86-fold greater potential to retain heat in the atmosphere compared to that of CO₂, calculated over a 20-year period (IEA, 2021). The global methane emissions are mainly driven by human activities such as livestock production, irrigated agriculture, oil and gas production, and landfill decomposition (IEA, 2021). Soil methane cycling is strongly dependent on the microbiota, since the biogenic source of this gas is methanogenic archaea. The biological consumption of methane is controlled by methanotrophic organisms, primarily bacteria. In soil, the balance between methanotrophic bacteria and methanogenic archaea is related to environmental conditions (i.e., moisture, temperature, soil density, and pH) and is sensitive to changes in agricultural management (Le Mer and Roger, 2001; Liu et al., 2007; Tian et al., 2015).

Methanotrophic bacteria in soil belong primarily to the *Gammaproteobacteria*, *Alphaproteobacteria*, *Verrucomicrobia*, and the phylum NC10 (Hanson and Hanson, 1996; Knief, 2015; Ettwig et al., 2009). The initial step of methane oxidation occurs through its conversion to methanol, which is mediated by the enzyme methane monooxygenase (MMO). A sub-unit of a common variant of this enzyme is coded for by the *pmoA* gene, which can be used as a methanotroph-specific marker for molecular studies.

Methanogenic archaea traditionally comprise members from eight orders within the phylum Euryarchaeota: *Methanopyrales*, *Methanobacteriales*, *Methanococcales*, *Methanomicrobiales*, *Methanocellales*, *Methanosarcinales*, *Methanomassiliicoccales*, and ‘*Candidatus* Methanophagales’ (Evans et al., 2019), with additional candidates in the phylum *Bathyarchaeota* (Kallistova et al., 2017). Methanogenesis is the final step of an anaerobic pathway that begins with the hydrolysis of organic polymers, fermentation of the resulting monomers and of initial fermentation products, and ends up in the production of CH₄ mostly from acetate, hydrogen, and CO₂. The final step in the methanogenesis pathway is facilitated through the action of the enzyme methyl-coenzyme M reductase, coded by the *mcrA* gene, which can be used as a methanogen-specific marker for molecular studies (Serrano-Silva et al., 2014).

The response of the soil microbial community to changes in land-use is not well understood (Nazaries et al., 2013; Tate, 2015), but previous studies have shown significant impacts on microbial diversity in the Amazon region (de Carvalho et al., 2016; Mendes et al., 2015; Navarrete et al., 2015; Rodrigues et al., 2013; Jesus et al., 2009). In well-managed pastures in the Amazon region, the grass root system can redistribute carbon to deeper layers, where it is less susceptible to decomposition (Fearnside and Imbrozio Barbosa, 1998). On the other hand, we can expect that degraded pastures with large bare soil areas can facilitate the release of carbon from the system, with superficial grassroots, higher loss of soil organic matter, and lower carbon stocks (Segnini et al., 2019). Proper management of pasture may involve several practices, such as soil acidity correction and continuous maintenance of grass cover to protect soil from erosion. These practices are particularly important in the Amazon region given the environmental extremes of this area. Altogether, the high soil acidity, high

rainfall, and high temperatures combined with exposure of the soil to equatorial solar radiation constitute factors that are associated with increased erosion and soil degradation (Demattê and Demattê, 1993).

The effect of soil liming on methane fluxes is still poorly understood, and studies in temperate forests show that liming can lead to both increases or decreases methane consumption (Wang et al., 2021; Borken and Brumme, 1997; Butterbach-Bahl et al., 2002). In wheat-focused agriculture liming has led to increased consumption of methane in soil (Hütsch et al., 1994). The increased methane consumption after liming was also observed in Mediterranean semiarid soils under lupine, wheat, and triticale, a hybrid of wheat and rye (Barton et al., 2013; García-Marco et al., 2016). However, for tropical soils little information is available regarding what influence liming has on methane production and consumption. In an assessment of greenhouse gas fluxes from soils under soybean cultivation in Brazil, the acidity correction presented no effect on methane fluxes (Lammel et al., 2018). Likewise, in a field experiment in Puerto Rico, soil consumption of atmospheric CH₄ in an intentionally acidified soil was about one-fourth of that at pH 6, and was not restored after liming (Mosier et al., 1998).

There is a growing consensus that the key to understanding major soil functions lies where plants and soil meet, in the rhizosphere (Lau and Lennon, 2011). The rhizosphere is a micro-environment with differentiated soil conditions and steep overlapping gradients, in which pH can be up to 2 units more acidic or more basic than the bulk soil. The rhizosphere can present heterogeneous concentrations of oxygen and moisture and can be enriched in root exudates (Philippot et al., 2013). These factors affect soil methane cycling not only by providing organic substrate for methanogenesis but also by promoting the oxidation of methane in the rhizosphere. Despite its known role in flooded rice soils (Frenzel et al., 1992), little is known about the impact of the rhizosphere on methane cycling in upland soils since these soils are commonly considered to be a sink of methane, not a source (Philippot et al., 2009). Thus, a better understanding of how the rhizosphere of land-intensive tropical pastures affects soil methane cycling can yield new strategies to mitigate greenhouse gas emissions related to cattle ranching.

The intensive land use in agriculture and cattle ranching in Amazonia can lead to soil and pasture degradation, and the presently somewhat degraded areas range from 50% to 70% of the total land (Dias-Filho, 2017). Within this context, this research aims to evaluate how the management of pastures can affect soil methane cycling. We hypothesized that CH₄ sinking is increased by liming soils, similar to that observed in soils from temperate regions, and by continuous grass coverage of *Urochloa brizantha* cv. Marandu, due to a reduction of methanogens in the rhizosphere. To test this hypothesis we combined 1) greenhouse experiments in which we controlled the environmental variables and 2) an *in-situ* study, in which we made measurements under natural conditions. First, we set up greenhouse experiments, where soil acidity was adjusted and grass was planted, the soil-air CH₄ fluxes were measured and shifts in the soil microbial community between bare soil and the rhizosphere of *Urochloa brizantha* cv. Marandu were determined. Second, we measured *in-situ* methane fluxes in soils from pasture field sites with and without grass cover, and compared the microbiota in bare pasture soil to that of the rhizosphere of grassy soil.

2. Materials and methods

2.1. Sampling

Experiments were performed with soils from both Western and Eastern Amazonia (Table S1). In the Western region (hereafter “Ariquemes”) sampling was carried out in April of 2017 at the Fazenda Nova Vida near

Ariquemes, RO (10°10'49.5" S, 62°49'23.9" W). While in the Eastern region (hereafter "Tapajós"), the samples were taken at the National Forest of Tapajós and the immediately surrounding pasture areas near Belterra, PA (3°07'53.8" S, 54°57'24.2" W), in August of 2019. The sampled soils were used in two rounds of greenhouse experiments at the Center for Nuclear Energy in Agriculture, Brazil (22°42'27.7" S, 47°38'41.0" W). In addition to the soil sampling a field study was performed, but only in the Tapajós region (detailed below).

Western Amazonia represents a region with a high degree of exploitation (de Moraes et al., 1996; Herpin et al., 2002; Reiners et al., 1994). The Fazenda Nova Vida region has fragments of primary forest and pastures of different ages. The sampled pasture area was established in 1972, and since then managed by cattle rotation, with the use of fire only to control eventual pests, mechanical removal of invasive trees, and at least one record of liming 15 years before the sampling. Soils sampled varied from average clay to sandy texture. The soil microbial community of the area were previously studied by Rodrigues et al. (2013) and Meyer et al. (2020).

Eastern Amazonia represents areas of more recent exploration. The Tapajós National Forest was sampled as a model of a conservation area and the pasture chosen is on a small property in Belterra, PA. The pasture used here was established between 1989 and 1994 and supported cattle at the time of sampling, had sparse signs of degradation, fire was applied when necessary to control invasive plants, and it has no history of liming. The soils sampled ranged from average clay to sandy texture.

During each expedition, we sampled 20–30 kg of soil from the upper 0–10 cm layer, gathered from 5 equidistant sampling points along 100–200 m at each site, from areas under primary forest and pastures cultivated with *Urochloa brizantha* cv. Marandu. The soils sampled for experiments were conserved in plastic bags partially closed with a paper filter in its opening, to ensure air exchange but little water loss. The plastic bags were transported to the Cell and Molecular Biology Laboratory at the University of Sao Paulo, CENA-USP, where aliquots were used for chemical analysis, and then were conserved under environmental temperature and shadow until the experiment start. In addition, soil sample aliquots were kept on ice at the end of each sampling day in the field and stored at –20 °C for future molecular analyses.

2.2. Greenhouse experiments

Sampled soil was homogenized, sieved (5 mm), and placed in clay pots with a volume of 10 L each, resulting in 10 cm high soil columns with 5 kg of soil per pot. The grass was raised from seeds of *Urochloa brizantha* cv. Marandú (BRSEEDS, Aracatuba, Sao Paulo, Brazil) in a subsample of the soil, and mature plants were transferred to the experimental clay pots at least 40 days after soil liming. The liming was performed by the addition of CaCO₃ to reach pH 6.5 (water), calculated for a base saturation of 70–75%. For each treatment, four pots were used to grow *Urochloa brizantha* cv. Marandu with 4 additional pots used as no-plant controls (bare soil), both at natural pH and with limed soils (4 pots × 2 soil types × 2 pH situations × 2 plant situations). At the beginning of the experiment, soil moisture was standardized to ~70% of the water retention capacity of the soil and adjusted every 2 to 4 days, taking as reference the weight variation after drying soil samples for 48 h at 75 °C. In the experiment with soil from Ariquemes, the plants were removed when they reached approximately 35 cm in height, and by shaking and with the help of a sterilized brush the rhizosphere soil was collected. Here we defined the rhizosphere as soil that remained attached to the roots even after vigorous plant shaking.

2.3. Field in-situ measurements

At two pastures in the Tapajós region, 100 m side squares were established and 4 points in the square corners, plus a point in the center, were selected to evaluate CH₄ fluxes and to sample soils for molecular and chemical analysis (Table S1). Those 5 points had grass coverage at the time, and before gas flux measurements with static gas collection chambers

the grass leaves were cut to their stems (2 cm above soil surface) and removed. Following chamber removal, the roots were collected, and the rhizospheric soil (defined below) was sampled and stored at –20 °C by the end of the day. Adjacent to each of the five selected sampling points, 1 m² plots without grass (bare soil) were used to measure methane fluxes and to collect soil samples for molecular and chemical analysis.

2.4. Determination of methane fluxes in soil

Gas fluxes in the experiment were measured after day 130 of the beginning of the experiment, the moment of the grass seeding, and lasted until day 238. The intervals among measurements were 130 to 137 (7 days), 148 to 158 (10 days), 158 to 214 (56 days), 214 to 226 (12 days) and 226 to 238 (12 days).

The measurements of CH₄ fluxes in both the field sampling sites and the greenhouse experiments were carried out using static gas collection chambers (20 cm in diameter and height, ~6 L inner volume). The chamber consisted of a metal circular ring, with a top lateral groove that was filled with water to fit the lid without gas leaking. Over a period of 10 min, measurements were taken at 10 s intervals using a portable gas analyzer (UGGA model 915–0011, Los Gatos Research, San Jose, CA, USA) and the first 12 points were excluded to reduce fluctuations in the headspace fluxes, defined after testing (data not shown). No fans were used for gas homogenizing, due to the relatively small volume of the chamber and the automatic pump of the portable gas analyzer that circulates the gas, mixing it continually. The laser calibration was verified by checking the graphs of optical absorption by laser frequency, according to the manufacturer recommendation. The methane values used were those already discounted of water vapor effect on optical absorption. This adjustment was performed by the software supplied with the equipment UGGA (model 915–0011, Los Gatos Research, San Jose, CA, USA). The daily flux of gases was estimated from the concentration in the chamber headspace. Daily flux (F, mass of gas m⁻².day⁻¹) was computed using the following equation (Ussiri et al., 2009):

$$F = \Delta g_{\text{as}} / \Delta t \times V / A \times k$$

Where $\Delta g_{\text{as}} / \Delta t$ is the rate of change in CH₄ concentration inside the chamber (i.e., mg CH₄-C); V is the chamber volume (m³); A is the surface area of the soil circumscribed by the chamber (m²) and k is the time conversion factor (1440 min day⁻¹). The cumulative gas emissions were calculated by linear interpolation of average emissions between two successive measurements and the sum of the results obtained over the entire study period. Finally, the data was expressed as differences in the cumulative CH₄ fluxes in relationship to the controls. To calculate the differences in the cumulative fluxes, we subtracted from the accumulated fluxes in the respective treatments (liming, grass coverage, and liming plus grass coverage) the control measurements of the average accumulated flux (bare soils).

2.5. Characterization of soil chemical properties

Approximately 600 g of soil were analyzed for their physical-chemical properties at the Laboratory of Chemical Analysis in the Soil Science Department at the Luiz de Queiroz College of Agriculture (ESALQ / USP) (van Raij et al., 2001). The soil attributes measured were: pH in CaCl₂; concentrations of phosphorus, potassium, calcium, and magnesium by extraction with ion exchange resin; aluminum by extraction of potassium chloride at 1 mol/L; potential acidity estimated by pH-SMP buffer test; organic matter by the dichromate-titrimetric method; boron by extraction with hot water; copper, iron, manganese and zinc extracted by the DTPA-TEA extractor (pH 7.3); and by calculating the sum of bases (BS); cation exchange capacity (CEC); base saturation (V%), and aluminum saturation (m%).

2.6. DNA extraction

DNA was extracted for molecular analyses from soils used in greenhouse experiments, previously originated from Ariquemes region and from soils of the field study in the Tapajós region. Total DNA was extracted from soil samples using the PowerLyzer PowerSoil DNA Isolation Kit (Qiagen, Hilden, Germany) from 250 mg of soil, according to the protocol provided by the manufacturer, except that after adding solution C1 the stirring time was extended to 15 min followed by 3 min centrifugation (Venturini et al., 2020). The amount and quality of the DNA extracted were analyzed in a Nanodrop 2000c spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA) at an optical density of 260 nm. The total DNA extracted was stored at -20°C .

2.7. Abundance of methane producers and oxidizers

The abundance of methanotrophs and methanogens in the soil was estimated by quantitative PCR. To this, DNA was extracted from soil samples at 40, 130 and 250 days of the experiment. Forty days marked the time after liming stabilization in the soil, 130 days the grass seeding and 250 days the end of the experiment.

Quantitative PCR (qPCR) was used to quantify the genes associated with methane cycling *mcrA* and *pmoA* (Table S2) in total soil DNA samples. For each gene, a standard curve was established spanning each order of magnitude from 10^1 to 10^7 copies of the gene. Target genes were previously obtained by PCR from genomic DNA of *Methanolinea mesofila* (DSMZ 23604) for the *mcrA* gene, and *Methylosinus sporium* (DSMZ 17706) for the gene *pmoA*, both obtained from the DSMZ (German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany). The qPCR was performed in triplicate for each sample on a StepOne Plus cycler (Thermo Fisher Scientific, Waltham, MA, USA), with a final volume of 10 μL , containing 5 μL of SYBR Green ROX qPCR (Thermo Fisher Scientific, MA, USA), 1 μL of each primer (5 pmol), 1 μL of soil DNA (adjusted to 10 ng/ μL), 0.8 μL of bovine serum albumin (20 mg / mL) (Sigma-Aldrich, San Luis, MO, USA), and 1.2 μL of sterile ultrapure water.

In order to minimize bias in the analysis between each qPCR plate run, gene abundance was quantified with the software LinRegPCR (Ramakers et al., 2003). Raw amplification data for each sample were used to calculate individual reaction efficiencies, and detection limits were established for each group of technical replicates. The data generated in arbitrary fluorescence units were converted to the number of copies of the genes using linear interpolation between the known quantities in the standard curve (5 best points out of 7) and the observed fluorescence measurements, using the curves of each plate as a reference for the respective samples.

2.8. Sequencing of 16S rRNA gene fragments

The composition of the microbial community was determined with high throughput sequencing (MiSeq Illumina platform with a 600c kit) of the V4 region of the 16S rRNA gene at the Functional Genomics Center of Luiz de Queiroz College of Agriculture. The V4 region was amplified with the primers 515F (Parada et al., 2016) and 806R (Apprill et al., 2015). This sequencing strategy was selected to match the highly diverse soil environment and for the size of the paired-end reads (average 300 bp). The DNA concentrations in the samples were adjusted to 10 ng/ μL using a Nanodrop 2000c spectrophotometer and the PCR reactions with 2.5 μL of $10\times$ buffer, 1 μL of 50 mM MgCl₂, 1 μL of 10 mM dNTPs, 0.5 μL of 10 μM forward and reverse primers, 0.5 μL of 5 U/ μL Taq Platinum – PCR (Thermo Fisher Scientific, Waltham, MA, USA) and water for PCR - 14 μL , in a total volume of 25 μL per reaction. Gene library preparation followed the conditions of 95°C for 3 min, followed by 25 cycles at 95°C for 30 s, 50°C for 30 s, 72°C for 30 s, and a final extension step at 72°C for 5 min. Subsequent DNA purification of the amplicon was performed using AMPure XP beads (Beckman Coulter, Brea, CA, USA) and verified on an agarose gel. Similarly, the adapters were added by synthesis, followed by another purification with AMPure XP beads and confirmed on an agarose gel electrophoresis.

The amplicon pool was normalized using quantification by qPCR with the KAPA Illumina quantification kit (Roche, Basel, Switzerland). The computational processing of these data was performed using QIIME2 2017.11 (Bolyen et al., 2019), with data quality control using the DADA2 tool (Callahan et al., 2017) and taxonomic identification of the sequences was performed using q2-feature-classifier (Bokulich et al., 2018) and the SILVA v.128 99% database (Quast et al., 2012).

2.9. Phylogenetic analyses

Because a few amplification sequence variants (ASVs) were grouped closest with the family Beijerinckiaceae, a family containing both methanotrophic and non-methanotrophic genera, we increased our phylogenetic resolution by analyzing phylogenetic trees containing only Beijerinckiaceae sequences that were created with two 16S rRNA primer pairs. One set of primers targeted the region V4, between 515F (Parada et al., 2016) and 806R (Apprill et al., 2015), and the second set of primers targeted the V3/V4 region, between 341F and 805R (Herlemann et al., 2011). For this last primers pair, the amplification protocol was identical to that described above, except the annealing temperature for the second primer pair was 55°C . The sequences were aligned, and trees were calculated using the software CLC Genomics Workbench 20.0 (QIAGEN, Aarhus, Denmark) at default parameters, and with a maximum likelihood model (PHYML function) with UPGMA (Unweighted Pair Group Method with Arithmetic mean) assuming common replacement frequencies to the bases (Kimura, 1980). The robustness of the final trees was tested with 1000 bootstrap replications. Reference sequences for members of the Beijerinckiaceae family were obtained from the curated database Ribosomal Database Project II version 11 (Cole et al., 2014) based on the criteria of high-quality reads with a length greater than 1200 bp and representing type strains. The only sequence available for the 16S rRNA gene of the methanotrophic bacterium USC α (Pratscher et al., 2018) was also added.

2.10. Statistical analysis

All comparative analyses between groups were performed with ANOVA followed by a Tukey Honestly Significant Difference (HSD) test and *p*-values calculated for a two-tailed distribution of the data using the package *agricolae* version 1.2–8 (R Core Team, 2013).

Significant explanatory variables of the methane fluxes were chosen by linear regression and model selection (backward) and by minimizing the Akaike Information Criterion (AIC). The statistical significance was assessed by 1000 permutations of the reduced model. The resulting significant explanatory variables were used to access their contribution to explaining the CH₄ fluxes, using the function *varpart* (Peres-Neto et al., 2006) in the vegan package (Oksanen et al., 2015). Statistical analyses were performed in software R (R Core Team, 2013).

The DEICODE tool (Martino et al., 2019) in the QIIME2 2019.10 was used to process the sequencing data. This tool can identify significant changes in the community based on relative abundance data. Next, the software QURRO (Fedarko et al., 2020) was used to assess shifts in the methane cycling community based on transformed abundance data (natural logarithm) and using a minimum of 10 occurrences per taxon.

3. Results

This study tested the effect of acidity correction by liming and the presence of a grass cover by *Urochloa brizantha* cv. Marandu on soil methane fluxes with soils from pasture and forest of different Amazon regions. The first experiment was performed with soils from western Amazon region (Ariquemes, RO) and the second with soils from eastern region (Tapajós, PA). In the Ariquemes experiment, liming resulted in a final pH of ~ 6.0 (CaCl₂, equivalent to pH 6.5 in H₂O), and an increase in calcium availability, as well a decrease in aluminum saturation (Table S3). Also, methane was consumed in both bare soils of forest and pasture at their respective natural pH values, with greater uptake in forest soils (Fig. 1). In the Tapajós

experiment, we observed methane emissions from bare soils from the pasture at natural pH, and methane uptake in bare soils from forest at natural pH (Fig. S1). When forest soils from Ariquemes had grass cover they exhibited the highest methane consumption (Fig. 1-a; $p = 0.059$), at values close to the naturally acidic forest soil, but significantly lower than both limed soils with or without grass cover (Fig. 1-a). The Tapajós soils showed a similar trend compared to Ariquemes soils (Fig. S1). Methane uptake in pasture soils increase by 35% on average when they have grass coverage (Fig. 1-b; $p = 0.001$). However, liming of pasture soils reduced their methane uptake (Fig. 1-b; $p = 0.001$) and turned forest soils from a methane sink into a methane source (Fig. S1-b; $p = 0.052$).

We compared our greenhouse observations to field methane flux measurements. *In situ* measurements of CH_4 fluxes were taken on two pastures in Belterra/PA, Tapajós region, during the end of wet season, at points with and without grass coverage. No significant differences were observed, but the trend is similar to that observed in the greenhouse experiments (Fig. S2; $p = 0.112$).

Molecular analyses were performed only with soils from the Ariquemes greenhouse experiment (Fig. 2), and from the field study in the Tapajós region (Fig. S3), due to logistic reasons. The rhizosphere community was evaluated only at the end of the experiment ($T_3 = 250$ days). During most of the experimental timeline we did not observe differences in the abundance of methanotrophs between pasture and forest soils (Fig. 2). Regarding methane producers, we observed a very low abundance in forest soils compared to pasture soils during the experimental duration (Fig. 2). The acidity correction shows a tendency to reduce methanotroph levels in forest soils after 250 days in the grass rhizosphere ($p = 0.339$) and in the bare soil ($p = 0.162$) (Fig. 2A). Pasture soils had between 100 and 1000-fold more methanogenic archaea than forest soils throughout the experiment, which did not change with acidity correction (Fig. 2B). The abundance of methanogenic archaea in the grass rhizosphere in pasture soils was reduced on average by 13 times compared to the bare soil (Fig. 2B; $p = 0.025$). No significant changes in methanotrophs abundance were recorded in the rhizosphere (Fig. 2A; $p = 0.263$). This reduction in methanogenic archaea in the grass rhizosphere was not observed in the field study (Fig. S3; $p = 0.186$).

To investigate the effects of acidity correction after liming and grass rhizosphere on specific groups of microorganisms, high throughput DNA sequencing of the 16S rRNA gene was performed. The results show fair sequencing depth, with rarefaction curves leveling off well below the minimal sequencing depth in soils from the Ariquemes experiment (Table S4, Fig. S4-a) and in soils from the field studies (Table S5, Fig. S4-b).

Considering only the community associated with methane cycling in the soil (identified with a minimum of 90% confidence), we filtered the groups recognized as methanotrophs (Knief, 2015) and methanogens (Evans et al., 2019) and observed results similar to those obtained in the quantification of gene copies in total DNA. The relative abundance of methanogens in forest soils was smaller than in pasture soils, while the relative abundance of methanotrophs was similar between forest and pasture soils (Fig. 3). The increase of methanogens in forest soils with acidity correction was not significant. A significant drop in methanotroph abundance was observed only for the combination of acidity correction and grass cover treatments (Fig. 3, $p = 0.024$). Methanotrophs in pasture soils did not change with liming or with the presence of grass cover, as previously observed in the quantification of *pmoA* gene copies (Fig. 2). However, the number of methanogens was significantly reduced in the grass rhizosphere, with ($p = 0.017$) or without ($p = 0.007$) acidity correction (Fig. 3). This last result was similar to that observed in the field, which, although not significant, points to a tendency to reduce methanogenic archaea of different groups in the grass rhizosphere (Fig. S5).

Finally, to understand which groups are associated with methane cycling in these soils, a detailed analysis was performed of all groups that presented sequences of the genera known to act in methane cycling (Knief, 2015; Evans et al., 2019) (Figs. 4 and 5). In pasture soils, the abundance of members of all methanogenic genera was lower when soil was grass-covered. Forest soils showed a low abundance of methanogenic archaea

belonging to *Methanosarcina* spp. compared to pasture soils. The Beijerinckiaceae family is abundant in these soils but it was not possible to identify the sequences at the genus level with the database SILVA v.128 (Quast et al., 2012). A new phylogenetic identification was performed comparing all the sequences annotated as Beijerinckiaceae in the RDP database, with the five amplification sequence variants (ASVs) from sequencing with primers 341F/805R and 8 ASVs from primers 515F/806R. In this analysis, only sequences of the 16S rRNA gene of this family were used as reference, and the results indicate that they are closely related to the methanotrophic clade, with more than 90% confidence (Figs. S6 and S7). Those Beijerinckiaceae are reduced in their relative abundance in forest soils with acidity correction (Fig. 5), without changes in pasture soil. No significant changes were observed compared to other methanotrophs.

To disentangle the effect of soil properties, microbial communities, and the presence of grass on the CH_4 fluxes we performed a variation partitioning analysis. This analysis showed that a large part of the explained variation of the methane fluxes (40%) was due to pH and other soil properties in the first days of the experiment (Fig. 6b). However, the grass biomass represented most of the explained variance at the later time-points (Fig. 6e-f). The effect of methanogens and methanotrophs abundance was minor at the beginning of the incubation (6%) but reached 13% of the explained variance at the last time point. Here we decided to separate pH from other soil properties when running this analysis as pH has been previously shown to be a major driver in CH_4 soil fluxes. Our results confirm this since at the beginning of the experiment the pH and other soil physical-chemical properties were the stronger explanatory variables of the CH_4 uptake capacity (Table 1). However, the contribution of the soil properties decreases through time while the presence of the grass and the microbial communities gain in explanatory power.

4. Discussion

4.1. Land-use change shifts soil methane fluxes

Deforestation of the Amazonian soils is often followed by the establishment of pastures. This forest-to-pasture conversion affects soil methane cycling, where forest soils that were previously acting as a methane sink now become sources of methane (Fernandes et al., 2002). The results obtained in the present study corroborate previous field studies (Meyer et al., 2020; Fernandes et al., 2002; Steudler et al., 1996; Goreau and De Mello, 1988), although the values obtained in our experiments cannot be directly compared to those previously reported, since the results are limited to a 10 cm surface layer of soil. The trend observed is the same recorded by Steudler et al. (1996), in that forests consume 2.74 more methane than pastures. In our experiments, these values were 0.6-fold in soils from Ariquemes in Western Amazonia, and 4.28-fold in soils from Tapajós in Eastern Amazonia. This discrepancy may be related to differences in soil microbial communities and chemical properties, but it can also be a consequence of conservation of the forest areas from which these soils originated. While in Ariquemes forests were fragmented and small samples from Tapajós originated from a contiguous forest in a conservation area. Forest fragmentation is known to be associated with increased greenhouse gas emissions (Laurance et al., 1998). Furthermore, the pastures sampled in Ariquemes have a history of long-term management, and the pasture soils sampled in Tapajós showed signs of degradation. As management affects the carbon stock in the soil (Fearnside and Imbrozio Barbosa, 1998), it can be expected that it will also affect methane cycling in the soil.

4.2. Land-use change impact of methanotrophs and methanogens

The forest-to-pasture conversion alters the physical-chemical properties of the soil, impacting the microorganisms that produce and consume methane. In field studies conducted in the same region in Ariquemes, a decrease in methanotrophic bacteria and an increase in methanogenic archaea were observed, in addition to changes in the composition of communities, which were attributed at least in part to changes in methane fluxes (Meyer et al.,




Fig. 1. Cumulative CH_4 fluxes in the experiment with a) forest and b) pasture soils of Ariquemes, with and without acidity correction and with and without grass coverage. Central lines indicate the mean and shaded areas the standard deviation. Sampling times in: 137, 148, 158, 214, 226 and 238 days after the beginning of the incubation.

2017; Meyer et al., 2020). Also, an increase in the activity of methanogens in pastures, compared to forest soils, was recorded in soils from the same region (Kroeger et al., 2021). Our results under controlled moisture conditions did not detect significant changes in the methanotrophic community nor changes in the relative abundance of specific methanotrophic groups. However, there was a significant increase in the abundance of methanogenic archaea in pastures compared to forest soils.

Pasture soils in the Amazonian region present a microbial community quite distinct from that observed in forest soils (de Carvalho et al., 2016; Rodrigues et al., 2013; Jesus et al., 2009). This is partly attributed to acidity reduction in the process of establishing pastures. Forest soils in Amazonia have pH values between 3.5 and 4.5 (Demattê and Demattê, 1993). pH is currently understood as one of the main drivers of microbial community structure in soils (Fierer and Jackson, 2006), and increasing it by liming is a strategy to improve fertility and reduce soil toxicity to plants

(Oliveira et al., 2003). This process becomes necessary in pasture management to counter the tendency for acidification of pasture soils over time with soil pH reaching values close to those observed in forest areas (de Moraes et al., 1996). In addition, degraded pastures, which can amount to more than 50% of pasture areas in Amazonia (Dias-Filho, 2017), also tend to need acidity correction for their restoration.

4.3. pH influence on methanotrophs and methanogens

Little is known about the effects of soil liming on the methane cycling process in tropical soils. It is known that the optimum growth pH of most cultivable methanotrophs and methanogens is neutral (Le Mer and Roger, 2001; Whittenbury et al., 1970), which is why soil pH represents an important explanatory variable for the distribution of methanotrophs. However, methane oxidation is observed in natural




Fig. 2. qPCR quantification of the genes (A) *pmoA* and (B) *mcrA* at different times after the beginning of the Ariquemes experiment. T1 = 40 days, T2 = 130 days and T3 = 250 days. Letters above each box plot indicate significant changes ($p < 0.05$). CN = copy number. Ns = Not significant.




Fig. 3. Changes in the logarithmic ratio between a) methanogenic and b) methanotrophic groups in relation to the whole community (16S rRNA gene) in the Ariqueles experiment. The dotted line indicates a calculated ratio of the minimum of 10 counts for each group. The more negative the number of the natural log, the lower the abundance in relation to the total community. Letters indicate significant changes within each treatment for the same land use and group of microorganisms (Tukey HSD; $p < 0.05$). ns = not significant.

environments across a wide pH range (Knief et al., 2003; Kolb, 2009; Nazaries et al., 2013). Our results indicate that the soil acidity correction for pH values close to 6.5 has different effects in pasture or forest soils, possibly because the forest soil has undergone a more intense pH correction, starting at 3.5–5.0 and ending at 6.5, while in pasture soils the change was from 4.5–5.5 to 6.5. In our forest soils from Ariqueles, we determined a decrease in methane uptake in response to liming, and

a shift from uptake to emission in forest soils from Tapajos. Yet, no significant differences were noticed in pasture soils. Thus, liming pasture soils may not impact methane emissions, but it helps to maintain the pH of these soils at values suitable for grass biomass productivity. For forest soils, we have shown that the reduction in acidity alone is enough to shift the soil from a methane sink to a source. This change in methane fluxes was not noticeable in the abundance of methanotrophic or




Fig. 4. Changes in the Log_e ratio between methanogenic microorganisms by genus (G) in relation to the total community in the Ariqueles experiment. All the identified genera are shown. The more negative the numbers, the lower the abundance. Letters indicate significant differences (Tukey HSD; $p < 0.05$).




Fig. 5. Changes in the Log_e ratio between methanotrophic microorganisms by genus (G) or family (Fa) in relation to the total community in the Ariqueemes experiment. The more negative the numbers, the lower the abundance. All the detected groups are presented. Letters indicate significant differences (Tukey HSD; $p < 0.05$).

methanogenic microorganisms by qPCR, despite a reduction in the relative abundance of methanotrophs that follows the acidity correction.

4.4. Diversity of microorganisms related to methane cycling

The identification of microorganisms based on short DNA sequences of the 16S rRNA gene, such as those generated in this study, is limited to the evolutionary information available in that fragment so it is not always feasible to identify the microorganisms at the genus level. Considering that the ability to oxidize methane is variable at the genus level in the Beijerinckiaceae family, identifying sequences at the family level is not enough to infer if they are methanotrophs. This family also includes generalist bacteria capable of using multiple carbon compounds as an energy source, and here Beijerinckiaceae are more abundant in forest soils than in pastures. Thus, identifying whether they are methanotrophs is relevant to understanding methane cycling in the forest-to-pasture conversion. The results demonstrate that the Beijerinckiaceae sequences observed in forest soils cluster together in phylogenetic trees. This cluster was observed on the two data sets with high support ($> 90\%$ in 1000 bootstraps) and also includes the methanotrophic USC α , which indicates that these sequences are potential methanotrophic Beijerinckiaceae.

The differences observed in methane fluxes after liming were not noticed in the abundance of producers and consumers. This discrepancy might be related to a reduction in the activity of forest soil to act as a methane sink after acidity correction, due to the lower availability of Fe and Cu which are necessary as cofactors for the activity of methane-monooxygenase (Semrau et al., 1995). Alternatively, this discrepancy might be due to limitations of the primers, drawn mostly with microbial references from temperate soils, but we applied them to tropical soils. Also, the difference between methane flux and shift in abundance of methane cycling microorganisms can be due to ammonia oxidizers, possibly oxidizing methane at a higher soil pH. The methanotrophs were potentially affected since acidity correction was followed by a reduction in the consumption of atmospheric methane by the soil (concentrations of ~ 1.8 ppm). Also, the duration of incubating soil from Ariqueemes for 250 days should be long enough to observe compensatory changes due to DNA replication, that should be detected in the DNA quantification analysis.

4.5. CH_4 sink capacity as a function of soil properties, grass cover and microbial communities

There is variability in the methane flux data, with pastures acting as methane sink and source, which is in fact a commonly reported final result. This observed variability means that pastures could seasonally or by location switch from being a methane source to temporarily becoming a methane sink (Fernandes et al., 2002; Steudler et al., 1996). Our initial hypothesis was that the methane sinking capacity of pasture soils would be related to intermediate moisture availability in the micro-environments of soil, since soil moisture is a determining factor for methane fluxes in pastures (Verchot et al., 2000). To eliminate moisture variation as a variable in the experiments we set the soil water contents at 70% of the holding capacity in the greenhouse experiments. The variability of pasture gas fluxes could also be explained by grass coverage, a factor associated with pasture management. The management of pastures can influence soil gas fluxes (Figueiredo et al., 2017), since it influences the carbon stocks in the soil (Fearnside and Imbrozio Barbosa, 1998), however the way ongoing pasture management can affect the microbial community remains an open question. Considering that management is performed with the goal of grass productivity, and greater aerial biomass is associated with greater root biomass, we expect that a larger root surface area in pasture would create a more interactive environment with the soil microbiota, and thus enable higher rhizosphere activity. The role of the rhizosphere on methane cycling in upland soils is still poorly understood, and even different plant species can influence the soil by increasing methane oxidation or production, depending on the type of soil or soil conditions (Praeg et al., 2017). In soils of the Ariqueemes experiment, we observed that plant cover will lead to a reduced methane flux in both forest and pasture soils compared to those with acidity correction. The methane flux rates with grass cover were similar to those of the original forest soil and tended to be higher than those of pasture without acidity correction. In soils from Tapajós experiment, the same trend was observed in forest soils, but likely due to the shorter duration of this experiment, there were no significant differences in the pasture.

When disentangling the contribution of different biotic and abiotic factors to CH_4 soil uptake capacity we found that its drivers change through




Fig. 6. Variation partitioning analysis to determine the drivers of CH₄ fluxes in the Ariquemes soils in time intervals from day 0 to day 7 (b), day 18 to 28 (c), 28–84 (d), 84–96 (e), and 96–108 (f). Variance was partitioned into four explanatory variables, soil physical-chemical properties (S), pH, abundance of methanotrophs and methanogens (G), plant biomass (P), and combinations of these potential predictors as exemplified in the outline (a). Geometric areas are proportional to the respective percentages of explained variation. The corners of the square depict the variation explained by each factor alone, while percentages of variation explained by interactions of two or all factors are indicated on the sides and in the middle of the square, respectively. All numbers represent percentages, graphically represented by the size of the respective hexagons. Only variance fractions $\geq 2\%$ are shown. The variables used for each variation partitioning are indicated in Table 1.

time, which could explain as previously discussed that soils might change from a source to a sink. While pH and other soil properties explained most of the variance at the beginning of the greenhouse experiment, the

Table 1
Selected explaining variables for the capacity of methane consumption determined with the variation partitioning analyses at five different time-intervals during the Ariquemes experiment.

Time interval	Soil properties ^a & pH	Gene abundances	Plant ^b
0–7	pH, OM, Cu, CEC	<i>mcrA</i>	NA
18–28	pH, Cu	<i>pmoA</i>	NA
28–84	pH, CEC	–	NA
84–96	–	–	Plant biomass
96–108	–	<i>mcrA</i>	Plant biomass

^a CEC corresponds to Cation Exchange Capacity, Cu to Copper, and OM to Organic Matter.

^b Plant biomass corresponds to the sum of the above and belowground dry weight.

abundance of microbial communities related to CH₄ fluxes and plant biomass explained most of the CH₄ uptake at the end of the experiment. These results suggest that our treatments (liming and planting grass) are changing the microbial communities and while the soil properties are initially the main variables explaining the CH₄ fluxes, after a couple of weeks the biotic factors are the main drivers of CH₄ fluxes in these soils. While a previous study showed that peak emissions of the greenhouse gas N₂O can be driven by the microorganisms related to the production and reduction of this greenhouse gas (Domeignoz-Horta et al., 2018), our results show how microorganisms related to methane cycling and plant cover play a role to understand the temporal dynamics of CH₄ uptake in soils. These results highlight the need for better characterizing microbial communities to increase our understanding of the relationship between abundance and diversity of microorganisms and their corresponding processes.

The results presented here demonstrate that soil acidity is an important factor for methane flux in tropical forest soils, since reducing acidity reduces the soil's capacity for methane consumption. In pastures, the effect of the acidity correction is less consequential compared to the presence of

grass coverage. This demonstrates that the correction of acidity in pastures, if combined with constant soil coverage with grass, would have little or no impact on methane emissions while improving soil structure and increasing nutrient availability, soil organic matter and grass productivity.

5. Conclusion

Our results suggest that the impact of methane emissions from pastures in the Amazonian region can be partially mitigated through pasture management, specifically by preventing soil exposure. The rhizosphere of *U. brizantha* cv. Marandu affects soil microbial communities by lowering the abundance of methanogenic archaea up to 10 times compared to the bare soil. The affected methanogens are composed of *Methanobacterium* spp., *Methanocella* spp., *Rice Cluster I*, and *Methanosarcina* spp. In addition, we demonstrate that the correction of acidity in pasture soils can reduce methane sequestration under atmospheric methane concentrations (high-affinity methanotrophs). Therefore, the level of acidity correction should be considered as a factor for additional emissions of greenhouse gases. In the acidic forest soils, an increase in pH reduced methane sequestration by more than 50%, thereby reversing the flux direction to turn forest soil from a methane sink into a source. Field studies with liming and a focus on the grass rhizosphere under seasonal conditions are needed to provide specific recommendations to policymakers and farmers.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2022.156225>.

CRediT authorship contribution statement

Leandro F. de Souza: Conceptualization, Methodology, Formal analysis, Investigation, Data curation, Project administration, Writing - original draft, Writing - review & editing. **Dasiel O. Alvares:** Conceptualization, Methodology, Formal analysis, Investigation, Writing - original draft, Writing - review & editing. **Luiz A. Domeignoz-Horta:** Conceptualization, Formal analysis, Writing - original draft, Writing - review & editing. **Fabio V. Gomes:** Methodology, Investigation. **Cassio S. Almeida:** Methodology, Investigation. **Luís F. Merloti:** Investigation, Writing - original draft. **Lucas W. Mendes:** Conceptualization, Writing - original draft, Writing - review & editing. **Fernando Dini-Andreote:** Conceptualization, Methodology, Funding acquisition, Writing - review & editing. **Brendan Bohannan:** Conceptualization, Funding acquisition, Writing - review & editing. **Jorge L.M. Rodrigues:** Conceptualization, Funding acquisition, Writing - review & editing. **Klaus Nüsslein:** Conceptualization, Methodology, Resources, Writing - original draft, Funding acquisition, Writing - review & editing, Supervision. **Siu Mui Tsai:** Conceptualization, Methodology, Resources, Writing - original draft, Funding acquisition, Writing - review & editing, Supervision.

Funding

This project was supported by the BIOTA FAPESP - Brazil, and NSF – Dimensions of Biodiversity - USA (2014/50320-4 and DEB 1442183) and by CNPq - Brazil (311008/2016-0). Additional funding in the form of scholarships were provided by FAPESP - Brazil (2018/09117-1), CNPq - Brazil (140953/2017-5), and CAPES - Brazil (001 and 88881.189492/2018-01).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors thank the owners and staff of Farm “Fazenda Nova Vida”, for logistical support and permission to work on their property. We also thank the private landowners: Aristeu, Bernardo e Elói for their support and access to their land. We would like to thank the Large-Scale

Biosphere-Atmosphere Program (LBA), coordinated by the National Institute for Amazon Research (INPA), for the use and availability of data for logistical support and infrastructure during field activities. Additionally, we are grateful to Prof. Plinio B. de Camargo, Henrique Cipriani (EMBRAPA-RO), Alexandre Pedrinho, and to Wagner Piccinini for assistance with fieldwork.

References

- Apprill, A., McNally, S., Parsons, R., Weber, L., 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat. Microb. Ecol.* 75 (2), 129–137. <https://doi.org/10.3354/ame01753>.
- Barton, L., Murphy, D.V., Butterbach-Bahl, K., 2013. Influence of crop rotation and liming on greenhouse gas emissions from a semi-arid soil. *Agric. Ecosyst. Environ.* 167, 23–32. <https://doi.org/10.1016/j.agee.2013.01.003>.
- Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon, M., Bolyen, E., Knight, R., Huttley, G.A., Caporaso, J.G., 2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* 6 (1). <https://doi.org/10.1186/s40168-018-0470-z>.
- Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M., Chase, J., Caporaso, J.G., 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* 37 (8), 852–857. <https://doi.org/10.1038/s41587-019-0209-9>.
- Borken, W., Brumme, R., 1997. Liming practice in temperate forest ecosystems and the effects on CO₂, N₂O and CH₄ fluxes. *Soil Use Manag.* 13, 251–257. <https://doi.org/10.1111/j.1475-2743.1997.tb00596.x>.
- Butterbach-Bahl, K., Rothe, A., Papen, H., 2002. Effect of tree distance on N₂O and CH₄ fluxes from soils in temperate forest ecosystems. *Plant Soil* 240 (1), 91–103. <https://doi.org/10.1023/A:1015828701885>.
- Callahan, B.J., McMurdie, P.J., Holmes, S.P., 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* 11 (12), 2639–2643. <https://doi.org/10.1038/ismej.2017.119>.
- de Carvalho, T.S., Barlow, J., Gardner, T.A., Soares, I.C., Tiedje, J.M., 2016. Land use intensification in the humid tropics increased both alpha and beta diversity of soil bacteria. *Ecol. Appl.* 97 (10), 2760–2771. <https://doi.org/10.1002/ecs.1513>.
- Cole, J.R., Wang, Q., Fish, J.A., Chai, B., McGarrell, D.M., Sun, Y., Brown, C.T., Porras-Alfaro, A., Kuske, C.R., Tiedje, J.M., 2014. Ribosomal database project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res.* 42, D633–D642. <https://doi.org/10.1093/nar/gkt1244>.
- Demattê, J.L.L., Demattê, J.A.M., 1993. Comparações entre as propriedades químicas de solos das regiões da floresta amazônica e do cerrado do Brasil central. *Sci. Agric.* 50 (2), 272–286. <https://doi.org/10.1590/S0103-90161993000200016>.
- Dias, L.C., Pimenta, F.M., Santos, A.B., Costa, M.H., Ladle, R.J., 2016. Patterns of land use, extensification, and intensification of Brazilian agriculture. *Glob. Chang. Biol.* 22 (8), 2887–2903.
- Dias-Filho, M.B., 2017. Degradação de pastagens: o que é e como evitar. *Embrapa, Brasília, DF, Brazil* 978-85-7035-688-8.
- Domeignoz-Horta, L.A., Philippot, L., Peyrard, C., Bru, D., Breuil, M.C., Bizouard, F., Justes, E., Mary, B., Leonard, J., Spor, A., 2018. Peaks of in situ N₂O emissions are influenced by N₂O producing and reducing microbial communities across arable soils. *Glob. Chang. Biol.* 24, 360–370. <https://doi.org/10.1111/gcb.13853>.
- Ettwig, K.F., van Alen, T., van de Pas-Schoonen, K.T., Jetten, M.S., Strous, M., 2009. Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum. *Appl. Environ. Microbiol.* 75, 3656–3662.
- Evans, P.N., Boyd, J.A., Leu, A.O., Woodcroft, B.J., Parks, D.H., Hugenholtz, P., Tyson, G.W., 2019. An evolving view of methane metabolism in the Archaea. *Nat. Rev. Microbiol.* 17, 219–232. <https://doi.org/10.1038/s41579-018-0136-7>.
- Fearnside, P.M., Imbrozio Barbosa, R., 1998. Soil carbon changes from conversion of forest to pasture in Brazilian Amazonia. *For. Ecol. Manag.* 108 (1–2), 147–166. [https://doi.org/10.1016/S0378-1127\(98\)00222-9](https://doi.org/10.1016/S0378-1127(98)00222-9).
- Fedarko, M.W., Martino, C., Morton, J.T., González, A., Rahman, G., Marotz, C.A., Minich, J.J., Allen, E.E., Knight, R., 2020. Visualizing 'omic feature rankings and log-ratios using Qurro. *NAR Genomics Bioinforma.* 2 (2), lqaa023. <https://doi.org/10.1093/nargab/lqaa023>.
- Fernandes, S.A.P., Bernoux, M., Cerri, C.C., Feigl, B.J., Piccolo, M.C., 2002. Seasonal variation of soil chemical properties and CO₂ and CH₄ fluxes in unfertilized and P-fertilized pastures in an Ultisol of the Brazilian Amazon. *Geoderma* 107 (3–4), 227–241. [https://doi.org/10.1016/S0016-7061\(01\)00150-1](https://doi.org/10.1016/S0016-7061(01)00150-1).
- Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial communities. *Proc. Natl. Acad. Sci. U. S. A.* 103 (3), 626–631. <https://doi.org/10.1073/pnas.0507535103>.
- Figueiredo, E.B., Jayasundara, S., Bordonal, R.O., Berchielli, T.T., Reis, R.A., Wagner-Riddle, C., La Scala, N., 2017. Greenhouse gas balance and carbon footprint of beef cattle in three contrasting pasture-management systems in Brazil. *J. Clean. Prod.* 142 (1), 420–431.
- Frenzel, P., Rothfuss, F., Conrad, R., 1992. Oxygen profiles and methane turnover in a flooded rice microcosm. *Biol. Fertil. Soils* 14 (2), 84–89. <https://doi.org/10.1007/BF00336255>.
- García-Marco, S., Abalos, D., Espejo, R., Vallejo, A., Mariscal-Sancho, I., 2016. No tillage and liming reduce greenhouse gas emissions from poorly drained agricultural soils in Mediterranean regions. *Sci. Total Environ.* 566, 512–520. <https://doi.org/10.1016/j.scitotenv.2016.05.117>.

- Goreau, T.J., De Mello, W.Z., 1988. Tropical deforestation: some effects on atmospheric chemistry. *Ambio* 17 (4), 275–281. <https://www.osti.gov/etdweb/biblio/6655174>.
- Hanson, R.S., Hanson, T.E., 1996. Methanotrophic bacteria. *Microbiol. Rev.* 60 (2), 439–471. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC239451/>.
- Herlemann, D.P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J.J., Andersson, A.F., 2011. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J.* 5 (10), 1571–1579. <https://doi.org/10.1038/ismej.2011.41>.
- Herpin, U., Cerri, C.C., Conceição Santana Carvalho, M., Markert, B., Enzweiler, J., Friese, K., Breulmann, G., 2002. Biogeochemical dynamics following land use change from forest to pasture in a humid tropical area (Rondônia, Brazil): a multi-element approach by means of XRF-spectroscopy. *Sci. Total Environ.* 286 (1–3), 97–109. [https://doi.org/10.1016/S0048-9697\(01\)00967-6](https://doi.org/10.1016/S0048-9697(01)00967-6).
- Hütsch, B.W., Webster, C.P., Powlson, D.S., 1994. Methane oxidation in soil as affected by land use, soil pH and N fertilization. *Soil Biol. Biochem.* 26 (12), 1613–1622. [https://doi.org/10.1016/0038-0717\(94\)90313-1](https://doi.org/10.1016/0038-0717(94)90313-1).
- IEA, International Energy Agency, 2021. Methane Tracker 2021. IEA, Paris. <https://www.iea.org/reports/methane-tracker-2021>.
- Jesus, E.C., Marsh, T., Tiedje, J., Moreira, F.M.S., 2009. Changes in land use alter the structure of bacterial communities in Western Amazon soils. *ISME J.* 3, 1004–1011. <https://doi.org/10.1038/ismej.2009.47>.
- Kallistova, A.Y., Merkel, A.Y., Tarnovskii, I.Y., Pimenov, N.V., 2017. Methane formation and oxidation by prokaryotes. *Microbiology* 86 (6), 671–691. <https://doi.org/10.1134/S0026261717060091>.
- Kimura, M., 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* 16 (2), 111–120. <https://doi.org/10.1007/BF01731581>.
- Knief, C., 2015. Diversity and habitat preferences of cultivated and uncultivated aerobic methanotrophic bacteria evaluated based on pmoA as molecular marker. *Terr. Microbiol.* 1346. <https://doi.org/10.3389/fmicb.2015.01346>.
- Knief, C., Lipski, A., Dunfield, P.F., 2003. Diversity and activity of methanotrophic bacteria in different upland soils. *Appl. Environ. Microbiol.* 69 (11), 6703–6714. <https://doi.org/10.1128/AEM.69.11.6703-6714.2003>.
- Kolb, S., 2009. The quest for atmospheric methane oxidizers in forest soils. *Environ. Microbiol. Rep.* 1 (5), 336–346. <https://doi.org/10.1111/j.1758-2229.2009.00047.x>.
- Kroeger, M.E., Meredith, L.K., Meyer, K.M., Webster, K.D., Camargo, P.B., de Souza, L.F., Tsai, S.M., Haren, J., Saleska, S., Bohannon, B.J.M., Rodrigues, J.L.M., Berenguer, E., Barlow, J., Nüsslein, K., 2021. Rainforest-to-pasture conversion stimulates soil methanogenesis across the Brazilian Amazon. *ISME J.* 15, 658–672. <https://doi.org/10.1038/s41396-020-00804-x>.
- Lammel, D.R., Barth, G., Ovaskainen, O., Cruz, L.M., Zanatta, J.A., Ryo, M., de Souza, E.M., Pedrosa, F.O., 2018. Direct and indirect effects of a pH gradient bring insights into the mechanisms driving prokaryotic community structures. *Microbiome* 6 (1), 106. <https://doi.org/10.1186/s40168-018-0482-8>.
- Lau, J.A., Lennon, J.T., 2011. Evolutionary ecology of plant–microbe interactions: soil microbial structure alters selection on plant traits. *New Phytol.* 192, 215–224. <https://doi.org/10.1111/j.1469-8137.2011.03790.x>.
- Laurance, W.F., Laurance, S.G., Delamonica, P., 1998. Tropical forest fragmentation and greenhouse gas emissions. *For. Ecol. Manag.* 110 (1–3), 173–180. [https://doi.org/10.1016/S0378-1127\(98\)00291-6](https://doi.org/10.1016/S0378-1127(98)00291-6).
- Le Mer, J., Roger, P., 2001. Production, oxidation, emission and consumption of methane by soils: a review. *Eur. J. Soil Biol.* 37 (1), 25–50. [https://doi.org/10.1016/S1164-5563\(01\)01067-6](https://doi.org/10.1016/S1164-5563(01)01067-6).
- Liu, C., Holst, J., Brüggemann, N., Butterbach-Bahl, K., Yao, Z., Yue, J., Han, S., Han, X., Krümmelbein, J., Horn, R., Zheng, X., 2007. Winter-grazing reduces methane uptake by soils of a typical semi-arid steppe in Inner Mongolia, China. *Atmos. Environ.* 41 (28), 5948–5958. <https://doi.org/10.1016/j.atmosenv.2007.03.017>.
- Margulis, S., 2003. Causes of Deforestation of the Brazilian Amazon. The World Bank <https://doi.org/10.1596/0-8213-5691-7>.
- Martino, C., Morton, J.T., Marotz, C.A., Thompson, L.R., Tripathi, A., Knight, R., Zengler, K., 2019. A novel sparse compositional technique reveals microbial perturbations. *MSystems* 4 (1). <https://doi.org/10.1128/msystems.00016-19>.
- Mendes, L.W., Tsai, S.M., Navarrete, A.A., de Hollander, M., van Veen, J.A., Kuramae, E.E., 2015. Soil-borne microbiome: linking diversity to function. *Microb. Ecol.* 70 (1), 255–265. <https://doi.org/10.1007/s00248-014-0559-2>.
- Meyer, K.M., Klein, A.M., Rodrigues, J.L.M., Nüsslein, K., Tringe, S.G., Mirza, B.S., Tiedje, J.M., Bohannon, B.J.M., 2017. Conversion of Amazon rainforest to agriculture alters community traits of methane-cycling organisms. *Mol. Ecol.* 26 (6), 1547–1556. <https://doi.org/10.1111/mec.14011>.
- Meyer, K.M., Morris, A.H., Webster, K., Klein, A.M., Kroeger, M.E., Meredith, L.K., Brændholt, A., Nakamura, F., Venturini, A., de Souza, L.F., Danielson, R., van Haren, J., Shek, K.L., de Camargo, P.B., Tsai, S.M., Dini-Andreote, F., Mauro, J., Barlow, J., Berenguer, E., Nüsslein, K., Saleska, S., Rodrigues, J.L.M., Bohannon, B.J.M., 2020. Belowground changes to community structure alter methane-cycling dynamics in Amazonia. *Environ. Int.* 145, 106131.
- de Moraes, J.F.L., Volkoff, B., Cerri, C.C., Bernoux, M., 1996. Soil properties under Amazon forest and changes due to pasture installation in Rondônia, Brazil. *Geoderma* 70 (1), 63–81. [https://doi.org/10.1016/0016-7061\(95\)00072-0](https://doi.org/10.1016/0016-7061(95)00072-0).
- Mosier, A.R., Delgado, J.A., Keller, M., 1998. Methane and nitrous oxide fluxes in an acid Oxisol in western Puerto Rico: effects of tillage, liming and fertilization. *Soil Biol. Biochem.* 30 (14), 2087–2098. [https://doi.org/10.1016/S0038-0717\(98\)00085-6](https://doi.org/10.1016/S0038-0717(98)00085-6).
- Navarrete, A.A., Tsai, S.M., Mendes, L.W., Faust, K., de Hollander, M., Cassman, N.A., Raes, J., van Veen, J.A., Kuramae, E.E., 2015. Soil microbiome responses to the short-term effects of Amazonian deforestation. *Mol. Ecol.* 24 (10), 2433–2448. <https://doi.org/10.1111/mec.13172>.
- Nazaries, L., Murrell, J.C., Millard, P., Baggs, L., Singh, B.K., 2013. Methane, microbes and models: fundamental understanding of the soil methane cycle for future predictions. *Environ. Microbiol.* 15 (9). <https://doi.org/10.1111/1462-2920.12149>.
- Oksanen, J., Blanchet, F.G., Kindt, R., Wagner, L.H., 2015. Vegan: Community ecology package. R package version 2.2-1. Available online at: <http://CRAN.R-project.org/package=vegan>.
- Oliveira, P.P.A., Boaretto, A.E., Trivelin, P.C.O., de Oliveira, W.S., Corsi, M., 2003. Liming and fertilization to restore degraded Brachiaria decumbens pastures grown on an entisol. *Sci. Agric.* 60 (1), 125–131.
- Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: assessing small sub-unit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ. Microbiol.* 18 (5), 1403–1414. <https://doi.org/10.1111/1462-2920.13023>.
- Peres-Neto, P.R., Legendre, P., Dray, S., Borcard, D., 2006. Variation partitioning of species data matrices: estimation and comparison of fractions. *Ecology* 87, 2614–2625.
- Philippot, L., Hallin, S., Börjesson, G., Baggs, E.M., 2009. Biochemical cycling in the rhizosphere having an impact on global change. *Plant Soil* 321, 61–81. <https://doi.org/10.1007/s11040-008-9796-9>.
- Philippot, Laurent, Raaijmakers, J.M., Lemanceau, P., van der Putten, W.H., 2013. Going back to the roots: the microbial ecology of the rhizosphere. *Nat. Rev. Microbiol.* 11 (11), 789–799. <https://doi.org/10.1038/nrmicro3109>.
- Praeg, N., Wagner, A.O., Illmer, P., 2017. Plant species, temperature, and bedrock affect net methane flux out of grassland and forest soils. *Plant Soil* 410 (1–2), 193–206. <https://doi.org/10.1007/s11040-016-2993-z>.
- Pratscher, J., Vollmers, J., Wiegand, S., Dumont, M.G., Kaster, A.-K., 2018. Unravelling the identity, metabolic potential and global biogeography of the atmospheric methane-oxidizing upland soil cluster α. *Environ. Microbiol.* <https://doi.org/10.1111/1462-2920.14036>.
- Quast, C., Priesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O., 2012. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* 41 (D1), D590–D596. <https://doi.org/10.1093/nar/gks1219>.
- R Core Team, 2013. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria URL <http://www.R-project.org/>.
- van Raij, B., Andrade, J.C., Cantarella, H., Quaggio, J.A., de, 2001. *Análise química para avaliação da fertilidade de solos tropicais* Campinas, Brazil.
- Ramakers, C., Ruijter, J.M., Lekanne Deprez, R.H., Moorman, A.F.M., 2003. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci. Lett.* 339 (1), 62–66. [https://doi.org/10.1016/S0304-3940\(02\)01423-4](https://doi.org/10.1016/S0304-3940(02)01423-4).
- Reiners, W.A., Bouwman, A.F., Parsons, W.F.J., Keller, M., 1994. Tropical rain forest conversion to pasture: changes in vegetation and soil properties. *Ecol. Appl.* 4 (2), 363–377. <https://doi.org/10.2307/1941940>.
- Rodrigues, J.L.M., Pellizari, V.H., Mueller, R., Baek, K., Paula, F.S., Mirza, B., Hamaoui, G.S., Tsai, S.M., Feigl, B., Tiedje, J.M., Bohannon, B.J.M., Nüsslein, K., Jesus, E., 2013. Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. *Proc. Natl Acad. Sci. USA* 110 (3), 988–993. <https://doi.org/10.1073/pnas.1220608110>.
- Segnini, A., Xavier, A.P., Otaviani-Junior, P.L., Oliveira, P.P.A., Pedrosa, A.F., Praes, M.F.M.M., Rodrigues, P.H.M., Milori, D.M.B.P., 2019. Soil carbon stock and humification in pastures under different levels of intensification in Brazil. *Sci. Agric.* 76 (1), 33–40. <https://doi.org/10.1590/1678-992x-2017-0131>.
- Semrau, J.D., Chistoserdov, A., Lebron, J., Costello, A., Davagnino, J., Kenna, E., Holmes, A.J., Finch, R., Murrell, J.C., Lidstrom, M.E., 1995. *Particulate methane monooxygenase genes in methanotrophs*. *J. Bacteriol.* 177, 3071–3079.
- Serrano-Silva, N., Sarria-Guzmán, Y., Dendooven, L., Luna-Guido, M., 2014. Methanogenesis and methanotrophy in soil: a review. *Pedosphere* 24 (3), 291–307. [https://doi.org/10.1016/S1002-0160\(14\)60016-3](https://doi.org/10.1016/S1002-0160(14)60016-3).
- Stauder, P.A., Melillo, J.M., Feigl, B.J., Neill, C., Piccolo, M.C., Cerri, C.C., 1996. Consequence of forest-to-pasture conversion on CH₄ fluxes in the Brazilian Amazon Basin. *J. Geophys. Res.-Atmos.* 101 (D13), 18547–18554. <https://doi.org/10.1029/96JD01551>.
- Tamas, I., Smirnova, A.V., He, Z., Dunfield, P.F., 2014. The (d)evolution of methanotrophy in the Beijerinckiaceae—a comparative genomics analysis. *ISME J.* 8 (2), 369–382. <https://doi.org/10.1038/ismej.2013.145>.
- Tate, K.R., 2015. Soil methane oxidation and land-use change – from process to mitigation. *Soil Biol. Biochem.* 80, 260–272. <https://doi.org/10.1016/j.soilbio.2014.10.010>.
- Tian, H., Chen, G., Lu, C., Xu, X., Ren, W., Zhang, B., Banger, K., Tao, B., Pan, S., Liu, M., Zhang, C., Bruhwiler, L., Wofsy, S., 2015. Global methane and nitrous oxide emissions from terrestrial ecosystems due to multiple environmental changes. *Ecosyst. Health Sustain.* 1 (1), 1–20. <https://doi.org/10.1890/EHS14-0015.1>.
- Ussiri, D.A.N., Lal, R., Jarecki, M.K., 2009. Nitrous oxide and methane emissions from long-term tillage under a continuous corn cropping system in Ohio. *Soil Tillage Res.* 104 (2), 247–255.
- Venturini, A.M., Nakamura, F.M., Gontijo, J.B., da França, C.A., Yoshiura, C.A., Mandro, J.A., Tsai, S.M., 2020. Robust DNA protocols for tropical soils. *Heliyon* 6 (5), e03830. <https://doi.org/10.1016/j.heliyon.2020.e03830>.
- Verchot, L.V., Davidson, E.A., Cattaneo, J.H., Ackerman, I.L., 2000. Land-use change and biogeochemical controls of methane fluxes in soils of Eastern Amazonia. *Ecosystems* 3 (1), 41–56. <https://doi.org/10.1007/s100210000009>.
- Wang, Y., Yao, Z., Zhang, Y., Zheng, X., Zhou, M., Yan, G., Wang, L., Werner, C., Butterbach-Bahl, K., 2021. Potential benefits of liming to acid soils on climate change mitigation and food security. *Glob. Chang. Biol.* <https://doi.org/10.1111/gcb.15607>.
- Whittenbury, R., Davies, S.L., Davey, J.F., 1970. Exospores and cysts formed by methane-utilizing Bacteria. *Microbiology* 61 (2), 219–226. <https://doi.org/10.1099/00221287-61-2-219>.