

FtsH2 and FtsH5: two homologous subunits use different integration mechanisms leading to the same thylakoid multimeric complex

Ricardo A. O. Rodrigues¹, Marcio C. Silva-Filho^{1,*} and Kenneth Cline^{2,*}

¹Departamento de Genética, Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo, Piracicaba, SP, Brazil, and

²Horticultural Sciences Department and Plant Molecular and Cellular Biology, University of Florida, Gainesville, FL, USA

Received 1 October 2010; revised 16 November 2010; accepted 25 November 2010; published online 10 January 2011.

*For correspondence (fax +55 19 3433 6706; e-mail mdcsilva@esalq.usp.br or kcline@ufl.edu).

SUMMARY

The *Arabidopsis* thylakoid FtsH protease complex is composed of FtsH1/FtsH5 (type A) and FtsH2/FtsH8 (type B) subunits. Type A and type B subunits display a high degree of sequence identity throughout their mature domains, but no similarity in their amino-terminal targeting peptide regions. In chloroplast import assays, FtsH2 and FtsH5 were imported and subsequently integrated into thylakoids by a two-step processing mechanism that resulted in an amino-proximal luminal domain, a single transmembrane anchor, and a carboxyl proximal stromal domain. FtsH2 integration into washed thylakoids was entirely dependent on the proton gradient, whereas FtsH5 integration was dependent on NTPs, suggesting their integration by Tat and Sec pathways, respectively. This finding was corroborated by *in organello* competition and by antibody inhibition experiments. A series of constructs were made in order to understand the molecular basis for different integration pathways. The amino proximal domains through the transmembrane anchors were sufficient for proper integration as demonstrated with carboxyl-truncated versions of FtsH2 and FtsH5. The mature FtsH2 protein was found to be incompatible with the Sec machinery as determined with targeting peptide-swapping experiments. Incompatibility does not appear to be determined by any specific element in the FtsH2 domain as no single domain was incompatible with Sec transport. This suggests an incompatible structure that requires the intact FtsH2. That the highly homologous type A and type B subunits of the same multimeric complex use different integration pathways is a striking example of the notion that membrane insertion pathways have evolved to accommodate structural features of their respective substrates.

Keywords: chloroplast, protease, protein transport, twin arginine, SecY, Tat.

INTRODUCTION

Thylakoidal FtsH proteins are bacterial-type metalloproteases that belong to the subfamily ATPases Associated with diverse cellular Activities (AAA⁺). In general, FtsH proteases have two transmembrane domains followed by a cytosolic or stromal ATPase module with Walker A and B motifs, a Zn⁺ binding region, and a proteolytic domain (Ito and Akiyama, 2005). FtsH proteases are known to degrade short-lived membrane proteins in a number of organisms. Studies in prokaryotes show that FtsH proteins form hexameric rings where the ATP binding motif faces the center of the ring (Niwa *et al.*, 2002). Although there are nine putative plastid FtsH family members in *Arabidopsis*, only one thylakoid FtsH complex has been detected in significant quantity, and this consists of FtsH1/FtsH5 (type A) and FtsH2/FtsH8 (type B)

subunits (Sakamoto *et al.*, 2003). Co-immunoprecipitation studies demonstrate that type A and type B subunits are members of the same complex (Sakamoto *et al.*, 2003; Yu *et al.*, 2004). Disruption of the *FtsH5* gene results in a variegated phenotype (*var1*); disruption of the *FtsH2* gene also results in a variegated phenotype (*var2*). The *var1* and *var2* phenotypes can be rescued by overexpression of FtsH1 and FtsH8, respectively (Yu *et al.*, 2004, 2005). Thus, only one of each isomer is required for correct function of the thylakoid FtsH complex (Zaltsman *et al.*, 2005).

All chloroplast FtsH proteins are encoded on nuclear genes and are synthesized in the cytosol. Cytosolically synthesized thylakoid proteins of chloroplasts are routed to their functional locations by several translocases (Cline

and Dabney-Smith, 2008). Most are initially imported into the plastid stroma by the Toc and Tic complexes in the outer and inner envelope membranes. The Sec, Tat, SRP/Alb3, and an unassisted pathway integrate the stromal intermediates into the thylakoid bilayer or transport them to the lumen (Cline and Dabney-Smith, 2008; Cline and Theg, 2007). Of particular interest are the Tat pathway (for twin arginine translocation), which transports fully folded protein substrates and relies only on the proton gradient, and the Sec pathway, which transports unfolded proteins through a SecYE channel by virtue of ATP binding and hydrolysis by the SecA translocation motor.

Here we analyzed the targeting mechanisms of two *Arabidopsis* FtsH members, FtsH5 and FtsH2, as representative members of type A and type B families. Both isoforms have an amino terminal stroma-targeting transit peptide followed by a hydrophobic sequence, which we show functions as a cleavable hydrophobic signal peptide rather than a transmembrane domain. We show that the Tat pathway integrates FtsH2, whereas the Sec pathway integrates FtsH5. Carboxyl proximal truncations suggest that targeting specificity resides in the signal peptide of each isoform. Transit peptide swapping experiments indicated

an incompatibility of the FtsH2 mature region with the Sec pathway, explaining the need for two different integration systems to target two highly homologous subunits. This finding demonstrates an intriguing twist on the biogenesis of membrane protein complexes, whereby two highly homologous subunits of the same multimeric complex are delivered to the membrane by different integration machineries.

RESULTS

Type A and type B FtsH subunits differ by the presence/absence of the twin arginine motif

The two FtsH type A family members share a high degree of homology throughout their entire peptide sequence; this is also true for type B FtsH family members, suggesting recent gene duplications (Figure S1b,c). By contrast, whereas FtsH type A and FtsH type B show identity in their stroma-exposed ATPase and protease domains, there is little homology between their chloroplast targeting peptides and the sequences flanking the amino proximal hydrophobic domain (Figure S1a). In particular, the type B proteins FtsH2 and FtsH8 possess a twin-arginine motif just before, and an A–X–A signal peptidase cleavage site consensus (where X is




Figure 1. FtsH type A and B differ by the presence/absence of the twin arginine motif.

The amino acid sequences of the N-terminal region of four *Arabidopsis* FtsH proteases are aligned.

The amino-acid sequences of the N-terminal region of four *Arabidopsis* FtsH precursors are aligned.

- Precursors to type A, FtsH5 and FtsH1, have a chloroplast-targeting domain followed by a hydrophobic domain and the tripeptide A-L-A, a cleavage motif for the thylakoid signal peptidase (Tpp). The latter two features are characteristic of a hydrophobic signal peptide.
- Precursors to type B, FtsH2 and FtsH8, possess a chloroplast targeting domain, a classical Tat motif (R-R-X-F-L-K), a hydrophobic domain, and an A-N-A Tpp cleavage site. The latter three features are characteristic of a Tat pathway signal peptide.
- Diagram of a generic (i.e. type A or type B) FtsH precursor protein. In the present work, the chloroplast targeting domain and the signal peptide combined are called the 'transit peptide' (TP). The luminal domain (L), transmembrane anchor (T), and stromal domain (S) are marked. In (a) and (b) of the figure, the amino-acid sequences start after the predicted Spp cleavage site. The arrows in sequence panels (a) and (b) are swapping points for fusions between FtsH5 and FtsH2 or truncation points (see below).

any amino acid) just after the first hydrophobic domain, suggesting that they are Tat-directed cleavable signal peptides (Figure 1b). For FtsH type A proteins the first hydrophobic domain also appears to be a cleavable signal peptide as evidenced by a relatively low hydrophobicity and an A–X–A cleavage site (Figure 1a). This sequence analysis suggests that the mature proteins for type A and type B FtsH proteases consist of an amino terminal lumen-facing domain (L), a transmembrane anchor (T), and a large stroma-facing catalytic domain (S) (Figure 1c). Because amino-acid sequence identity of family members within a type was very high, even in the signal peptide and transit peptide regions (Figure S1b,c) we selected FtsH5 and FtsH2 for the type A and type B groups, respectively, for *in vitro* assays to determine their paths to the thylakoids.

pFtsH2 and pFtsH5 are localized *in vitro* by a two-step pathway

In vitro translated precursor to FtsH2 (pFtsH2; 74 kDa) and precursor to FtsH5 (pFtsH5; 75 kDa) were incubated with intact pea chloroplasts. The precursors were imported into the organelle and processed to a mature form of 65 kDa for mFtsH2 and 67 kDa for mFtsH5 (Figure 2a,b, lane 2). Protease treatment of recovered chloroplasts confirmed that the mature forms were inside of the chloroplasts (lane 3). Fractionation of the recovered chloroplasts showed that both mature proteins were present in the thylakoid fraction (T, lane 5) rather than the soluble stromal fraction (S, lane 4).

To verify the proper integration into the membrane, aliquots of the thylakoid fraction were extracted with

100 mM NaOH, 200 mM Na₂CO₃, or treated with thermolysin. The FtsH2 protein was resistant to extraction with NaOH (Figure 2a, TN, lane 7) and thermolysin treatment produced a partial degradation product of approximately 13 kDa (Figure 2a, lane 14), which is the expected size of the luminal tail plus one transmembrane domain. That the protease resistance of the FtsH2 fragment is due to protection by the membrane bilayer was verified by treatment of thylakoids with thermolysin plus 1% Triton X-100, which resulted in complete degradation (Figure S2). These characteristics are in agreement with those observed for the previously characterized pepper chromoplast FtsH protein Pftf (Summer *et al.*, 2000) (Figure 2a). For the following experiments, the presence of the characteristic protease protected fragment or resistance to NaOH was taken as an indicator of proper integration of FtsH2.

In vitro translation of FtsH5 either with a homemade wheat germ system or a TNT coupled Transcription and Translation Kit (Promega) produced both the 75 kDa full length precursor protein and an additional C-terminally truncated protein of approximately 25 kDa (Figure S3). Production of C-terminally truncated proteins (early quitters) from large mRNAs is not unusual. This truncated precursor appeared to be imported into the chloroplasts and processed to approximately 17 kDa, which is very close to the migration of the expected protease-protected product. For this reason, resistance to extraction with Na₂CO₃ was used to assess integration of FtsH5 into membrane. Imported and thylakoid localized FtsH5 was fully resistant to extraction with the alkaline 200 mM Na₂CO₃ (Figure 2b, TCO, lane 7),




Figure 2. pFtsH2 and pFtsH5 are imported into isolated chloroplasts, processed to an intermediate form, and integrated into thylakoids.

Radiolabeled *in vitro* translated pFtsH2, pPftf, (a) pFtsH5 or pOE33 (b) (lane 1, tp) were incubated with intact isolated pea chloroplasts in a reaction containing 5 mM ATP and approximately 100 μ E m⁻² sec⁻¹ light at 25°C for 20 min, in the absence or presence of nigericin and valinomycin (for pFtsH2 and pPftf) or sodium azide (for pFtsH5 and pOE33), as depicted above the panels. Intact chloroplasts were recovered from the reaction (C, lanes 2 and 8) and treated with thermolysin (C+, lanes 3 and 9). Untreated intact chloroplasts were fractionated into stroma (S, lanes 4 and 10) and thylakoids. Thylakoids aliquots were washed with import buffer (T, lanes 5 and 11), treated with thermolysin (T+, lanes 6 and 12) or treated with 100 mM NaOH or 200 mM Na₂CO₃ (TN or TCO, respectively, lanes 7 and 13). Samples were analyzed by SDS-PAGE on 7.5% gels and fluorography, except for thermolysin treated thylakoids, which were analyzed on 12.5% gels (a, lanes 14 and 15). Translation products (tp, lanes 1) represent 2% and all other samples represent 5% of the amount present in each assay. The precursor, intermediate, and mature forms of the proteins are designated p, i, and m, respectively, on the right side of the panels. Mr values in parentheses next to the various species were estimated from their migration compared with standard marker proteins.

although not resistant to extraction with 100 mM NaOH (data not shown). Resistance to 200 mM Na₂CO₃ is a generally accepted criterion for a membrane anchored proteins, although some membrane proteins are resistant to the harsher 100 mM NaOH treatment. It's not clear why FtsH2, but not FtsH5, is resistant to NaOH, as both have strongly hydrophobic transmembrane anchors as determined by Membrane Protein Explorer. Nevertheless, resistance to 200 mM Na₂CO₃ was taken as an indicator of proper integration of FtsH5.

A two-step processing mechanism for localization of FtsH2 and FtsH5 is suggested by the prediction of sequentially cleaved targeting peptides, i.e. cleavage by the stromal peptidase produces an intermediate sized stromal form; cleavage by the thylakoid signal peptidase produces a smaller integrated mature protein. This was tested by import in the presence of inhibitors characteristic for the Tat and SecA/SecYE pathways. For pFtsH2, the chloroplast import reaction was conducted in the presence of ionophores nigericin and valinomycin, which dissipates the proton motive force, the driving force for the Tat pathway. This prevented FtsH2 membrane integration (Figure 2a, lanes 13 and 15) and resulted in accumulation of an intermediate sized form (lane 10) that was found both in the stromal fraction and associated with but not integrated into the membranes. Nigericin/valinomycin has a limited and variable effect on the Sec pathway (Yuan and Cline, 1994) and had virtually no effect on pFtsH5 integration (data not shown and Figure 3, lane 6). However, azide, which inhibits SecA of the Sec pathway (Yuan and Cline, 1994), partially inhibited integration of FtsH5 (Figure 2b, lane 13), resulting in accumulation of an intermediate-sized form largely associated


with the membrane fraction (lane 11) and which was largely extracted by Na₂CO₃. Interestingly, the truncated FtsH5 protein accumulated an intermediate-sized form in the stroma and membranes of azide-treated chloroplasts, and the membrane-associated form was extracted by Na₂CO₃ (Figure S3). Thylakoid transport of the control SecA/SecYE substrate pOE33 was also inhibited by sodium azide and accumulated the intermediate iOE33 (Figure 2b).

The results demonstrate that pFtsH2 and pFtsH5 possess bipartite transit peptides with a chloroplast-targeting domain that is removed after import into the stroma and a signal peptide that is removed after integration into thylakoids. Combined with the protease and alkaline extraction data in Figure 2 and Figure S3, this result supports a model in which both FtsH2 and FtsH5 have a single transmembrane anchor and a large carboxyl-proximal stroma-exposed catalytic domain.

pFtsH2 requires the proton gradient whereas pFtsH5 requires NTPs for integration

To further examine the requirements specific for thylakoid integration, pFtsH2 and pFtsH5 were directly assayed for integration into isolated thylakoids and compared with the Tat substrate pOE23 and the SecA/SecY substrate pOE33 (Figure 3). In the presence of stromal extract, light to generate a proton gradient, and ATP, both proteins were integrated into the membrane (lane 1) as assessed by the partial protection against thermolysin post-treatment (for FtsH2) or resistance to extraction with 200 mM Na₂CO₃ (for FtsH5). FtsH2 integration was dependent on the proton motive force as it was inhibited by the ionophores nigericin and valinomycin in the light (lane 6) and in the dark was

Figure 3. pFtsH2 integration into isolated thylakoids requires the thylakoidal proton motive force whereas FtsH5 integration requires NTPs. The requirements for integration into isolated pea thylakoids was examined by incubating radiolabeled pFtsH2, pFtsH5, pOE23 or pOE33 for 30 min at 25°C under the conditions depicted above the panels. These include the absence or presence of stromal extract, light, 5 mM ATP, nigericin and valinomycin (N/V), sodium azide (SA), and apyrase. The recovered thylakoids were treated with thermolysin (for pFtsH2, pOE23 and pOE33) or extracted with 200 mM Na₂CO₃ (for pFtsH5). The samples were analyzed by SDS-PAGE and fluorography on 12.5% gels. Bands in the fluorograms were quantified using Image J software and the amounts in each band relative to the band in lane 1 of each panel are depicted below the panels.



ATP-dependent, via reverse action of the ATP synthase (Figure 3, lanes 4 and 5). In the presence of light FtsH2 integration occurred without stromal extract (Figure 3, lanes 2 and 7) and without NTPs, which were scrubbed from the reaction with apyrase (lanes 3, and 7), and in presence of sodium azide (lane 8). Identical requirements were exhibited for transport of the Tat substrate pOE23. pFtsH5 integration was dependent on NTPs (lanes 3 and 7) and reduced in samples treated with sodium azide (lane 8). The integration was unaffected by ionophores (lane 6). pFtsH5 requirements are similar to the requirements for transport of the SecA/SecY substrate pOE33, with two notable differences; FtsH5 integration was virtually unaffected by the absence of stromal extract, the main source of SecA (lane 2), whereas pOE33 transport was significantly reduced, and FtsH5 integration was reduced by sodium azide whereas pOE33 transport was eliminated (lane 8). This finding might indicate a smaller dependence of FtsH5 than pOE33 on SecA, or possibly that an unidentified NTPase serves as a translocation motor for FtsH5 integration.

Antibody to Hcf106 inhibits pFtsH2 integration; antibody to SecY inhibits pFtsH5 integration

The involvement of Tat and Sec translocation machinery in FtsH integration was tested by antibody inhibition assays, whereby thylakoids were pretreated with antibodies to components of the three thylakoid pathways (Mori and Cline, 2001; Mori *et al.*, 1999). As shown in Figure 4, treatment of thylakoids with antibody to the Tat component Hcf106 prevented integration of FtsH2 (lane 2), whereas treatment of thylakoids with antibody to chloroplast SecY




Figure 4. Pre-treating thylakoids with Hcf106 antibody prevents integration of pFtsH2, whereas pre-treating with SecY antibody prevents pFtsH5 integration. For antibody inhibition, pea thylakoids were incubated without (–) or with anti-Hcf106 IgG (1 mg ml⁻¹), anti-SecY serum, or anti-Alb3 IgG (1 mg ml⁻¹) for 1 h on ice. Thylakoids were washed with import buffer, supplemented with stromal extract, and incubated with radiolabeled pFtsH2, pFtsH5, pOE23, pOE33 or pLHCP and 5 mM ATP in the light at 25°C for 30 min. The samples were then treated with thermolysin (for pFtsH2, pOE23, pOE33 and pLHCP) or 200 mM Na₂CO₃ (for pFtsH5). Samples were analyzed by SDS-PAGE and fluorography on 12.5% gels. FtsH2-DP is the thermolysin degradation product of the integrated protein and LHCP-DP is the characteristic thermolysin degradation product of integrated LHCP.

(lane 3) inhibited the integration of FtsH5 (lane 3). Integration of either protein was unaffected by antibodies to the chloroplast SRP pathway component Alb3 (lane 4). The effects on antibody treatments on the control proteins pOE23 (Tat), pOE33 (SecA/SecYE), and pLHCP (SRP/Alb3) confirmed the efficacy and specificity of the treatments. As an additional test for pathway specificity, the integration pathways of FtsH2 and FtsH5 was queried by an *in organello* competition assay, wherein a pre-import incubation accumulates thylakoid saturating concentrations of either a Tat substrate (iOE23) or a Sec substrate (iOE33) before challenge with radiolabeled precursor proteins. As expected, the Tat substrate competed for FtsH2 localization, whereas the SecA/SecYE substrate competed for FtsH5 localization (Figure S4).

The amino-proximal regions of pFtsH2 and pFtsH5 are sufficient for thylakoid integration

Previous work in many laboratories has shown that targeting specificity to SecA/SecYE or Tat pathways is governed by the signal peptide. However, a previous attempt by us to characterize targeting determinants *in vivo* suggested that the mature domain of FtsH5 was important for thylakoid targeting (unpublished results). Specifically GFP was fused in frame to the amino terminal sequences of FtsH2 and FtsH5 from the amino terminus to the second hydrophobic domain carboxyl flanking sequence. Although pFtsH2-GFP was correctly localized to thylakoids in tobacco transgenic plants, pFtsH5-GFP accumulated in the stromal compartment. In order to clarify whether or not all of the targeting specificity resides in the amino terminal domains of pFtsH2 and pFtsH5, truncated versions of pFtsH2 (pFtsH2Δ) and pFtsH5 (pFtsH5Δ), containing the amino terminal sequences but lacking the stroma-facing enzymatic domain, were prepared by *in vitro* transcription and translation. As shown in Figure 5, both truncated precursors, pFtsH2Δ and pFtsH5Δ, were able to correctly localize in an *in vitro* chloroplast import assay. pFtsH2Δ was imported into chloroplasts (left panel, C, C+) and inserted into thylakoids, as confirmed by protection against protease treatment (T, T+) and NaOH extraction (TN). pFtsH5Δ was imported into chloroplasts (right panel, C, C+) and inserted into thylakoids, as confirmed by protection against protease treatment (T, T+) and Na₂CO₃ extraction (TCO). These results demonstrate that all necessary requirements for import of pFtsH2 and pFtsH5 into isolated chloroplasts and subsequent integration into thylakoids reside in the N-terminal domain.

The pFtsH2 mature domain is incompatible for integration by the Sec-pathway

As an initial test for the underlying molecular basis for dual integration pathways for homologous proteins of the same multimeric complex, we prepared chimeric precursor proteins, in which the transit peptide (including the signal peptide) of one precursor protein was fused to the mature




Figure 5. The N-terminal regions of pFtsH2 and pFtsH5 are able to insert into thylakoid membranes.

Radiolabeled pFtsH2Δ or pFtsH5Δ (as diagrammed below the panels) was incubated with intact pea chloroplasts in a reaction containing 5 mM ATP and light at 25°C for 20 min. Intact chloroplasts were recovered (C) and treated with thermolysin (C+). Untreated intact chloroplasts were fractionated into stroma (S) and thylakoids. Thylakoid aliquots were washed with import buffer (T), treated with thermolysin (T+) or extracted with 100 mM NaOH (TN). For pFtsH5Δ, a thylakoid aliquot was extracted with 200 mM Na₂CO₃ (TCO). Samples were analyzed by SDS-PAGE on 12.5% gels and fluorography.

domain of the other, producing TP2Mat5 and TP5Mat2 (see Figure 1a,b), points (a). As shown in Figure 6(a) the chimeric precursor TP2Mat5 (76 kDa) was imported into chloroplasts, localized to thylakoids, and processed to a mature form (67 kDa, lane 2). Because *in vitro* translation of TP2Mat5 only produces the full length precursor protein, protease treatment of thylakoids was used to verify membrane integration. The protease treated membranes produced a protected fragment of approximately 17 kDa (lane 11), confirming integration. Essentially the same results were obtained in a parallel assay but with the chloroplasts pre-treated with sodium azide (lanes 8–10 and 13). By contrast, when the chloroplasts were treated with nigericin and valinomycin (N/V), the imported protein accumulated as intermediate sized precursor (approximately 72 kDa) in the stromal fraction (lane 6) and thylakoids, which was completely degraded by protease treatment (lane 12). These results indicate that the pFtsH2 transit peptide and signal peptide are able to direct import and integration of the FtsH5 mature domain by the Tat pathway.

The chimeric precursor TP5Mat2 (73 kDa) was imported into chloroplasts and processed to a 70 kDa form (Figure 6, lane 2), which is slightly larger than the 65 kDa band expected after a thylakoid processing step. Interestingly, the processed form remained exclusively associated with the stromal fraction (lane 3). The same result was obtained with the nigericin/valinomycin treated chloroplasts (lanes 5–7), and sodium azide treated chloroplasts (lanes 8–10). This result indicates that the pFtsH5 transit peptide was not able to direct integration of the FtsH2 mature region into thylakoids, suggesting that the FtsH2 mature region has a conformational or sequence-based incompatibility with the Sec machinery.

The FtsH2 luminal tail, transmembrane anchor, and stromal exposed regions are all required for incompatibility

There are three different domains of the mature FtsH2 that could potentially cause incompatibility, the luminal tail, the transmembrane anchor, and the large catalytic stromal domain. Three different fusion precursor proteins were




Figure 6. The FtsH2 mature domain is incompatible for transport by the Sec-pathway.

Radiolabeled TP2Mat5 (a) or TP5Mat2 (b) as diagrammed below the panels was incubated with intact pea chloroplasts in a reaction containing 5 mM ATP and light at 25°C for 20 min, in the presence or absence of nigericin and valinomycin (N/V) or sodium azide (SA). Intact chloroplasts were recovered from reaction and treated with thermolysin (C+). Untreated intact chloroplasts were fractionated into stroma (S) and thylakoids. Thylakoid aliquots were washed with import buffer (T) or treated with thermolysin (T+). Samples were analyzed by SDS-PAGE and fluorography on 7.5% gels except for the T+ samples, which were analyzed on a 12.5% gel. Arrows on diagrams denote swapping points between FtsH5 and FtsH2 sequences.




Figure 7. Both the FtsH2 lumen exposed- and stroma exposed-domains are required for incompatibility with the Sec pathway. Radiolabeled TP5Mat2Δ, TPL5S2, and TPL5TS2 were incubated with intact isolated pea chloroplasts in a reaction containing 5 mM ATP in light at 25°C for 20 min. Intact chloroplasts were recovered and treated with thermolysin (C+). Untreated intact chloroplasts were fractionated into stroma (S) and thylakoids. Thylakoids aliquots were washed with import buffer (T), treated with thermolysin (T+), or extracted with 200 mM Na₂CO₃ (Na₂CO₃) or with 100 mM NaOH (NaOH). Samples were analyzed by SDS-PAGE and fluorography. The transit peptide (TP), luminal domain (L), and transmembrane domain (T) are labeled 2 or 5 to signify their origin from FtsH2 or FtsH5, respectively.

constructed to test these domains for incompatibility: TP5Mat2Δ, which has the same swapping point of TP5Mat2 but lacks the stroma-exposed domain to test the FtsH2 luminal (L) and transmembrane (T) domains (Figure 7a); TPL5S2, in which the FtsH5 sequence through the transmembrane domain is fused to the FtsH2 stromal domain (Figure 7b); and TPL5TS2 in which the FtsH5 transit peptide and luminal tail are fused to the FtsH2 transmembrane domain and stromal domain (Figure 7c). As shown in Figure 7, all three fusion proteins were imported into chloroplasts (lanes 2), proteolytically processed, and localized to thylakoids (lanes 4). Each protein was properly integrated into thylakoids as determined by either partial protease protection (Figure 7a, lane 5) or by resistance to 100 mM

NaOH (Figure 7c, lane 6) or resistance to 200 mM Na₂CO₃ (Figure 7b, lane 5), depending on the source of the transmembrane anchor in the fusion protein. This result indicates that no single FtsH2 domain or combination of two FtsH2 domains prevent integration by the Sec machinery. Perhaps there is a folded structure consisting of the all domains that prevents access by the Sec machinery.

DISCUSSION

Three classes of bacterially conserved ATP-dependent proteases in chloroplasts are the Clp, Lon, and FtsH proteases [for review see (Adam *et al.*, 2006; Sakamoto, 2006)]. Clp and Lon are located in the stroma and FtsH is located in the membrane. All possess the ATP-binding motif but differ in their catalytic domains for proteolysis. The FtsH protease was originally described in *E. coli* and shown to degrade short-lived proteins (for review see Ito and Akiyama, 2005). FtsH, as well as orthologous FtsH proteins in different organisms, have two transmembrane domains, followed by an ATPase domain, and a zinc-binding motif, which serves as the catalytic site of the protease. These sites face the central pore of hexamers, access to which is controlled by the ATPase domain. The ATPase domain is also thought to translocate the polypeptide through the protease site as well as to dislocate membrane protein substrates out of the bilayer (Ito and Akiyama). Arabidopsis has 12 FtsH genes (FtsH1-12). Eight of the 12 FtsHs (FtsH1, 2, 5, 6, 7, 8, 9, 12) appear to reside in chloroplasts based on GFP fusion localizations (Sakamoto *et al.*, 2003), three appear to be mitochondrial (FtsH3, 4, 10), and FtsH11 is dual targeted to chloroplasts and mitochondria (Urantowka *et al.*, 2005). Chloroplast targeted FtsH1/FtsH5 (type A), FtsH2/FtsH8 (type B), and FtsH7/FtsH9 are closely related pairs based on sequence identity and genetic analysis. It is now clear that type A subunits and type B subunits combine to assemble a single type of multimeric complex (Yu *et al.*, 2004, 2005). Type A and type B are the most abundant plastid FtsHs.

Our original intention was to include FtsH7/FtsH9 pair in this study because FtsH7 has been detected in a proteomics study of the plastid envelope (Ferro *et al.*, 2003) and appears to possess two transmembrane anchors, rather than the single anchors demonstrated here for FtsH2 and FtsH5. In preliminary work, we determined that pFtsH7 and pFtsH9 were imported into isolated chloroplasts and localized to the membrane fraction in an alkaline extraction resistant form (Rodrigues *et al.*, unpublished). This finding confirms that FtsH7/9 are indeed plastid membrane proteins. Unfortunately, our attempts to define their routing pathways were unsuccessful.

The routing and integration pathways for type A and the type B subunits were much more amenable to *in vitro* studies. Our data indicate that both proteins are localized by a two-step pathway in which the precursor is imported into

the plastid stroma where the chloroplast targeting domain of the transit peptide is removed, exposing a thylakoid targeting signal peptide that directs thylakoid integration. Following integration into thylakoids, the signal peptide is removed, producing mature proteins with amino-proximal luminal tails of 118 (FtsH5) and 84 (FtsH2) residues, single transmembrane anchors, and large catalytic stromal domains. The topology of these domains was verified by protease treatment and alkaline extraction (Figure 2, Figure S2 and S3). From the stroma, FtsH5 and FtsH2 are targeted to separate membrane integration machineries. The plastid Sec machinery integrates FtsH5, whereas the Tat machinery integrates FtsH2. That FtsH5 is integrated by the Sec pathway was demonstrated by the requirement for ATP, inhibition by the SecA inhibitor azide, competition by the Sec pathway substrate iOE33, and inhibition by pretreatment with antibodies to SecY. By analogy with the operation of the *E. coli* Sec system (Osborne *et al.*, 2005; Driessen and Nouwen, 2008), we envision that SecA engages the hydrophobic signal peptide, which serves as a start-transfer sequence, and by a processive insertion of unfolded peptide segments through the SecYE channel, SecA delivers the luminal domain across the thylakoid membrane. The transmembrane anchor, which would function as a stop transfer sequence, arrests further transport and is laterally released to the lipid bilayer.

Evidence that the Tat pathway integrates FtsH2 includes the PMF as sole energy requirement, competition by the Tat substrate iOE23, and inhibition by pretreatment with antibodies to the Tat component Hcf106. Previous work in our lab showed that a *Capsicum* FtsH protease, Pftf, implicated in chromoplast vesicle fusion, is also integrated by the Tat pathway (Summer *et al.*, 2000). The defining feature of Tat-targeted FtsH proteins is the presence of a twin arginine motif. The consensus Tat motif for thylakoid proteins is R-R-X-Hyd-Hyd, where X is any amino acid and Hyd is a hydrophobic amino acid. FtsH2 and FtsH8 possess Tat motifs that fit the more stringent bacterial Tat motif, R-R-X-F-L-K, with a phenylalanine at position RR+2. The RR+2 F is important for transport efficiency in *E. coli* (Stanley *et al.*, 2000) and can increase the binding affinity of thylakoid precursors that normally lack phenylalanine at this position (Gerard and Cline, 2007). A brief survey of the plant sequence database indicates that, in general, plants and green algae possess a Tat-targeted FtsH protein. FtsH2 is one of only a handful of membrane proteins integrated by the Tat machinery (Summer *et al.*, 2000; Molik *et al.*, 2001; Hatzixanthis *et al.*, 2003). The manner by which Tat integrates membrane proteins is unclear because Tat transports fully folded protein domains with charged and hydrophilic surfaces (for review, see (Cline and Theg, 2007)). Experiments with artificial substrates showed that even a small exposed hydrophobic domain can terminate Tat transport (Richter *et al.*, 2007), apparently causing disassembly of the

transient translocase complex and release of the hydrophobic segment to the lipid bilayer.

Signal peptide swapping experiments with other substrates have shown that the Tat pathway can transport Sec substrate mature domains with moderate efficiency, but that the Sec pathway is nearly unable to transport Tat substrate mature domains (Henry *et al.*, 1997). A similar situation exists for the FtsH proteases. The chimeric precursor TP2Mat5 was readily imported into chloroplasts and integrated into thylakoids by the Tat pathway. By contrast, TP5Mat2 was imported into plastids but incapable of integration. A tightly folded structure and/or the presence of basic residues near signal peptidase cleavage site appear to be two main reasons for incompatibility of Tat substrates on the Sec pathway (Bogsch *et al.*, 1997; Hynds *et al.*, 1998). We attempted to narrow the relevant incompatibility regions of FtsH2 by making chimeric constructs between FtsH5 and FtsH2. However, we found that no individual domain of FtsH2 was sufficient to abort Sec-mediated integration, suggesting that the entire FtsH2 mature domain presents a structure that interferes with either the recognition by or access to the Sec machinery.

Sakamoto (2006) notes that the multiplicity of FtsH isoforms is associated with photosynthetic organisms rather than non-photosynthetic organisms. Four FtsH isomers have been identified in the cyanobacterium *Synechocystis* sp. PCC6803. Of interest is that one of these isomers (sll1463) contains a twin arginine immediately preceding the first hydrophobic region, which is followed by a putative cleavage site (A-P-A). Thus the presence of chloroplast FtsH isomers integrated by the Tat pathway may be due to a gene duplication and pathway segregation that occurred in the photosynthetic prokaryote progenitor of chloroplasts. It is not unusual for a photosynthetic complex to contain subunits targeted by different thylakoid translocation pathways. For example plant chloroplast photosystem 2 contains subunits translocated by all four known thylakoid pathways. However, the situation for thylakoid FtsH proteins, where homologous subunits of the same complex take different integration pathways enroute to their functional form is highly intriguing. The specific factors that dictate the use of mechanistically diverse machinery may yet yield insight into the operations of Tat and Sec pathways for post-translational integration of membrane proteins.

EXPERIMENTAL PROCEDURES

In-silico analysis

Arabidopsis FtsH sequences were obtained from TAIR database (<http://www.arabidopsis.org>). Stromal processing sites were predicted using ChloroP (<http://www.cbs.dtu.dk/services/ChloroP>) and transmembrane domains by TMpred (http://www.ch.embnet.org/software/TMPRED_form.html). Binary alignments were obtained using SIM and LALNVIEW (<http://www.expasy.ch/tools/sim-prot.html>).

Preparation of precursor proteins

Clones for pLHCP, pOE23 and pOE33 were previously described (Cline *et al.*, 1993a). Nucleotide sequences encoding the full-length FtsH2 and FtsH5 cDNAs were amplified from clones kindly provided by Dr. W. Sakamoto (Okayama University), using primers containing *Bam*H1 (forward) and *Pst*I (reverse) restriction sites. Truncated versions for FtsH2 and FtsH5 were amplified with a forward *Bam*H1-containing primer and an internal primer containing a stop codon and *Pst*I site. Swappings between FtsH2 and FtsH5 were prepared by the splicing method (Horton *et al.*, 1989), i.e. by first amplifying the appropriate N- and C-terminal regions, purifying the resulting PCR products, and then PCR amplifying the spliced product using the outside forward *Bam*H1 and reverse *Pst*I primers. Truncated versions and swapping points are indicated in Figure 1. All PCR products were digested with *Bam*H1 and *Pst*I and ligated into *Bam*H1/*Pst*I-digested pGEM-4Z plasmid and transformed into XL1-Blue cells. All the resulting clones were sequenced entirely on both strands. Radiolabeled precursors were produced by transcription with SP6 polymerase (Promega) and translation with a homemade wheat germ translation system in the presence of 3 H-leucine (Cline, 1986). Translation products were diluted with one volume of 60 mM leucine in 2x import buffer (IB; 1x = 50 mM HEPES/KOH, pH 8.0, 0.33 M sorbitol) prior to use.

Chloroplast import and thylakoid integration

Intact chloroplasts were isolated from 9- to 10-day-old pea seedlings (*Pisum sativum* cv. Laxton's Progress 9) as described (Cline, 1986). Chlorophyll concentrations were determined according to (Arnon, 1949). Chloroplast lysate was prepared from intact chloroplasts and washed thylakoids and stroma prepared from lysate (Cline *et al.*, 1993b). Import of radiolabeled precursors into pea chloroplasts or transport/integration into washed thylakoids or chloroplast lysate (0.33 mg chlorophyll ml $^{-1}$ or equivalent) and 5 mM MgATP (unless otherwise stated) was conducted at 25°C in 70–100 μ E m $^{-2}$ sec $^{-1}$ white light (Cline *et al.*, 1993b) for the time indicated in the figure legends. Where indicated, chloroplasts, lysates, or thylakoids were preincubated with nigericin (0.75 μ M final concentration) and valinomycin (1.5 μ M final concentration), or sodium azide (10 mM final concentration) on ice for 10 min. Reactions were initiated by the addition of translation products equivalent to one-sixth the assay volume and terminated by transfer to an ice bath. Intact chloroplasts were recovered, with or without protease treatment, by centrifugation through a 35% Percoll cushion. For fractionation, recovered chloroplasts were lysed by resuspension in 20 mM HEPES, KOH, pH 8, and incubation on ice for 5 min. The thylakoid membranes were separated from the stromal fraction by centrifugation for 8 min at 3200 g . Thylakoids were washed with import buffer, extracted with 100 mM NaOH, extracted with 200 mM Na₂CO₃, or treated with thermolysin as described (Summer *et al.*, 2000). Thermolysin post-treatment of chloroplasts or thylakoids was with 1 μ g thermolysin per microgram chlorophyll for at least 40 min at 4°C followed by washing in import buffer containing 14 mM EDTA. Samples were analyzed by SDS-PAGE and fluorography. On the fluorograms, translation products represent 2% and all other samples represent 5% of the amounts present in each assay. Mr values for bands on gels were estimated from their migration compared to standard curves for marker proteins (EZ Run pre-stained protein ladder, Fisher Scientific).

In organello competition

In organello competition for Tat and Sec pathways was conducted essentially as described (Cline *et al.*, 1993b). Unlabeled inclusion

bodies of pOE23 or pOE33 were dissolved in fresh 10 M urea, 10 mM DTT for 3 h at room temperature. Isolated chloroplasts, 5 mM MgATP and 1.5 mM DTT were preincubated with solubilized competitors for 7 min in the light at 25°C. Competitors were aliquotted from stocks, such that the final competitor concentration was 0.75 μ M pOE23, 0.75 μ M pOE33, or no competitor, and the urea concentration was 0.25 M in all assays. Radiolabeled precursors for pFtsH2, pFtsH5, pOE23 or pOE33 were then added (1/6 volume), and the incubation was continued for an additional 10 min and then transferred to ice. Chloroplasts were repurified by centrifugation through Percoll cushions. Recovered chloroplasts were lysed in 10 mM HEPES/KOH, pH 8.0, and 10 mM MgCl₂ buffer, and separated into stroma and thylakoids by centrifugation at 3200 g for 8 min. An aliquot of the thylakoid fraction was treated with 0.2 mg ml $^{-1}$ thermolysin 40 min at 4°C (FtsH2) or extracted with 200 mM Na₂CO₃ (FtsH5).

Antibody inhibition of thylakoid protein integration

Antibody inhibition of protein transport was conducted as described (Mori *et al.*, 1999). Briefly, washed thylakoids were suspended in import buffer containing 10 mM MgCl₂ plus 3% BSA at 1 mg of chlorophyll ml $^{-1}$. The suspension was adjusted with 20 mM Hepes/KOH, pH 8, to a final chlorophyll concentration of 0.33 mg ml $^{-1}$ and antibody concentrations as indicated in figure legends. After 1 h on ice, thylakoids were recovered by centrifugation at 3200 g for 8 min and washed with import buffer containing 10 mM MgCl₂. Aliquots of pretreated thylakoids (equivalent to 25 μ g chlorophyll) were supplemented with stromal extract equivalent to 50 μ g chlorophyll of intact chloroplasts and Mg-ATP (5 mM final concentration) and then incubated with radiolabeled proteins pFtsH2, pFtsH5, pOE23, pOE33 and pLHCP in a final volume of 75 μ l. Reactions were conducted at 25°C for 30 min in the light and terminated by transfer to ice. Thylakoids, recovered by centrifugation, were treated with 0.2 mg ml $^{-1}$ thermolysin for 40 min at 4°C or were extracted with 200 mM Na₂CO₃ (FtsH5).

ACKNOWLEDGEMENTS

We thank Dr Wataru Sakamoto for the FtsH cDNA clones. We thank Cassie Aldridge, Jose Celedon, Xianyue Ma, and Michael McCaffery for critical review of the manuscript, and Michael McCaffery for excellent technical assistance. This work was supported in part by National Institutes of Health grant R01 GM46951 to KC and Fundação de Amparo à Pesquisa do Estado de São Paulo grant 08/52067-3 to MCSF. MCSF is also a research fellow of CNPq. RAOR is a recipient of CAPES (BEX 0375/09-8) and FAPESP (2007/57806-6) fellowships.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Figure S1. Sequence alignment of chloroplast FtsH proteins.

Figure S2. pFtsH2 is completely degraded by thermolysin without protection by the thylakoid bilayer.

Figure S3. *In vitro* translation of pFtsH5 RNA produces a truncated precursor in addition to the full-length pFtsH5.

Figure S4. pFtsH2 integration is competed *in organello* by a Tat precursor while FtsH5 is competed by a Sec precursor.

Please note: As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

REFERENCES

Adam, Z., Rudella, A. and van Wijk, K.J. (2006) Recent advances in the study of Clp, FtsH and other proteases located in chloroplasts. *Curr. Opin. Plant Biol.* **9**, 234–240.

Arnon, D.I. (1949) Copper enzymes in isolated chloroplasts. Polyphenoloxidase in Beta vulgaris. *Plant Physiol.* **24**, 1–15.

Bogsch, E., Brink, S. and Robinson, C. (1997) Pathway specificity for a delta pH-dependent precursor thylakoid lumen protein is governed by a 'Sec-avoidance' motif in the transfer peptide and a 'Sec-incompatible' mature protein. *EMBO J.* **16**, 3851–3859.

Cline, K. (1986) Import of proteins into chloroplasts. Membrane integration of a thylakoid precursor protein reconstituted in chloroplast lysates. *J. Biol. Chem.* **261**, 14804–14810.

Cline, K. and Dabney-Smith, C. (2008) Plastid protein import and sorting: different paths to the same compartments. *Curr. Opin. Plant Biol.* **11**, 585–592.

Cline, K. and Theg, S.M. (2007) The Sec and Tat protein translocation pathways in chloroplasts. In *Molecular Machines Involved in Protein Transport across Cellular Membranes*, Vol. XXV (Dalbey, R.E., Tamanai, F. and Koehler, C.M., eds). London: Elsevier, pp. 463–492.

Cline, K., Henry, R., Li, C. and Yuan, J. (1993a) Multiple pathways for protein transport into or across the thylakoid membrane. *EMBO J.* **12**, 4105–4114.

Driessens, A.J. and Nouwen, N. (2008) Protein translocation across the bacterial cytoplasmic membrane. *Annu. Rev. Biochem.* **77**, 643–667.

Ferro, M., Salvi, D., Brugiere, S., Miras, S., Kowalski, S., Louwagie, M., Garin, J., Joyard, J. and Rolland, N. (2003) Proteomics of the chloroplast envelope membranes from *Arabidopsis thaliana*. *Mol. Cell Proteomics*, **2**, 325–345.

Gerard, F. and Cline, K. (2007) The thylakoid proton gradient promotes an advanced stage of signal peptide binding deep within the Tat pathway receptor complex. *J. Biol. Chem.* **282**, 5263–5272.

Hatzixanthitis, K., Palmer, T. and Sargent, F. (2003) A subset of bacterial inner membrane proteins integrated by the twin-arginine translocase. *Mol. Microbiol.* **49**, 1377–1390.

Henry, R., Carrigan, M., McCaffrey, M., Ma, X. and Cline, K. (1997) Targeting determinants and proposed evolutionary basis for the Sec and the Delta pH protein transport systems in chloroplast thylakoid membranes. *J. Cell Biol.* **136**, 823–832.

Horton, R.M., Hunt, H.D., Ho, S.N., Pullen, J.K. and Pease, L.R. (1989) Engineering hybrid genes without the use of restriction enzymes: gene splicing by overlap extension. *Gene*, **77**, 61–68.

Hynds, P.J., Robinson, D. and Robinson, C. (1998) The sec-independent twin-arginine translocation system can transport both tightly folded and malfolded proteins across the thylakoid membrane. *J. Biol. Chem.* **273**, 34868–34874.

Ito, K. and Akiyama, Y. (2005) Cellular functions, mechanism of action, and regulation of FtsH protease. *Annu. Rev. Microbiol.* **59**, 211–231.

Molik, S., Karnauchov, I., Weidlich, C., Herrmann, R.G. and Klosgen, R.B. (2001) The Rieske Fe/S protein of the cytochrome b6/f complex in chloroplasts: missing link in the evolution of protein transport pathways in chloroplasts? *J. Biol. Chem.* **276**, 42761–42766.

Mori, H. and Cline, K. (2001) Post-translational protein translocation into thylakoids by the Sec and Deltaph-dependent pathways. *Biochim. Biophys. Acta*, **1541**, 80–90.

Mori, H., Summer, E.J., Ma, X. and Cline, K. (1999) Component specificity for the thylakoidal Sec and Delta pH-dependent protein transport pathways. *J. Cell Biol.* **146**, 45–56.

Niwa, H., Tsuchiya, D., Makyio, H., Yoshida, M. and Morikawa, K. (2002) Hexameric ring structure of the ATPase domain of the membrane-integrated metalloprotease FtsH from *Thermus thermophilus* HB8. *Structure*, **10**, 1415–1423.

Osborne, A.R., Rapoport, T.A. and van den Berg, B. (2005) Protein translocation by the Sec61/SecY channel. *Annu. Rev. Cell Dev. Biol.* **21**, 529–550.

Richter, S., Lindenstrauss, U., Lucke, C., Bayliss, R. and Bruser, T. (2007) Functional Tat transport of unstructured, small, hydrophilic proteins. *J. Biol. Chem.* **282**, 33257–33264.

Sakamoto, W. (2006) Protein degradation machineries in plastids. *Annu. Rev. Plant Biol.* **57**, 599–621.

Sakamoto, W., Zaltsman, A., Adam, Z. and Takahashi, Y. (2003) Coordinated regulation and complex formation of yellow variegated1 and yellow variegated2, chloroplastic FtsH metalloproteases involved in the repair cycle of photosystem II in *Arabidopsis* thylakoid membranes. *Plant Cell*, **15**, 2843–2855.

Stanley, N.R., Palmer, T. and Berks, B.C. (2000) The twin arginine consensus motif of Tat signal peptides is involved in Sec-independent protein targeting in *Escherichia coli*. *J. Biol. Chem.* **275**, 11591–11596.

Summer, E.J., Mori, H., Settles, A.M. and Cline, K. (2000) The thylakoid delta pH-dependent pathway machinery facilitates RR-independent N-tail protein integration. *J. Biol. Chem.* **275**, 23483–23490.

Urantowka, A., Knorpp, C., Olczak, T., Kolodziejczak, M. and Janska, H. (2005) Plant mitochondria contain at least two i-AAA-like complexes. *Plant Mol. Biol.* **59**, 239–252.

Yu, F., Park, S. and Rodermel, S.R. (2004) The *Arabidopsis* FtsH metalloprotease gene family: interchangeability of subunits in chloroplast oligomeric complexes. *Plant J.* **37**, 864–876.

Yu, F., Park, S. and Rodermel, S.R. (2005) Functional redundancy of AtFtsH metalloproteases in thylakoid membrane complexes. *Plant Physiol.* **138**, 1957–1966.

Yuan, J. and Cline, K. (1994) Plastocyanin and the 33 kDa subunit of the oxygen-evolving complex are transported into thylakoids with similar requirements as predicted from pathway specificity. *J. Biol. Chem.* **269**, 18463–18467.

Zaltsman, A., Ori, N. and Adam, Z. (2005) Two types of FtsH protease subunits are required for chloroplast biogenesis and Photosystem II repair in *Arabidopsis*. *Plant Cell*, **17**, 2782–2790.